

Jaromír Duda

Factorable congruences and factorable congruence blocks on powers of a finite algebra

Czechoslovak Mathematical Journal, Vol. 42 (1992), No. 1, 89–93

Persistent URL: <http://dml.cz/dmlcz/128309>

Terms of use:

© Institute of Mathematics AS CR, 1992

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

FACTORABLE CONGRUENCES AND FACTORABLE
CONGRUENCE BLOCKS ON POWERS OF A FINITE ALGEBRA

JAROMÍR DUDA, Brno

(Received January 7, 1991)

1. INTRODUCTION

R. Willard has proved in [4] that any power A^n , $n \geq 2$, of a finite k -element algebra A , $k \geq 2$, has factorable congruences whenever the power $A^{k^3+k^2-k}$ has the same property. In this paper the exponent $k^3 + k^2 - k$ is reduced to $3k^2 - 2k$. Further, it is shown that the factorability of congruence blocks on the power A^{2k^2-k} ensures this property on any power A^n , $n \geq 2$.

2. FACTORABLE CONGRUENCES

Definition 1. Let A_1, \dots, A_n , $n \geq 2$, be algebras of the same type. We say that the product $B = A_1 \times \dots \times A_n$ has *factorable congruences* whenever $\Theta = \Theta_1 \times \dots \times \Theta_n$ holds for any congruence Θ on B where $\Theta_1, \dots, \Theta_n$ are congruences on A_1, \dots, A_n , respectively.

Notation 1. Let A_1, \dots, A_n , $n \geq 2$, be algebras of the same type, $B = A_1 \times \dots \times A_n$. Elements $\langle a_1, \dots, a_n \rangle$, $\langle b_1, \dots, b_n \rangle$, ... of B are denoted by \bar{a} , \bar{b} , Further, denote

$$\sigma(B) = \{ \langle \bar{a}, \bar{b}, \bar{c}, \bar{d} \rangle \in B^4; \text{ for each } i \leq n \text{ either } \langle a_i, b_i \rangle = \langle c_i, d_i \rangle \text{ or } a_i = b_i \}$$

and

$$\gamma(B) = \left\{ \langle \bar{a}, \bar{b}, \bar{c}, \bar{d} \rangle \in B^4; \text{ for each } i \leq n \text{ either } \langle a_i, b_i \rangle = \langle c_i, d_i \rangle \right. \\ \left. \text{or } a_i = b_i, c_i = d_i \text{ or } a_i = b_i = d_i \right\}.$$

Notation 2. Let B be an algebra, $c, d \in B$. Then the symbol $\Theta_B(c, d)$ denotes the principal congruence on B generated by the pair $\langle c, d \rangle$.

Lemma 1. Let $A_1, \dots, A_n, n \geq 2$, be algebras of the same type, $B = A_1 \times \dots \times A_n$. The following conditions are equivalent:

- (1) B has factorable congruences;
- (2) $\langle \bar{a}, \bar{b}, \bar{c}, \bar{d} \rangle \in \sigma(B)$ implies $\langle \bar{a}, \bar{b} \rangle \in \Theta_B(\bar{c}, \bar{d})$ for any elements $\bar{a}, \bar{b}, \bar{c}, \bar{d} \in B$;
- (3) $\langle \bar{a}, \bar{b}, \bar{c}, \bar{d} \rangle \in \gamma(B)$ implies $\langle \bar{a}, \bar{b} \rangle \in \Theta_B(\bar{c}, \bar{d})$ for any elements $\bar{a}, \bar{b}, \bar{c}, \bar{d} \in B$.

Proof. (1) \Leftrightarrow (2): See [4; Lemma 4.3, p. 339].

(2) \Rightarrow (3) is trivial since $\gamma(B) \subseteq \sigma(B)$;

(3) \Rightarrow (2): Let $\langle \bar{a}, \bar{b}, \bar{c}, \bar{d} \rangle \in \sigma(B)$. Then $\langle a_i, b_i \rangle = \langle c_i, d_i \rangle, i \in I$, and $a_i = b_i, i \in J$, for some disjoint index sets $I, J, I \cup J = \{1, \dots, n\}$.

(a) Introduce a new quadruple $\langle \bar{a}', \bar{b}', \bar{c}', \bar{d}' \rangle \in B^4$ by the rule

$$\langle a'_i, b'_i, c'_i, d'_i \rangle = \begin{cases} \langle a_i, b_i, c_i, d_i \rangle & \text{for } i \in I \\ \langle d_i, d_i, c_i, d_i \rangle & \text{for } i \in J. \end{cases}$$

Then $\langle \bar{a}', \bar{b}', \bar{c}', \bar{d}' \rangle \in \gamma(B)$ and so $\langle \bar{a}', \bar{b}' \rangle \in \Theta_B(\bar{c}', \bar{d}')$, by hypothesis (3).

(b) Further, introduce a quadruple $\langle \bar{a}'', \bar{b}'', \bar{c}'', \bar{d}'' \rangle \in B^4$ via

$$\langle a''_i, b''_i, c''_i, d''_i \rangle = \begin{cases} \langle a_i, b_i, c_i, d_i \rangle & \text{for } i \in I \\ \langle a_i, a_i, d_i, d_i \rangle & \text{for } i \in J. \end{cases}$$

Since evidently $\langle \bar{a}'', \bar{b}'', \bar{c}'', \bar{d}'' \rangle \in \gamma(B)$ we have $\langle \bar{a}'', \bar{b}'' \rangle \in \Theta_B(\bar{c}'', \bar{d}'')$, by (3) again.

Moreover $\langle \bar{a}', \bar{b}' \rangle = \langle \bar{c}'', \bar{d}'' \rangle, \langle \bar{c}', \bar{d}' \rangle = \langle \bar{c}, \bar{d} \rangle, \langle \bar{a}'', \bar{b}'' \rangle = \langle \bar{a}, \bar{b} \rangle$, and thus $\langle \bar{a}, \bar{b} \rangle = \langle \bar{a}'', \bar{b}'' \rangle \in \Theta_B(\bar{c}'', \bar{d}'') = \Theta_B(\bar{a}', \bar{b}') \subseteq \Theta_B(\bar{c}', \bar{d}') = \Theta_B(\bar{c}, \bar{d})$, i.e. $\langle \bar{a}, \bar{b} \rangle \in \Theta_B(\bar{c}, \bar{d})$, as required. \square

Lemma 2. Let B, C be algebras of the same type, φ a homomorphism from B to C . Then $\langle a, b \rangle \in \Theta_B(c, d)$ implies $\langle \varphi(a), \varphi(b) \rangle \in \Theta_C(\varphi(c), \varphi(d))$ for any elements $a, b, c, d \in B$.

Proof. Applying the binary scheme, see [2; Thm 1, p. 41], to the relation formula $\langle a, b \rangle \in \Theta_B(c, d)$ we obtain

$$\begin{aligned} a &= t_1(c, d, b_1, \dots, b_m), \\ t_i(d, c, b_1, \dots, b_m) &= t_{i+1}(c, d, b_1, \dots, b_m), \quad 1 \leq i < n, \\ b &= t_n(d, c, b_1, \dots, b_m) \end{aligned}$$

for some elements $b_1, \dots, b_m \in B$ and suitable terms t_1, \dots, t_n . Then

$$\begin{aligned} \varphi(a) &= t_1(\varphi(c), \varphi(d), \varphi(b_1), \dots, \varphi(b_m)), \\ t_i(\varphi(d), \varphi(c), \varphi(b_1), \dots, \varphi(b_m)) &= t_{i+1}(\varphi(c), \varphi(d), \varphi(b_1), \dots, \varphi(b_m)), \quad 1 \leq i < n, \\ \varphi(b) &= t_n(\varphi(d), \varphi(c), \varphi(b_1), \dots, \varphi(b_m)), \end{aligned}$$

which means that $\langle \varphi(a), \varphi(b) \rangle \in \Theta_C(\varphi(c), \varphi(d))$, see [2] again. \square

Notation 3. Let C be an algebra, $p_1, p_2, p_3, p_4: C^4 \rightarrow C$ canonical projections, and S a subset of C^4 . Then $p_1^S, p_2^S, p_3^S, p_4^S$ denote the restrictions of p_1, p_2, p_3, p_4 , respectively to S .

Theorem 1. Let C be a finite algebra. The following conditions are equivalent:

- (1) C^n has factorable congruences for any $n \geq 2$;
- (2) $C^{\gamma(C)}$ has factorable congruences.

Proof. We use the arguments from [4; Lemma 4.4, p. 339]: Let $(\bar{a}, \bar{b}, \bar{c}, \bar{d})$ be an arbitrary quadruple from $\gamma(C^n)$, $n \geq 2$. It is a routine to verify that

- (a) $\langle p_1^{\gamma(C)}, p_2^{\gamma(C)}, p_3^{\gamma(C)}, p_4^{\gamma(C)} \rangle \in \gamma(C^{\gamma(C)})$;
- (b) the correspondence $\varphi: g \mapsto \langle g(a_1, b_1, c_1, d_1), \dots, g(a_n, b_n, c_n, d_n) \rangle$ is homomorphism from $C^{\gamma(C)}$ to C^n which sends $p_1^{\gamma(C)}, p_2^{\gamma(C)}, p_3^{\gamma(C)}, p_4^{\gamma(C)}$ to $\bar{a}, \bar{b}, \bar{c}, \bar{d}$, respectively.

Now, by hypothesis (2) the algebra $C^{\gamma(C)}$ has factorable congruences and so (a) implies $\langle p_1^{\gamma(C)}, p_2^{\gamma(C)} \rangle \in \Theta_{C^{\gamma(C)}}(p_3^{\gamma(C)}, p_4^{\gamma(C)})$. Applying the homomorphism φ to this principal congruence formula we obtain $(\bar{a}, \bar{b}) \in \Theta_{C^n}(\bar{c}, \bar{d})$ which proves (1), see Lemma 1 again. \square

Corollary 1. Let C be a finite k -element algebra, $k \geq 2$. The following conditions are equivalent:

- (1) C^n has factorable congruences for any $n \geq 2$;
- (2) C^{3k^2-2k} has factorable congruences.

Proof. Evidently $\text{card } \gamma(C) = 3k^2 - 2k$ whenever $\text{card } C = k$. \square

FACTORABLE CONGRUENCE BLOCKS

Definition 2. Let A_1, \dots, A_n , $n \geq 2$, be algebras of the same type. A subset S of $B = A_1 \times \dots \times A_n$ is said to be *factorable* whenever $S = S_1 \times \dots \times S_n$ for some subsets $S_i \subseteq A_i$, $i \leq n$.

Further, we say that B has factorable congruence blocks whenever any congruence block on B is factorable.

Lemma 3. Let A_1, \dots, A_n , $n \geq 2$, be algebras of the same type, S a subset of $B = A_1 \times \dots \times A_n$. The following conditions are equivalent:

- (1) S is factorable;
- (2) $\bar{c}, \bar{d} \in S$ implies $\bar{a} \in S$ where $a_i \in \{c_i, d_i\}$, $i \leq n$.

Proof. (1) \Rightarrow (2): Let $\bar{c}, \bar{d} \in S = S_1 \times \dots \times S_n$. Then $c_i, d_i \in S_i$, $i \leq n$, and thus also $a_i \in S_i$, $i \leq n$, for $a_i \in \{c_i, d_i\}$, $i \leq n$. Altogether, $\bar{a} = \langle a_1, \dots, a_n \rangle \in S_1 \times \dots \times S_n = S$ as required.

(2) \Rightarrow (1): Denote $S_i = pr_i S$, $i \leq n$. Evidently the inclusion $S \subseteq S_1 \times \dots \times S_n$ holds. Conversely, let $\bar{s} = \langle s_1, \dots, s_n \rangle \in S_1 \times \dots \times S_n$. Then there are elements $\langle a_{i1}, \dots, a_{in} \rangle \in S$, $i \leq n$, such that $a_{ii} = s_i$, $i \leq n$, by the definition of subsets S_i , $i \leq n$. Now from $\langle a_{11}, \dots, a_{1n} \rangle, \langle a_{21}, \dots, a_{2n} \rangle \in S$ we obtain $\langle s_1, s_2, a_{23}, \dots, a_{2n} \rangle = \langle a_{11}, a_{22}, a_{23}, \dots, a_{2n} \rangle \in S$, by hypothesis (2). Repeating this process we find that $\bar{s} = \langle s_1, \dots, s_n \rangle \in S$, which proves the factorability of C \square

Notation 4. Let A_1, \dots, A_n , $n \geq 2$, be algebras of the same type, $B = A_1 \times \dots \times A_n$. Denote by

$$\beta(B) = \{ \langle \bar{a}, \bar{b}, \bar{c}, \bar{d} \rangle \in B^4, \text{ for each } i \leq n \text{ either } \langle a_i, b_i \rangle = \langle c_i, d_i \rangle \\ \text{or } a_i = b_i = d_i \}.$$

Lemma 4. Let A_1, \dots, A_n , $n \geq 2$, be algebras of the same type, $B = A_1 \times \dots \times A_n$. The following conditions are equivalent:

- (1) B has factorable congruence blocks;
- (2) $\langle \bar{a}, \bar{b}, \bar{c}, \bar{d} \rangle \in \beta(B)$ implies $\langle \bar{a}, \bar{b} \rangle \in \Theta_B(\bar{c}, \bar{d})$ for any elements $\bar{a}, \bar{b}, \bar{c}, \bar{d} \in B$.

Proof. (1) \Rightarrow (2): Let $\langle \bar{a}, \bar{b}, \bar{c}, \bar{d} \rangle \in \beta(B)$. Then $\bar{b} = \bar{d}$ and $a_i \in \{c_i, d_i\}$, $i \leq n$. Evidently $\bar{c}, \bar{d} \in [\bar{d}]\Theta_B(\bar{c}, \bar{d})$ and thus also $\bar{a} \in [\bar{d}]\Theta_B(\bar{c}, \bar{d})$, by Lemma 3. In other words, we have $\langle \bar{a}, \bar{b} \rangle = \langle \bar{a}, \bar{d} \rangle \in \Theta_B(\bar{c}, \bar{d})$.

(2) \Rightarrow (1): Let S be an arbitrary congruence block on B and let $\bar{c}, \bar{d} \in S$. Consider an element $\bar{a} = \langle a_1, \dots, a_n \rangle$ such that $a_i \in \{c_i, d_i\}$, $i \leq n$. Then $\langle \bar{a}, \bar{d}, \bar{c}, \bar{d} \rangle \in \beta(B)$ and so $\langle \bar{a}, \bar{d} \rangle \in \Theta_B(\bar{c}, \bar{d})$, by hypothesis (2). This means that $\bar{a} \in [\bar{d}]\Theta_B(\bar{c}, \bar{d}) \subseteq S$ and so S is factorable, see Lemma 3. \square

Theorem 2. Let C be a finite algebra. The following conditions are equivalent:

- (1) C^n has factorable congruence blocks for any $n \geq 2$;
- (2) $C^{\beta(C)}$ has factorable congruence blocks.

Proof goes along the same lines as in Theorem 1 and hence can be omitted. \square

Corollary 2. Let C be a finite k -element algebra, $k \geq 2$. The following conditions are equivalent:

- (1) C^n has factorable congruence blocks;
- (2) C^{2k^2-k} has factorable congruence blocks.

Proof. We have $\text{card } \beta(C) = 2k^2 - k$ whenever $\text{card } C = k$. \square

References

- [1] *S. Burris, R. Willard*: Finitely many primitive positive clones, *Proc. Amer. Math. Soc.* **101** (1987), 427–430.
- [2] *J. Duda*: On two schemes applied to Mal'cev type theorems, *Ann. Univ. Sci. Budapest, Sect. Math.* **26** (1983), 39–45.
- [3] *J. Duda*: Varieties having directly decomposable congruence classes, *Čas. Pěst. Matem.* **111** (1986), 394–403.
- [4] *R. Willard*: Congruence lattices of powers of an algebra, *Algebra Univ.* **26** (1989), 332–340.

Author's address: 616 00 Brno, Kroftova 21, Czechoslovakia.