

Radhey S. Singh; Renu Shrivastava
Rings which have projective coflat module

Czechoslovak Mathematical Journal, Vol. 42 (1992), No. 3, 491–496

Persistent URL: <http://dml.cz/dmlcz/128346>

Terms of use:

© Institute of Mathematics AS CR, 1992

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

RINGS WHICH HAVE PROJECTIVE COFLAT MODULE

R. S. SINGH and RENU SHRIVASTAVA, Sagar

(Received January 30, 1991)

INTRODUCTION AND PRELIMINARIES

In [1] R. R. Colby has defined a ring R , as left (right) *IF* if every injective left (right) R -module is flat. In [2] R. F. Damiano has defined *FC* ring over which left (right) flat and left (right) coflat modules are precisely the same. Induced by these paper we have defined a ring R as left (right) *PC* if every left (right) projective R -module is coflat. A ring R is quasi-frobenius ring if R as a right R -module is noetherian and injective. In [3] U. Chase characterised a left perfect ring R as a ring over which every left flat R -module is projective.

Through-out this paper R will be understood a ring with unity, and every module over R will be unitary. Every right (left) R -module M is denoted by M_R (${}_R M$) and dual of M is denoted by ${}_R M^*$ (M_R^*), i.e. $M^* = \text{Hom}_R(M, R)$. S is the R -endomorphisms ring of M , i.e.

$$S = \text{Hom}_R(M, M) = \text{End}_R(M).$$

Trace of an R -module M on M is denoted by $T_M(M)$ and defined as

$$T_M(M) = \left\{ \sum \text{Im } f : f \in \text{Hom}_R(M, M) \right\}.$$

Definition. A ring R is right (left) *PC* ring if every right (left) projective R -module is coflat. A ring R is *PC* ring if it is both right as well as left *PC* ring.

Obvioulsy every *FC* ring and every *QF* ring is *PC* ring.

Theorem 1. For a ring R following statements are equivalent.

1. R is right *PC* ring.
2. Every right free module over R is coflat.
3. For every right free R module M , trace of M on M is coflat.

Proof. $1 \Rightarrow 2$ Let R is right PC ring and M is a free module over R . Then obviously it is coflat, since every free module is projective and over right PC ring every right projective module is coflat.

$2 \Rightarrow 3$ Let every free right R -module M is coflat. But M is a flat module [\because Free \Rightarrow Projective \Rightarrow flat] hence ${}_s M \otimes_R R \cong {}_s M$ is a flat module. Therefore by [4, Lemma 1.3(2)] M generates all kernels of homomorphism $M^n \rightarrow M, n \in \mathbb{N}$, which implies by [4, Lemma 1.3(1)] that the mapping $\emptyset'(M): \text{Hom}_R(M, M) \otimes_s M \rightarrow T_M(M)$ is an isomorphism or

$$S \otimes_s M_R \text{ is isomrphic to } T_M(M)$$

i.e. M_R is isomrphic to $T_M(M)$ and so $T_M(M)$ is coflat.

$3 \Rightarrow 1$ Let for every free right R -module $M, T_M(M)$ is coflat. Since M is also flat, by the same argument as in $2 \Rightarrow 3$ we have $T_M(M)$ is isomorphic to M . Which implies very free right R -module M is coflat. Now let P be any projective right R module. Then P is a direct summand of a free module, say M . But M is coflat so by [2, Theorem 1.8] every direct summand of M is coflat i.e. P is coflat. Hence R is right PC ring. \square

This theorem is true for left PC ring R and left free R -module M also.

Example 1. [2, R. F. Damiano, example 2.7]

Let $R = \mathbb{Z} \oplus \mathbb{Q}/\mathbb{Z}$ with multiplication defined by $(n_1, q_1) \cdot (n_2, q_2) = (n_1 n_2, n_1 q_2 + n_2 q_1), n_i \in \mathbb{Z}, q_i \in \mathbb{Q}/\mathbb{Z}$. Then R is a commutative ring with Jacobson radical

$$J(R) = \{(n, q) \mid n = 0\}.$$

More-over it is clear that every finitely generated ideal of R is principal. Hence R is PC ring, but $R/J(R) \cong \mathbb{Z}$ is not PC ring, as the homomorphism $f: n\mathbb{Z} \rightarrow \mathbb{Z}$ via $n\nu \rightarrow \nu$ can not extend over \mathbb{Z} , which implies \mathbb{Z} over \mathbb{Z} is not coflat.

Example 2. A left PC ring which is not ring PC .

(R. F. Damiano [2, Example 2.8]). Let R be an algebra over a field F with basis $\{1, e_0, e_1, e_2, \dots, x_1, x_2, \dots\}$ where for all i, j

$$\begin{aligned} e_i e_j &= \delta_{i,j} e_j, \\ x_i e_j &= \delta_{i,j+1} x_i, \\ e_i x_j &= \delta_{i,j} x_j, \\ x_i x_j &= 0. \end{aligned}$$

It is easy to see that R is left coherent and that every R -homomorphism

$$f: {}_R I \rightarrow {}_R R$$

extends to one over R . Then R is left coflat, and hence R is left PC ring.

However R is not right coflat since the homomorphism $x_i R \rightarrow e_o R$.

Via $x_i r \rightarrow e_o r$ can not be extended over R . Thus every projective right R -module is not coflat. Therefore R is not right PC ring.

Theorem 2. *If R is right PC ring for a finitely generated projective right R -module P following are true.*

1. $S = \text{Hom}_R(P, P)$ is coflat.
2. $P^* = \text{Hom}_R(P, R)$ is coflat.

Proof. (1) Let $o \rightarrow I \rightarrow S$ be an exact sequence with I finitely generated right ideal of S . Now P_R is projective so P_R is flat. Hence ${}_s P \otimes_R R \cong {}_s P$ is flat. So we have an exact sequence $o \rightarrow I \otimes_s P \rightarrow S \otimes_s P$ with $I \otimes_s P$ finitely generated. Again P_R is coflat module [$\because R$ is right PC ring]. So consider the following commutative diagram

$$\begin{array}{ccccc} \text{Hom}_R(S \otimes_s P, P) & \longrightarrow & \text{Hom}_R(I \otimes_s P, P) & \longrightarrow & 0 \\ \downarrow \sim & & \downarrow \sim & & \\ \text{Hom}_s(S, \text{Hom}(P, P)) & \longrightarrow & \text{Hom}_s(I, \text{Hom}(P, P)) & \longrightarrow & 0 \end{array}$$

The exactness of bottom row implies that $\text{Hom}_R(P, P)$ is coflat.

(2) Similar to the proof of (1). □

Above theorem is true for the left PC ring also.

Theorem 3. *If R is any right PC ring, the ring of endomorphisms of a finitely generated projective right R -module P is also PC ring.*

Proof. Let S be the ring of endomorphism of P i.e. $S = \text{Hom}_R(P, P)$. Let Q be any left (right) projective module over S . Then by the dual basis lemma choose a generating set $\{x_i\}_{i \in I}$ for Q and $\{f_i\}_{i \in I} \subset \text{Hom}_s(Q, S)$ such that for each $x \in Q$, $f_i(x) = 0$ for all but a finite number of i and $x = \sum_{i \in I} f_i(x)x_i \left(\sum_{i \in I} x_i f_i(x) \right)$. For each $i \in I$ define $g_i: Q \rightarrow Sx_i$ ($x_i S$) by $g_i(x) = f_i(x)x_i \left(x_i f_i(x) \right)$, $x \in Q$. Then g_i is an endomorphism of Q . Therefore $g_i(Q)$ is a direct summand of Q , hence a direct summand of Sx_i ($x_i S$). Thus $g_i(Q)$ is cyclic.

Then $g_i(Q) \cong Se$ (eS) for any nonzero idempotent of S .

But R is right PC ring so by theorem 2 S is coflat as S module. Hence Se (eS) is coflat ($\because Se(eS)$ is direct summand of S and [2, Theorem 1.8]).

Thus each $g_i(Q)$ is coflat.

But $Q = \sum_{i \in I} g_i(Q)$.

Hence by [2, Theorem 1.8] Q is coflat and $S0$ S is a PC ring. □

Corollary 1. *If R is a right PC ring then for a non-zero idempotent $e \in R$, eRe is a PC ring.*

Proof. For a nonzero idempotent $e \in R$, eR is a finitely generated projective right R -module. Hence $\text{End}_R(eR)$ is a PC ring by theorem 3. But

$$eRe \cong \text{End}_R(eR).$$

So that eRe is a PC ring. □

Corollary 2. *Let R be a right PC ring and M is an right R module having a PC endomorphisms ring. Then every direct summand of M also has a PC endomorphism ring.*

Proof. Let $S = \text{Hom}_R(M, M)$ and N a direct summand of M . Let e is the projection onto N . Then e is an idempotent of S .

Now $\text{Hom}_R(N, N) = eSe$, and by corollary 1 eSe is a PC ring. Hence $\text{Hom}_R(N, N)$ is a PC ring. □

Corollary 3. *If R is a ring and $n > 0$ is an integer, let $M_n(R)$ denote the ring of $n \times n$ matrices over R . Then R is a right PC ring iff $M_n(R)$ is PC ring.*

Proof. Let R is right PC ring. Now R^n is a finitely generated projective module, hence $\text{Hom}_R(R^n, R^n)$ is PC ring by theorem 3. But $M_n(R) \cong \text{Hom}_R(R^n, R^n)$, so $M_n(R)$ is PC ring.

Conversely let $M_n(R)$ is PC ring. Then $\text{Hom}_R(R^n, R^n) \cong M_n(R)$ is PC ring. Since R_R is a direct summand of R^n_R , by corollary 2 $\text{Hom}_R(R, R)$ is PC ring. Which implies that R is right PC ring. □

Theorem 4. *For a left perfect ring R followings are equivalent.*

1. R is left PC ring.
2. R is left FC ring.
3. Every left coflat module is projective.

Proof. $1 \Rightarrow 2$ Let R is left PC ring, then every projective left R -module is coflat. But R is left perfect ring where every left flat module is projective. Hence every left flat module is coflat, and so R is left FC ring [2, Theorem 2.4].

$2 \Rightarrow 3$ Let R is left FC ring. Then every left coflat module is flat. But R is left perfect, so every left flat module is projective. Hence every coflat module is projective.

$3 \Rightarrow 1$ Let every left coflat module projective.

Then every left coflat module is flat.

$\Rightarrow R$ is left FC ring [2, Theorem 2.4].

$\Rightarrow R$ is left PC ring [\because Every left FC ring is left PC ring]. □

Example. A PC ring which is not FC ring. In [5] Rosenberg and Zelinsky gave an example of a quasi-frobenius, hence FC ring R and a nonzero idempotent $e \in R$ such that eRe is not QF . Thus, since eRe is artinian, it is not FC ring. But eRe is PC ring, (since, R is a PC ring and eR is a finitely generated projective module which implies that $eRe \cong \text{End}_R(eR)$ is a PC ring by theorem 3).

Theorem 5. If R is right noetherian ring following statements are equivalent.

1. R is right PC ring.
2. R is QF ring.
3. R is right IF ring.
4. R is FC ring.

Proof. $1 \Rightarrow 2$ Let R is right PC ring. Then every right projective module is coflat. Hence R_R is coflat. But R is right noetherian hence by [2, Corollary 1.10] R_R is injective. Thus R_R is noetherian and coflat so that R is QF ring.

$2 \Rightarrow 3$ Let R is QF ring. Then every right injective R -module is projective or every right injective R -module is flat. Hence R is right IF ring.

$3 \Rightarrow 4$ Let R is right IF ring. Then every right injective R -module is flat. As R is right noetherian every right coflat module is right injective, so every coflat module is flat. Hence R is FC ring [2, Theorem 2.4].

$4 \Rightarrow 1$ Let R is FC ring, then every right flat module is coflat. Hence every right projective module is right coflat and so R is right PC ring. □

Example. A PC ring which is not QF ring.

Let K is a field and for any set A

$$R = K^A = \prod_{i \in A} K_i, \quad K_i = K \quad \forall i \in A.$$

Then R is a commutative regular ring.

Or R is a PC ring. But R is not QF ring. Since $\bigoplus_{i \in A} R_{ij}R_i = R$ is projective but not injective as R is not noetherian ring.

References

- [1] *R. R. Colby*: Rings which have flat-injective modules, *Journal of Algebra* 35 (1975), 239–252.
- [2] *R. F. Damiano*: Coflat rings and modules, *Pacific Journal of Mathematics* 81 (1979), 349–369.
- [3] *S. U. Chase*: Direct products of modules, *Trans. Amer. Math. Soc.* 97 (1960), 457–473.
- [4] *B. Zimmermann-Huisgen*: Endomorphism rings of self generators, *Pacific Journal of Math.* 61 (1975), 587–602.
- [5] *A. Rosenberg and D. Zelinsky*: Annihilators, *Portugalies Mathematica* 20 (1961), 53–65.

Authors' address: Department of Mathematics and Statistics, Dr. Harisingh Gour Vishwavidyalaya, Sagar (M. P.), 470003, (Formerly: University of Sagar), India.