

Svatoslav Staněk

On a class of functional boundary value problems for second-order functional differential equations with parameter

Czechoslovak Mathematical Journal, Vol. 43 (1993), No. 2, 339–348

Persistent URL: <http://dml.cz/dmlcz/128403>

Terms of use:

© Institute of Mathematics AS CR, 1993

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON A CLASS OF FUNCTIONAL BOUNDARY VALUE PROBLEMS
FOR SECOND-ORDER FUNCTIONAL DIFFERENTIAL EQUATIONS
WITH PARAMETER

SVATOSLAV STANĚK, Olomouc

(Received December 18, 1991)

In this paper sufficient conditions concerning only operators Q , F are given for the functional differential equation

$$y''(t) - Q[y, y'](t) \cdot y(t) = F[y, y', \mu](t)$$

depending on the parameter μ to admit, for a suitable value of μ , a solution y satisfying functional boundary conditions

$$\alpha_1(y(t_1) - y(t)|J_1) = 0, \quad y(t_2) = 0, \quad \alpha_2(y(t_3) - y(t)|J_2) = 0,$$

where $-\infty < t_1 < t_2 < t_3 < \infty$, α_i are continuous functionals and $y(t)|J_i$ denotes the restriction of y to $J_i = \langle t_i, t_{i+1} \rangle$ ($i = 1, 2$). Next, sufficient conditions are given under which the above equation has, for a suitable value of the parameter μ , a bounded solution y on the halfline (t_1, ∞) and $\alpha_1(y(t_1) - y(t)|J_1) = 0$, $y(t_2) = 0$.

1. INTRODUCTION

Let $-\infty < t_1 < t_2 < t_3 < \infty$, $-\infty < a < b < \infty$, $J = \langle t_1, t_3 \rangle$, $J_1 = \langle t_1, t_2 \rangle$, $J_2 = \langle t_2, t_3 \rangle$, $I = \langle a, b \rangle$ and X (X_1 ; X_2) be the Banach space of the C^0 -functions on J (J_1 ; J_2) with the norm $\|y\| = \max\{|y(t)|; t \in J\}$ ($\|y\|_1 = \max\{|y(t)|; t \in J_1\}$; $\|y\|_2 = \max\{|y(t)|; t \in J_2\}$). Consider the functional differential equation

$$(1) \quad y''(t) - Q[y, y'](t) \cdot y(t) = F[y, y', \mu](t),$$

depending on a parameter μ . Here $Q: X \times X \rightarrow X$, $F: X \times X \times I \rightarrow X$ are continuous operators, $Q[y, z](t) > 0$ on J for all $[y, z] \in X \times X$.

Let $\alpha_i: X_i \rightarrow R$ ($i = 1, 2$) be continuous increasing (i.e. $\alpha_i(x) < \alpha_i(y)$ for all $x, y \in X_i$, $x(t) < y(t)$ for $t \in J_i - \{t_{2i-1}\}$, $x(t_{2i-1}) = y(t_{2i-1}) = 0$) functionals, $\alpha_i(0) = 0$. The purpose of this paper is to obtain using the Schauder linearization technique and the Schauder fixed point theorem, sufficient conditions imposed on the operators Q, F under which equation (1) admits, for a suitable value of the parameter μ , a solution y satisfying the functional boundary conditions

$$(2) \quad \alpha_1(y(t_1) - y(t)|J_1) = 0, \quad y(t_2) = 0, \quad \alpha_2(y(t_3) - y(t)|J_2) = 0,$$

where $y(t)|J_i$ ($i = 1, 2$) denotes the restriction of y to the interval J_i .

In Section 4, we use BVP (1)–(2) to consider bounded solutions of (1) on the halfline (t_1, ∞) satisfying the functional boundary conditions

$$\alpha_1(y(t_1) - y(t)|J_1) = 0, \quad y(t_2) = 0.$$

The paper generalizes the author's results in [1]–[3] and, in a special case, also his results in [4]. In [1] the existence of solutions of (1) satisfying for example the boundary conditions $y(t_1) - y(t_2) = y(t_3) = y(t_4) - y(t_5) = 0$ ($-\infty < t_1 < t_2 < t_3 < t_4 < t_5 < \infty$) was studied.

In [2] sufficient conditions for the existence (and uniqueness) of solutions of the differential equation

$$(3) \quad y'' - q(t)y = f(t, y, y', \mu)$$

satisfying the boundary conditions

$$(4) \quad y(t_1) = y(t_2) = y(t_3) = 0$$

($-\infty < t_1 < t_2 < t_3 < \infty$) was established.

In [4] the author considered the functional differential equation

$$y''(t) - q(t)y(t) = f(t, y(t), y(h_0(t)), y'(t), y'(h_1(t)), \mu)$$

with boundary conditions

$$\sum_{i=1}^m \alpha_i y(t_i) = 0, \quad y(c) = 0, \quad \sum_{j=1}^n \beta_j y(x_j) = 0$$

($\alpha_i > 0, \beta_j > 0$ constants, $a = t_1 < \dots < t_m < c < x_n < \dots < x_1 = b$).

In [3]—among other—sufficient conditions for the boundedness of solutions of (3) on a halfline (t_1, ∞) satisfying the boundary conditions $y(t_1) = y(t_2) = 0$ ($t_2 > t_1$) were obtained.

A functional boundary value problem depending on one parameter was studied also in [5]. In this paper the retarded functional differential equation

$$y'' - q(t)y = f(t, y_t, \mu)$$

with boundary conditions (4) was considered.

2. NOTATION, LEMMAS

Let $\varphi \in C^1(J)$ and let u_φ, v_φ be the solutions of the differential equation

$$(5) \quad y'' = Q[\varphi, \varphi'](t)y,$$

$u_\varphi(t_2) = 0, u'_\varphi(t_2) = 1, v_\varphi(t_2) = 1, v'_\varphi(t_2) = 0$. For $(t, s) \in J \times J$ define $r(t, s; \varphi)$ and $r'_1(t, s; \varphi)$ by

$$\begin{aligned} r(t, s; \varphi) &= u_\varphi(t)v_\varphi(s) - u_\varphi(s)v_\varphi(t) \quad (= -r(s, t; \varphi)), \\ r'_1(t, s; \varphi) &= u'_\varphi(t)v_\varphi(s) - u_\varphi(s)v'_\varphi(t) \quad (= \frac{\partial}{\partial t}r(t, s; \varphi)). \end{aligned}$$

Then $r(t, s; \varphi) > 0$ for all $t_1 \leq s < t \leq t_3$, $r(t, s; \varphi) < 0$ for all $t_1 \leq t < s \leq t_3$, $r'_1(t, s; \varphi) > 1$ for all $(t, s) \in J \times J$ and $t \neq s$, $r'_1(t, t; \varphi) = 1$ for all $t \in J$ (for the proof, see e.g. [2]).

Lemma 1. Assume $\varphi \in C^1(J)$, $h \in C^0(J \times I)$, $h(t, \cdot)$ is increasing on I for each fixed $t \in J$ and

$$(6) \quad h(t, a)h(t, b) \leq 0 \quad \text{for all } t \in J.$$

Then there is a unique $\mu_0 \in I$ such that the differential equation

$$(7) \quad y'' = Q[\varphi, \varphi'](t)y + h(t, \mu)$$

with $\mu = \mu_0$ admits a solution y satisfying (2). Moreover, this solution y is unique.

Proof. The function $y(t; \mu, c)$ defined on $J \times I \times R$ by

$$y(t; \mu, c) = c u_\varphi(t) + \int_{t_2}^t r(t, s; \varphi)h(s, \mu) ds$$

is the general solution of (7) vanishing at the point $t = t_2$. Since

$$\begin{aligned} y(t_1; \mu, c) - y(t; \mu, c) &= c(u_\varphi(t_1) - u_\varphi(t)) + \\ &+ \int_{t_2}^t [r(t_1, s; \varphi) - r(t, s; \varphi)] h(s, \mu) ds + \int_t^{t_1} r(t_1, s; \varphi) h(s, \mu) ds, \\ y(t_3; \mu, c) - y(t; \mu, c) &= c(u_\varphi(t_3) - u_\varphi(t)) + \\ &+ \int_{t_2}^t [r(t_3, s; \varphi) - r(t, s; \varphi)] h(s, \mu) ds + \int_t^{t_3} r(t_3, s; \varphi) h(s, \mu) ds \end{aligned}$$

and $u_\varphi(t_1) - u_\varphi(t) < 0$ on (t_1, t_3) , $u_\varphi(t_3) - u_\varphi(t) > 0$ on (t_1, t_3) , $r(t_1, s; \varphi) - r(t, s; \varphi) = r'_1(\xi, s; \varphi)(t_1 - t) < 0$ for $(t, s) \in J \times J$, $t \neq t_1$ (where ξ lies between t_1 and t), $r(t_3, s; \varphi) - r(t, s; \varphi) = r'_1(\eta, s; \varphi)(t_3 - t) > 0$ for $(t, s) \in J \times J$, $t \neq t_3$ (where η lies between t_3 and t), we see that the functions $p_i: I \times R \rightarrow R$, $p_i(\mu, c) = \alpha_i(y(t_{2i-1}; \mu, c) - y(t; \mu, c)|J_i)$ ($i = 1, 2$) are continuous on $I \times R$, $p_i(\cdot, c)$ are increasing on I for each fixed $c \in R$, $p_1(\mu, \cdot)$ ($p_2(\mu, \cdot)$) is decreasing (increasing) on R for each fixed $\mu \in I$. Finally, one can check that $\lim_{c \rightarrow -\infty} p_1(\mu, c) > 0$, $\lim_{c \rightarrow \infty} p_1(\mu, c) < 0$, $\lim_{c \rightarrow -\infty} p_2(\mu, c) < 0$, $\lim_{c \rightarrow \infty} p_2(\mu, c) > 0$ for each fixed $\mu \in I$. Hence there are unique functions $c_i: I \rightarrow R$ ($i = 1, 2$) such that

$$p_i(\mu, c_i(\mu)) = 0 \quad \text{for all } \mu \in I \text{ and } i = 1, 2,$$

and $c_1(\mu)$ ($c_2(\mu)$) is increasing (decreasing) on I .

To prove that c_i ($i = 1, 2$) are continuous functions on I we suppose there are sequences $\{\mu'_n\}$, $\{\mu''_n\}$ from I such that $\lim_{n \rightarrow \infty} \mu'_n = \lim_{n \rightarrow \infty} \mu''_n = \mu_0$ and $\lim_{n \rightarrow \infty} c_i(\mu'_n) = \lambda_1$, $\lim_{n \rightarrow \infty} c_i(\mu''_n) = \lambda_2$, $\lambda_1 < \lambda_2$, for some $i \in \{1, 2\}$. Then $0 = \lim_{n \rightarrow \infty} p_i(\mu'_n, c_i(\mu'_n)) = p_i(\mu_0, \lambda_1)$, $0 = \lim_{n \rightarrow \infty} p_i(\mu''_n, c_i(\mu''_n)) = p_i(\mu_0, \lambda_2)$, which is a contradiction to $p_i(\mu_0, \lambda_1) \neq p_i(\mu_0, \lambda_2)$.

It remains to prove the existence of a unique $\mu_0 \in I$ such that $c_1(\mu_0) = c_2(\mu_0)$. Since $h(t, a) \leq 0$, $h(t, b) \geq 0$ on J (cf. (6)) we have $y(t_1; a, 0) - y(t; a, 0) \leq 0$, $y(t_1; b, 0) - y(t; b, 0) \geq 0$ for $t \in (t_1, t_2)$, $y(t_3; a, 0) - y(t; a, 0) \leq 0$, $y(t_3; b, 0) - y(t; b, 0) \geq 0$ for $t \in (t_2, t_3)$, and then $p_i(a, 0) \leq 0$, $p_i(b, 0) \geq 0$ ($i = 1, 2$). Using the fact that $p_1(a, \cdot)$, $p_1(b, \cdot)$ ($p_2(a, \cdot)$, $p_2(b, \cdot)$) are decreasing (increasing) on R and $p_i(a, c_i(a)) = 0$, $p_i(b, c_i(b)) = 0$ ($i = 1, 2$), we get $c_1(a) \leq 0$, $c_1(b) \geq 0$, $c_2(a) \geq 0$, $c_2(b) \leq 0$, therefore $c_1(a) - c_2(a) \leq 0$, $c_1(b) - c_2(b) \geq 0$. Since $c_1(\mu) - c_2(\mu)$ is continuous increasing on I , the equation $c_1(\mu) - c_2(\mu) = 0$ has a unique solution on I . \square

Next, we will suppose that there exist positive constants r_0, r_1 such that the operators Q, F satisfy the following assumptions:

- (H₁) $|F[y, y', \mu](t)| \leq r_0 \cdot Q[y, y'](t)$ for all $t \in J$ and $[y, y', \mu] \in D \times I$,
 where $D = \{[y, y']; y \in C^1(J), \|y^{(i)}\| \leq r_i \text{ for } i = 0, 1\}$;
- (H₂) $F[y, y', \mu_1](t) < F[y, y', \mu_2](t)$ for all $t \in J$
 and $[y, y'] \in D, \mu_1, \mu_2 \in I, \mu_1 < \mu_2$;
- (H₃) $F[y, y', a](t) \cdot F[y, y', b](t) \leq 0$ for all $t \in J$ and $[y, y'] \in D$;
- (H₄) $\min\{(A + r_0 B)\tau, 2\sqrt{r_0}\sqrt{A + r_0 B}\} \leq r_1$,
 where $A = \sup\{\|F[y, y', \mu]\|; [y, y', \mu] \in D \times I\}$,
 $B = \sup\{\|Q[y, y']\|; [y, y'] \in D\}$, $\tau = \max\{t_2 - t_1, t_3 - t_2\}$.

Lemma 2. *Let assumptions (H₁)–(H₄) be fulfilled for positive constants r_0, r_1 and let $\varphi \in C^1(J), \|\varphi^{(i)}\| \leq r_i$ ($i = 0, 1$). Then there exists a unique $\mu_0 \in I$ such that the equation*

$$(8) \quad y'' = Q[\varphi, \varphi'](t)y + F[\varphi, \varphi', \mu](t)$$

with $\mu = \mu_0$ admits a (then unique) solution y satisfying (2) and, moreover,

$$(9) \quad \|y^{(i)}\| \leq r_i \quad \text{for } i = 0, 1.$$

Proof. Setting $h(t, \mu) = F[\varphi, \varphi', \mu](t)$ for $(t, \mu) \in J \times I$, the function h fulfils the assumptions of Lemma 1 and hence there is a unique $\mu_0 \in I$ such that equation (8) with $\mu = \mu_0$ admits a (then unique) solution y satisfying (2).

Now we prove $\|y\| \leq r_0$. Let $|y(\xi)| = \|y\| > r_0$ for some $\xi \in J$. If $\xi \in (t_1, t_3)$ then the function $y \cdot \text{sign } y(\xi)$ has a local maximum at the point $t = \xi$, which contradicts $y''(\xi) \cdot \text{sign } y(\xi) > 0$. The last inequality follows from assumption (H₁). Hence $\xi \in \{t_1, t_3\}$. If $\xi = t_1$ ($\xi = t_3$) then due to $y(t_2) = 0$ and assumption (H₁) we have $(y(t_1) - y(t)) \text{ sign } y(t_1) > 0$ for all $t \in (t_1, t_2)$ ($(y(t_3) - y(t)) \cdot \text{sign } y(t_3) > 0$ for all $t \in (t_2, t_3)$), which contradicts $\alpha_1(y(t_1) - y(t)|_{J_1}) = 0$ ($\alpha_2(y(t_3) - y(t)|_{J_2}) = 0$). Thus $\|y\| \leq r_0$.

Since $\alpha_i(y(t_{2i-1}) - y(t)|_{J_i}) = 0$, α_i are increasing functionals and $\alpha_i(0) = 0$ ($i = 1, 2$), there exist $\xi_1 \in (t_1, t_2)$, $\xi_2 \in (t_2, t_3)$ such that $y(t_{2i-1}) - y(\xi_i) = 0$ and therefore $y'(\eta_i) = 0$ for some $\eta_1 \in (t_1, \xi_1)$, $\eta_2 \in (\xi_2, t_3)$. For the next part of the proof of the inequality $\|y'\| \leq r_1$ see e.g. [2] and [4]. \square

3. EXISTENCE THEOREM

Theorem 1. *Assume assumptions (H₁)–(H₄) are fulfilled for positive constants r_0 and r_1 . Then there exists $\mu_0 \in I$ such that equation (1) with $\mu = \mu_0$ admits a solution y satisfying (2) and (9).*

Proof. Let Y be the Banach space of the C^1 -functions on J with the norm $\|y\|_Y = \|y\| + \|y'\|$ for $y \in Y$ and $K = \{y; y \in Y, \|y^{(i)}\| \leq r_i \text{ for } i = 0, 1\}$. K is a bounded convex closed subset of Y . Let $\varphi \in K$. By Lemma 2 there is a unique $\mu_0 \in I$ such that equation (8) with $\mu = \mu_0$ admits a (then unique) solution y satisfying (2) and $y \in K$. Setting $T(\varphi) = y$ we obtain an operator $T: K \rightarrow K$. To prove Theorem 1 it is sufficient to show that T has a fixed point.

First we prove that T is a continuous operator. Let $\{y_n\} \subset K$ be a convergent sequence, $\lim_{n \rightarrow \infty} y_n = y$ and let $z_n = T(y_n)$, $z = T(y)$. Then there are sequences $\{\mu_n\} \subset I$, $\{c_n\} \subset R$ and $\mu_0 \in I$, $c_0 \in R$ such that we have (see the proof of Lemma 1)

$$z_n(t) = c_n u_{y_n}(t) + \int_{t_2}^t r(t, s; y_n) F[y_n, y'_n, \mu_n](s) ds \text{ for all } t \in J \text{ and } n \in N,$$

$$z(t) = c_0 u_y(t) + \int_{t_2}^t r(t, s; y) F[y, y', \mu_0](s) ds \text{ for all } t \in J,$$

and

$$\alpha_1(z_n(t_1) - z_n(t)|J_1) = 0, \quad z_n(t_2) = 0, \quad \alpha_2(z_n(t_3) - z_n(t)|J_2) = 0 \text{ for all } n \in N,$$

$$\alpha_1(z(t_1) - z(t)|J_1) = 0, \quad z(t_2) = 0, \quad \alpha_2(z(t_3) - z(t)|J_2) = 0.$$

The sequence $\{c_n\}$ is bounded since $\lim_{n \rightarrow \infty} y_n = y$ and $\|z_n\| \leq r_0$ for all $n \in N$. If $\{c_n\}$ is not convergent there are convergent subsequences $\{c_{k_n}\}$, $\{c_{r_n}\}$ and convergent subsequences $\{\mu_{k_n}\}$, $\{\mu_{r_n}\}$ of $\{\mu_n\}$ such that $\lim_{n \rightarrow \infty} c_{k_n} = c^{(1)}$, $\lim_{n \rightarrow \infty} c_{r_n} = c^{(2)}$, $\lim_{n \rightarrow \infty} \mu_{k_n} = \mu^{(1)}$, $\lim_{n \rightarrow \infty} \mu_{r_n} = \mu^{(2)}$, $c^{(1)} < c^{(2)}$ and $\mu^{(1)}, \mu^{(2)}$ are either equal or not. Then

$$(k_1(t) :=) \lim_{n \rightarrow \infty} z_{k_n}(t) = c^{(1)} u_y(t) + \int_{t_2}^t r(t, s; y), F[y, y', \mu^{(1)}](s) ds,$$

$$(k_2(t) :=) \lim_{n \rightarrow \infty} z_{r_n}(t) = c^{(2)} u_y(t) + \int_{t_2}^t r(t, s; y) F[y, y', \mu^{(2)}](s) ds$$

uniformly on J and

$$(10) \quad \alpha_1(k_i(t_1) - k_i(t)|J_1) = 0, \quad k_i(t_2) = 0,$$

$$\alpha_2(k_i(t_3) - k_i(t)|J_2) = 0 \quad \text{for } i = 1, 2.$$

The equalities ($i = 1, 2$)

$$\begin{aligned}
 k_i(t_1) - k_i(t) &= c^{(i)}(u_y(t_1) - u_y(t)) + \int_{t_2}^t (r(t_1, s; y) - r(t, s; y)) \\
 &\quad \times F[y, y', \mu^{(i)}](s) ds + \int_t^{t_1} r(t_1, s; y) F[y, y', \mu^{(i)}](s) ds, \\
 k_i(t_3) - k_i(t) &= c^{(i)}(u_y(t_3) - u_y(t)) + \int_{t_2}^t (r(t_3, s; y) - r(t, s; y)) \\
 &\quad \times F[y, y', \mu^{(i)}](s) ds + \int_t^{t_3} r(t_3, s; y) F[y, y', \mu^{(i)}](s) ds
 \end{aligned}$$

imply (see the proof of Lemma 1)

$$\begin{aligned}
 k_1(t_1) - k_1(t) &> k_2(t_1) - k_2(t) \quad \text{for } t \in (t_1, t_2) \quad \text{and } \mu^{(1)} \geq \mu^{(2)}, \\
 k_2(t_3) - k_2(t) &> k_1(t_3) - k_1(t) \quad \text{for } t \in (t_2, t_3) \quad \text{and } \mu^{(1)} \leq \mu^{(2)},
 \end{aligned}$$

which contradicts (10). Hence $\{c_n\}$ is convergent, and let $\lim_{n \rightarrow \infty} c_n = c^*$. If $\{\mu_n\}$ is not convergent there are convergent subsequences $\{\mu_{j_n}\}$, $\{\mu_{i_n}\}$, $\lim_{n \rightarrow \infty} \mu_{j_n} = \lambda^{(1)}$, $\lim_{n \rightarrow \infty} \mu_{i_n} = \lambda^{(2)}$, $\lambda^{(1)} < \lambda^{(2)}$. Then

$$\begin{aligned}
 (p_1(t) :=) \lim_{n \rightarrow \infty} z_{j_n}(t) &= c^* u_y(t) + \int_{t_2}^t r(t, s; y) F[y, y', \lambda^{(1)}](s) ds, \\
 (p_2(t) :=) \lim_{n \rightarrow \infty} z_{i_n}(t) &= c^* u_y(t) + \int_{t_2}^t r(t, s; y) F[y, y', \lambda^{(2)}](s) ds
 \end{aligned}$$

uniformly on J and

$$\begin{aligned}
 (11) \quad \alpha_1(p_i(t_1) - p_i(t)|_{J_1}) &= 0, \quad p_i(t_2) = 0, \\
 \alpha_2(p_i(t_3) - p_i(t)|_{J_2}) &= 0 \quad \text{for } i = 1, 2.
 \end{aligned}$$

As above we may verify

$$\begin{aligned}
 p_2(t_1) - p_2(t) &> p_1(t_1) - p_1(t) \quad \text{for all } t \in (t_1, t_2), \\
 p_2(t_3) - p_2(t) &> p_1(t_3) - p_1(t) \quad \text{for all } t \in (t_2, t_3),
 \end{aligned}$$

which contradicts (11). Hence $\{\mu_n\}$ is convergent, and let $\lim_{n \rightarrow \infty} \mu_n = \mu^*$. Then

$$(z^*(t) :=) \lim_{n \rightarrow \infty} z_n(t) = c^* u_y(t) + \int_{t_2}^t r(t, s; y) F[y, y', \mu^*](s) ds$$

uniformly on J , and consequently, z^* is a solution of the differential equation

$$w'' = Q[y, y'](t) w + F[y, y', \mu^*](t)$$

and

$$\alpha_1(z^*(t_1) - z^*(t)|J_1) = 0, \quad z^*(t_2) = 0, \quad \alpha_2(z^*(t_3) - z^*(t)|J_2) = 0.$$

By Lemma 2 it is necessary that $z = z^*$ and $\mu_0 = \mu^*$. Since $\lim_{n \rightarrow \infty} z_n^{(i)}(t) = z^{(i)}(t)$ uniformly on J for $i = 0, 1$, we have $z = \lim_{n \rightarrow \infty} z_n = \lim_{n \rightarrow \infty} T(y_n) = T(y)$ and therefore T is a continuous operator. Let $\varphi \in K$ and $T(\varphi) = y$. Then the equality

$$y''(t) = Q[\varphi, \varphi'](t) y(t) + F[\varphi, \varphi', \mu_0](t)$$

holds on J for some $\mu_0 \in I$, thus $\|y''\| \leq A + r_0 B$ ($:= r_2$) and $K \subset L = \{y; y \in C^2(J), \|y^{(i)}\| \leq r_i \text{ for } i = 0, 1, 2\}$. Since L is a compact subset of Y , K is a relative compact subset of Y .

By the Schauder fixed point theorem there is a fixed point of T . This completes the proof. \square

Remark 1. If $\alpha_1(z) = \alpha_2(z) = z(t_2)$, then Theorem 1 in [2] and Theorem 1 in [4] (where $m = n = 1$) follow from Theorem 1.

Let $t_1 < x_1 < t_2 < x_2 < t_3$. If $\alpha_1(z) = z(x_1)$, $\alpha_2(z) = z(x_2)$, then Theorem 1 in [1] follows from Theorem 1.

Example 1. Consider the functional differential equation

$$(12) \quad y''(t) = y(t) \exp \{ |y(y'(t))| \} + \frac{1}{2} \cos (t + y'(y(t))) + \mu$$

on the interval $J = \langle 0, t_3 \rangle$, where $t_3 \geq 2\sqrt{1+e}$. Let $t_2 \in (0, t_3)$. Assumptions (H_1) – (H_4) are fulfilled with $r_0 = 1$, $r_1 = 2\sqrt{1+e}$ and $I = \langle -\frac{1}{2}, \frac{1}{2} \rangle$. Let $\alpha_1(z) = \int_0^{t_2} z^3(s) ds$ for $z \in C^0((0, t_2))$ and $\alpha_2(z) = \max \{ z(t); t \in \langle t_2, \frac{1}{2}(t_2 + t_3) \rangle \}$ for $z \in C^0(\langle t_2, t_3 \rangle)$. Then by Theorem 1 there is $\mu_0 \in \langle -\frac{1}{2}, \frac{1}{2} \rangle$ such that equation (12) with $\mu = \mu_0$ admits a solution y satisfying

$$\int_0^{t_2} (y(t_1) - y(s))^3 ds = 0, \quad y(t_2) = 0, \quad \max \{ y(t_3) - y(t); t \in \langle t_2, \frac{1}{2}(t_2 + t_3) \rangle \} = 0$$

and

$$\|y\| \leq 1, \quad \|y'\| \leq 2\sqrt{1+e}.$$

4. BOUNDED SOLUTIONS ON A HALFLINE

In this section BVP (1)–(2) is applied to the investigation of bounded solutions of a functional differential equation of type (1) with functional boundary conditions

$$(13) \quad \alpha_1(y(t_1) - y(t)|J_1) = 0, y(t_2) = 0.$$

Let Y be the space of bounded C^0 -functions on the halfline $\langle t_1, \infty \rangle$ with the topology of uniform convergence on compact subintervals of $\langle t_1, \infty \rangle$. Consider the functional differential equation

$$(14) \quad y''(t) - U[y, y'](t)y(t) = V[y, y', \mu](t),$$

where $U: Y \times Y \rightarrow Y$, $V: Y \times Y \times I \rightarrow Y$ are continuous operators, $U[y, z](t) > 0$ for all $t \geq t_1$ and $[y, z] \in Y \times Y$. Further we shall assume that there exists an increasing sequence $\{x_n\} \subset R$, $x_1 > t_2$, $\lim_{n \rightarrow \infty} x_n = \infty$ such that the functions $U[y, z](t)$, $V[y, z, \mu](t)$ are defined on $\langle t_1, x_n \rangle$ only by the restrictions of y, z to the interval $\langle t_1, x_n \rangle$ ($n = 1, 2, \dots$), that is

$$U: Y_n \times Y_n \rightarrow Y_n, \quad V: Y_n \times Y_n \times I \rightarrow Y_n \quad (n = 1, 2, \dots),$$

where Y_n is the Banach space of the C^0 -functions on $\langle t_1, x_n \rangle$ with the sup norm. The differential equation $y'' - q(t, y, y')y = f(t, y, y', \mu)$, where $q \in C^0(\langle t_1, \infty \rangle \times R^2)$, $f \in C^0(\langle t_1, \infty \rangle \times R^2 \times I)$, is a special case of (14).

Suppose there are positive constants r_0, r_1 such that the operators U, V satisfy the following assumptions:

- (C₁) $|V[y, y', \mu](t)| \leq r_0 U[y, y'](t)$ for all $t \geq t_1$ and $[y, y', \mu] \in H \times I$, where $H = \{[y, y']; y \in C^1(\langle t_1, \infty \rangle), |y^{(i)}(t)| \leq r_i \text{ for } t \geq t_1, i = 0, 1\}$;
- (C₂) $V[y, y', \mu_1](t) < V[y, y', \mu_2](t)$ for all $t \geq t_1$, $[y, y'] \in H$ and $\mu_1, \mu_2 \in I$, $\mu_1 < \mu_2$;
- (C₃) $V[y, y', a](t)V[y, y', b](t) \leq 0$ for all $t \geq t_1$ and $[y, y'] \in H$;
- (C₄) $2\sqrt{r_0}\sqrt{A + r_0B} \leq r_1$, where $A = \sup_{t \geq t_1} \{ \sup |V[y, y', \mu](t)|; [y, y', \mu] \in H \times I \}$, $B = \sup_{t \geq t_1} \{ \sup |U[y, y'](t)|; [y, y'] \in H \}$.

Lemma 3. *Assume assumptions (C₁)–(C₄) are fulfilled with positive constants r_0, r_1 . Then for any x_n ($n = 1, 2, \dots$) there exists a $\mu_n \in I$ such that equation (14) with $\mu = \mu_n$ admits a solution y_n defined on the interval $\langle t_1, x_n \rangle$ and satisfying the boundary conditions*

$$(15) \quad \alpha_1(y_n(t_1) - y_n(t)|J_1) = 0, \quad y_n(t_2) = 0, \quad y_n(x_n) = 0 \quad (n = 1, 2, \dots),$$

and, moreover,

$$(16) \quad \begin{aligned} |y_n(t)| &\leq r_0, & |y'_n(t)| &\leq r_1, \\ |y''_n(t)| &\leq A + r_0 B & \text{for } t \in \langle t_1, x_n \rangle, & \quad (n = 1, 2, \dots). \end{aligned}$$

Proof. The proof follows immediately from Theorem 1 if we set $t_3 = x_n$ and $\alpha_2(z) = z(t_2)$. The last inequality in (16) is evident. \square

Theorem 2. Assume assumptions (C_1) – (C_4) are fulfilled with positive constants r_0, r_1 . Then there exists a $\mu_0 \in I$ such that equation (14) with $\mu = \mu_0$ admits a solution y satisfying (13) and

$$(17) \quad |y(t)| \leq r_0, \quad |y'(t)| \leq r_1 \quad \text{for } t \geq t_1.$$

Proof. According to Lemma 3 there exists a sequence $\{y_n\}$ of solutions of equation (14) with $\mu = \mu_n (\in I)$ on the intervals $\langle t_1, x_n \rangle$ satisfying (15) and (16). Using the Ascoli-Arzelà theorem, a diagonal process of Cantor and the fact that $\{\mu_n\}$ is a bounded sequence, we may assume without loss of generality that $\{y_n(t)\}$ and $\{y'_n(t)\}$ are locally uniformly convergent on $\langle t_1, \infty \rangle$ and $\{\mu_n\}$ is convergent. Setting $\lim_{n \rightarrow \infty} y_n(t) = y(t)$ for $t \in \langle t_1, \infty \rangle$ and $\lim_{n \rightarrow \infty} \mu_n = \mu_0$, then y is a solution of equation (14) with $\mu = \mu_0$ satisfying (13) and (17). \square

Example 2. Consider the functional differential equation

$$(18) \quad y''(t) = 6\pi y(t) \exp\{|y(t + (\sin t)^2)|\} + \ln(e + |y'(\sqrt{t})|) \arctan t + (1 + y^2(t))\mu.$$

The assumptions of Theorem 2 are satisfied with $t_1 \geq 1$, $r_0 = 1$, $r_1 = e^3$ and $I = \langle -2\pi, 0 \rangle$. Therefore there exists a $\mu_0 \in \langle -2\pi, 0 \rangle$ such that equation (18) with $\mu = \mu_0$ has a solution y defined on $\langle t_1, \infty \rangle$, and (13) and $|y(t)| \leq 1$, $|y'(t)| \leq e^3$ for $t \geq t_1$ hold.

References

- [1] S. Staněk: On a class of five-point boundary value problems in second-order functional differential equations with parameter, *Acta Math. Hungar.*, to appear.
- [2] S. Staněk: Three-point boundary value problem for nonlinear second-order differential equations with parameter, *Czech. Math. J.* 42 (117) (1992), 241–256.
- [3] S. Staněk: On the boundedness of solutions of nonlinear second-order differential equations with parameter, *Arch. Math. (Brno)* 27 (1991), 229–241.
- [4] S. Staněk: Multi-point boundary value problem for a class of functional differential equations with parameter, *Math. Slovaca* 42 (1992), no. 1, 85–96.
- [5] S. Staněk: Three-point boundary value problem of retarded functional differential equation of the second order with parameter, *Acta UPO, Fac. rer. Nat.* 97 Math. XXIX. (1990), 107–121.

Author's address: Department of Math. Analysis, Palacký University, tř. Svobody 26, 771 46 Olomouc, Czech Republic.