

Mohammad Ismail

A topological characterization of stationary sets

Czechoslovak Mathematical Journal, Vol. 43 (1993), No. 4, 765–767

Persistent URL: <http://dml.cz/dmlcz/128435>

Terms of use:

© Institute of Mathematics AS CR, 1993

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A TOPOLOGICAL CHARACTERIZATION OF STATIONARY SETS

MOHAMMAD ISMAIL, Slippery Rock

(Received May 25, 1992)

If X is a topological space and $x \in X$, then let $t(x, X)$ denote the ‘tightness of X at x ’. Recall that $t(x, X)$ is the smallest cardinal number τ such that if $A \subseteq X$ and $x \in \overline{A}$, then there exists $B \subseteq A$ such that $|B| \leq \tau$ and $x \in \overline{B}$. The ‘tightness’ of X is defined as

$$t(X) = \sup\{t(x, X) : x \in X\}.$$

Let κ denote an uncountable regular cardinal with the usual order topology. A subset S of κ is called ‘stationary’ if for every closed unbounded (cub) subset C of κ , $C \cap S \neq \varnothing$. Stationary sets, with the subspace topology, are interesting topological spaces and their properties have been extensively studied in such articles as [F] and [ν DL]. The aim of this article is to prove the following topological characterization of stationary sets.

Theorem. *For a subset S of κ , the following are equivalent:*

(a) S is stationary

(b) *If X is a compact Hausdorff space and for each $x \in X$, $t(x, X) < \kappa$, then for any continuous function $f: S \rightarrow X$, there exists a unique point $x \in X$ such that $|f^{-1}(x)| = \kappa$.*

(c) *If X is a compact Hausdorff space which contains a subspace homeomorphic to S , then there exists $x \in X$ such that $t(x, X) \geq \kappa$.*

Proof. (a) \rightarrow (b) Let S be stationary and let $f: S \rightarrow X$ be a continuous function, where X is a compact Hausdorff space and for each $x \in X$, $t(x, X) < \kappa$.

For each $\beta < \kappa$, let $S_\beta = \{\alpha \in S : \alpha \geq \beta\}$ and let $Y_\beta = \overline{f(S_\beta)}$. Then $\{Y_\beta : \beta < \kappa\}$ is a decreasing family of nonempty closed subsets of X . Therefore, $\bigcap_{\beta < \kappa} Y_\beta \neq \varnothing$. Let $x \in \bigcap_{\beta < \kappa} Y_\beta$.

Let us show that $|f^{-1}(x)| = \kappa$. Assume the contrary and let $|f^{-1}(x)| < \kappa$. Then there exists $\gamma < \kappa$ such that $f^{-1}(x) \cap S_\gamma = \varnothing$. Then for each $\alpha \in S_\gamma$, there exists

a neighborhood U_α of $f(\alpha)$ in X such that $x \notin \overline{U_\alpha}$. By the continuity of f , there exists $\beta_\alpha < \alpha$ such that $f((\beta_\alpha, \alpha] \cap S) \subseteq U_\alpha$. Then $x \notin f((\beta_\alpha, \alpha] \cap S)$. Since the map $\alpha \rightarrow \beta_\alpha$ is a regressive map of the stationary set S_γ into κ , by Fodor's theorem for stationary sets (otherwise known as the Pressing Down Lemma) (See [F] or [J, Theorem 22, p. 59]), there is a stationary subset A of S_γ and a $\beta < \kappa$ such that $\beta_\alpha = \beta$, for each $\alpha \in A$. Thus $x \notin f((\beta, \alpha] \cap S)$, for each $\alpha \in A$. Therefore, if $B \subseteq \overline{S_\beta}$ and $|B| < \kappa$, then, since there exists $\alpha \in A$ such that $f(B) \subseteq f((\beta, \alpha] \cap S)$, $x \notin f(B)$. This is a contradiction since $x \in Y_\beta = \overline{f(S_\beta)}$ and $t(x, X) < \kappa$. Hence $|f^{-1}(x)| = \kappa$.

Note that the point x is unique. Indeed, if $y \in X$ and $y \neq x$, then since $f^{-1}(x)$ and $f^{-1}(y)$ are disjoint closed subsets of S , they cannot be both unbounded. Hence $|f^{-1}(y)| < \kappa$.

(b) \rightarrow (c) is obvious

(c) \rightarrow (a)

Let S be a nonstationary subset of κ . Then $S \subseteq \kappa \setminus C$, for some cub subset C of κ . We will show that $\kappa \setminus C$, and hence S , can be embedded in a compact Hausdorff space X such that $t(x, X) < \kappa$, for each $x \in X$.

Let $\{P_\alpha : \alpha \in J\}$ be the collection of all maximal convex subsets of $\kappa \setminus C$. Note that P_α 's are disjoint clopen subsets of $\kappa \setminus C$ and $\kappa \setminus C = \bigcup_{\alpha \in J} P_\alpha$. Hence $\kappa \setminus C = \oplus \{P_\alpha : \alpha \in J\}$, the disjoint sum of P_α 's. Since each P_α is a bounded subset of κ , it can be embedded in a compact Hausdorff space Y_α such that $t(y, Y_\alpha) < \kappa$, for each $y \in Y_\alpha$. For example, if $P_\alpha \subseteq [0, \beta]$, for some $\beta < \kappa$, then $[0, \beta]$ is one such space. Let Y be the disjoint sum of Y_α 's. Then Y is a locally compact Hausdorff space and $\kappa \setminus C$ is embedded in Y . Let $X = Y \cup \{x\}$ be the one-point compactification of Y . Note that $t(x, X) \leq \omega_0 < \kappa$ and for each $y \in Y$, $t(y, X) = t(y, Y) < \kappa$. Hence X is the desired compact Hausdorff space. \square

From (c) \rightarrow (a), it follows that if $\kappa = \nu^+$ is a successor cardinal, then every nonstationary subset of κ can be embedded in a compact Hausdorff space X such that $t(X) \leq \nu < \kappa$. Also, if $\kappa = \omega_1$, each nonstationary subset of κ can be embedded in a disjoint sum of first countable compact Hausdorff spaces. The one-point compactification of such a disjoint sum is sequential. We, therefore, obtain the following.

Theorem. *If κ is a successor cardinal (respectively, if $\kappa = \omega_1$), then a subset T of κ is nonstationary if and only if T can be embedded in a compact Hausdorff space X such that $t(X) < \kappa$ (respectively, X is sequential).*

The following example shows that the above theorem is not true for regular limit cardinals.

Example. Let κ be an uncountable regular limit cardinal, and let C be the set of all limit cardinals less than κ . Then C is a club subset of κ . Let $T = \kappa \setminus C$. Then T is nonstationary. Let us show that $t(T) = \kappa$.

Let ν be an arbitrary cardinal less than κ and let $A = \{\alpha \in T : \nu < \alpha < \nu^+\}$. Then $\nu^+ \in \bar{A}$ in T , but for any subset B of A such that $|B| \leq \nu$, $\nu^+ \notin \bar{B}$ in T . Therefore, $t(T) \geq \nu^+$. Since ν is arbitrary, it follows that $t(T) = \kappa$. Hence, T cannot be embedded in any space of tightness less than κ .

Example. For every uncountable regular cardinal κ , there exists a first countable countably compact Hausdorff space of cardinality κ which cannot be embedded in a compact Hausdorff space of tightness less than κ . Let $S = \{\alpha < \kappa : cf(\alpha) = \omega_0\}$. Then S is such a space. Note that S is a stationary subset of κ .

References

- [F] *W. G. Fleissner: Applications of Stationary sets in Topology, Surveys in General Topology, Academic Press, 1980, pp. 163–193.*
- [J] *T. Jeck: Set Theory, Academic Press, 1978.*
- [ν DL] *E. K. Van Douwen and D. J. Lutzer: On the Classification of Stationary Sets, Michigan Math. J. 26 (1979), 47–64.*

Author's address: Department of Mathematics, Slippery Rock University, Slippery Rock, PA 16057, U.S.A.