

Ivan Chajda; M. Kotle

Subdirectly irreducible and congruence distributive  $q$ -lattices

*Czechoslovak Mathematical Journal*, Vol. 43 (1993), No. 4, 635–642

Persistent URL: <http://dml.cz/dmlcz/128438>

## Terms of use:

© Institute of Mathematics AS CR, 1993

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

SUBDIRECTLY IRREDUCIBLE  
AND CONGRUENCE DISTRIBUTIVE  $Q$ -LATTICES

I. CHAJDA, M. KOTRLE, Olomouc

(Received February 13, 1992)

By a  $q$ -lattice (see [3]) we mean an algebra  $A = (A; \vee, \wedge)$  with two binary operations satisfying the following identities:

(associativity):	$a \vee (b \vee c) = (a \vee b) \vee c,$	$a \wedge (b \wedge c) = (a \wedge b) \wedge c,$
(commutativity):	$a \vee b = b \vee a,$	$a \wedge b = b \wedge a,$
(weak absorption):	$a \vee (a \wedge b) = a \vee a,$	$a \wedge (a \vee b) = a \wedge a,$
(weak idempotence):	$a \vee (b \vee b) = a \vee b,$	$a \wedge (b \wedge b) = a \wedge b,$
(equalization):	$a \vee a = a \wedge a.$	

A  $q$ -lattice  $A$  is called *distributive* if it satisfies the distributive identity

$$a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$$

for each  $a, b, c$  from  $A$ . A  $q$ -lattice  $A$  is *bounded* if there exist elements 0 and 1 of  $A$  such that

$$a \wedge 0 = 0 \quad \text{and} \quad a \vee 1 = 1$$

for each  $a \in A$ .

An element  $a$  of a  $q$ -lattice  $A$  is called an *idempotent* if  $a \vee a = a$  (and, by equalization, also  $a \wedge a = a$ ). The set of all idempotents of  $A$  is called the *skeleton* of  $A$ . It is clear that the skeleton of  $A$  is a sub- $q$ -lattice of  $A$  which is the maximal sublattice contained in  $A$ .

A non-singleton subset  $C$  of a  $q$ -lattice  $A$  is called a *cell* of  $A$  if  $a, b \in C$  implies  $a \vee a = b \vee b$  and  $C$  is a maximal subset of  $A$  with respect to this property.

Evidently, a  $q$ -lattice  $A$  is a lattice if and only if it has no cell, i.e. if  $A$  is equal to its skeleton. Every cell  $C$  of  $A$  has just one idempotent.

Evidently, every cell  $D$  of a  $q$ -lattice  $A$  is a sub- $q$ -lattice of  $A$ . If  $A$  is a cell, then the skeleton of  $A$  is a singleton.

Distributive and/or bounded  $q$ -lattices were investigated in [4]. Let us notice that the distributive identity is equivalent to its dual; on the other hand, the foregoing identities for 0 and 1 do not imply  $a \vee 0 = a$  and  $1 \wedge a = a$  but only the weaker laws  $a \vee 0 = a \vee a$  and  $a \wedge 1 = a \wedge a$ .

By the foregoing definitions, the class of all distributive  $q$ -lattices as well as the class of all bounded distributive lattices form varieties. Therefore, it makes sense to look for SI-members of these varieties. Although  $q$ -lattices look rather similar to lattices, these varieties have another number of SI-members than the variety of (bounded) distributive lattices.

**Theorem 1.** *The variety  $D$  of all distributive  $q$ -lattices has exactly two non-trivial SI-members, namely those visualized in Fig. 1 as  $B$  and  $C$ .*



Fig. 1

Before proceeding to proof, let us remark that every  $q$ -lattice  $A = (A; \vee, \wedge)$  can be viewed as a quasiordered set  $(A; Q)$ , where the quasiorder  $Q$  on  $A$  is induced by  $\vee$  (or  $\wedge$ ) as follows (see e.g. [3], [4]):

$$\langle a, b \rangle \in Q \quad \text{iff} \quad a \vee b = b \vee b$$

(or, equivalently,  $\langle a, b \rangle \in Q$  iff  $a \wedge b = a \wedge a$ ). Henceforth, we can visualize this quasiorder  $Q$  in the diagrams of  $q$ -lattices by oriented arrows; i. e.  $\langle a, b \rangle \in Q$  iff there exists an oriented path from  $a$  to  $b$  consisting of arrows.

**Proof of Theorem 1.** Since both  $B$  and  $C$  are two-element  $q$ -lattices, they are subdirectly irreducible. Hence it remains to prove that any other non-trivial distributive  $q$ -lattice  $A$  different from  $B, C$  is subdirectly reducible.

(i) If  $A$  has no cell, then  $A$  is a lattice. In the case of  $A \neq B, C$ ,  $A$  is subdirectly reducible by [2].

(ii) Let  $D$  be a cell of  $A$ .

(a) Let  $A$  contain an element  $a \notin D$ . Denote by  $d$  the idempotent of  $D$  and

$$A_1 = A - (D - \{d\}).$$

Then  $A_1$  and  $D$  are sub- $q$ -lattices of  $A$  and  $\text{card } A_1 > 1$ ,  $\text{card } D > 1$ . Introduce a mapping  $\alpha: A \rightarrow A_1 \times D$  by the rule

$$\begin{aligned}\alpha(x) &= \langle x, d \rangle \quad \text{for } x \in A - (D - \{d\}), \\ \alpha(x) &= \langle d, x \rangle \quad \text{for } x \in D.\end{aligned}$$

It is clear that  $\alpha$  is an injection and  $\text{pr}_1\alpha(A) = A_1$ ,  $\text{pr}_2\alpha(A) = D$ . Prove that  $\alpha$  is a homomorphism:

if  $x \in A_1$ ,  $y \in D$ , then

$$\begin{aligned}\alpha(x \vee y) &= \alpha(x \vee d) = \langle x \vee d, d \rangle, \\ \alpha(x) \vee \alpha(y) &= \langle x, d \rangle \vee \langle d, y \rangle = \langle x \vee d, d \rangle;\end{aligned}$$

if  $x, y \in A_1$ , then

$$\alpha(x \vee y) = \langle x \vee y, d \rangle = \langle x, d \rangle \vee \langle y, d \rangle = \alpha(x) \vee \alpha(y);$$

if  $x, y \in D$ , then

$$\alpha(x \vee y) = \alpha(d) = \langle d, d \rangle = \langle d, x \rangle \vee \langle d, y \rangle = \alpha(x) \vee \alpha(y).$$

Dually this can be shown for the meet. Hence  $A$  is subdirectly reducible.

(b) Suppose  $A = D$ . If  $A \neq C$ , there exist elements  $a, b$  of  $D$  such that  $a \neq b \neq d \neq a$ . Put  $A_1 = A - \{b\}$  and  $A_2 = \{d, b\}$ . Thus  $\text{card } A_1 > 1$ ,  $\text{card } A_2 > 1$ . Introduce a mapping  $\alpha: A \rightarrow A_1 \times A_2$  as follows:

$$\begin{aligned}\alpha(x) &= \langle x, d \rangle \quad \text{for } x \in A_1, \\ \alpha(x) &= \langle d, x \rangle \quad \text{for } x \in A_2.\end{aligned}$$

Evidently,  $\alpha$  is an injection and  $\text{pr}_1\alpha(A) = A_1$ ,  $\text{pr}_2\alpha(A) = A_2$ . Prove that  $\alpha$  is a homomorphism:

if  $x \in A_1$ ,  $y \in A_2$ , then

$$\begin{aligned}\alpha(x \vee y) &= \alpha(d) = \langle d, d \rangle, \\ \alpha(x) \vee \alpha(y) &= \langle x, d \rangle \vee \langle d, y \rangle = \langle d, d \rangle;\end{aligned}$$

if  $x, y \in A_1$ , then

$$\alpha(x \vee y) = \alpha(d) = \langle d, d \rangle = \langle x, d \rangle \vee \langle y, d \rangle = \alpha(x) \vee \alpha(y);$$

if  $x, y \in A_2$  then

$$\alpha(x \vee y) = \alpha(d) = \langle d, d \rangle = \langle d, x \rangle \vee \langle d, y \rangle = \alpha(x) \vee \alpha(y).$$

Dually this can be done for  $\wedge$ , i.e.  $A$  is a subdirect product of  $A_1, A_2$ . □

**Theorem 2.** *The class of all bounded distributive  $q$ -lattices with  $0 \neq 1$  has exactly three nontrivial SI-members, namely  $B$  (in Fig. 1),  $C_1, C_2$  (in Fig. 2).*




Fig. 2

**Proof.** As was already mentioned,  $B$  is subdirectly irreducible. Since the lattices of congruences not collapsing  $0$  and  $1$  of  $C_1, C_2$  are three-element chains, see Fig. 3, also  $C_1, C_2$  are subdirectly irreducible in this class.




Fig. 3

We have to prove that if  $A$  is a bounded distributive  $q$ -lattice different from  $B, C_1, C_2$  then  $A$  is subdirectly reducible in this class.

- (A) If  $A$  has no cell than this was done by G. Birkhoff in [2].
- (B) If  $A$  has at least two cells, say  $D_1, D_2$ , then clearly  $D_1 \cap D_2 = \emptyset$ . Put

$$\Theta_1 = D_1 \times D_1 \cup \omega, \quad \Theta_2 = D_2 \times D_2 \cup \omega$$

where  $\omega$  denotes the identity relation on  $A$ . It can be easily shown that  $\Theta_1, \Theta_2$  are congruences on  $A$  and  $\Theta_1 \cap \Theta_2 = \omega$ ; thus, by the Birkhoff Theorem [2],  $A$  is subdirectly reducible.

- (C) It remains to deal with the case when  $A$  has just one cell  $D$ .

(i) Suppose that the skeleton of  $A$  contains just two elements, namely  $0$  and  $1$ . Let  $0 \in D$ . Since  $A$  is not isomorphic with  $C_2$ , it means that  $D$  contains at least two non-idempotent elements  $a, b$ , i.e.  $a \neq 0 \neq b \neq a$ . We can put

$$A_1 = \{0, 1, a\}, \quad A_2 = A - \{a\}.$$

It is easy to see that both  $A_1, A_2$  are bounded distributive  $q$ -lattices (moreover,  $A_1 \simeq C_2$ ). Define  $\alpha: A \rightarrow A_1 \times A_2$  as follows:

$$\begin{aligned} \alpha(0) &= \langle 0, 0 \rangle, & \alpha(1) &= \langle 1, 1 \rangle, \\ \alpha(a) &= \langle a, 0 \rangle; \\ \alpha(x) &= \langle 0, x \rangle \text{ for } x \in D, x \neq a. \end{aligned}$$

We can see that  $\alpha$  is an injection and  $\text{pr}_1\alpha(A) = A_1$ ,  $\text{pr}_2\alpha(A) = A_2$ . It remains to prove that  $\alpha$  is a homomorphism. It is almost evident in the case  $z, y \in A_1$  that  $\alpha(z \vee y) = \alpha(z) \vee \alpha(y)$  and  $\alpha(z \wedge y) = \alpha(z) \wedge \alpha(y)$ , and analogously for  $z, y \in A_2$ . Suppose  $z \in A_1 - A_2$ ,  $y \in A_2 - A_1$ . Then  $z = a$  and  $y \in D$ ,  $y \neq a$ ,  $y \neq 0$ . We have

$$\begin{aligned}\alpha(z) \vee \alpha(y) &= \alpha(a) \vee \alpha(y) = \langle a, 0 \rangle \vee \langle 0, x \rangle = \langle 0, 0 \rangle, \\ \alpha(z \vee y) &= \alpha(0) = \langle 0, 0 \rangle \quad \text{and} \\ \alpha(z) \wedge \alpha(y) &= \langle a, 0 \rangle \wedge \langle 0, x \rangle = \langle 0, 0 \rangle = \alpha(0) = \alpha(z \wedge y).\end{aligned}$$

Consequently,  $A$  is isomorphic to a subdirect product of  $A_1, A_2$ .

(ii) If the skeleton of  $A$  contains just two elements (0 and 1) and  $1 \in D$ , where  $D$  is the unique cell of  $A$ , the proof is dual to that of (i).

(iii) Let the skeleton of  $A$  have more than two elements. We have three cases:

(a) Suppose there exists an idempotent  $d \in A$  with  $0 \neq d \neq 1$  and  $d \in D$ .

Put

$$A_1 = \{x; \langle x, d \rangle \in Q\} \quad \text{and} \quad A_2 = \{x; \langle d, x \rangle \in Q\}$$

for the induced quasiorder  $Q$ . Define  $\alpha: A \rightarrow A_1 \times A_2$  as follows:

$$\begin{aligned}\alpha(x) &= \langle x \wedge d, x \vee d \rangle \quad \text{for } x \notin D \quad \text{and} \\ \alpha(x) &= \langle x, x \rangle \quad \text{for } x \in D.\end{aligned}$$

Since  $x \notin D$  is an idempotent of  $A$  (because  $A$  has just one cell  $D$ ), it is easy to verify that  $\alpha$  is an injective homomorphism satisfying  $\text{pr}_1\alpha(A) = A_1$ ,  $\text{pr}_2\alpha(A) = A_2$ , thus  $A$  is isomorphic to a subdirect product of  $A_1, A_2$ .

(b) Suppose there exists an idempotent  $d \in A$  with  $0 \neq d \neq 1$  and  $0 \in D$ .

Put

$$A_1 = \{x; \langle x, d \rangle \in Q\}, \quad A_2 = \{x; \langle d, x \rangle \in Q\}$$

and introduce a mapping  $\alpha: A \rightarrow A_1 \times A_2$  by

$$\begin{aligned}\alpha(x) &= \langle x \wedge d, x \vee d \rangle \quad \text{for } x \notin D, \\ \alpha(x) &= \langle x, d \rangle \quad \text{for } x \in D.\end{aligned}$$

We can easily verify that  $\alpha$  is an injective homomorphism with  $\text{pr}_i\alpha(A) = A_i$  ( $i = 1, 2$ ), thus  $A$  is a subdirect product of  $A_1, A_2$ .

(c) The last case with  $d \in A$ ,  $0 \neq d \neq 1$ ,  $1 \in D$  is dual to (b), only  $\alpha$  is defined for  $x \in D$  by  $\alpha(x) = \langle d, x \rangle$ .  $\square$

**Corollary.** Every non-trivial distributive  $q$ -lattice  $A$  is a subdirect product of  $q$ -lattices  $B$  and  $C$ . Every bounded distributive  $q$ -lattice  $A$  with  $0 \neq 1$  is a subdirect product of  $q$ -lattices  $B, C_1, C_2$ .

It is well-known than for any lattice  $L$ , the congruence lattice  $\text{Con } L$  is distributive, see e.g. [1]. We can ask if a similar result is also valid for  $q$ -lattices. It is easy to show that the answer is negative in the general case. More precisely, we can state

**Lemma.** Let  $C$  be a  $q$ -lattice which is a cell. Then  $\text{Con } C \simeq \Pi_n$ , where  $n = \text{card } C$  and  $\Pi_n$  is the partition lattice of the set of cardinality  $n$ .

The proof is trivial since every equivalence on  $C$  is a congruence.

**Theorem 3.** Let  $A$  be a  $q$ -lattice which has just one  $n$ -element cell  $C$ , let  $S$  be the skeleton of  $A$ . Then  $\text{Con } A \simeq \Pi_n \times \text{Con } S$ .

**Proof.** (a) If  $\Theta_1 \in \text{Con } S$  and  $\Theta_2 \in \text{Con } C \simeq \Pi_n$  and  $d$  is the only idempotent of  $C$  (i.e.  $\{d\} = S \cap C$ ), then clearly

$$\Theta_1 \cup \Theta_2 \cup \{[d]_{\Theta_1} \cup [d]_{\Theta_2}\}^2 \in \text{Con } A.$$

(b) If  $\Theta \in \text{Con } A$ , put  $\Theta_1 = \Theta \cap S^2$ ,  $\Theta_2 = \Theta \cap C^2$ .

Evidently,  $\Theta = \Theta_1 \cup \Theta_2 \cup \{[d]_{\Theta_1} \cup [d]_{\Theta_2}\}^2$ . Hence each  $\Theta \in \text{Con } A$  is of the above mentioned form, i.e. it is uniquely determined by some  $\Theta_1 \in \text{Con } S$  and  $\Theta_2 \in \text{Con } C$ , i.e. the mapping

$$h: \Theta \rightarrow \langle \Theta_2, \Theta_1 \rangle$$

is a bijection of  $\text{Con } A$  onto  $\Pi_n \times \text{Con } S$ . It is easy to show that  $h$  is an isomorphism. □

**Theorem 4.** For a  $q$ -lattice  $A$ , the congruence lattice  $\text{Con } A$  is distributive if and only if  $A$  contains at most one cell with at most 2 elements.  $\text{Con } A$  is modular if and only if  $A$  contains at most one cell with at most 3 elements.

**Proof.** If  $A$  has no cell, then  $A$  is a lattice and  $\text{Con } A$  is distributive, see [1].

If  $A$  contains just one  $n$ -element cell then, by Theorem 3,  $\text{Con } A \simeq \Pi_n \times \text{Con } S$ , where  $S$  is the skeleton of  $A$ . However,  $\Pi_n$  is distributive if and only if  $n \leq 2$ ,  $\Pi_n$  is modular if and only if  $n \leq 3$  (see e.g. Ex. 5 of Par. 9, Ch. IV in [1]). Since  $\text{Con } S$  is distributive, we arrive at the statement.

On the contrary, suppose  $A$  has at least two cells  $C_1, C_2$ . Let  $a_i$  be an idempotent of  $C_i$  and  $b_1 \in C_1, b_1 \neq a_1, b_2 \in C_2, b_2 \neq a_2$ . Then clearly  $\langle a_1, a_2 \rangle = \langle b_1 \vee b_2 \rangle \vee \langle b_1 \vee b_2 \rangle$ , i.e.

$$\Theta(a_1, a_2) \subseteq \Theta(b_1, b_2).$$

But  $\langle b_1, b_2 \rangle \notin \Theta(a_1, a_2)$ , i.e.  $\Theta(a_1, a_2) \neq \Theta(b_1, b_2)$ .

(i) If  $a_1 < a_2$  then the congruences

$$\Theta(a_1, b_1), \Theta(a_1, a_2), \Theta(b_1, b_2), \Theta(a_1, b_1) \wedge \Theta(a_1, a_2), \Theta(a_1, b_1) \vee \Theta(b_1, b_2)$$

form a sublattice of  $\text{Con } A$  isomorphic to  $N_5$ , see Figs. 4 and 5.




Fig. 4




Fig. 5

(ii) If  $a_1 \parallel a_2$ , then the congruences

$$\Theta(a_1, a_1 \wedge a_2), \Theta(a_1, a_2), \Theta(b_1, b_2),$$

$$\Psi = \Theta(b_1, a_1 \wedge a_2) \vee \Theta(b_2, a_1 \vee a_2),$$

$$\Phi = \psi \vee \Theta(b_1, b_2)$$



form a sublattice of  $\text{Con } A$  isomorphic with  $N_5$  again, see Figs. 6 and 7. □




Fig. 6




Fig. 7

*References*

- [1] *G. Birkhoff*: Lattice Theory, Publ. AMS, 3<sup>rd</sup> edition (1973).
- [2] *G. Birkhoff*: Subdirect unions in universal algebras, Bull. AMS 50 (1944), 764–768.
- [3] *I. Chajda*: Lattices in quasiordered sets, Acta Palack. Univ. Olomouc 31 (1992), 6–12.
- [4] *I. Chajda*: An algebra of quasiordered logic, submitted to Math. Bohemica.

*Author's address*: Dept. of Algebra and Geometry, Sci. Faculty, Palacký University, Tomkova 38, 779 00 Olomouc, Czech Republic.