

Ján Jakubík

On iterated limits of subsets of a convergence ℓ -group

Mathematica Bohemica, Vol. 126 (2001), No. 1, 53–61

Persistent URL: <http://dml.cz/dmlcz/133921>

Terms of use:

© Institute of Mathematics AS CR, 2001

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON ITERATED LIMITS OF SUBSETS OF
A CONVERGENCE ℓ -GROUP

JÁN JAKUBÍK, Košice

(Received February 12, 1999)

Abstract. In this paper we deal with the relation

$$\lim_{\alpha} \lim_{\alpha} X = \lim_{\alpha} X$$

for a subset X of G , where G is an ℓ -group and α is a sequential convergence on G .

Keywords: convergence ℓ -group, disjoint subset, direct product, lexico extension

MSC 2000: 06F15, 22C05

For a convergence ℓ -group (shorter: cl-group) we apply the same notation and definitions as in [4] with the distinction that now we do not assume the commutativity of the group operation.

Let (G, α) be a cl-group (where G is an ℓ -group and α is a convergence on G). For $X \subseteq G$ the symbol $\lim_{\alpha} X$ has the usual meaning. X will be said to be regular with respect to (G, α) if the relation

$$\lim_{\alpha} \lim_{\alpha} X = \lim_{\alpha} X$$

is valid.

An ℓ -group G will be called absolutely regular, if whenever (G, α) is a convergence ℓ -group and H is an ℓ -subgroup of G , then H is regular with respect to (G, α) .

We denote by F the class of all ℓ -groups K such that each disjoint subset of K is finite; such ℓ -groups were studied in [1] (cf. also [2] and [6]).

Supported by grant VEGA 2/5125/99.

In the present paper we prove that each ℓ -group belonging to F is absolutely regular.

This generalizes a result from [5] concerning ℓ -groups which can be represented as direct products of a finite number of linearly ordered groups.

1. PRELIMINARIES

In the whole paper G is an ℓ -group; the group operation is written additively, but we do not assume commutativity of this operation.

For the notion of convergence $\alpha \in \text{conv } G$ we apply the same definition as in [4] with the distinction that to the conditions for α used in [4] we add the following one:

(*) α is a normal subset of $(G^N)^+$ (i.e., if $s \in (G^N)^+$, then $s + \alpha = \alpha + s$).

The corresponding convergence ℓ -group will be denoted by (G, α) .

If X is a nonempty subset of G , then by $\lim_{\alpha} X$ we denote the set of all $g \in G$ such that there exists a sequence $(x_n) \in X$ with $x_n \rightarrow_{\alpha} g$.

It is easy to verify that

- (i) if X is an ℓ -subgroup of G , then $\lim_{\alpha} X$ is an ℓ -subgroup of G as well;
- (ii) if X is convex in G , then the same holds for $\lim_{\alpha} X$.

We shall often apply the following rule:

If $x_n \rightarrow_{\alpha} g$ and $x_n \leq g$ for each $n \in N$, then $\bigvee_{n \in N} x_n = g$ (and dually).

A subset Y of G is called disjoint if $Y \subseteq G^+$ and $y_1 \wedge y_2 = 0$ whenever y_1 and y_2 are distinct elements of G .

The direct product of ℓ -groups G_1, G_2, \dots, G_k is defined in the usual way; it will be denoted by $G_1 \times G_2 \times \dots \times G_k$.

If H is a convex ℓ -subgroup of G such that $g > h$ for each $g \in G^+ \setminus H$ and each $h \in H$, then G is said to be a lexico extension of H ; we express this fact by writing $G = \langle H \rangle$. For the properties of the lexico extension cf., e.g., [2].

2. AUXILIARY RESULTS

Let (G, α) be a cl-group.

2.1. Lemma. *Let (x_n) be a sequence in G , $x_n \leq x_{n+1}$ for each $n \in N$, $g \in G$, $x_n \rightarrow_{\alpha} g$. Then $\bigvee_{n \in N} x_n = g$.*

P r o o f. If there exists a subsequence (x_n^1) of (x_n) such that $x_n^1 \leq g$ for each $n \in N$, then $\bigvee_{n \in N} x_n^1 = g$, and hence we have also $\bigvee_{n \in N} x_n = g$. If such a subsequence

(x_n^1) does not exist, then there is a subsequence (x_n^2) of (x_n) such that for each $n \in N$, either $x_n^2 > g$ or x_n^2 is incomparable with g . Hence $x_n^2 \vee g > g$ for each $n \in N$. Thus we obtain

$$(*_1) \quad x_n^2 \vee g \rightarrow_\alpha g$$

and

$$g < x_n^2 \vee g \leq x_n^2 \vee g \quad \text{for each } n \in N,$$

so that the relation $(*_1)$ cannot be valid. □

2.2. Lemma. *Let H be an ℓ -subgroup of the ℓ -group G . Suppose that H can be represented as a lexico extension $H = \langle A \rangle$ with $A \neq \{0\}$. Then*

$$\lim_\alpha H = \bigcup_{h \in H} \lim_\alpha (h + A).$$

Moreover, if $h_1, h_2 \in H$ and $h_1 \notin h_2 + A$, then

$$\lim_\alpha (h_1 + A) \cap \lim_\alpha (h_2 + A) = \emptyset.$$

P r o o f. For $h \in H$ we put $\bar{h} = h + A$. If $h_1, h_2 \in H$ and $h_1 \notin h_2 + A$, then from the properties of the lexico extension we infer that either

$$(i) \quad h'_1 < h'_2 \text{ for each } h'_1 \in h_1 + A \text{ and each } h'_2 \in h_2 + A,$$

or

$$(ii) \quad h'_2 < h'_1 \text{ for each } h'_1 \in h_1 + A \text{ and each } h'_2 \in h_2 + A.$$

Let $g \in G$ and suppose that there exists a sequence (h_n) in H such that $h_n \rightarrow_\alpha g$.

a) First suppose that there exist $h_1 \in H$ and a subsequence (h'_n) of (h_n) such that $h'_n \in h_1 + A$ for each $n \in N$. Then $h'_n \rightarrow_\alpha g$, whence $g \in \lim_\alpha (h_1 + A)$.

b) Now suppose that the assumption from a) is not valid. Then there exists a subsequence (h'_n) of (h_n) such that, whenever $n(1)$ and $n(2)$ are distinct positive integers, then

$$h'_{n(1)} + A \neq h'_{n(2)} + A.$$

Thus in view of the relations (i) and (ii) above, if $n(1)$ and $n(2)$ are distinct, then either $h'_{n(1)} < h'_{n(2)}$ or $h'_{n(1)} > h'_{n(2)}$. This implies that there exists a subsequence (h''_n) of (h'_n) such that either

$$h''_n < h''_{n+1} \quad \text{for each } n \in N,$$

or

$$h''_n > h''_{n+1} \quad \text{for each } n \in N.$$

Suppose that the first case occurs (in the second case we apply a dual argument). We have $h''_n \rightarrow_\alpha g$ and thus according to 2.1 the relation

$$\bigvee_{n \in N} h''_n = g$$

is valid.

If there exists $n(1) \in N$ such that $h''_{n(1)} + A = g + A$, then $h''_{n(1)+1} > g$, which is a contradiction. Hence

$$h''_{n(1)} + A \neq g + A \quad \text{for each } n(1) \in N.$$

Since $A \neq \{0\}$, there exists $a \in A$ with $a > 0$. Then

$$h''_n < g - a \quad \text{for each } n \in N,$$

which is impossible. Thus we have verified that the condition from a) must be valid. Therefore

$$\bigcup_{h \in H} \lim_\alpha (h + A) \subseteq \lim_\alpha H \subseteq \bigcup_{h \in H} \lim_\alpha (h + A),$$

which proves the first assertion of the lemma.

c) Let g be as above; we have shown that there is $h_1 \in H$ such that $g \in \lim_\alpha (h_1 + A)$. Let $h_2 \in H$, $h_1 \notin h_2 + A$. By way of contradiction, suppose that $g \in \lim_\alpha (h_2 + A)$. Hence there exists a sequence (h_n^2) in $h_2 + A$ such that $h_n^2 \rightarrow_\alpha g$. At the same time, there exists a sequence (h_n^1) in $h_1 + A$ such that $h_n^1 \rightarrow_\alpha g$. Let a be as above. If (i) is valid, then

$$h_n^1 + a < h_n^2 \quad \text{for each } n \in N,$$

thus $g + a \leq g$, which is a contradiction. In the case when (ii) is valid we proceed dually. \square

2.3. Lemma. *Let H be as in 2.2. Then $\lim_\alpha H = \langle \lim_\alpha A \rangle$.*

Proof. We obviously have $\lim_\alpha A \subseteq \lim_\alpha H$ and thus $\lim_\alpha A$ is an ℓ -subgroup of $\lim_\alpha H$. Let $h_1, h_2 \in \lim_\alpha A$, $h \in \lim_\alpha H$, $h_1 \leq h \leq h_2$. Then there exist sequences $(h_n^1), (h_n^2)$ in A and (h'_n) in H such that

$$h_n^1 \rightarrow_\alpha h_1, \quad h_n^2 \rightarrow_\alpha h_2, \quad h'_n \rightarrow_\alpha h.$$

Put $(h'_n \vee h_n^1) \wedge h_n^2 = h''_n$. Then $h''_n \in A$ for each $n \in N$ and

$$h''_n \rightarrow_\alpha (h \vee h_1) \wedge h_2 = h,$$

whence $h \in \lim_\alpha A$. Thus $\lim_\alpha A$ is a convex subset of $\lim_\alpha H$.

Let $h \in (\lim_\alpha H)^+ \setminus \lim_\alpha A$. In view of 2.2 there exist $h^1 \in H$ and a sequence (h_n) in $h^1 + A$ such that $h_n \rightarrow_\alpha h$. Moreover, h^1 does not belong to A . Since $h \in G^+$, without loss of generality we can suppose that all h_n belong to G^+ . Further, 2.2 yields that there is a subsequence (h_n^1) of (h_n) such that for each $n \in N$ the relation $h_n^1 \notin A$ is valid. Thus $h_n^1 > a$ for each $a \in A$. Therefore $h \geq a$; since $h \notin A$ we obtain that $h > a$ for each $a \in A$.

If $a' \in \lim_\alpha A$, then there exists a sequence (a_n) in A with $a_n \rightarrow_\alpha a'$. Thus $h > a_n$ for each $n \in N$, hence $h \geq a'$. Since $h \notin \lim_\alpha A$ we get $h > a'$ for each $a' \in \lim_\alpha A$. Therefore $\lim_\alpha H = \langle \lim_\alpha A \rangle$. \square

2.4. Corollary. *If H is as in 2.2 and if A is regular with respect to (G, α) , then H is regular with respect to (G, α) .*

2.5. Corollary. *Let H be an ℓ -group, $H = \langle A \rangle$, $A \neq \{0\}$ and suppose that A is absolutely regular. Then H is absolutely regular.*

2.6. Proposition. *Let A be an ℓ -group which can be represented as a direct product of a finite number of linearly ordered groups. Suppose that $A \neq \{0\}$ and $H = \langle A \rangle$. Then H is absolutely regular.*

P r o o f. This is a consequence of 2.6 and of Theorem 3.6, [3]. \square

2.7. Lemma. *Let H be an ℓ -subgroup of G such that*

- (i) *H can be represented as a direct product $H_1 \times H_2 \times \dots \times H_k$;*
- (ii) *there are ℓ -subgroups A_i of H_i such that $H_i = \langle A_i \rangle$, $H_i \neq A_i \neq \{0\}$ ($i = 1, 2, \dots, k$).*

Then $\lim_\alpha H = \lim_\alpha H_1 \times \dots \times \lim_\alpha H_k$.

P r o o f. Let $i \in \{1, 2, \dots, k\}$. In view of 2.3,

$$\lim_\alpha H_i = \langle \lim_\alpha A_i \rangle.$$

Now we proceed by induction with respect to k . For $k = 1$ the assertion is trivial. Let $k > 1$. Consider an element $g \in \lim_\alpha H$ with $g > 0$. Then there exists a sequence (z_n) in H such that $z_n \rightarrow_\alpha g$ and $z_n > 0$ for each $n \in N$.

a) First we prove that g cannot be an upper bound of the set H . In fact, if $g \geq h$ for each $h \in H$, then $g \geq z_n$ for each $n \in N$, whence $g = \bigvee_{n \in N} z_n$ and thus $g = \sup H$. There exists $h_0 \in H$ with $h_0 > 0$. Then $h + h_0 \in H$ for each $h \in H$, yielding that $h + h_0 \leq g$. Hence $h \leq g - h_0 < g$ for each $h \in H$, which is a contradiction.

b) For $h \in H$ and $i \in I$ we denote by $h(H_i)$ the component of h in H_i . If $h \geq 0$, then

$$h = h(H_1) + h(H_2) + \dots + h(H_n) = h(H_1) \vee h(H_2) \vee \dots \vee h(H_n).$$

Thus in view of a) there exists $i_0 \in \{1, 2, \dots, k\}$ such that g fails to be an upper bound of the set H_{i_0} . Without loss of generality we can suppose that $i_0 = k$. Therefore there exists $x_0 \in H_k^+$ such that $x_0 \not\leq g$.

We have

$$z_n \wedge x_0 = (z_n(H_1) \vee z_n(H_2) \vee \dots \vee z_n(H_k)) \wedge x_0 = z_n(H_k) \wedge x_0 \in H_k$$

(since $z_n(H_i) \wedge x_0 = 0$ for $i = 1, 2, \dots, k-1$). Then

$$z_n(H_k) \wedge x_0 \rightarrow g \wedge x_0,$$

whence $g \wedge x_0 \in \lim_{\alpha} H_k \subseteq \lim_{\alpha} H$.

For each $h^k \in H_k$ we denote $\overline{h^k} = h^k + A_k$. Further we put

$$\overline{H}_k = \{\overline{h^k} : h^k \in H_k\}.$$

If $\overline{h_1^k}$ and $\overline{h_2^k}$ are distinct elements of \overline{H}_k and $h_1^k < h_2^k$, then we put $\overline{h_1^k} < \overline{h_2^k}$. In this way \overline{H}_k turns out to be a linearly ordered set.

Consider the sequence $(z_n(\overline{H}_k))$. If there existed a subsequence (\overline{y}_n) of $(z_n(\overline{H}_k))$ such that $\overline{y}_n > \overline{x}_0$ for each $n \in N$, then we would have $g \geq x_0$, which is a contradiction. Hence there is a subsequence (\overline{y}_n) of $(z_n(\overline{H}_k))$ such that $\overline{y}_n \leq \overline{x}_0$ for each $n \in N$.

Since $H_k \neq A_k$ there exists $x'_0 \in H_k$ such that $\overline{x}_0 < \overline{x'_0}$. We can replace \overline{x}_0 by $\overline{x'_0}$ and then the previous considerations remain valid. Moreover, $\overline{y}_n < \overline{x'_n}$ for each $n \in N$. We have $y_n = z_n^1(H_k)$, where (z_n^1) is a subsequence of (z_n) . Thus

$$z_n^1(H_k) < x'_0 \quad \text{for each } n \in N,$$

and $z_n^1(H_k) \wedge x'_0 \rightarrow_{\alpha} g \wedge x'_0$. Hence $z_n^1(H_k) \rightarrow_{\alpha} g \wedge x'_0$. This yields that

$$z'_n - z'_n(H_k) \rightarrow_{\alpha} g - (g \wedge x'_0).$$

Since

$$z'_n - z'_n(H_k) = z'_n(H_1) + z'_n(H_2) + \dots + z'_n(H_{k-1}) \in H_1 \times \dots \times H_{k-1},$$

in view of the induction hypothesis we obtain

$$g - (g \wedge x'_0) \in \lim_{\alpha} H_1 \times \lim_{\alpha} H_2 \times \dots \times \lim_{\alpha} H_{k-1}.$$

Denote

$$\lim_{\alpha} H_1 \times \lim_{\alpha} H_2 \times \dots \times \lim_{\alpha} H_{k-1} = Y_{k-1}.$$

It is easy to verify that if $y_{k-1} \in (Y_{k-1})^+$ and $y_k \in (\lim_{\alpha} H_k)^+$, then

$$y_{k-1} \wedge y_k = 0.$$

Further, we obviously have

$$0 \in (Y_{k-1})^+ \cap (\lim_{\alpha} H_k)^+.$$

Let Y be the sublattice of the lattice G^+ generated by the set

$$(Y_{k-1})^+ \cup (\lim_{\alpha} H_k)^+.$$

Since the lattice G^+ is distributive, we obtain

$$Y = \{y_{k-1} \vee y_k : y_{k-1} \in (Y_{k-1})^+ \text{ and } y_k \in (\lim_{\alpha} H_k)^+\}.$$

Thus in view of Lemma 3.4 in [5] we get

$$(1) \quad Y = (Y_{k-1})^+ \times Y_k^+,$$

where Y_k^+ is the underlying lattice of the lattice ordered semigroup $(\lim_{\alpha} H_k)^+$.

For $A, B \subseteq G$ we put

$$A - B = \{a - b : a \in A \text{ and } b \in B\}.$$

Clearly

$$\lim_{\alpha} H_k = Y_k^+ - Y_k^+.$$

Therefore according to (1) and by applying Theorem 2.9 in [3] we obtain

$$\begin{aligned} \lim_{\alpha} H &= Y - Y = ((Y_{k-1})^+ - (Y_{k-1})^+) \times (Y_k^+ - Y_k^+) = Y_{k-1} \times \lim_{\alpha} H_k \\ &= \lim_{\alpha} H_1 \times \lim_{\alpha} H_2 \times \dots \times \lim_{\alpha} H_{k-1} \times \lim_{\alpha} H_k. \end{aligned}$$

□

2.8. Lemma. *Let H and H_1, H_2, \dots, H_k be as in 2.7. Further suppose that all A_i ($i = 1, 2, \dots, k$) are regular with respect to (G, α) . Then $\lim_{\alpha} H$ can be represented in the form*

$$\lim_{\alpha} H = \langle \lim_{\alpha} A_1 \rangle \times \langle \lim_{\alpha} A_2 \rangle \times \dots \times \langle \lim_{\alpha} A_k \rangle$$

and all $\lim_{\alpha} A_i$ ($i = 1, 2, \dots, k$) are regular with respect to (G, α) .

P r o o f. The first assertion is a consequence of 2.7 and 2.3; the latter is obvious. □

2.9. Lemma. *Let H and H_1, H_2, \dots, H_k be as in 2.8. Then H is regular with respect to (G, α) .*

P r o o f. In view of 2.3, 2.7 and 2.8 we have

$$\begin{aligned} \lim_{\alpha} \lim_{\alpha} H &= \lim_{\alpha} \langle \lim_{\alpha} A_1 \rangle \times \dots \times \lim_{\alpha} \langle \lim_{\alpha} A_k \rangle \\ &= \langle \lim_{\alpha} \lim_{\alpha} A_1 \rangle \times \dots \times \langle \lim_{\alpha} \lim_{\alpha} A_k \rangle \\ &= \langle \lim_{\alpha} A_1 \rangle \times \dots \times \langle \lim_{\alpha} A_k \rangle = \lim_{\alpha} H. \end{aligned}$$

□

2.10. Corollary. *Let H and H_i ($i = 1, 2, \dots, k$) be ℓ -groups such that the conditions (i) and (ii) from 2.7 are valid. Further suppose that all A_i ($i = 1, 2, \dots, k$) are absolutely regular. Then H is absolutely regular.*

3. ON ℓ -GROUPS BELONGING TO F

In this section we assume that H is an ℓ -group belonging to the class F and that $H \neq \{0\}$.

It follows from the results of [1] concerning the structure of ℓ -groups belonging to the class F that there exist a positive integer n and finite systems F_1, F_2, \dots, F_n of convex nonzero subgroups of H such that

- (i) $F_1 = \{A_1^1, A_2^1, \dots, A_{n(1)}^1\}$, all ℓ -groups A_i^1 ($i = 1, \dots, n(1)$) are linearly ordered and $A_{i(1)}^1 \cap A_{i(2)}^1 = \{0\}$ whenever $i(1), i(2)$ are distinct elements of the set $\{1, 2, \dots, n(1)\}$.
- (ii) If $k > 1$, then $F_k = \{A_1^k, A_2^k, \dots, A_{n(k)}^k\}$ such that
 - (ii1) $A_{i(1)}^k \cap A_{i(2)}^k = \{0\}$ whenever $i(1), i(2)$ are distinct elements of the set $\{1, 2, \dots, n(k)\}$;

- (ii) if $i \in \{1, 2, \dots, n(k)\}$, then either A_i^k is equal to an element of F_{k-1} , or there are $B_1, B_2, \dots, B_{t(i)} \in F_{k-1}$ such that $t(i) \geq 2$ and $A_i^k = \langle B_1 \times B_2 \times \dots \times B_{t(i)} \rangle$.
- (iii) $F_n = \{H\}$.

3.1. Lemma. *Let us apply the above notation and let $k \in \{1, 2, \dots, n\}$. Then all ℓ -groups of the system F_k are absolutely regular.*

P r o o f. We proceed by induction with respect to k . For $k = 1$, this is a consequence of Theorem 3.6 in [5]. Suppose that $k > 1$ and that the assertion is valid for $k - 1$. Then 2.10 yields that the elements of F_k are absolutely regular. \square

As a corollary we obtain

3.2. Theorem. *Each ℓ -group belonging to F is absolutely regular.*

If an ℓ -group H is a direct product of a finite number of linearly ordered groups, then H belongs to F . Hence 3.2 generalizes Theorem 3.6 from [5].

References

- [1] *P. Conrad*: The structure of a lattice ordered group with a finite number of disjoint elements. Michigan Math. J. 7 (1960), 171–180.
- [2] *P. Conrad*: Lattice Ordered Groups. Lecture Notes, Tulane University, 1970.
- [3] *J. Jakubík*: Direct decompositions of partially ordered groups, II. Czechoslovak Math. J. 11 (1961), 490–515. (In Russian.)
- [4] *J. Jakubík*: Sequential convergences in ℓ -groups without Urysohn’s axiom. Czechoslovak Math. J. 42 (1992), 101–116.
- [5] *J. Jakubík*: Closed convex ℓ -subgroups and radical classes of convergence ℓ -groups. Math. Bohem. 122 (1997), 301–315.
- [6] *V. M. Kopytov, N. Ya. Medvedev*: The Theory of Lattice Ordered Groups. Kluwer Academic Publishers, Dordrecht-Boston-London, 1994.

Author’s address: *Ján Jakubík*, Matematický ústav SAV, Grešákova 6, 040 01 Košice, Slovakia.