

Jan Kühr

Pseudo BL -algebras and $DR\ell$ -monoids

Mathematica Bohemica, Vol. 128 (2003), No. 2, 199–208

Persistent URL: <http://dml.cz/dmlcz/134040>

Terms of use:

© Institute of Mathematics AS CR, 2003

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

PSEUDO BL -ALGEBRAS AND $DR\ell$ -MONOIDS

JAN KÜHR, Olomouc

(Received March 1, 2002)

Abstract. It is shown that pseudo BL -algebras are categorically equivalent to certain bounded $DR\ell$ -monoids. Using this result, we obtain some properties of pseudo BL -algebras, in particular, we can characterize congruence kernels by means of normal filters. Further, we deal with representable pseudo BL -algebras and, in conclusion, we prove that they form a variety.

Keywords: pseudo BL -algebra, $DR\ell$ -monoid, filter, polar, representable pseudo BL -algebra

MSC 2000: 06F05, 03G25

1. CONNECTIONS BETWEEN PSEUDO BL -ALGEBRAS AND $DR\ell$ -MONOIDS

Recently, pseudo BL -algebras were introduced by A. Di Nola, G. Georgescu and A. Iorgulescu in [3] as a noncommutative extension of Hájek's BL -algebras (see [6]).

An algebra $\mathfrak{A} = (A, \vee, \wedge, \odot, \rightarrow, \rightsquigarrow, 0, 1)$ of type $\langle 2, 2, 2, 2, 0, 0 \rangle$ is called a *pseudo BL -algebra* iff $(A, \vee, \wedge, 0, 1)$ is a bounded lattice, $(A, \odot, 1)$ is a monoid and the following conditions are satisfied for all $x, y, z \in A$:

- (1) $x \odot y \leq z$ iff $x \leq y \rightarrow z$ iff $y \leq x \rightsquigarrow z$,
- (2) $x \wedge y = (x \rightarrow y) \odot x = x \odot (x \rightsquigarrow y)$,
- (3) $(x \rightarrow y) \vee (y \rightarrow x) = (x \rightsquigarrow y) \vee (y \rightsquigarrow x) = 1$.

By [3, Corollary 3.29], pseudo BL -algebras satisfying the identity

$$(x \rightsquigarrow 0) \rightarrow 0 = (x \rightarrow 0) \rightsquigarrow 0 = x$$

are the duals of pseudo MV -algebras.

In the same way, (noncommutative) $DR\ell$ -monoids extend Swamy's $DR\ell$ -semi-groups which were introduced in [12] as a common generalization of abelian ℓ -groups and Brouwerian algebras.

An algebra $\mathfrak{A} = (A, +, 0, \vee, \wedge, \rightarrow, \leftarrow)$ of type $\langle 2, 0, 2, 2, 2, 2 \rangle$ is a *dually residuated lattice ordered monoid*, or simply a *DR ℓ -monoid*, iff

- (1) $(A, +, 0, \vee, \wedge)$ is an ℓ -monoid, that is, $(A, +, 0)$ is a monoid, (A, \vee, \wedge) is a lattice and, for any $x, y, s, t \in A$, the following distributive laws are satisfied:

$$\begin{aligned} s + (x \vee y) + t &= (s + x + t) \vee (s + y + t), \\ s + (x \wedge y) + t &= (s + x + t) \wedge (s + y + t); \end{aligned}$$

- (2) for any $x, y \in A$, $x \rightarrow y$ is the least $s \in A$ such that $s + y \geq x$, and $x \leftarrow y$ is the least $t \in A$ such that $y + t \geq x$;
- (3) \mathfrak{A} fulfils the identities

$$\begin{aligned} ((x \rightarrow y) \vee 0) + y &\leq x \vee y, \quad y + ((x \leftarrow y) \vee 0) \leq x \vee y, \\ x \rightarrow x &\geq 0, \quad x \leftarrow x \geq 0. \end{aligned}$$

Note that the inequalities $x \rightarrow x \geq 0$ and $x \leftarrow x \geq 0$ can be omitted, and the condition (2) is equivalent to the system of identities (see [10])

$$\begin{aligned} (x \rightarrow y) + y &\geq x, \quad y + (x \leftarrow y) \geq x, \\ x \rightarrow y &\leq (x \vee z) \rightarrow y, \quad x \leftarrow y \leq (x \vee z) \leftarrow y, \\ (x + y) \rightarrow y &\leq x, \quad (y + x) \leftarrow y \leq x. \end{aligned}$$

In [11], mutual relationships between *BL*-algebras and bounded representable commutative *DR ℓ -monoids* are described.

Theorem 1.1. *Let $\mathfrak{A} = (A, \vee, \wedge, \odot, \rightarrow, \rightsquigarrow, 0, 1)$ be a pseudo *BL*-algebra. If we set*

$$\begin{aligned} x + y &:= x \odot y, \quad x \vee_d y := x \wedge y, \quad x \wedge_d y := x \vee y, \\ x \rightarrow y &:= y \rightarrow x, \quad x \leftarrow y := y \rightsquigarrow x, \quad 0_d := 1, \quad 1_d := 0 \end{aligned}$$

for any $x, y \in A$, then $\mathfrak{A}_d = (A, +, 0_d, \vee_d, \wedge_d, \rightarrow, \leftarrow)$ is a bounded *DR ℓ -monoid* with the greatest element 1_d . In addition, this *DR ℓ -monoid* satisfies the identities

$$(*) \quad \begin{aligned} (x \rightarrow y) \wedge_d (y \rightarrow x) &= 0_d, \\ (x \leftarrow y) \wedge_d (y \leftarrow x) &= 0_d. \end{aligned}$$

Proof. Since $(A, \odot, 1, \vee, \wedge)$ is an ℓ -monoid, by [3, Propositions 3.3, 3.9], so is $(A, +, 0_d, \vee_d, \wedge_d)$. The rest follows directly by the definitions. Note that if a *DR ℓ -monoid* \mathfrak{A}_d contains the greatest element 1_d then 0_d is its least element, by [8, Theorem 1.2.3]. \square

In view of Theorem 1.1, it is easily seen that in the definition of a pseudo BL -algebra, the condition (1) can be equivalently replaced by the following identities:

$$\begin{aligned}(x \rightarrow y) \odot x &\leq y, \quad x \odot (x \rightsquigarrow y) \leq y, \\ x \rightarrow y &\geq x \rightarrow (y \wedge z), \quad x \rightsquigarrow y \geq x \rightsquigarrow (y \wedge z), \\ y \rightarrow (x \odot y) &\geq x, \quad y \rightsquigarrow (y \odot x) \geq x.\end{aligned}$$

Consequently, pseudo BL -algebras form a variety of algebras of type $\langle 2, 2, 2, 2, 2, 0, 0 \rangle$. This variety is arithmetical; in accordance with [8, Theorem 3.1.1], the Pixley term of the variety of pseudo BL -algebras can be taken as follows:

$$p(x, y, z) = ((x \rightsquigarrow y) \rightarrow z) \wedge ((z \rightsquigarrow y) \rightarrow x) \wedge (x \vee z).$$

Theorem 1.2. *Let $\mathfrak{A} = (A, +, 0, \vee, \wedge, \rightarrow, \leftarrow)$ be a $DR\ell$ -monoid with the greatest element 1. For any $x, y \in A$ set*

$$\begin{aligned}x \odot y &:= x + y, \quad x \vee_d y := x \wedge y, \quad x \wedge_d y := x \vee y, \\ x \rightarrow y &:= y \leftarrow x, \quad x \rightsquigarrow y := y \leftarrow x, \quad 0_d := 1, \quad 1_d := 0.\end{aligned}$$

Then $\mathfrak{A}_d = (A, \vee_d, \wedge_d, \odot, \rightarrow, \rightsquigarrow, 0_d, 1_d)$ is a pseudo BL -algebra if and only if \mathfrak{A} satisfies ().*

Proof. In any $DR\ell$ -monoid we have

$$x \vee y = ((y \leftarrow x) \vee 0) + x = x + ((y \leftarrow x) \vee 0).$$

Since \mathfrak{A} is bounded, that is, $0 \leq x \leq 1$ for any $x \in A$, it follows that

$$x \wedge_d y = (x \rightarrow y) \odot x = x \odot (x \rightsquigarrow y).$$

The rest is obvious. □

Let \mathcal{PBL} be the category of pseudo BL -algebras, that is, the category whose objects are pseudo BL -algebras and morphisms are homomorphisms of pseudo BL -algebras. Let $\mathcal{DR}\mathcal{L}_{1(*)}$ be the category of bounded $DR\ell$ -monoids satisfying (*). Its morphisms are homomorphisms of $DR\ell$ -monoids which preserve also 1, thus in the sequel, bounded $DR\ell$ -monoids are regarded as algebras $(A, +, 0, \vee, \wedge, \rightarrow, \leftarrow, 1)$ of type $\langle 2, 0, 2, 2, 2, 2, 0 \rangle$.

Theorem 1.3. *The categories \mathcal{PBL} and $\mathcal{DR}\mathcal{L}_{1(*)}$ are equivalent.*

Proof. Theorems 1.1 and 1.2 enable us to define a functor $\mathcal{F}: \mathcal{PBL} \rightarrow \mathcal{DR}\mathcal{L}_{1(*)}$ as follows: (i) $\mathcal{F}(\mathfrak{A}) = \mathfrak{A}_d$ for any pseudo BL -algebra \mathfrak{A} , and (ii) $\mathcal{F}(h) = h$ for any pseudo BL -homomorphism h . It is easy to see that \mathcal{F} is really a categorical equivalence. □

2. FILTERS

According to [3], a subset F of a pseudo BL -algebra \mathfrak{A} with the following properties is said to be a *filter* of \mathfrak{A} :

- (F1) $1 \in F$;
- (F2) $\forall x, y \in F; x \odot y \in F$;
- (F3) $\forall x \in F \forall y \in A; x \leq y \implies y \in F$.

For any subset $M \subseteq A$, the intersection of all filters containing M is called a *filter generated by M* and denoted by $[M]$. It is clear that

$$[M] = \{x \in A; x \geq a_1 \odot \dots \odot a_n \text{ for some } a_1, \dots, a_n \in M \text{ and } n \geq 1\},$$

and if we write briefly $[a]$ for $[\{a\}]$ then

$$[a] = \{x \in A; x \geq a^n \text{ for some } n \geq 1\}.$$

In Section 1, we have already proved that $DR\ell$ -monoids include the duals of pseudo BL -algebras. It is obvious that $F \subseteq A$ is a filter of a pseudo BL -algebra \mathfrak{A} iff it is an ideal of the induced bounded $DR\ell$ -monoid \mathfrak{A}_d , that is,

- (I1) $0_d \in F$;
- (I2) $\forall x, y \in F; x + y \in F$;
- (I3) $\forall x \in F \forall y \in A; x \geq_d y \implies y \in F$.

Ideals of noncommutative $DR\ell$ -monoids were studied in [9]. Considering the above facts, we immediately obtain the following results.

Proposition 2.1. *The set of all filters of any pseudo BL -algebra \mathfrak{A} , ordered by set inclusion, is an algebraic Brouwerian lattice. For any filters F, G of \mathfrak{A} , the relative pseudocomplement of F with respect to G is given by*

$$F * G = \{a \in A; a \vee x \in G \text{ for all } x \in F\}.$$

Let \mathfrak{A} be a pseudo BL -algebra and $X \subseteq A$. The set

$$X^\perp = \{a \in A; a \vee x = 1 \text{ for any } x \in X\}$$

is called the *polar* of X . For any $x \in A$ we write x^\perp instead of $\{x\}^\perp$.

A subset X of A is a *polar* in \mathfrak{A} iff $X = Y^\perp$ for some $Y \subseteq A$.

Proposition 2.2 [3, Propositions 4.38, 4.39]. *For all subsets X, Y of a pseudo BL -algebra \mathfrak{A} , (i) X^\perp is a filter of \mathfrak{A} , (ii) $X \subseteq X^{\perp\perp}$, (iii) $X \subseteq Y$ implies $Y^\perp \subseteq X^\perp$, (iv) $X^\perp = X^{\perp\perp\perp}$.*

Proposition 2.3. For any subset X of a pseudo BL -algebra \mathfrak{A} , X is a polar in \mathfrak{A} iff $X = X^{\perp\perp}$.

Proof. Let $X = Y^\perp$; then $X^{\perp\perp} = Y^{\perp\perp\perp} = Y^\perp = X$. □

By Proposition 2.1, the pseudocomplement of a filter F is

$$F^* = \{a \in A; a \vee x = 1 \text{ for any } x \in F\}.$$

Moreover, it is clear that $F^\perp = F^*$ whenever F is a filter, and conversely, any polar is the pseudocomplement of some filter; in fact, $X = (X^\perp)^*$. Thus the polars in any pseudo BL -algebra are precisely the pseudocomplements in the lattice of its filters. Therefore, by the Glivenko-Frink Theorem, we directly obtain

Theorem 2.4. The set of all polars in any pseudo BL -algebra, ordered by set inclusion, is a complete Boolean algebra.

A filter F of a pseudo BL -algebra \mathfrak{A} is said to be *normal* iff it satisfies the following condition for each $x, y \in A$:

$$x \rightarrow y \in F \iff x \rightsquigarrow y \in F.$$

Proposition 2.5. For any filter F , the following conditions are equivalent:

- (i) F is normal;
- (ii) $x \odot F = F \odot x$ for each $x \in A$.

Proposition 2.6. If F and G are normal filters of \mathfrak{A} then

$$F \vee G = \{x \in A; x \geq a \odot b \text{ for some } a \in F, b \in G\}.$$

In addition, $F \vee G$ is a normal filter. Consequently, normal filters of any pseudo BL -algebra form a complete sublattice of the lattice of all its filters.

Theorem 2.7. In any pseudo BL -algebra, there is a one-to-one correspondence between the normal filters and the congruence relations. In fact, F corresponds to $\Theta(F)$ defined by

$$\langle x, y \rangle \in \Theta(F) = \Theta_1(F) \iff (x \rightarrow y) \wedge (y \rightarrow x) \in F,$$

or equivalently,

$$\langle x, y \rangle \in \Theta(F) = \Theta_2(F) \iff (x \rightsquigarrow y) \wedge (y \rightsquigarrow x) \in F.$$

As proved in [3], and in general for noncommutative $DR\ell$ -monoids in [9], if F is not a normal filter then the binary relations defined in the previous theorem, $\Theta_1(F)$ and $\Theta_2(F)$, are two distinct congruence relations on the distributive lattice $\mathfrak{L}(\mathfrak{A}) = (A, \vee, \wedge, 0, 1)$. In the quotient lattices $\mathfrak{L}(\mathfrak{A})/\Theta_1(F)$ and $\mathfrak{L}(\mathfrak{A})/\Theta_2(F)$ we have

$$(2.1) \quad [x]\Theta_1(F) \leq [y]\Theta_1(F) \iff x \rightarrow y \in F$$

and

$$(2.2) \quad [x]\Theta_2(F) \leq [y]\Theta_2(F) \iff x \rightsquigarrow y \in F,$$

respectively.

Let \mathfrak{A} be a pseudo BL -algebra. A filter F of \mathfrak{A} is said to be *prime* if it is a finitely meet-irreducible element in the lattice of filters of \mathfrak{A} .

By [3, Theorem 4.28], for any filter F of a pseudo BL -algebra \mathfrak{A} and for each ideal I of the lattice $\mathfrak{L}(\mathfrak{A})$, if $F \cap I = \emptyset$ then there exists a prime filter P of \mathfrak{A} with $F \subseteq P$ and $P \cap I = \emptyset$. Consequently, every proper filter is the intersection of all prime filters including it. In particular, the intersection of all prime filters is equal to $\{1\}$.

Theorem 2.8. *For any filter F of a pseudo BL -algebra \mathfrak{A} , the following conditions are equivalent:*

- (i) F is prime;
- (ii) for all filters G, H of \mathfrak{A} , $G \cap H \subseteq F$ implies $G \subseteq F$ or $H \subseteq F$;
- (iii) for any $x, y \in A$, $x \vee y \in F$ implies $x \in F$ or $y \in F$;
- (iv) for any $x, y \in A$, $x \vee y = 1$ implies $x \in F$ or $y \in F$;
- (v) for any $x, y \in A$, $x \rightarrow y \in F$ or $y \rightarrow x \in F$;
- (vi) for any $x, y \in A$, $x \rightsquigarrow y \in F$ or $y \rightsquigarrow x \in F$;
- (vii) $\mathfrak{L}(\mathfrak{A})/\Theta_1(F)$ is totally ordered;
- (viii) $\mathfrak{L}(\mathfrak{A})/\Theta_2(F)$ is totally ordered;
- (ix) the set of all filters including F is totally ordered under set inclusion.

Remark. The equivalence of (iii), (v), (vi), (vii) and (viii) is due to [3, Proposition 4.25].

Proof. (i) \Rightarrow (ii): Using the distributivity of the lattice of filters, $G \cap H \subseteq F$ implies $F = F \vee (G \cap H) = (F \vee G) \cap (F \vee H)$, whence $F = F \vee G$ or $F = F \vee H$, that is, $F \supseteq G$ or $F \supseteq H$.

(ii) \Rightarrow (iii): Obviously, $x \vee y \in F$ yields $[x] \cap [y] = [x \vee y] \subseteq F$. Hence, by (ii), $[x] \subseteq F$ or $[y] \subseteq F$ and thus $x \in F$ or $y \in F$.

(iii) \Rightarrow (iv): This is evident since $1 \in F$.

(iv) \Rightarrow (v) and (iv) \Rightarrow (vi): By the definition of a pseudo BL -algebra,

$$(x \rightarrow y) \vee (y \rightarrow x) = (x \rightsquigarrow y) \vee (y \rightsquigarrow x) = 1,$$

which implies the assertion by (iv).

(v) \Rightarrow (vii) and (vi) \Rightarrow (viii): This is obvious from (2.1) and (2.2), respectively.

(vii) \Rightarrow (ix): If $F \subseteq G, H$ and neither $G \subseteq H$ nor $H \subseteq G$ then there exist $a, b \in A$ with $a \in G \setminus H$ and $b \in H \setminus G$. For instance, let $a \rightarrow b \in F$. Then $b \geq a \wedge b = (a \rightarrow b) \odot a \in G$, whence $b \in G$; a contradiction. Similarly (viii) \Rightarrow (ix).

(ix) \Rightarrow (i): $F = G \cap H$ entails $F = G$ or $F = H$, because either $G \subseteq H$ or $H \subseteq G$. \square

3. REPRESENTABLE PSEUDO BL -ALGEBRAS

Proposition 3.1. *If P is a minimal prime filter of a pseudo BL -algebra \mathfrak{A} then $A \setminus P$ is a maximal ideal of the lattice $\mathfrak{L}(\mathfrak{A})$.*

Proof. By Zorn's Lemma, there is a maximal ideal I of $\mathfrak{L}(\mathfrak{A})$ with $A \setminus P \subseteq I$. (Since P is also a prime filter of $\mathfrak{L}(\mathfrak{A})$, it follows that $A \setminus P$ is a prime ideal of $\mathfrak{L}(\mathfrak{A})$ which is included in some maximal (prime) ideal.) We will show that $I = A \setminus P$. Denote $Q = \bigcup \{a^\perp; a \in I\}$. We claim that $P = Q$.

If $x \in a^\perp$ for some $a \in I$, then $x \vee a = 1$ and $x \notin I$. Indeed, if $x \in I$ then $x \vee a \neq 1$ since $x \vee a = 1$ would mean $I = A$. Thus $x \in A \setminus I \subseteq A \setminus (A \setminus P) = P$, whence $a^\perp \subseteq A \setminus I \subseteq P$ and consequently, $Q \subseteq A \setminus I \subseteq P$.

We shall now prove that Q is a prime filter of \mathfrak{A} . (F1): Since any principal polar a^\perp contains 1, so does Q . (F2): If $x, y \in Q$, that is, $x \in a^\perp, y \in b^\perp$ for some $a, b \in I$, then $a \vee b \in I$ and

$$(x \odot y) \vee a \vee b \geq (x \vee a \vee b) \odot (y \vee a \vee b) = 1 \odot 1 = 1.$$

Therefore $x \odot y \in (a \vee b)^\perp \subseteq Q$. (F3): It is obvious since a^\perp is a filter of \mathfrak{A} for each $a \in I$.

To prove that Q is prime, suppose $x \vee y = 1$ and $x \notin Q$, that is, $x \vee a \neq 1$ for all $a \in I$. If $x \notin I$ then the ideal in the lattice $\mathfrak{L}(\mathfrak{A})$ generated by $I \cup \{x\}$, $(I \cup \{x\}]$, is proper, i.e., $A \setminus P \subseteq I \subset (I \cup \{x\}] \neq A$, since $(I \cup \{x\}] = A$ would entail $1 \leq x \vee a$ for some $a \in I$; a contradiction. Hence $x \in I$ and thus $y \in x^\perp \subseteq Q$, proving that Q is prime.

However, P is a minimal prime filter of \mathfrak{A} ; thus $Q \subseteq A \setminus I \subseteq P$ yields $Q = A \setminus I = P$ as claimed. Therefore $I = A \setminus P$. \square

Corollary 3.2. *If P is a minimal prime filter then*

$$P = \bigcup \{a^\perp; a \notin P\}.$$

Proof. By the proof of the previous proposition, $P = \bigcup \{a^\perp; a \in I\}$, where $I = A \setminus P$. \square

A pseudo BL -algebra is said to be *representable* if it is a subdirect product of linearly ordered pseudo BL -algebras.

By Theorems 2.7 and 2.8, subdirect representations by totally ordered pseudo BL -algebras are associated with families of normal prime filters whose intersections are precisely $\{1\}$. Therefore it is obvious that every BL -algebra is representable (see also [11]). In contrast, for pseudo BL -algebras, this assertion fails.

The following results generalize the similar properties of pseudo MV -algebras, [4, Theorem 2.20], [1, Theorem 5.9], and [2, Theorem 6.11].

Theorem 3.3. *For any pseudo BL -algebra \mathfrak{A} , the following statements are equivalent.*

- (i) \mathfrak{A} is representable.
- (ii) There exists a family $\{P_i\}_{i \in I}$ of normal prime filters of \mathfrak{A} such that

$$\bigcap_{i \in I} P_i = \{1\}.$$

- (iii) Any polar of \mathfrak{A} is a normal filter of \mathfrak{A} .
- (iv) Any principal polar is a normal filter.
- (v) Any minimal prime filter is normal.

Proof. As argued above, the equivalence of (i) and (ii) is clear.

(i) \Rightarrow (iii): Suppose that \mathfrak{A} is a subdirect product of linearly ordered pseudo BL -algebras $\{\mathfrak{A}_i\}_{i \in I}$. Observe that

$$(3.1) \quad x \vee y = 1 \text{ iff } \{i \in I; x_i \neq 1_i\} \cap \{i \in I; y_i \neq 1_i\} = \emptyset$$

for all $x, y \in A$, since \mathfrak{A}_i are totally ordered.

Let now P be a polar in \mathfrak{A} , i.e. $P = P^{\perp\perp}$. Let $x \in A, a \in P$ and $y \in P^\perp$. Then $x \odot a \leq x$ implies $x \odot a = (x \odot a) \wedge x = (x \rightarrow (x \odot a)) \odot x$. Further, $\{i \in I; x_i \rightarrow (x_i \odot a_i) \neq 1_i\} \subseteq \{i \in I; a_i \neq 1_i\}$. Indeed, if $a_i = 1_i$ then $x_i \rightarrow (x_i \odot a_i) = x_i \rightarrow (x_i \odot 1_i) = x_i \rightarrow x_i = 1_i$. Hence we obtain

$$\{i \in I; x_i \rightarrow (x_i \odot a_i) \neq 1_i\} \cap \{i \in I; y_i \neq 1_i\} \subseteq \{i \in I; a_i \neq 1_i\} \cap \{i \in I; y_i \neq 1_i\} = \emptyset$$

by (3.1), since $a \in P$ and $y \in P^\perp$. Therefore $(x \rightarrow (x \odot a)) \vee y = 1$, and thus $x \rightarrow (x \odot a) \in P^{\perp\perp} = P$. Hence $x \odot a = (x \rightarrow (x \odot a)) \odot x \in P \odot x$, proving $x \odot P \subseteq P \odot x$.

(iii) \Rightarrow (iv): Obvious.

(iv) \Rightarrow (v): By Corollary 3.2, $P = \bigcup\{a^\perp; a \notin P\}$ for any minimal prime filter P . If $x \rightarrow y \in P$ then there is $a \notin P$ with $x \rightarrow y \in a^\perp$ which is a normal filter, and hence $x \rightsquigarrow y \in a^\perp \subseteq P$. Summarizing, $x \rightarrow y \in P$ iff $x \rightsquigarrow y \in P$.

(v) \Rightarrow (i): Since any prime filter contains a minimal prime filter and the intersection of all prime filters of \mathfrak{A} is obviously $\{1\}$, so does the intersection of minimal prime filters. Thus, by (ii), \mathfrak{A} is representable. \square

Theorem 3.4. *A pseudo BL-algebra is representable if and only if it satisfies the identities*

$$(3.2) \quad (y \rightarrow x) \vee (z \rightsquigarrow ((x \rightarrow y) \odot z)) = 1,$$

$$(3.3) \quad (y \rightsquigarrow x) \vee (z \rightarrow (z \odot (x \rightsquigarrow y))) = 1.$$

Consequently, the class of representable pseudo BL-algebras is a variety.

Proof. In any linearly ordered pseudo BL-algebra \mathfrak{A} , either $y \rightarrow x = 1$ or $x \rightarrow y = 1$ (and also $y \rightsquigarrow x = 1$ or $x \rightsquigarrow y = 1$), and so it is easy to verify that \mathfrak{A} fulfils (3.2) and (3.3). Therefore the part “only if” is obvious.

Conversely, suppose that (3.2) and (3.3) are satisfied by \mathfrak{A} . In view of Theorem 3.3 (iv), it suffices to prove that any principal polar x^\perp is a normal filter of \mathfrak{A} .

Let $y \in x^\perp$, that is, $y \vee x = 1$. Observe that in this case

$$x = 1 \rightarrow x = (y \vee x) \rightarrow x = (y \rightarrow x) \wedge (x \rightarrow x) = (y \rightarrow x) \wedge 1 = y \rightarrow x$$

and similarly $y = x \rightarrow y$. Hence, by (3.2),

$$x \vee (z \rightsquigarrow (y \odot z)) = (y \rightarrow x) \vee (z \rightsquigarrow ((x \rightarrow y) \odot z)) = 1,$$

thus $z \rightsquigarrow (y \odot z) \in x^\perp$. Further, $y \odot z \leq z$ implies $y \odot z = (y \odot z) \wedge z = z \odot (z \rightsquigarrow (y \odot z)) \in z \odot x^\perp$, which shows $x^\perp \odot z \subseteq z \odot x^\perp$. The other inclusion follows similarly by (3.3). \square

References

- [1] *A. Dvurečenskij*: On pseudo MV -algebras. *Soft Computing* 5 (2001), 347–354.
- [2] *A. Dvurečenskij*: States on pseudo MV -algebras. *Studia Logica* 68 (2001), 301–327.
- [3] *A. Di Nola, G. Georgescu, A. Iorgulescu*: Pseudo BL -algebras: Part I. Preprint.
- [4] *G. Georgescu, A. Iorgulescu*: Pseudo MV -algebras. *Mult. Val. Logic* 6 (2001), 95–135.
- [5] *G. Grätzer*: General Lattice Theory. Birkhäuser, Berlin, 1998.
- [6] *P. Hájek*: Basic fuzzy logic and BL -algebras. *Soft Computing* 2 (1998), 124–128.
- [7] *P. Hájek*: Metamathematics of Fuzzy Logic. Kluwer, Amsterdam, 1998.
- [8] *T. Kovář*: A general theory of dually residuated lattice ordered monoids. Ph.D. thesis, Palacký Univ., Olomouc, 1996.
- [9] *J. Kühr*: Ideals of noncommutative $DR\ell$ -monoids. Manuscript.
- [10] *J. Rachůnek*: A non-commutative generalization of MV -algebras. *Czechoslovak Math. J.* 52 (2002), 255–273.
- [11] *J. Rachůnek*: A duality between algebras of basic logic and bounded representable $DR\ell$ -monoids. *Math. Bohem.* 126 (2001), 561–569.
- [12] *K. L. N. Swamy*: Dually residuated lattice ordered semigroups. *Math. Ann.* 159 (1965), 105–114.

Author's address: Jan Kühr, Department of Algebra and Geometry, Faculty of Sciences, Palacký University, Tomkova 40, 779 00 Olomouc, Czech Republic, e-mail: kuhr@inf.upol.cz.