

Elango Roja; Mallasamudram Kuppusamy Uma; Ganesan Balasubramanian
Urysohn's lemma, gluing lemma and contraction* mapping theorem for fuzzy
metric spaces

Mathematica Bohemica, Vol. 133 (2008), No. 2, 179--185

Persistent URL: <http://dml.cz/dmlcz/134052>

Terms of use:

© Institute of Mathematics AS CR, 2008

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

URYSOHN'S LEMMA, GLUING LEMMA AND CONTRACTION*
MAPPING THEOREM FOR FUZZY METRIC SPACES

E. ROJA, M. K. UMA, Salem, G. BALASUBRAMANIAN, Chennai

(Received November 11, 2006)

Abstract. In this paper the concept of a fuzzy contraction* mapping on a fuzzy metric space is introduced and it is proved that every fuzzy contraction* mapping on a complete fuzzy metric space has a unique fixed point.

Keywords: fuzzy contraction mapping, fuzzy continuous mapping

MSC 2000: 54A40, 03E72

1. INTRODUCTION

The theory of fuzzy sets was introduced by Zadeh in 1965 [7]. Since then many authors (Zi-ke 1982 [8], Erceg 1979 [1], George and Veeramani 1994 [2], Kaleva and Seikkala 1984 [5]) have introduced the concept of a fuzzy metric space in different ways. In this paper we follow the definition of a metric space given by George and Veeramani [2] since the topology induced by the fuzzy metric according to the definition of George and Veeramani [2] is Hausdorff. Motivated by the concept of a metric space, Urysohn's lemma and gluing lemma are studied. Based on the concept of a fuzzy contraction mapping [6], the fuzzy contraction* mapping theorem is established.

2. PRELIMINARIES

Definition 1 [4]. A binary operation $*$: $[0, 1] \times [0, 1] \rightarrow [0, 1]$ is a continuous t -norm if $*$ satisfies the following conditions:

1. $*$ is associative and commutative,
2. $*$ is continuous,

3. $a * 1 = a$ for all $a \in [0, 1]$,
4. $a * b \leq c * d$ whenever $a \leq c$ and $b \leq d$, ($a, b, c, d \in [0, 1]$).

Definition 2 [2]. The triple $(X, M, *)$ is said to be a fuzzy metric space if X is an arbitrary set, $*$ is a continuous t -norm and M is a fuzzy set on $X^2 \times (0, \infty)$ satisfying the following conditions:

1. $M(x, y, t) > 0$,
2. $M(x, y, t) = 1$ if and only if $x = y$,
3. $M(x, y, t) = M(y, x, t)$,
4. $M(x, y, t) * M(y, z, s) \leq M(x, z, t + s)$, $x, y, z \in X$ and $t, s > 0$,
5. $M(x, y, \cdot): X^2 \times (0, \infty) \rightarrow [0, 1]$ is continuous, $x, y, z \in X$ and $t, s > 0$.

Remark 1 [2]. $M(x, y, t)$ can be thought of as the degree of nearness between x and y with respect to t . We identify $x = y$ with $M(x, y, t) = 1$, for $t > 0$ and $M(x, y, t) = 0$ with $x = \infty$ or $y = \infty$.

Remark 2 [2]. In a fuzzy metric space $(X, M, *)$, whenever $M(x, y, t) > 1 - r$ for x, y in X , $t > 0$, $0 < r < 1$, we can find a t_0 , $0 < t_0 < 1$ such that $M(x, y, t_0) > 1 - r$.

Definition 3 [4]. A sequence $\{x_n\}$ in a fuzzy metric space $(X, M, *)$ is said to be a Cauchy sequence if for each ε , $0 < \varepsilon < 1$ and $t > 0$ there exists $n_0 \in \mathbb{N}$ such that $M(x_n, x_m, t) > 1 - \varepsilon$ for all $n, m \geq n_0$.

Definition 4 [2]. Let $(X, M, *)$ be a fuzzy metric space. We define the open ball $B(x, r, t)$ with centre $x \in X$ and radius r , $0 < r < 1$, $t > 0$ as

$$B(x, r, t) = \{y \in X: M(x, y, t) > 1 - r\}.$$

Definition 5 [4]. Let $(X, M, *)$ be a fuzzy metric space. Define $T = \{A \subset X: x \in A \text{ if and only if there exist } r, t > 0, 0 < r < 1 \text{ such that } B(x, r, t) \subset A\}$. Then T is topology on X . This topology is called the topology induced by the fuzzy metric.

Then by Theorem 3.11 of (George and Veeramani 1994 [2]) we know that a sequence $x_n \rightarrow x$ (x_n converges to x) if and only if $M(x_n, x, t) \rightarrow 1$ as $n \rightarrow \infty$.

Definition 6 [2]. A fuzzy metric space is said to be complete if every Cauchy sequence is convergent.

Notation. $M_A(x, y, t)$ denotes the degree of nearness between x and y with respect to t when $x, y \in A$.

3. URYSOHN'S LEMMA AND GLUING LEMMA

Proposition 1 (Urysohn's Lemma). *Let $(X, M, *)$ be a fuzzy metric space. Let T be a topology on X induced by the fuzzy metric. Let A and B be distinct members of r . Then there exists a fuzzy continuous function $f: X \rightarrow [0, 1]$ such that $f = 0$ on A and $f = 1$ on B .*

Proof. Define a function $f: X \rightarrow [0, 1]$ by

$$f(x) = \frac{1 - M_A(x, x, t)}{M_B(x, x, t) - M_A(x, x, t)}.$$

Note that $M_B(x, x, t) - M_A(x, x, t) \neq 0$ for any $x \in X$. If $x \in A$, $M_A(x, x, t) = 1$, then $f(x) = 0$. If $x \in B$, $M_B(x, x, t) = 1$, then $f(x) = 1 - M_A(x, x, t)/1 - M_A(x, x, t) = 1$. Since $M(x, y, t)$ is fuzzy continuous (George and Veeramani 1994 [2]), f is fuzzy continuous. □

Proposition 2 (Gluing Lemma). *Let $(X, M, *)$ and $(Y, M, *)$ be two fuzzy metric spaces. Let $U_i, i \in I$ be members of fuzzy induced topology T on X such that $\bigcup_{i \in I} U_i = X$. Assume that there exists a fuzzy continuous function [3] $f_i: U_i \rightarrow Y$ for each $i \in I$ with the property that $f_i(x) = f_j(x)$ for all $x \in U_i \cap U_j$ and $i, j \in I$. Then the function $f: X \rightarrow Y$ defined by $f(x) = f_i(x)$ if $x \in U_i$ is well defined and fuzzy continuous on X .*

Proof. Let $x, y \in X$. Since f_i is continuous for given $r \in (0, 1), t > 0$ we can find $r_0 \in (0, 1), t/4 > 0$ such that $M(x, y, t_0) > 1 - r_0$ implies $M(f_i(x), f_i(y), t/2) > 1 - r$. Now $M(x, y, t/4) > 1 - r_0$. Let $x \in U_i, y \in U_j$ for some $i \neq j$. Let $x_i \in U_i \cap U_j$. Then

$$\begin{aligned} M(f(x), f(y), t/2) &> M(f(x), f(x_i), t/4) * M(f(x_i), f(y), t/4) \\ &= M(f_i(x), f_i(x_i), t/4) * M(f_j(x_i), f_j(y), t/4) \\ &> (1 - r) * (1 - r) = 1 - r. \end{aligned}$$

Therefore f is fuzzy continuous. □

4. FUZZY CONTRACTION * MAPPING

Definition 7. Let $(X, M, *)$ be a fuzzy metric space. A function $f: X \rightarrow X$ is called a fuzzy contraction* mapping if $M(x, y, t) \geq 1 - (1 - r^2)$ for all $0 < 1 - r^2 < 1$. Then $M(f(x), f(y), t) \geq 1 - (1 - r_0^2)$ for each $x, y \in X$ for some $1 - r_0^2 < 1 - r^2, 1$.

Example 1. Consider the fuzzy metric space $(\mathbb{R}, M, *)$, where \mathbb{R} is the set of all real numbers and $M(x, y, t) = t/(t + |x - y|)$. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ and define $f(x) = x/2$. Then $M(x, y, t) = t/(t + |x - y|) \geq 1 - (1 - r^2)$, $t > 0$, $0 < 1 - r^2 < 1$ where $1 - r^2 \geq |x - y|/(t + |x - y|)$. Then

$$\begin{aligned} M(f(x), f(y), t) &= \frac{t}{t + |(x/2) - (y/2)|} \\ &= \frac{1 - (|(x/2) - (y/2)|)}{t + |(x/2) - (y/2)|} \geq 1 - (1 - r_0^2) \end{aligned}$$

where

$$1 - r_0^2 \geq \frac{|(x/2) - (y/2)|}{t + |(x/2) - (y/2)|}.$$

Further,

$$\begin{aligned} (1 - r^2) - (1 - r_0^2) &\geq \frac{|x - y|}{t + |x - y|} - \frac{(|(x/2) - (y/2)|)}{t + |(x/2) - (y/2)|} \\ &\geq \frac{|x - y|}{t + |x - y|} - \frac{\frac{1}{2}|x - y|}{t + \frac{1}{2}|x - y|} \\ &\geq \frac{|x - y|(t + \frac{1}{2}|x - y|) - \frac{1}{2}|x - y|(t + |x - y|)}{(t + |x - y|)(t + \frac{1}{2}|x - y|)} \\ &\geq \frac{|x - y|t - \frac{1}{2}(|x - y|t)}{(t + |x - y|)(t + \frac{1}{2}|x - y|)} \\ &\geq \frac{(t/2)|x - y|}{(t + |x - y|)(t + \frac{1}{2}|x - y|)} = 0, \end{aligned}$$

which implies that f is a fuzzy contraction* by Definition 7.

Definition 8. A mapping from a fuzzy metric space X to a fuzzy metric space Y is said to be fuzzy continuous* if for given $1 - r^2$, $t > 0$, $0 < 1 - r^2 < 1$ we can find $1 - r_0^2 \in (0, 1)$, $t_0 > 0$ such that $M(x, y, t_0) > 1 - (1 - r_0^2)$ implies $M(f(x), f(y), t/2) > 1 - (1 - r^2)$.

Proposition 3. Every fuzzy contraction* mapping on a fuzzy metric space is fuzzy continuous*.

Proof. Let $f: X \rightarrow X$ be a fuzzy contraction* mapping. Therefore for $x, y \in X$, given $1 - r^2 \in (0, 1)$, $t > 0$, we can find $1 - r_0^2 \in (0, 1)$, $t/4 > 0$ such that $1 - r^2 = (1 - (1 - r_0^2)) * (1 - (1 - r_0^2))$. Now $M(x, y, t/4) > 1 - (1 - r_0^2)$ implies $M(f(x), f(y), t/4) > 1 - (1 - s^2) > 1 - (1 - r_0^2)$ where $1 - s^2 \in (0, (1 - r_0^2))$ (since f is a fuzzy contraction* mapping). Let $x_1 \in X$. Then

$$\begin{aligned} M(f(x), f(y), t/2) &> M(f(x), f(x_1), t/4) * M(f(x_1), f(y), t/4) \\ &> (1 - (1 - r_0^2)) * (1 - (1 - r_0^2)) > (1 - (1 - r^2), (1 - r^2) \in (0, 1) \end{aligned}$$

which implies that f is a fuzzy continuous* mapping. □

Remark 3. The converse need not be true as the following example shows.

Example 2. Consider the fuzzy metric space $(\mathbb{R}, M, *)$ [2] where \mathbb{R} is the set of all real numbers and

$$M(x, y, t) = \frac{t}{t + |x - y|}.$$

Let $f: \mathbb{R} \rightarrow \mathbb{R}$ and define $f(x) = x^2$. Then

$$M(x, y, t) = \frac{t}{t + |x - y|} \geq 1 - (1 - r^2)$$

where $(1 - r^2) \geq |x - y|/(t + |x - y|)$. Then

$$\begin{aligned} M(f(x), f(y), t/2) &= \frac{(t/2)}{(t/2) + |x^2 - y^2|} = \frac{t}{t + 2(|x^2 - y^2|)} \\ &\geq 1 - (1 - r_0^2) \text{ where } 1 - r_0^2 \geq \frac{2|x^2 - y^2|}{t + 2(|x^2 - y^2|)} \end{aligned}$$

which implies that f is a fuzzy continuous* mapping. However, $M(f(x), f(y), t) = t/(t + |x^2 - y^2|) \geq 1 - (1 - s^2)$ where $1 - s^2 \geq |x^2 - y^2|/(t + |x^2 - y^2|)$ since

$$\begin{aligned} (1 - s^2) - (1 - r^2) &\geq \frac{|x^2 - y^2|}{t + |x^2 - y^2|} - \frac{|x - y|}{t + |x - y|} \\ &\geq \frac{(t + |x - y|)|x^2 - y^2| - (t + |x^2 - y^2|)|x - y|}{(t + |x^2 - y^2|)(t + |x - y|)} \\ &\geq \frac{t(|x^2 - y^2| - |x - y|)}{(t + |x^2 - y^2|)(t + |x - y|)} \begin{cases} \geq 0 & \text{if } x, y \text{ are integers} \\ \leq 0 & \text{if } x, y \text{ are not integers} \end{cases} \end{aligned}$$

and consequently, f is not a fuzzy contraction* mapping.

Proposition 4. *Every fuzzy contraction* mapping on a complete fuzzy metric space [2] has a unique fixed point.*

Proof. Let f be a fuzzy contraction* mapping on a complete fuzzy metric space $(X, M, *)$. □

Uniqueness part. If possible let $x_0 \neq y_0$ be two fixed points of f . Then we have

$$\begin{aligned} x_0 &= f^1(x_0) = f^2(x_0) = f^3(x_0) = \dots = f^n(x_0), \\ y_0 &= f^1(y_0) = f^2(y_0) = f^3(y_0) = \dots = f^n(y_0) \quad \text{for each } n \in \mathbb{N}. \end{aligned}$$

Now

$$\begin{aligned} M(x_0, y_0, t) &= M(f^n(x_0), f^n(y_0), t) \geq 1 - (1 - r^2)/k^n \\ &> M(x_0, y_0, t) \quad (= 1 - (1 - r^2)) \end{aligned}$$

where $k > 1$, a contradiction, hence $x_0 = y_0$. Therefore the fixed points are unique.

Existence part. Let $x_1 = f(x_0)$, $x_2 = f(x_1)$, \dots , $x_n = f(x_{n-1}) = f^{n-1}(x_1)$. Then

$$\begin{aligned} M(x_n, x_{n+1}, t) &= M(f^{n-1}(x_1), f^{n-1}(x_2), t) \geq 1 - (1 - r^2)/k^{n-1} \\ &\geq 1 - \frac{1}{1 - s^2} \quad \text{for some } \frac{1}{1 - s^2} \in (0, 1). \end{aligned}$$

Therefore,

$$(A) \quad M(x_n, x_{n+1}, t) \geq 1 - \frac{1}{1 - s^2}.$$

For a given $t' = (m - n)t > 0$, $\varepsilon > 0$, choose n_0 such that $1/n_0 < \varepsilon$. Then for $m \geq n \geq n_0$,

$$\begin{aligned} M(x_n, x_m, t') &\geq M(x_n, x_{n+1}, t) * M(x_{n+1}, x_{n+2}, t) * \dots * M(x_{m-1}, x_m, t) \\ &\geq (1 - (1 - s^2)^{-1}) * (1 - (1 - s^2)^{-1}) * \dots * (1 - (1 - s^2)^{-1}) \\ &\geq 1 - \frac{1}{n} \quad \text{for some } \frac{1}{n} \in (0, 1) \geq 1 - \varepsilon \end{aligned}$$

and hence $\{x_n\}$ is a Cauchy sequence. Since X is a complete metric space, this sequence converges to, say, $z_0 \in X$. Now we assert that z_0 is a fixed point of f . Consider $n \in \mathbb{N}$ for $0 < 1 - r^2 < 1$, $t > 0$. Then we have

$$\begin{aligned} M(f(z_0), z_0, t) &\geq M(f(z_0), f(x_0), t/n + 1) * M(f(x_0), f^2(x_0), t/n + 1) * \dots \\ &\quad * M(f^n(x_0), z_0, t/n + 1), \end{aligned}$$

and since f is a fuzzy contraction* mapping, this is for $k > 1$ and $1/(1 - s_n^2) \in (0, 1)$ greater than or equal to

$$\begin{aligned} & (1 - (1 - s_n^2)) * (1 - (1 - r^2)) * (1 - (1 - r^2)/k) * \dots \\ & * (1 - (1 - r^2)/k^{n-1}) * M(f^n(x_0), z_0, t/n + 1) \\ & \geq (1 - (1 - r^2)/k^{n+p}) * M(f^n(x_0), z_0, t/n + 1) \end{aligned}$$

for some $p \in \mathbb{N}$. Taking limit on both sides as $n \rightarrow \infty$ we obtain

$$\begin{aligned} \lim_{n \rightarrow \infty} M(f(z_0), z_0, t) & \geq \lim_{n \rightarrow \infty} (1 - (1 - r^2)/k^{n+p}) * \lim_{n \rightarrow \infty} M(f^n(x_0), z_0, t/n + 1) \\ & \Rightarrow M(f(z_0), z_0, t) \geq 1 * 1 \Rightarrow f(z_0) = z_0. \end{aligned}$$

Acknowledgement. The authors thank the referee for his/her comments and suggestions.

References

- [1] *M. A. Erceg*: Metric spaces in fuzzy set theory. *J. Math. Anal. Appl.* *69* (1979), 205–230. zbl
- [2] *A. George, P. Veeramani*: On some results in fuzzy metric spaces. *Fuzzy Sets Syst.* *64* (1994), 395–399. zbl
- [3] *A. George, P. Veeramani*: Some theorems in fuzzy metric spaces. *J. Fuzzy Math.* *3* (1995), 933–940. zbl
- [4] *A. George, P. Veeramani*: On some results of analysis for fuzzy metric spaces. *Fuzzy Sets Syst.* *90* (1997), 365–368. zbl
- [5] *O. Kaleva, S. Seikkala*: On fuzzy metric spaces. *Fuzzy Sets Syst.* *12* (1984), 215–229. zbl
- [6] *G. Rajendran, G. Balasubramanian*: Fuzzy contraction mapping theorem for fuzzy metric spaces. *Bull. Calcutta Math. Soc.* *94* (2002), 453–458. zbl
- [7] *L. A. Zadeh*: Fuzzy sets. *Inform and Control* *8* (1965), 338–353. zbl
- [8] *Zi-Ke, Deng*: Fuzzy pseudo metric spaces. *J. Math. Anal. Appl.* *86* (1982), 74–95. zbl

Authors' addresses: *E. Roja, M. K. Uma*, Department of Mathematics, Sri Sarada College for Women, Salem-636 016, Tamilnadu, Indie, e-mail: rpbalan@sancharnet.in; *G. Balasubramanian*, Ramanujan Institute for Advanced Study in Mathematics, University of Madras, Chennai-600 005, Tamilnadu, Indie.