

Ladislav Nebeský

The induced paths in a connected graph and a ternary relation determined by them

*Mathematica Bohemica*, Vol. 127 (2002), No. 3, 397–408

Persistent URL: <http://dml.cz/dmlcz/134072>

## Terms of use:

© Institute of Mathematics AS CR, 2002

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

THE INDUCED PATHS IN A CONNECTED GRAPH AND  
A TERNARY RELATION DETERMINED BY THEM

LADISLAV NEBESKÝ, Praha

(Received October 5, 2000)

*Abstract.* By a ternary structure we mean an ordered pair  $(X_0, T_0)$ , where  $X_0$  is a finite nonempty set and  $T_0$  is a ternary relation on  $X_0$ . By the underlying graph of a ternary structure  $(X_0, T_0)$  we mean the (undirected) graph  $G$  with the properties that  $X_0$  is its vertex set and distinct vertices  $u$  and  $v$  of  $G$  are adjacent if and only if

$$\{x \in X_0; T_0(u, x, v)\} \cup \{x \in X_0; T_0(v, x, u)\} = \{u, v\}.$$

A ternary structure  $(X_0, T_0)$  is said to be the B-structure of a connected graph  $G$  if  $X_0$  is the vertex set of  $G$  and the following statement holds for all  $u, x, y \in X_0$ :  $T_0(x, u, y)$  if and only if  $u$  belongs to an induced  $x - y$  path in  $G$ . It is clear that if a ternary structure  $(X_0, T_0)$  is the B-structure of a connected graph  $G$ , then  $G$  is the underlying graph of  $(X_0, T_0)$ . We will prove that there exists no sentence  $\sigma$  of the first-order logic such that a ternary structure  $(X_0, T_0)$  with a connected underlying graph  $G$  is the B-structure of  $G$  if and only if  $(X_0, T_0)$  satisfies  $\sigma$ .

*Keywords:* connected graph, induced path, ternary relation, finite structure

*MSC 2000:* 05C38, 03C13

INTRODUCTION

The letters  $i, j, k, m$  and  $n$  will be reserved for denoting integers.

By a graph we mean here a graph in the sense of [2], i.e. a finite undirected graph without loops or multiple edges. If  $G$  is a graph, then  $V(G)$  and  $E(G)$  denote its vertex set and its edge set, respectively.

Let  $G$  be a graph, let  $v_0, \dots, v_n \in V(G)$ , and let

$$P: v_0, \dots, v_n$$

---

Research supported by Grant Agency of the Czech Republic, grant No. 401/98/0383.

be a path in  $G$ . We say that  $P$  is an *induced path* in  $G$  if  $v_i v_j \notin E(G)$  for all  $i, j \in \{0, \dots, n\}$  such that  $|i - j| \neq 1$ . Note that instead of the term “induced path” the term “minimal path” is sometimes used. If  $G$  is a connected graph, then we say that  $P$  is a *geodesic* in  $G$ , if  $d(v_0, v_n) = n$ , where  $d$  denotes the distance function of  $G$ . Instead of the term “geodesic” the term “shortest path” is sometimes used.

Let  $P$  and  $P'$  be induced paths in a graph  $G$ ; we will say that  $P$  and  $P'$  are disjoint if no vertex of  $G$  belongs both to  $P$  and to  $P'$ ; we will say that  $P$  and  $P'$  are non-adjacent in  $G$  if there exists no pair of vertices  $u$  and  $u'$  such that  $u$  belongs to  $P$ ,  $u'$  belongs to  $P'$  and  $u$  and  $u'$  are adjacent in  $G$ .

## PART 1

By a *ternary structure* we mean an ordered pair  $(X_0, T_0)$ , where  $X_0$  is a *finite* nonempty set and  $T_0$  is a ternary relation on  $X_0$ .

Let  $(X_1, T_1)$  and  $(X_2, T_2)$  be ternary structures. By a *partial isomorphism* from  $(X_1, T_1)$  to  $(X_2, T_2)$  we mean such an injective mapping  $q$  that  $\text{Def}(q) \subseteq X_1$ ,  $\text{Im}(q) \subseteq X_2$  and

$$T_1(x, u, y) \text{ if and only if } T_2(q(x), q(u), q(y))$$

for all  $u, x, y \in \text{Def}(q)$ . (Note that the notion of a partial isomorphism from a ternary structure to a ternary structure is a special case of the notion of a partial isomorphism in the sense of [4], p. 15). Let  $(X_0, T_0)$  be a ternary structure. By the *pseudointerval function* of  $(X_0, T_0)$  we mean the mapping  $J$  of  $X_0 \times X_0$  into  $2^{X_0}$  defined as follows:

$$J(x, y) = \{u \in X_0; T_0(x, u, y)\}$$

for all  $x, y \in X_0$ .

Let  $(X_0, T_0)$  be a ternary structure, and let  $J$  denote its pseudointerval function. By the *underlying graph* of  $(X_0, T_0)$  we mean the graph  $G$  defined as follows:  $V(G) = X_0$  and

$$E(G) = \{uv; u, v \in X_0, u \neq v \text{ and } J(u, v) \cup J(v, u) = \{u, v\}\}.$$

We will say that  $(X_0, T_0)$  is *connected* if its underlying graph is connected.

Let  $G$  be a connected graph, and let  $\mathbf{P}_0$  be a subset of the set of all paths in  $G$ . By the  $\mathbf{P}_0$ -*structure* of  $G$  we mean the ternary structure  $(X_0, T_0)$  such that  $X_0 = E(G)$  and

$T_0(x, u, y)$  if and only if

there exists an  $x - y$  path  $P$  in  $G$  such that  $P \in \mathbf{P}_0$  and  $u$  belongs to  $P$

for all  $u, x, y \in X_0$ . Let  $(X_0, T_0)$  be the  $\mathbf{P}_0$ -structure of  $G$ . If  $\mathbf{P}_0$  is the set of all paths in  $G$ , the set of all induced paths in  $G$ , or the set of all geodesics in  $G$ , then we say that  $(X_0, T_0)$  is the A-structure of  $G$ , the B-structure of  $G$ , or the  $\Gamma$ -structure of  $G$ , respectively.

Let  $G$  be a connected graph, and let  $d$  denote its distance function. By the  $\Sigma$ -structure of  $G$  we mean the ternary structure  $(X_0, T_0)$  such that  $X_0 = V(G)$  and

$$T_0(x, u, y) \text{ if and only if } d(x, u) = 1 \text{ and } d(u, y) = d(x, y) - 1$$

for all  $u, x, y \in X_0$ .

Let  $(X_0, T_0)$  be a ternary structure, and let  $\mathbf{Z}$  stand for A, B,  $\Gamma$  or  $\Sigma$ . We say that  $(X_0, T_0)$  is a  $\mathbf{Z}$ -structure if there exists a connected graph  $G$  such that  $(X_0, T_0)$  is the  $\mathbf{Z}$ -structure of  $G$ .

Let  $(T_0, X_0)$  be a ternary structure, and let  $J$  denote its pseudointerval function. We will say that  $(X_0, T_0)$  satisfies condition C1, C1', C2 or C3 if

- (C1)  $J(x, x) = \{x\}$  for all  $x \in X_0$ ,
- (C1')  $J(x, x) = \emptyset$  for all  $x \in X_0$ ,
- (C2)  $J(x, y) = J(y, x)$  for all  $x, y \in X_0$ , or
- (C3)  $x \in J(x, y)$  for all  $x, y \in X_0$ ,

respectively. It is obvious that all A-structures, B-structures and  $\Gamma$ -structures satisfy conditions C1, C2 and C3 and that all  $\Sigma$ -structures satisfy condition C1'.

Let  $\mathbf{Z}$  stand for B,  $\Gamma$  or  $\Sigma$ . It is easy to see that if  $(X_0, T_0)$  is a  $\mathbf{Z}$ -structure, then it is the  $\mathbf{Z}$ -structure of exactly one connected graph, namely of the underlying graph of  $(X_0, T_0)$ . This means that all B-structures, all  $\Gamma$ -structures and all  $\Sigma$ -structures are connected. However, this is not the case with A-structures. The underlying graph of the A-structure of a complete graph with at least three vertices has no edges.

Let  $(X_0, T_0)$  be a ternary structure, and let  $J$  denote its pseudointerval function. We will say that  $(X_0, T_0)$  is *scant* if (a) it satisfies conditions C1 and C2, and (b) the following statement holds for all distinct  $x, y \in X_0$ : if  $J(x, y) \neq \{x, y\}$ , then  $J(x, y) = X_0$ . Clearly, every scant ternary structure is determined by its underlying graph. It is not difficult to see that if the  $\Gamma$ -structure of a connected graph  $G$  is scant, then the diameter of  $G$  does not exceed two. This is not the case with B-structures. It is obvious that the B-structure of every cycle is scant. Thus, for every  $n \geq 3$  there exists a connected graph  $G$  of diameter  $n$  such that the B-structure of  $G$  is scant.

Let  $(X_0, T_0)$  be a ternary structure, let  $J$  denote its pseudointerval function, and let  $G$  denote the underlying graph of  $(X_0, T_0)$ . If  $J$  satisfies conditions C1, C2 and C3, then  $J$  is a transit function on  $G$  in the sense of Mulder [7]. Recall that if  $(X_0, T_0)$

is a  $\Gamma$ -structure or a B-structure, then it is respectively the  $\Gamma$ -structure or the B-structure of  $G$ . If  $(X_0, T_0)$  is a  $\Gamma$ -structure, then  $J$  is called the interval function of  $G$ ; cf. Mulder [6], where the interval function of a connected graph was studied widely. If  $(X_0, T_0)$  is a B-structure, then  $J$  is called the induced path function or the minimal path function on  $G$  in [7]. The induced path function on a connected graph was studied by Duchet [3] and by Morgana and Mulder [5].

The pseudointerval functions of A-structures were characterized in Changat, Klavžar and Mulder [1] while the pseudointerval functions of  $\Gamma$ -structures were characterized by the present author in [8], [10] and [12]. These characterizations can be reformulated easily as characterizations of A-structures and of  $\Gamma$ -structures by a finite set of axioms or, more strictly, by a unique axiom.

The result obtained for  $\Sigma$ -structures by the present author in [9] and [11] is not too strong:  $\Sigma$ -structures were characterized as connected ternary structures satisfying a finite set of axioms. This result could be reformulated as follows: there exists an axiom  $\sigma$  in a language of the first order logic such that a connected ternary structure  $(X_0, T_0)$  is a  $\Sigma$ -structure if and only if  $(X_0, T_0)$  satisfies  $\sigma$ .

In the present paper we will prove that a similar result does not hold for B-structures. To prove this, we will need a certain portion of mathematical logic; for precise formulations and further details the reader is referred to Ebbinghaus and Flum [4], p. 1–12. (Especially, the explanation of the term “satisfy”, which will be used in Theorem 1, can be found in [4], p. 6).

Let  $T$  be the symbol for a ternary relation. By an atomic formula of the first-order logic of vocabulary  $\{T\}$  (shortly: by an atomic formula) we mean an expression

$$x = y,$$

where  $x$  and  $y$  are variables, or an expression

$$T(x, u, y),$$

where  $u$ ,  $x$  and  $y$  are variables. The formulae of the first-order logic of vocabulary  $\{T\}$  (shortly: the formulae) will be defined as follows:

- every atomic formula is a formula;
- if  $\alpha$  is a formula, then  $\neg\alpha$  is a formula;
- if  $\alpha_1$  and  $\alpha_2$  are formulae, then  $\alpha_1 \vee \alpha_2$  is a formula;
- if  $\alpha$  is a formula and  $x$  is a variable, then  $\exists x\alpha$  is a formula;
- no other expressions are formulae.

Following [4] we define the *quantifier rank*  $qr(\alpha)$  of a formula  $\alpha$ :

if  $\alpha$  is atomic, then  $qr(\alpha) = 0$ ;

if  $\alpha$  is  $\neg\beta$ , where  $\beta$  is a formula, then  $qr(\alpha) = qr(\beta)$ ;

if  $\alpha$  is  $\beta_1 \vee \beta_2$ , where  $\beta_1$  and  $\beta_2$  are formulae, then  $qr(\alpha) = \max(qr(\beta_1), qr(\beta_2))$ ;

if  $\alpha$  is  $\exists x\beta$ , where  $\beta$  is a formula and  $x$  is a variable, then  $qr(\alpha) = qr(\beta) + 1$ .

The most important formulae are sentences: a formula  $\alpha$  is called a sentence if for every atomic subformula  $\beta$  of  $\alpha$ , every variable belonging to  $\beta$  is in the scope of the corresponding quantifier.

The next theorem, which is a special case of Fraïssé's Theorem, will be an important tool for us:

**Theorem 1.** *Let  $(X_1, T_1)$  and  $(X_2, T_2)$  be ternary structures, and let  $n \geq 1$ . Then the following statements (A) and (B) are equivalent:*

(A)  *$(X_1, T_1)$  and  $(X_2, T_2)$  satisfy the same sentences  $\sigma$  with  $qr(\sigma) \leq n$ .*

(B) *There exist nonempty sets  $\mathbf{Q}_0, \dots, \mathbf{Q}_n$  of partial isomorphisms from  $(X_1, T_1)$  to  $(X_2, T_2)$  such that for each  $m, 1 \leq m < n$ , we have*

(I) *for every  $q \in \mathbf{Q}_{m+1}$  and every  $x \in X_1$  there exists  $r \in \mathbf{Q}_m$  such that  $q \subseteq r$  and  $x \in \text{Def}(r)$ ;*

(II) *for every  $q \in \mathbf{Q}_{m+1}$  and every  $x \in X_2$  there exists  $r \in \mathbf{Q}_m$  such that  $q \subseteq r$  and  $x \in \text{Im}(r)$ .*

For the proof of Fraïssé's Theorem (and further closely related results) the reader is referred to [4], Chapter 1.

## PART 2

Assume that an infinite sequence

$$u_0, w_0, u_1, w_1, u_2, w_2, \dots$$

of mutually distinct vertices is given.

Let  $k \geq 3$ . By  $F_k$  we denote the graph with vertices

$$u_0, w_0, u_1, w_1, \dots, u_{6k-1}, w_{6k-1}$$

and with edges

$$\begin{aligned}
 &u_0u_1, u_1u_2, \dots, u_{3k-2}u_{3k-1}, u_{3k-1}u_0, \\
 &u_{3k}u_{3k+1}, u_{3k+1}u_{3k+2}, \dots, u_{6k-2}u_{6k-1}, u_{6k-1}u_{3k}, \\
 &w_0w_1, w_1w_2, \dots, w_{3k-2}w_{3k-1}, w_{3k-1}w_0, \\
 &w_{3k}w_{3k+1}, w_{3k+1}w_{3k+2}, \dots, w_{6k-2}w_{6k-1}, w_{6k-1}w_{3k}, \\
 &u_0w_0, u_1w_1, \dots, u_{6k-1}w_{6k-1}, \\
 &u_0u_{3k}, u_ku_{4k}, u_{2k}u_{5k}.
 \end{aligned}$$

A diagram of  $F_3$  is presented in Fig. 1.




Fig. 1.

**Lemma 1.** *Let  $k \geq 3$ . Then the B-structure of  $F_k$  is scant.*

*Proof.* Let  $x \in V(F_k)$ . Then there exists exactly one  $i$ ,  $0 \leq i \leq 6k - 1$ , such that  $x = u_i$  or  $x = w_i$ ; we define  $\text{ind}(x) = i$ . For every  $y \in V(F_k)$  we define  $y^L$  and  $y^R$  as follows:

- if  $\text{ind}(y) \in \{0, k, 2k, 3k, 4k, 5k\}$ , then  $y^L = y^R = u_{\text{ind}(y)}$ ;
- if  $jk < \text{ind}(y) < (j+1)k$ , where  $j \in \{0, 1, 3, 4\}$ , then  $y^L = u_{jk}$  and  $y^R = u_{(j+1)k}$ ;
- if  $2k < \text{ind}(y) < 3k$ , then  $y^L = u_{2k}$  and  $y^R = u_0$ ;
- if  $5k < \text{ind}(y) < 6k$ , then  $y^L = u_{5k}$  and  $y^R = u_{3k}$ .

Let  $J$  denote the pseudointerval function of the B-system of  $F_k$ . Consider arbitrary  $x, y \in V(F_k)$  such that  $d(x, y) \geq 2$ , where  $d$  denotes the distance function of  $F_k$ . We want to prove that  $J(x, y) = V(F_k)$ .

Denote  $V_1 = \{v \in V(F_k); 0 \leq \text{ind}(v) \leq 3k - 1\}$  and  $V_2 = V(F_k) \setminus V_1$ . Without loss of generality we assume that  $x \in V_1$ . We distinguish two cases.

C a s e 1. Let  $y \in V_1$ . It is clear that  $V_1 \subseteq J(x, y)$  and

$$V_2 \subseteq J(u_0, u_k) \cap J(u_k, u_{2k}) \cap J(u_{2k}, u_0).$$

Recall that  $d(x, y) \geq 2$ . We can see that there exist  $x_1 \in \{x^L, x^R\}$  and  $y_1 \in \{y^L, y^R\}$  such that  $x_1 \neq y_1$  and there exist an induced  $x - x_1$  path  $P_x$  in  $F_k$  and an induced  $y_1 - y$  path  $P_y$  in  $F_k$  with the property that  $P_x$  and  $P_y$  are disjoint and non-adjacent in  $F_k$ . This implies that  $J(x, y) = V(F_k)$ .

C a s e 2. Let  $y \in V_2$ . We distinguish two subcases.

S u b c a s e 2.1. Let  $d(x, y) = 2$ . Then  $x \in \{u_0, u_k, u_{2k}\}$  or  $y \in \{u_{3k}, u_{4k}, u_{5k}\}$ . Without loss of generality we assume that  $x = u_0$ . Then  $y = w_{3k}$  or  $y = u_{3k+1}$  or  $y = u_{6k-1}$ .

First, let  $y = w_{3k}$ . Consider the following five sequences:

$$\begin{aligned} &u_0, u_{3k}, w_{3k}; \\ &u_0, u_1, \dots, u_{k-1}, u_k, u_{4k}, u_{4k-1}, \dots, u_{3k+1}, u_{3k}, w_{3k}; \\ &u_0, u_{3k-1}, u_{3k-2}, \dots, u_{k+1}, u_k, u_{4k}, u_{4k+1}, \dots, u_{6k-2}, u_{6k-1}, u_{3k}, w_{3k}; \\ &u_0, w_0, w_1, \dots, w_{k-1}, w_k, u_k, u_{4k}, w_{4k}, w_{4k-1}, \dots, w_{3k+1}, w_{3k}; \\ &u_0, w_0, w_{3k-1}, w_{3k-2}, \dots, w_k, u_k, u_{4k}, w_{4k}, w_{4k+1}, \dots, w_{6k-1}, w_{3k}. \end{aligned}$$

Each vertex of  $F_k$  belongs to at least one of these sequences. Moreover, each of these sequences is an induced  $x - y$  path in  $F_k$ . Thus  $J(x, y) = V(F_k)$ .

Now, let  $y \neq w_{3k}$ . Without loss of generality we assume that  $y = u_{3k+1}$ . Consider the following five sequences:

$$\begin{aligned} &u_0, u_{3k}, u_{3k+1}; \\ &u_0, u_1, \dots, u_{k-1}, u_k, u_{4k}, u_{4k-1}, \dots, u_{3k+1}; \\ &u_0, u_{3k-1}, \dots, u_{k+1}, u_k, u_{4k}, u_{4k+1}, \dots, u_{6k-2}, u_{6k-1}, w_{6k-1}, w_{3k}, w_{3k+1}, u_{3k+1}; \\ &u_0, w_0, w_1, \dots, w_{k-1}, w_k, u_k, u_{4k}, w_{4k}, w_{4k-1}, \dots, w_{3k+1}, u_{3k+1}; \\ &u_0, w_0, w_{3k-1}, w_{3k-2}, \dots, w_k, u_k, u_{4k}, w_{4k}, w_{4k+1}, \dots, w_{6k-1}, w_{3k}, w_{3k+1}, u_{3k+1}. \end{aligned}$$

Again, each vertex of  $F_k$  belongs to at least one of these sequences and each of these sequences is an induced  $x - y$  path in  $F_k$ . Thus  $J(x, y) = V(F_k)$ .

S u b c a s e 2.2. Let  $d(x, y) \geq 3$ . Then there exist  $x_2 \in \{x^L, x^R\}$  and  $y_2 \in \{y^L, y^R\}$  such that  $d(x_2, y) \geq 3$  and  $d(x, y_2) \geq 3$ . Define  $x^* = u_{\text{ind}(x_2)+3k}$  and  $y^* = u_{\text{ind}(y_2)-3k}$ . Obviously,  $d(x^*, y) \geq 2$  and  $d(x, y^*) \geq 2$ . It is clear that  $V_1 \subseteq J(x, y^*)$  and  $V_2 \subseteq J(x^*, y)$ . This implies that  $J(x, y) = V(F_k)$ .

The proof is complete. □

Let  $k > 2$ . By  $F'_k$  we denote the graph with vertices

$$u_0, w_0, u_1, w_1, \dots, u_{6k-1}, w_{6k-1}$$

and with edges

$$\begin{aligned} &u_0u_1, u_1u_2, \dots, u_{2k-2}u_{2k-1}, u_{2k-1}u_0, \\ &u_{2k}u_{2k+1}, u_{2k+1}u_{2k+2}, \dots, u_{4k-2}u_{4k-1}, u_{4k-1}u_{2k}, \\ &u_{4k}u_{4k+1}, u_{4k+1}u_{4k+2}, \dots, u_{6k-2}u_{6k-1}, u_{6k-1}u_{4k}, \\ &w_0w_1, w_1w_2, \dots, w_{2k-2}w_{2k-1}, w_{2k-1}w_0, \\ &w_{2k}w_{2k+1}, w_{2k+1}w_{2k+2}, \dots, w_{4k-2}w_{4k-1}, w_{4k-1}w_{2k}, \\ &w_{4k}w_{4k+1}, w_{4k+1}w_{4k+2}, \dots, w_{6k-2}w_{6k-1}, w_{6k-1}w_{4k}, \\ &u_0w_0, u_1w_1, \dots, u_{6k-1}w_{6k-1}, \\ &u_ku_{2k}, u_{3k}u_{4k}, u_{5k}u_0. \end{aligned}$$

A diagram of  $F'_3$  is presented in Fig. 2.




Fig. 2.

**Lemma 2.** *Let  $k \geq 3$ . Then the B-structure of  $F'_k$  is not scant.*

*Proof.* Let  $J$  denote the pseudointerval function of the B-structure of  $F'_k$ . Since  $J(u_{k-1}, u_{k+1}) \neq V(F'_k)$ , the result follows.  $\square$

**Lemma 3.** Let  $n \geq 1$  and  $k > 2^{n+1}$ . Assume that  $(X_1, T_1)$  and  $(X_2, T_2)$  are scant ternary structures such that the underlying graph of  $(X_1, T_1)$  is  $F_k$  and the underlying graph of  $(X_2, T_2)$  is  $F'_k$ . Then  $(X_1, T_1)$  and  $(X_2, T_2)$  satisfy the same sentences  $\sigma$  with  $qr(\sigma) \leq n$ .

*Proof.* Put  $U = \{u_0, u_1, \dots, u_{6k-1}\}$ ,  $U^b = \{u_0, u_k, u_{2k}, u_{3k}, u_{4k}, u_{5k}\}$ ,  $W = \{w_0, w_1, \dots, w_{6k-1}\}$  and  $W^b = \{w_0, w_k, w_{2k}, w_{3k}, w_{4k}, w_{5k}\}$ . Obviously,  $X_1 = U \cup W = X_2$ .

If  $x, y \in U \cup W$ , then we will write  $x \sim y$  if and only if  $x, y \in U$  or  $x, y \in W$ . We define  $u_i^\diamond = w_i$  and  $w_i^\diamond = u_i$  for all  $i$ ,  $0 \leq i \leq 6k - 1$ . Thus  $(x^\diamond)^\diamond = x$  for each  $x \in U \cup W$  and  $y^\diamond \sim z^\diamond$  if and only if  $y \sim z$  for all  $y, z \in U \cup W$ . We define  $[x] = x$  for every  $x \in U$  and  $[x] = x^\diamond$  for every  $x \in W$ .

By  $F^*$  we mean  $F_k$  or  $F'_k$ . Let  $d^*$  denote the distance function of  $F^*$ . Define

$$e^*(x, y) = d^*([x], [y]) \text{ for all } x, y \in U \cup W.$$

Obviously,  $e^*(x, y) = 0$  if and only if  $x = y$  or  $x^\diamond = y$  for all  $x, y \in U \cup W$ .

Recall that  $k > 2^{n+1}$ . Consider an arbitrary  $x \in U \cup W$  and denote  $D(x) = \{y \in U^b \cup W^b; e^*(x, y) \leq 2^n\}$ ; it is easy to see that  $|D(x)| \leq 4$  and if  $D(x) \neq \emptyset$ , then the subgraph of  $F^*$  induced by  $D(x)$  is a path of length either one or three.

Consider arbitrary  $x, y \in U \cup W$  such that  $e^*(x, y) \leq 2^n$ . It is easy to see that (i) every  $x - y$  geodesic in  $F^*$  contains at most two vertices in  $U^b$ ; (ii) if at least one  $x - y$  geodesic in  $F^*$  contains two vertices in  $U^b$ , then every  $x - y$  geodesic in  $F^*$  contains two vertices in  $U^b$  and these two vertices are adjacent in  $F^*$ . We will write  $f^*(x, y) = 1$  if every  $x - y$  geodesic in  $F^*$  contains at most one vertex in  $U^b$  and  $f^*(x, y) = 2$  otherwise.

For every  $m$ ,  $0 \leq m \leq n$  and for all  $x, y \in U \cup W$  we define

$$\begin{aligned} e_m^*(x, y) &= e^*(x, y) \text{ if } e^*(x, y) \leq 2^m, \\ e_m^*(x, y) &= \infty \text{ if } e^*(x, y) > 2^m. \end{aligned}$$

Consider an arbitrary  $m$ ,  $0 \leq m < n$ . We see that

(1) if  $e_{m+1}^*(x, y) = \infty$  and  $e_m^*(y, z) < \infty$ , then  $e_m^*(x, z) = \infty$  for all  $x, y, z \in U \cup W$ .

We will write  $e$ ,  $e_m$  and  $f$  instead of  $e^*$ ,  $e_m^*$  and  $f^*$  respectively if  $F^*$  is  $F_k$ , and  $e'$ ,  $e'_m$  and  $f'$  instead of  $e^*$ ,  $e_m^*$  and  $f^*$  respectively if  $F^*$  is  $F'_k$ .

Recall that  $(X_1, T_1)$  and  $(X_2, T_2)$  are scant. We denote by PART the set of all partial isomorphisms  $p$  from  $F_k$  to  $F'_k$  such that  $U^b \cup W^b \subseteq \text{Def}(p)$ ,

$$p(x) \sim x \text{ for all } x \in \text{Def}(p),$$

and

$$\begin{aligned} p(u_0) &= u_0, p(w_0) = w_0, p(u_k) = u_k, p(w_k) = w_k, p(u_{2k}) = u_{4k}, p(w_{2k}) = w_{4k}, \\ p(u_{3k}) &= u_{5k}, p(w_{3k}) = w_{5k}, p(u_{4k}) = u_{2k}, p(w_{4k}) = w_{2k}, \\ p(u_{5k}) &= u_{3k} \text{ and } p(w_{5k}) = w_{3k}. \end{aligned}$$

Obviously, there exists exactly one  $p_0 \in \text{PART}$  such that  $\text{Def}(p_0) = U^b \cup W^b$ .

For every  $m$ ,  $0 \leq m \leq n$ , we denote by  $\mathbf{Q}_m$  the set of all  $q \in \text{PART}$  such that  $|\text{Def}(q)| \leq 12 + n - m$  and that  $e'_m(q(x), q(y)) = e_m(x, y)$  for all  $x, y \in \text{Def}(q)$ .

It is clear that  $\mathbf{Q}_n = \{p_0\}$ . As follows from the definition,  $\mathbf{Q}_n \subseteq \dots \subseteq \mathbf{Q}_0$ .

Consider an arbitrary  $m$ ,  $0 \leq m < n$ . We need to show that conditions (I) and (II) (of Theorem 1) hold.

Consider an arbitrary  $q \in \mathbf{Q}_{m+1}$  and an arbitrary  $x \in U \cup W$ . If  $x \in \text{Def}(q)$ , we put  $r = q$ . Assume that  $x \notin \text{Def}(q)$ . Then  $x \notin U^b \cup W^b$ . We distinguish two cases.

Case 1. Assume that there exists  $y \in \text{Def}(q)$  such that  $e_m(x, y) < \infty$ . Without loss of generality we assume that  $e_m(x, y) \leq e_m(x, y_0)$  for every  $y_0 \in \text{Def}(q)$ .

First, let  $e_m(x, y) = 0$ . Since  $x \notin \text{Def}(q)$ , we have  $y = x^\diamond$ . We put  $x' = (q(y))^\diamond$ .

Now, we assume that  $e_m(x, y) > 0$ . We distinguish four subcases.

Subcase 1.1. Assume that

$$(2) \quad \begin{aligned} &\text{there exists } z \in \text{Def}(q) \text{ such that} \\ &e_m(x, z) < \infty \text{ and } e(y, z) = e_m(y, x) + e_m(x, z). \end{aligned}$$

Without loss of generality we assume that  $e_m(x, z) \leq e_m(x, z_0)$  for every  $z_0 \in \text{Def}(q)$  such that  $e_m(x, z_0) < \infty$  and  $e(y, z_0) = e_m(y, x) + e_m(x, z_0)$ . Since  $e_m(x, y) > 0$ , it is obvious that  $e_m(x, z) > 0$ . Since  $e_m(x, y) < \infty$  and  $e_m(x, z) < \infty$ , we get  $e_{m+1}(y, z) < \infty$ . Since  $y, z \in \text{Def}(q)$ , we have  $e'_{m+1}(q(y), q(z)) = e_{m+1}(y, z)$ . There exists exactly one  $x' \in (U \cup W) \setminus \text{Im}(q)$  such that  $e'(q(y), q(z)) = e'_m(q(y), x') + e_m(x', q(z))$  and  $x' \sim x$ .

Subcase 1.2. Assume (2) does not hold and

$$(3) \quad \begin{aligned} &\text{there exists } z \in \text{Def}(q) \text{ such that} \\ &0 < e_{m+1}(y, z) < \infty, f(y, z) = 1 \text{ and } e(x, z) = e_m(x, y) + e_{m+1}(y, z). \end{aligned}$$

Without loss of generality we assume that  $e_{m+1}(y, z) \leq e_{m+1}(y, z_0)$  for every  $z_0 \in \text{Def}(q)$  such that  $0 < e_{m+1}(y, z_0) < \infty, f(y, z_0) = 1$  and  $e(x, z_0) = e_m(x, y) + e_{m+1}(y, z_0)$ . Since  $y, z \in \text{Def}(q)$ , we get  $e'_{m+1}(q(y), q(z)) = e_{m+1}(y, z)$ . There exists exactly one  $x' \in (U \cup W) \setminus \text{Im}(q)$  such that  $e'(x', q(z)) = e'_m(x', q(y)) + e'_{m+1}(q(y), q(z))$  and  $x' \sim x$ .

S u b c a s e 1.3. Assume (2) and (3) do not hold and

- (4) there exists  $z \in \text{Def}(q)$  such that  
 $0 < e_{m+1}(y, z) < \infty, f(y, z) = 2$  and  $e(x, z) = e_m(x, y) + e_{m+1}(y, z)$ .

Without loss of generality we assume that  $e_{m+1}(y, z) \leq e_{m+1}(y, z_0)$  for every  $z_0 \in \text{Def}(q)$  such that  $0 < e_{m+1}(y, z_0) < \infty, f(y, z_0) = 2$  and  $e(x, z_0) = e_m(x, y) + e_{m+1}(y, z_0)$ . It is easy to see that  $y, z \in U^b \cup W^b$  and  $e(y, z) = 1$ . Since  $y, z \in \text{Def}(q)$ , we get  $q(y), q(z) \in U^b \cup W^b$  and  $e'(q(y), q(z)) = 1$ . There exist exactly two vertices belonging to  $(U \cup W) \setminus \text{Im}(q)$ , say vertices  $v_1$  and  $v_2$ , such that  $e'(v_j, q(z)) = e'_m(v_j, q(y)) + 1$  and  $v_j \sim x$  for  $j = 1, 2$ . Consider an arbitrary  $x' \in \{v_1, v_2\}$ .

S u b c a s e 1.4. Assume (2), (3) and (4) do not hold. Then there exists no  $z \in \text{Def}(q)$  such that  $0 < e_{m+1}(y, z) \leq e_m(x, y) + 2^m$ . Thus there exists no  $z \in \text{Def}(q)$  such that  $0 < e'_{m+1}(q(y), q(z)) \leq e_m(x, y) + 2^m$ . This implies that there exist exactly two vertices belonging to  $(U \cup W) \setminus \text{Im}(q)$ , say vertices  $v_1$  and  $v_2$ , such that  $e'_m(v_j, q(y)) = e_m(x, y)$  and  $v_j \sim x$  for  $j = 1, 2$ . Consider an arbitrary  $x' \in \{v_1, v_2\}$ .

C a s e 2. Assume that  $e_m(x, y) = \infty$  for every  $y \in \text{Def}(q)$ . There exists  $x' \in (U \cup W) \setminus \text{Im}(q)$  such that  $x' \sim x$  and  $e'_m(x', q(y)) = \infty$  for every  $y \in \text{Def}(q)$ .

Define  $r = q \cup \{(x, x')\}$ . If we take (1) into account, we can see that  $r \in \mathbf{Q}_m$ . Thus condition (I) holds.

The fact that condition (II) holds can be proved analogously. Applying Theorem 1, we obtain the result of the lemma.  $\square$

R e m a r k. The introduction of functions  $e_m^*$  in the proof of Lemma 3 is a modification of one of the ideas in Example 1.3.5 of [4].

**Theorem 2.** *There exists no sentence  $\sigma$  of the first-order logic of vocabulary  $\{T\}$  such that a connected ternary structure is a B-structure if and only if it satisfies  $\sigma$ .*

P r o o f. Combining Lemmas 1, 2 and 3, we get the theorem.  $\square$

Note that Theorem 2 can be reformulated as follows: There exists no finite set  $S$  of sentences of first-order logic of vocabulary  $\{T\}$  such that a connected ternary structure is a B-structure if and only if it satisfies each sentence in  $S$ .

#### References

- [1] *M. Changat, S. Klavžar, H. M. Mulder*: The all-path transit function of a graph. Czechoslovak Math. J. 52 (2001), 439–448.
- [2] *G. Chartrand, L. Lesniak*: Graphs & Digraphs. Third edition. Chapman & Hall, London, 1996.
- [3] *P. Duchet*: Convex sets in graphs, II. Minimal path convexity. J. Combinatorial Theory, Series B 44 (1998), 307–316.

- [4] *H.-D. Ebbinghaus, J. Flum*: Finite Model Theory. Springer, Berlin, 1995.
- [5] *M. A. Morgana, H. M. Mulder*: The induced path convexity, betweenness and svelte graphs. To appear in Discrete Mathematics.
- [6] *H. M. Mulder*: The interval function of a graph. MC-tract 132, Mathematisch Centrum, Amsterdam, 1980.
- [7] *H. M. Mulder*: Transit functions on graphs. In preparation.
- [8] *L. Nebeský*: A characterization of the interval function of a connected graph. Czechoslovak Math. J. *44* (1994), 173–178.
- [9] *L. Nebeský*: Geodesics and steps in a connected graph. Czechoslovak Math. J. *47* (1997), 149–161.
- [10] *L. Nebeský*: Characterizing the interval function of a connected graph. Math. Bohem. *123* (1998), 137–144.
- [11] *L. Nebeský*: A theorem for an axiomatic approach to metric properties of connected graphs. Czechoslovak Math. J. *50* (2000), 121–133.
- [12] *L. Nebeský*: The interval function of a connected graph and a characterization of geodetic graphs. Math. Bohem. *126* (2001), 247–254.

*Author's address*: *L. Nebeský*, Univerzita Karlova v Praze, Filozofická fakulta, nám. Jana Palacha 2, 116 38 Praha 1, Czech Republic, e-mail: [ladislav.nebesky@ff.cuni.cz](mailto:ladislav.nebesky@ff.cuni.cz).