

Alfonz Haviar; Gabriela Monoszová  
Constructions of cell algebras

*Mathematica Bohemica*, Vol. 130 (2005), No. 1, 89–100

Persistent URL: <http://dml.cz/dmlcz/134224>

## Terms of use:

© Institute of Mathematics AS CR, 2005

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

## CONSTRUCTIONS OF CELL ALGEBRAS

ALFONZ HAVIAR, GABRIELA MONOSZOVÁ, Banská Bystrica

(Received April 22, 2004)

*Abstract.* A construction of cell algebras is introduced and some of their properties are investigated. A particular case of this construction for lattices of nets is considered.

*Keywords:* Plonka sum, cell algebra, prelattice

*MSC 2000:* 08A05, 06F99

## 1. INTRODUCTION

There are many ways how to construct a “new” algebra from algebras of the same type. The relationship between the resulting algebra and the original ones depends on the construction. For instance, the direct product  $\prod_{i \in I} \mathcal{A}_i$  of algebras of the same type is an algebra satisfying the identities which hold in all algebras  $\mathcal{A}_i$ ,  $i \in I$ . On the other hand, the Plonka sum  $\sum_{i \in I} \mathcal{A}_i$  [9] satisfies only the regular identities which hold in all algebras  $\mathcal{A}_i$ ,  $i \in I$ . A less known construction was introduced by Hecht in [7]. The algebra he constructed preserves only identities of the type

$$(1.1) \quad \begin{aligned} f(r(x_1, \dots, x_n), p_2(x_1, \dots, x_n), \dots, p_k(x_1, \dots, x_n)) \\ = f(r(x_1, \dots, x_n), q_2(x_1, \dots, x_n), \dots, q_k(x_1, \dots, x_n)) \end{aligned}$$

and all their consequences, where  $f$  is a  $k$ -ary operational symbol and  $r$ ,  $p_i$ ,  $q_i$ ,  $i = 2, \dots, k$ , are polynomials of variables  $x_1, \dots, x_n$ .

We introduce a construction of algebras which is similar both to Plonka sums and Hecht’s construction.

---

The authors were supported by the grant VEGA 2/2060/22 of the Slovak Grant Agency.

## 2. CELL ALGEBRAS

Throughout the paper we assume that all algebras considered are of a given type  $\tau$ . By  $F$  we denote the set of all operational symbols of the type  $\tau$ , i.e.  $F = \{f_t; t \in \tau\}$ . We write  $f_t^{(A)}$  for the realization of  $f_t$  on a set  $A$ . We often denote briefly by  $f$  an operational symbol and also its realization (when no confusion can arise).

Let  $\mathcal{A} = (A, F)$  be an algebra of a type  $\tau$ . For each element  $a \in A$  let an algebra  $\mathcal{B}_a = (B_a, F)$  of the type  $\tau$  be given and let  $B_a \cap B_b = \emptyset$  if  $a \neq b$ . Moreover, for each  $k$ -ary ( $k \geq 1$ ) operation  $f \in F$  let  $\mathcal{S}^{(f)}$  be a system of mappings with the following property:

$$(2.1) \quad \begin{aligned} & \text{if } f(a_1, \dots, a_k) = a \quad \text{for } f \in F \text{ and } a_1, \dots, a_k \in A, \\ & \text{then there exists a mapping} \\ & \varphi_{a_i, a}^{(f)}: B_{a_i} \rightarrow B_a \quad \text{from } \mathcal{S}^{(f)} \text{ for each } i \in \{1, \dots, k\}. \end{aligned}$$

Let us denote  $S^{(F)} = \{S^{(f)}; f \in F\}$ .

**Definition 1.** Let  $\mathcal{A} = (A, F)$  be an algebra of the type  $\tau$ , let  $\mathcal{B}_a = (B_a, F)$ ,  $a \in A$ , be a system of algebras of the same type  $\tau$  and  $S^{(F)}$  a system of mappings satisfying (2.1). By the cell algebra with the basic algebra  $\mathcal{A}$ , the cells  $\mathcal{B}_a$ ,  $a \in A$  and with the system  $S^{(F)}$  we mean the algebra of the type  $\tau$  with the carrier  $M = \bigcup_{a \in A} B_a$  and the operations  $f^{(M)}$  defined on  $M$  as follows:

1. if  $f \in F$  is a  $k$ -ary operational symbol,  $k \geq 1$ ,  $x_1 \in B_{a_1}, \dots, x_k \in B_{a_k}$  and  $f(a_1, \dots, a_k) = a$  then

$$(2.2) \quad f^{(M)}(x_1, \dots, x_k) = f^{(B_a)}(\varphi_{a_1, a}^{(f)}(x_1), \dots, \varphi_{a_k, a}^{(f)}(x_k));$$

2. if  $f$  is a nullary operational symbol and  $f^{(A)} = c$  then  $f^{(M)} = f^{(B_c)}$ .

We denote it by  $\mathcal{A}(\mathcal{B}_a; a \in A)$  or briefly by  $\mathcal{A}(\mathcal{B})$ .

The next construction is described in [7]. Let  $\mathcal{A} = (A, F)$  be an algebra of the type  $\tau$ ,  $\{S_a; a \in A\}$  a family of pairwise disjoint nonvoid sets and  $\varphi_{a, \bar{a}}^{(f)}: S_a \rightarrow S_{\bar{a}}$  a family of mappings for all  $a \in A$ ,  $f \in F$ ,  $\bar{a} \in \{b \in A; b = f(a, a_1, \dots, a_{k-1}) \text{ for some } a_1, \dots, a_{k-1} \in A\}$ . For a  $k$ -ary operational symbol  $f$ ,  $k \geq 1$ , the operation  $f^{(M)}$  on  $M = \bigcup_{a \in A} S_a$  is defined by

$$(2.2a) \quad f^{(M)}(x_1, \dots, x_k) = \varphi_{a_1, a}^{(f)}(x_1),$$

where  $x_1 \in S_{a_1}, \dots, x_k \in S_{a_k}$ ,  $f(a_1, \dots, a_k) = a$ . If for each  $a \in A$ ,  $f \in F$  we define an operation  $f^{(S_a)}$  on  $S_a$  by  $f^{(S_a)}(x_1, \dots, x_k) = x_1$ , we get an algebra  $\mathcal{B}_a = (S_a, F)$

of the same type  $\tau$  and the identity (2.2) is of the form (2.2a). So, the algebra constructed in [7] is a special case of a cell algebra.

If we do not require any additional conditions for the system of mappings  $\mathcal{S}^{(F)}$  (analogously to [9]) then the algebra  $\mathcal{A}(\mathcal{B})$  has no close relationship to algebras  $\mathcal{A}$  and  $\mathcal{B}_a$ . However, there are some identities preserved by the construction of cell algebras.

**Theorem 2.** *Let  $\mathcal{A}(\mathcal{B}) = (M, F)$  be a cell algebra with a basic algebra  $\mathcal{A} = (A, F)$ , cells  $\mathcal{B}_a = (B_a, F)$ ,  $a \in A$ , and let the system  $\mathcal{S}^{(F)}$  satisfy (2.1) and moreover  $S^{(f)} = S^{(g)}$  for every  $f, g \in F$ . If the basic algebra  $\mathcal{A}$  and also every cell  $\mathcal{B}_a$  satisfies an identity*

$$(2.3) \quad f(x_1, \dots, x_m) = g(y_1, \dots, y_n)$$

where  $f, g$  are  $m$ -ary and  $n$ -ary operational symbols,  $m \geq 1$ ,  $n \geq 1$ , then the identity (2.3) holds in the cell algebra  $\mathcal{A}(\mathcal{B})$ , too.

**Proof.** Let  $x_1 \in B_{a_1}, \dots, x_m \in B_{a_m}$ ,  $y_1 \in B_{b_1}, \dots, y_n \in B_{b_n}$  and  $f(a_1, \dots, a_m) = a$ ,  $g(b_1, \dots, b_n) = b$ . By assumption we have  $a = b$ , and moreover

$$\begin{aligned} f^{(M)}(x_1, \dots, x_m) &= f^{(B_a)}(\varphi_{a_1, a}^{(f)}(x_1), \dots, \varphi_{a_m, a}^{(f)}(x_m)) \\ &= g^{(B_a)}(\varphi_{b_1, a}^{(g)}(y_1), \dots, \varphi_{b_n, a}^{(g)}(y_n)) = g^{(M)}(y_1, \dots, y_n). \end{aligned}$$

□

**Corollary 3.** *If a basic algebra  $\mathcal{A}$  and each cell  $\mathcal{B}_a$  ( $a \in A$ ) is an abelian groupoid, then the cell algebra  $\mathcal{A}(\mathcal{B})$  is also an abelian groupoid.*

Common identities of more complicated type than (2.3) are not preserved by the cell algebra construction. For example, let us consider an identity of the type

$$(2.4) \quad f(p(x_1, \dots, x_m), x_2, \dots, x_m) = g(y_1, \dots, y_n),$$

where  $f, g$  are operational symbols of the type  $\tau$  and  $p$  is a term of the type  $\tau$  which is not a projection. There exist a basic algebra  $\mathcal{A} = (A, F)$ , cells  $\mathcal{B}_a = (B_a, F)$ ,  $a \in A$  and a system of mappings  $\mathcal{S}^{(F)}$  such that the identity (2.4) holds in  $\mathcal{A}$  and in each cell  $\mathcal{B}_a$ ,  $a \in A$ , but (2.4) does not hold in the cell algebra  $\mathcal{A}(\mathcal{B}) = (M, F)$ . Assume  $x_1 \in B_{a_1}, \dots, x_m \in B_{a_m}$ ,  $y_1 \in B_{b_1}, \dots, y_n \in B_{b_n}$  and  $p^{(A)}(a_1, \dots, a_m) = a_0$ ,  $f^{(A)}(a_0, a_2, \dots, a_m) = a = g^{(A)}(b_1, \dots, b_n)$ . We get

$$\begin{aligned} f^{(M)}(p^{(M)}(x_1, \dots, x_m), x_2, \dots, x_m) \\ = f^{(B_a)}(\varphi_1(p^{(B_{a_0})}(x'_1, \dots, x'_m)), \varphi_{a_2, a}^{(f)}(x_2), \dots, \varphi_{a_m, a}^{(f)}(x_m)) \end{aligned}$$

where  $x'_1, \dots, x'_m$  are some elements and  $\varphi_1$  is a mapping depending not only on  $x_1, \dots, x_m$  but also on the term  $p$ . The result depends on the system of the maps  $\mathcal{S}^{(f)}$ , too. A special case of (2.4) is, for example, the identity

$$(2.4a) \quad f(h(x, y), y) = g(x, y),$$

where  $f, g, h$  are binary operational symbols. Let the realizations of these operational symbols in the basic algebra  $\mathcal{A}$  and also in each cell  $\mathcal{B}_a$ ,  $a \in A$ , satisfy the identity

$$h(x, y) = g(x, y) = y, \quad f(t, y) = t.$$

Then the identity (2.4a) is satisfied in the basic algebra and also in every cell. If  $x \in B_a$ ,  $y \in B_b$  in the cell algebra  $\mathcal{A}(\mathcal{B})$  we get

$$(2.5) \quad f^{(M)}(h^{(M)}(x, y), y) = \varphi_{b,b}^{(f)}(\varphi_{b,b}^{(h)}(y)), \quad g^{(M)}(x, y) = \varphi_{b,b}^{(g)}(y)$$

and so the identity (2.4a) need not be satisfied in  $\mathcal{A}(\mathcal{B})$ .

A class of identities preserved by the cell algebra construction can be increased by assuming some suitable conditions for mappings  $\varphi_{a,b}$  (analogous to the conditions for Plonka sums). First, let us consider algebras with one binary operation  $f$ . Which conditions are necessary for  $S^{(f)}$  in order that  $f$  satisfies the associative law or the idempotency?

Let a basic algebra  $\mathcal{A} = (A, f)$  and every cell  $\mathcal{B}_a = (B_a, f)$  be semigroups, i.e. let

$$(2.6) \quad f(f(x, y), z) = f(x, f(y, z))$$

hold in  $\mathcal{A}$  and in every cell  $\mathcal{B}_a$ ,  $a \in A$ . Let us assume that the realizations  $f^{(A)}$  and  $f^{(B_a)}$ ,  $a \in A$ , satisfy the identity

$$f(x, y) = x$$

(i.e.  $\mathcal{A}$  and  $\mathcal{B}_a$ ,  $a \in A$ , are left-zero semigroups). Let mappings  $\varphi_{a,b}^{(f)}$  be given for every  $a, b \in A$ . Take elements  $x \in B_{a_1}$ ,  $y \in B_{a_2}$ ,  $z \in B_{a_3}$  and let  $f(a_1, a_2) = a_0$ ,  $f(a_0, a_3) = a$ ,  $f(a_2, a_3) = a_4$  (by assumption  $f(a_1, a_4) = a$ ). Putting the elements considered to the left-hand side of the identity (2.6) we get (for the realization  $f^{(M)}$  of the cell algebra)

$$\begin{aligned} f^{(M)}(f^{(M)}(x, y), z) &= f^{(B_a)}(\varphi_{a_0,a}^{(f)}(f^{(B_{a_0})}(\varphi_{a_1,a_0}^{(f)}(x), \varphi_{a_2,a_0}^{(f)}(y))), \varphi_{a_3,a}^{(f)}(z)) \\ &= \varphi_{a_0,a}^{(f)}(\varphi_{a_1,a_0}^{(f)}(x)). \end{aligned}$$

Analogously, putting the elements to the right-hand side of (2.6) we get

$$f^{(M)}(x, f^{(M)}(y, z)) = \varphi_{a_1, a}^{(f)}(x).$$

Thus (2.6) holds in the cell algebra  $\mathcal{A}$  if

$$\varphi_{a_0, a}^{(f)}(\varphi_{a_1, a_0}^{(f)}(x)) = \varphi_{a_1, a}^{(f)}(x).$$

Hence

$$(2.6a) \quad \varphi_{b, c}^{(f)} \circ \varphi_{a, b}^{(f)} = \varphi_{a, c}^{(f)}$$

is a necessary condition for the associative law to hold in this case. The use of (2.6a) requires that  $\varphi_{a, b}^{(f)}$  be homomorphisms (analogously to [9]).

**Theorem 4.** *Let a basic algebra  $\mathcal{A} = (A, f)$  and every cell  $\mathcal{B}_a = (B_a, f)$ ,  $a \in A$ , be semigroups. If  $\mathcal{S}^{(f)}$  is a family of homomorphisms satisfying (2.1) and (2.6a) then the cell algebra  $\mathcal{A}(\mathcal{B})$  is also a semigroup.*

*Proof.* Consider as above  $x \in B_{a_1}$ ,  $y \in B_{a_2}$ ,  $z \in B_{a_3}$ . If  $f(a_1, a_2) = a_0$ ,  $f(a_0, a_3) = a$ ,  $f(a_2, a_3) = a_4$  we get

$$\begin{aligned} f^{(M)}(f^{(M)}(x, y), z) &= f^{(B_a)}(\varphi_{a_0, a}^{(f)}(f^{(B_{a_0})}(\varphi_{a_1, a_0}^{(f)}(x), (\varphi_{a_2, a_0}^{(f)}(y))), \varphi_{a_3, a}^{(f)}(z)) \\ &= f^{(B_a)}(f^{(B_{a_0})}(\varphi_{a_0, a}^{(f)}(\varphi_{a_1, a_0}^{(f)}(x)), \varphi_{a_0, a}^{(f)}(\varphi_{a_2, a_0}^{(f)}(y))), \varphi_{a_3, a}^{(f)}(z)) \\ &= f^{(B_a)}(f^{(B_a)}(\varphi_{a_1, a}^{(f)}(x), \varphi_{a_2, a}^{(f)}(y)), \varphi_{a_3, a}^{(f)}(z)), \end{aligned}$$

and similarly

$$f^{(M)}(x, f^{(M)}(y, z)) = f^{(B_a)}(\varphi_{a_1, a}^{(f)}(x), f^{(B_a)}(\varphi_{a_2, a}^{(f)}(y), \varphi_{a_3, a}^{(f)}(z))).$$

Since  $(B_a, f)$  is a semigroup, it follows that

$$f^{(M)}(f^{(M)}(x, y), z) = f^{(M)}(x, f^{(M)}(y, z)).$$

□

Let a basic algebra  $(A, f)$  and every cell  $(B_a, f)$ ,  $a \in A$ , be idempotent groupoids, i.e. let

$$(2.7) \quad f(x, x) = x.$$

By taking an element  $x \in B_a$  we get (in the cell algebra  $\mathcal{A}(\mathcal{B})$ )

$$f^{(M)}(x, x) = f^{(B_a)}(\varphi_{a,a}^{(f)}(x), \varphi_{a,a}^{(f)}(x)) = \varphi_{a,a}^{(f)}(x).$$

Hence the identity 2.7 holds if

$$(2.7a) \quad \varphi_{a,a}^{(f)} = \text{id} = \Delta_{B_a}$$

for each element  $a$  from the set  $A$ .

**Theorem 5.** *Let a basic algebra  $\mathcal{A} = (A, f)$  and every cell  $\mathcal{B}_a = (B_a, f)$  be bands (idempotent semigroups) or monoids. If  $\mathcal{S}^{(f)}$  is a family of homomorphisms satisfying (2.1), (2.6a) and (2.7a) then the cell algebra is also a band or a monoid, respectively.*

**Proof.** If  $\mathcal{A}$  and every cell are bands and the conditions concerning  $\mathcal{S}^{(f)}$  are fulfilled then  $\mathcal{A}(\mathcal{B})$  is also a band by Theorem 4 and the above considerations.

Let  $\mathcal{A}$  and each cell  $\mathcal{B}_a$ ,  $a \in A$  be monoids. We denote by 1 the neutral element in  $\mathcal{A}$  and by  $1_a$  the neutral element in  $\mathcal{B}_a$ . We are going to show that the element  $1_1$  is the neutral element in the cell algebra  $\mathcal{A}(\mathcal{B}) = (M, f)$ . For  $x \in B_a$  we get

$$f^{(M)}(x, 1_1) = f^{(B_a)}(\varphi_{a,a}^{(f)}(x), \varphi_{1,a}^{(f)}(1_1)) = f^{(B_a)}(\varphi_{a,a}^{(f)}(x), 1_a) = \varphi_{a,a}^{(f)}(x) = x$$

(a homomorphic image of a neutral element is a neutral element and  $a.1 = a$ ). Analogously,  $f^{(M)}(1_1, x) = x$ .  $\square$

When a basic algebra  $\mathcal{A} = (A, f)$  is a group, for each  $a, b \in A$  there exist elements  $x, y \in A$  for which  $f(x, a) = b$  and  $f(a, y) = b$ . It follows that for each homomorphism  $\varphi_{a,b}^{(f)} \in S^{(f)}$  there exists a homomorphism  $\varphi_{b,a}^{(f)} \in S^{(f)}$ . Moreover, if (2.6a) and (2.7a) hold, we have

$$\varphi_{a,b}^{(f)} \circ \varphi_{b,a}^{(f)} = \varphi_{a,a}^{(f)} = \text{id},$$

therefore  $\varphi_{a,b}^{(f)}$  and  $\varphi_{b,a}^{(f)}$  are bijections of  $B_a$  onto  $B_b$  and conversely. So,  $\varphi_{a,b}^{(f)}$  and  $\varphi_{b,a}^{(f)}$  are inverse isomorphisms. The next theorem shows that if a basic algebra and every cell are groups then one can obtain as cell algebras only direct products of groups.

**Theorem 6.** Let  $\mathcal{A}, \mathcal{B}, \mathcal{B}_a, a \in A$ , be algebras of the type  $\tau$  and for every  $a \in A$  let there exist an isomorphism  $\varphi_a: \mathcal{B}_a \rightarrow \mathcal{B}$ . If  $\mathcal{S}^{(F)}$  is a family of isomorphisms  $\varphi_{a,b}: \mathcal{B}_a \rightarrow \mathcal{B}_b$  for every  $a, b \in A$  (i.e.  $\mathcal{S}^{(f)} = \mathcal{S}^{(g)}$  for any  $f, g \in F$ ), then the cell algebra  $\mathcal{A}(\mathcal{B})$  is isomorphic to the direct product  $\mathcal{A} \times \mathcal{B}$ .

**Proof.** Without loss of generality we can assume that for each  $a, b \in A$  we have  $\varphi_b \circ \varphi_{a,b} = \varphi_a$  where  $\varphi_b, \varphi_a$  are isomorphisms of the cells  $\mathcal{B}_b, \mathcal{B}_a$  onto algebra  $\mathcal{B}$ . We are going to show that the mapping

$$\varphi: M \rightarrow A \times B$$

defined by

$$\varphi(x) = [a, \varphi_a(x)] \text{ if } x \in B_a$$

is an isomorphism of the cell algebra  $\mathcal{A}(\mathcal{B})$  onto the direct product  $\mathcal{A} \times \mathcal{B}$ . Evidently  $\varphi$  is a bijection. If  $f$  is a  $k$ -ary operational symbol,  $x_1 \in B_{a_1}, \dots, x_k \in B_{a_k}$ ,  $f^{(A)}(a_1, \dots, a_k) = a$  then

$$\begin{aligned} \varphi(f^{(M)}(x_1, \dots, x_k)) &= [a, \varphi_a(f^{(M)}(x_1, \dots, x_k))] \\ &= [f^{(A)}(a_1, \dots, a_k), \varphi_a(f^{(B_a)}(\varphi_{a_1,a}(x_1), \dots, \varphi_{a_k,a}(x_k)))] \\ &= [f^{(A)}(a_1, \dots, a_k), f^{(B)}(\varphi_a(\varphi_{a_1,a}(x_1)), \dots, \varphi_a(\varphi_{a_k,a}(x_k)))] \\ &= [f^{(A)}(a_1, \dots, a_k), f^{(B)}(\varphi_{a_1}(x_1), \dots, \varphi_{a_k}(x_k))] \\ &= f^{(A \times B)}([a_1, \varphi_{a_1}(x_1)], \dots, [a_k, \varphi_{a_k}(x_k)]) \\ &= f^{(A \times B)}(\varphi(x_1), \dots, \varphi(x_k)). \end{aligned}$$

□

**Theorem 7.** Let a basic algebra  $\mathcal{A}$  and every cell  $\mathcal{B}_a, a \in A$ , be algebras of the type  $\tau$ . Let  $\mathcal{S}^{(F)}$  be a family of homomorphisms  $\varphi_{a,b}: B_a \rightarrow B_b$  such that (2.1), (2.6a) and (2.7a) hold and moreover  $\mathcal{S}^{(f)} = \mathcal{S}^{(g)}$  for all operations  $f, g \in F$  (i.e. the family  $\mathcal{S}^{(F)}$  does not depend on operations). If an identity

$$p(x_1, \dots, x_n) = q(x_1, \dots, x_n)$$

holds in  $\mathcal{A}$  and also in each  $\mathcal{B}_a$  then it holds in the cell algebra  $\mathcal{A}(\mathcal{B})$ , too.

**Proof.** First, we will show that

$$p^{(M)}(x_1, \dots, x_n) = p^{(B_a)}(\varphi_{a_1,a}(x_1), \dots, \varphi_{a_n,a}(x_n))$$

if  $p(x_1, \dots, x_n)$  is an arbitrary term of the type  $\tau$ ,  $x_1 \in B_{a_1}, \dots, x_n \in B_{a_n}$  and  $p^{(A)}(a_1, \dots, a_n) = a$ . We prove it by induction with respect to the number of operational symbols in the term  $p(x_1, \dots, x_n)$ .

Let

$$p(x_1, \dots, x_n) = f(p_1(x_1, \dots, x_n), \dots, p_k(x_1, \dots, x_n))$$

where  $f$  is a  $k$ -ary operational symbol. Let  $x_1 \in B_{a_1}, \dots, x_n \in B_{a_n}$ ,  $p_i^{(A)}(a_1, \dots, a_n) = b_i$  for  $i = 1, 2, \dots, k$ . By induction hypothesis we have

$$(2.8) \quad p_i^{(M)}(x_1, \dots, x_n) = p_i^{(B_{b_i})}(\varphi_{a_1, b_i}(x_1), \dots, \varphi_{a_n, b_i}(x_n))$$

for  $i = 1, 2, \dots, k$ . Let  $p^{(A)}(a_1, \dots, a_n) = a$ , i.e.  $f^{(A)}(b_1, \dots, b_k) = a$ . We get

$$\begin{aligned} p^{(M)}(x_1, \dots, x_n) &= f^{(B_a)}(\varphi_{b_1, a}(p_1^{(B_{b_1})}(x_1, \dots, x_n)), \dots, \varphi_{b_k, a}(p_k^{(B_{b_k})}(x_1, \dots, x_n))) \\ &= f^{(B_a)}(\varphi_{b_1, a}(p_1^{(B_{b_1})}(\varphi_{a_1, b_1}(x_1), \dots, \varphi_{a_n, b_1}(x_n))), \dots, \\ &\quad \varphi_{b_k, a}(p_k^{(B_{b_k})}(\varphi_{a_1, b_k}(x_1), \dots, \varphi_{a_n, b_k}(x_n)))) \\ &= f^{(B_a)}(p_1^{(B_a)}(\varphi_{b_1, a}(\varphi_{a_1, b_1}(x_1)), \dots, \varphi_{b_1, a}(\varphi_{a_n, b_1}(x_n))), \dots, \\ &\quad p_k^{(B_a)}(\varphi_{b_k, a}(\varphi_{a_1, b_k}(x_1)), \dots, \varphi_{b_k, a}(\varphi_{a_n, b_k}(x_n)))) \\ &= f^{(B_a)}(p_1^{(B_a)}(\varphi_{a_1, a}(x_1), \dots, \varphi_{a_n, a}(x_n)), \dots, \\ &\quad p_k^{(B_a)}(\varphi_{a_1, a}(x_1), \dots, \varphi_{a_n, a}(x_n))) \\ &= p^{(B_a)}(\varphi_{a_1, a}(x_1), \dots, \varphi_{a_n, a}(x_n)). \end{aligned}$$

Therefore under the above mentioned assumptions we obtain

$$\begin{aligned} p^{(M)}(x_1, \dots, x_n) &= p^{(B_a)}(\varphi_{a_1, a}(x_1), \dots, \varphi_{a_n, a}(x_n)) \\ &= q^{(B_a)}(\varphi_{a_1, a}(x_1), \dots, \varphi_{a_n, a}(x_n)) = q^{(M)}(x_1, \dots, x_n). \end{aligned}$$

□

**Corollary 8.** *Let a basic algebra  $\mathcal{A} = (A, f)$  and every cell  $\mathcal{B}_a = (B_a, f)$ ,  $a \in A$ , be groupoids. Let  $\mathcal{S}^{(f)}$  be a family of homomorphisms  $\varphi_{a,b}: B_a \rightarrow B_b$  such that (2.1), (2.6a) and (2.7a) hold. If an identity*

$$p(x_1, \dots, x_n) = q(x_1, \dots, x_n)$$

holds in  $\mathcal{A}$  and also in each  $\mathcal{B}_a$  then it holds in the groupoid  $\mathcal{A}(\mathcal{B})$ , too.

### 3. $N$ -SKEW LATTICES

In this section we will give a characterization of  $N$ -skew lattices using the construction of cell algebras.

An algebra  $(L, \wedge, \vee)$  of the type  $(2, 2)$  is called a noncommutative lattice if the binary operations  $\wedge$  and  $\vee$  are associative, idempotent and satisfy some absorption identities.

M. D. Gerhards has investigated noncommutative lattices satisfying the identities

$$(3.1) \quad x \wedge (x \vee y) = x \quad \& \quad (y \wedge x) \vee x = x$$

and

$$(3.2) \quad (z \vee y \vee x) \wedge (x \vee y) = y \vee x \quad \& \quad (y \wedge x) \vee (x \wedge y \wedge z) = x \wedge y$$

which are called prelattices (fastverbands). In [3] it is shown that every prelattice is the direct product of a lattice and a nest. In [2] M. D. Gerhards characterized prelattices as relational structures. Recall that a nest is an algebra  $(L, \wedge, \vee)$  of the type  $(2, 2)$  satisfying the identities

$$(3.3) \quad x \wedge y = x \quad \& \quad y \vee x = x.$$

V. Slavík investigated prelattices in [11] and varieties of prelattices in [12].

M. Yamada and N. Kimura in [13] investigated idempotent semigroups (bands) satisfying the identity  $xyz = xzy$  and showed that they are semilattices of trivial algebras (seminests). In [6] A. Haviar introduced a larger class of noncommutative lattices, so-called  $N$ -skew lattices, which can be characterized as relational systems, too.  $N$ -skew lattices are noncommutative lattices satisfying the identity (3.1) and the identities

$$(3.4) \quad x \wedge (y \wedge z) = x \wedge (z \wedge y) \quad \& \quad (z \vee y) \vee x = (y \vee z) \vee x.$$

**Theorem 9.** *An algebra  $(L, \wedge, \vee)$  of the type  $(2, 2)$  is an  $N$ -skew lattice if and only if  $(L, \wedge, \vee)$  is isomorphic to a cell algebra  $\mathcal{A}(\mathcal{B})$  in which the basic algebra  $\mathcal{A}$  is a lattice, every cell  $\mathcal{B}_a$ ,  $a \in A$ , is a nest and the system of mappings  $\varphi_{b,a}^{(\wedge)}: B_b \rightarrow B_a$  and  $\varphi_{a,b}^{(\vee)}: B_a \rightarrow B_b$  for each  $a \leq b$ ,  $a, b \in A$ , satisfies the conditions (2.6a) and (2.7a).*

*Proof.* a) Let  $\mathcal{L} = (L, \wedge, \vee)$  be an  $N$ -skew lattice. We define a relation  $\Theta$  on  $L$  as follows:

$$a \Theta b \iff a \wedge b = a \quad \& \quad b \wedge a = b.$$

The relation  $\Theta$  is a congruence relation of  $\mathcal{L}$ , the algebra  $\mathcal{L}/\Theta$  is a lattice (a modification of  $\mathcal{L}$  in the variety of lattices) and every block  $a\Theta = B_a$  is a nest (see [11]).

For  $a\Theta \leq b\Theta$  we define mappings

$$\varphi_{b\Theta, a\Theta}^{(\wedge)}: b\Theta \rightarrow a\Theta \quad \text{and} \quad \varphi_{a\Theta, b\Theta}^{(\vee)}: a\Theta \rightarrow b\Theta$$

by

- (i)  $\forall x \in b\Theta \quad \varphi_{b\Theta, a\Theta}^{(\wedge)}(x) = x \wedge a,$   
(ii)  $\forall x \in a\Theta \quad \varphi_{a\Theta, b\Theta}^{(\vee)}(x) = b \vee x.$

Let  $a_1 \in a\Theta$  and  $b_1 \in b\Theta$ . Since  $x \wedge a_1 = x \wedge a_1 \wedge a = x \wedge a \wedge a_1 = x \wedge a$  (by (3.4)) and similarly  $b_1 \vee x = b \vee x$ , the mappings  $\varphi_{b\Theta, a\Theta}^{(\wedge)}$  and  $\varphi_{a\Theta, b\Theta}^{(\vee)}$  are defined correctly. (Moreover, the mappings  $\varphi_{b\Theta, a\Theta}^{(\wedge)}$  and  $\varphi_{a\Theta, b\Theta}^{(\vee)}$  are homomorphisms because  $a\Theta$  and  $b\Theta$  are nests.)

If  $a\Theta \leq b\Theta \leq c\Theta$  then

$$\varphi_{b\Theta, a\Theta}^{(\wedge)}(\varphi_{c\Theta, b\Theta}^{(\wedge)}(x)) = \varphi_{b\Theta, a\Theta}^{(\wedge)}(x \wedge b) = (x \wedge b) \wedge a = x \wedge (a \wedge b) = x \wedge a = \varphi_{c\Theta, a\Theta}^{(\wedge)}(x)$$

and

$$\varphi_{a\Theta, a\Theta}^{(\wedge)}(x) = x \wedge a = x$$

and dually for  $\varphi_{a\Theta, b\Theta}^{(\vee)}$ , hence the mappings  $\varphi_{b\Theta, a\Theta}^{(\wedge)}$  and  $\varphi_{a\Theta, b\Theta}^{(\vee)}$  satisfy the conditions (2.6a) and (2.7a).

Let  $\mathcal{S}^{(\wedge)}$  and  $\mathcal{S}^{(\vee)}$  be systems of mappings

$$\mathcal{S}^{(\wedge)} = \{\varphi_{b\Theta, a\Theta}^{(\wedge)}; a\Theta \leq b\Theta\}, \quad \mathcal{S}^{(\vee)} = \{\varphi_{a\Theta, b\Theta}^{(\vee)}; a\Theta \leq b\Theta\}.$$

Denote by  $\sqcap$  and  $\sqcup$  the operations of a cell algebra with the basic algebra  $\mathcal{L}/\Theta$ , cells  $B_a = a\Theta$ ,  $a\Theta \in L/\Theta$  and systems of mappings  $\mathcal{S}^{(\wedge)}$ ,  $\mathcal{S}^{(\vee)}$ . For any elements  $x, y \in \bigcup_{a \in L} B_a = M$  we get

$$x \sqcap y = \varphi_{x\Theta, x \wedge y\Theta}^{(\wedge)}(x) \wedge \varphi_{y\Theta, x \wedge y\Theta}^{(\wedge)}(y) = (x \wedge (x \wedge y)) \wedge (y \wedge (x \wedge y)) = x \wedge y$$

and dually  $x \sqcup y = x \vee y$ .

b) Conversely, let  $\mathcal{A}(\mathcal{B})$  be a cell algebra for which the basic algebra  $\mathcal{A}$  is a lattice  $(A, \wedge, \vee)$ , let each cell  $B_a$ ,  $a \in A$ , be a nest and for every  $a \leq b$  let the mappings

$$\varphi_{b, a}^{(\wedge)}: B_b \rightarrow B_a, \quad \varphi_{a, b}^{(\vee)}: B_a \rightarrow B_b$$

satisfy the conditions (2.6a) and (2.7a).

The operations of the basic algebra as well as those of every cell are associative, idempotent and the mappings  $\varphi_{b,a}^{(\wedge)}, \varphi_{a,b}^{(\vee)}$  are homomorphisms, hence by Theorem 5 the operations of the cell algebra  $\mathcal{A}(\mathcal{B})$  are also associative and idempotent. By Corollary 8 the operations of the cell algebra  $\mathcal{A}(\mathcal{B})$  satisfy the identity (3.4), too.

For any elements  $x \in B_a, y \in B_b$  we get

$$\begin{aligned} x \sqcap (x \sqcup y) &= x \sqcap (\varphi_{a,a \vee b}^{(\vee)}(x) \vee \varphi_{b,a \vee b}^{(\vee)}(y)) = x \sqcap \varphi_{b,a \vee b}^{(\vee)}(y) \\ &= \varphi_{a,a \wedge (a \vee b)}^{(\wedge)}(x) \wedge \varphi_{a \vee b, a \wedge (a \vee b)}^{(\wedge)}(\varphi_{b,a \vee b}^{(\vee)}(y)) = \varphi_{a,a}^{(\wedge)}(x) = x \end{aligned}$$

and dually  $(y \sqcap x) \sqcup x = x$ . □

Now let us assume that the basic algebra of a cell algebra is a distributive lattice. A lattice is distributive if it satisfies the identity  $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$  which is satisfied in every nest, too.

A slight change in the proof of Theorem 9 enables us to show the next statement.

**Theorem 10.** *An algebra  $(L, \wedge, \vee)$  of the type  $(2, 2)$  is a distributive  $N$ -skew lattice if and only if  $(L, \wedge, \vee)$  is isomorphic to a cell algebra  $\mathcal{A}(\mathcal{B})$  in which the basic algebra  $\mathcal{A}$  is a distributive lattice, each cell  $\mathcal{B}_a, a \in A$ , is a nest and the system of mappings  $\varphi_{b,a}^{(\wedge)}: B_b \rightarrow B_a, \varphi_{a,b}^{(\vee)}: B_a \rightarrow B_b$  for every  $a \leq b, a, b \in A$ , satisfies the conditions (2.6a) and (2.7a).*

### References

- [1] *S. Burris, H. P. Sankappanavar:* A Course in Universal Algebra. Springer, New York, 1981.
- [2] *M. D. Gerhardt:* Schrägverbände und Quasiordnungen. Math. Ann. 181 (1969), 65–73.
- [3] *M. D. Gerhardt:* Zerlegungshomomorphismen in Schrägverbänden. Arch. Math. 21 (1970), 116–122.
- [4] *E. Graczyńska:* On the sums of double systems of some universal algebras. Bull. de L'Acad. Polon. Sci. Math. Astronom. Phys. 23 (1975), 509–513.
- [5] *G. Grätzer:* General Lattice Theory. Academic-Verlag, Berlin, 1978.
- [6] *A. Haviar:* N-Schrägverbände und Quasiordnungen. Matematický časopis 23 (1973), 240–248.
- [7] *T. Hecht:* Constructions of non-commutative algebras. Lectures in Universal Algebra (eds. L. Szabó, A. Szendrei), North-Holland, Amsterdam (1986), 177–187.
- [8] *J. Leech:* Recent developments in the theory of skew lattices. Semigroup Forum 53 (1996), 7–24.
- [9] *J. Plonka:* On a method of construction of abstract algebras. Fund. Math. 61 (1967), 183–189.
- [10] *J. Plonka:* Representations of algebras from varieties defined by some regular identities with nullary operation symbols. Algebra Universalis 33 (1995), 441–457.
- [11] *V. Slavík:* On skew lattices I. Comment. Math. Univ. Carolin. 14 (1973), 73–85.
- [12] *V. Slavík:* On skew lattices II. Comment. Math. Univ. Carolin. 14 (1973), 493–506.

- [13] *M. Yamada, N. Kimura*: Note on idempotent semigroups II. Proc. Japan. Acad. 34 (1958), 110–112.

*Authors' addresses:* *Alfonz Haviar*, Department of Mathematics, Faculty of Natural Sciences, Matej Bel University, Tajovského 40, 974 01 Banská Bystrica, Slovakia, e-mail: [haviar@fpv.umb.sk](mailto:haviar@fpv.umb.sk); *Gabriela Monoszová*, Department of Mathematics, Faculty of Natural Sciences, Matej Bel University, Tajovského 40, 974 01 Banská Bystrica, Slovakia, e-mail: [monosz@fpv.umb.sk](mailto:monosz@fpv.umb.sk).