

Applications of Mathematics

Wiktor Oktaba

Densities of determinant ratios, their moments and some simultaneous confidence intervals in the multivariate Gauss-Markoff model

Applications of Mathematics, Vol. 40 (1995), No. 1, 47--54

Persistent URL: <http://dml.cz/dmlcz/134277>

Terms of use:

© Institute of Mathematics AS CR, 1995

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

DENSITIES OF DETERMINANT RATIOS, THEIR MOMENTS
AND SOME SIMULTANEOUS CONFIDENCE INTERVALS
IN THE MULTIVARIATE GAUSS-MARKOFF MODEL

WIKTOR OKTABA, Lublin

(Received May 10, 1993)

Summary. The following three results for the general multivariate Gauss-Markoff model with a singular covariance matrix are given or indicated. 1° determinant ratios as products of independent chi-square distributions, 2° moments for the determinants and 3° the method of obtaining approximate densities of the determinants.

Keywords: products of independent chi-squares, multivariate general linear Gauss-Markoff model, Wishart distribution, singular covariance matrix, set of linear parametric functions, moment of determinant ratio, approximate simultaneous confidence interval

AMS classification: 62H10

1. INTRODUCTION

We are interested in the multivariate general linear Gauss-Markoff model MGM

$$(1.1) \quad (U, XB, \sigma^2 \Sigma \otimes V),$$

with a known matrix $V \geq 0$, where \otimes is the symbol of the Kronecker product of matrices, B is a matrix of unknown fixed parameters, X is a given known design matrix, U is a random matrix of observations with the expected value $\varepsilon(U) = XB$ and with a fixed nonsingular matrix Σ , σ^2 is an unknown positive scalar. Let $T = V + XMX'$ where $M = M'$ is an arbitrary matrix such that $R(X) \subset R(T)$. The symbol $R(X)$ is used to denote the vector space spanned on the columns of the matrix X .

Let $A \sim W_p(n, \Sigma)$ denote that a $p \times p$ random matrix A is distributed as the p -dimensional Wishart distribution with n degrees of freedom and a covariance matrix Σ . Determinant ratios as products of independent chi-square distributions under

the assumption $V^*W_p(n, \Sigma)$, where V^* and $\Sigma > 0$ are $p \times p$ matrices, have been given by Srivastava and Khatri [9] p. 82,3.38; Rao [7] p. 540, X; Muirhead [4] p. 447.

By applying this result and that given by Oktaba and Kieloch [6] for MGM three theorems are established (Sec. 2).

Moments of determinant ratios $|V^*||\Sigma|^{-1}$ under the assumption $V^*W_p(n, \Sigma)$ can be found in Anderson [1] or Rao ([7], p. 540). Their forms in the case of the MGM model are deduced by using results of Oktaba and Kieloch (loc. cit) (Sec. 3).

In very many papers and books the confidence intervals are given, e.g. S.N. Roy [9], Anderson [1]; Browkovicz and Linnik [2]; Muirhead [4]; Rao [7], Srivastava and Khatri [9]; Wani and Kabe [10], and some others. Using the procedure of Srivastava and Khatri, the method of approximate density of determinant rations is suggested. In this way we get a generalization of Wani and Kabe results [10] from the standard model $(U, XB, \Sigma \otimes I)$ to the MGM model. It creates a possibility for constructing an approximate simultaneous confidence region for a pair: a set of parametric estimable functions L^*B and the determinant $|\sigma^2\Sigma|$.

2. DENSITIES OF DETERMINANT RATIOS IN MGM MODEL

The following theorem (Srivastava and Khatri [9] pp. 82, 3.38; Rao [7], p. 540, X; Muirhead [4], p. 447) will be used in this section.

Theorem 2.1. *Let us consider a $p \times p$ matrix $V^*W_p(n, \Sigma)$, where Σ is $p \times p$ matrix, $p \times p$ matrix $\Sigma > 0$. Then the determinant*

$$(2.1) \quad |W| = \frac{|V^*|}{|\Sigma|} = z_1 \cdot z_2 \dots z_p = \chi_n^2 \chi_{n-1}^2 \dots \chi_{n-p+1}^2,$$

where $z_i, i = 1, \dots, p$ are independently distributed as χ_{n-i+1}^2 with $n - i + 1$ degrees of freedom.

Let $r(A)$ denote the rank of the matrix A . The symbol $(A:B)$ is reserved for the matrix involving two submatrices A and B .

The additivity of central p -dimensional Wishart distributions is stated in the following theorem.

Theorem 2.2 (Srivastava and Khatri [9] p. 82; Muirhead [4] p. 446). *If $p \times p$ matrices V_1^*, \dots, V_r^* are stochastically independent and $V_i^*W_p(\nu_i, \Sigma), i = 1, \dots, r$, then*

$$(2.2) \quad \Sigma_{i=1}^r V_i^*W_p \left(\sum_{i=1}^n \nu_i, \Sigma \right).$$

Theorems 2.3, 2.4 and 2.5 of Oktaba and Kieloch [6] will be applied in this section.

Let L^*B be a set of linear estimable parametric functions, where L^* is an $a \times m$ matrix. The following quadratic forms are of interest:

$$(2.3) \quad S_H = (L^*\hat{B} - \psi)'L^-(L^*\hat{B} - \psi) = (L^*\hat{B} - \psi)'(L^*C_4L^*)^-(L^*\hat{B} - \psi),$$

$$(2.4) \quad S_e = U'C_1U,$$

where

$$(2.5) \quad \hat{B} = (X'T^-X)^-X'T^-U = C_2'U = C_3U,$$

$$(2.6) \quad \begin{cases} C_1 = T^- - T^-X(X'T^-X)^-X'T^- = T^-(I - XC_3) \\ C_2' = C_3 = (X'T^-X)^-X'T^-, \end{cases}$$

$$(2.7) \quad L = (L^*C_3)V(L^*C_3)' = L^*C_4L^*$$

(Oktaba [5], (2.1), 179).

The symbol L^- is reserved for any choice of the g -inverse, i.e. the following relation holds:

$$(2.8) \quad LL^-L = L.$$

We recall that

$$\begin{pmatrix} V & X \\ X' & 0 \end{pmatrix}^- = \begin{pmatrix} C_1 & C_2 \\ C_3 & -C_4 \end{pmatrix}.$$

Theorem 2.3 (Oktaba and Kieloch [6]). *If*

$$(2.9) \quad U \sim N_{np}(XB, \sigma^2\Sigma \otimes V),$$

where $N_{np}(\cdot, \cdot)$ denotes an np -variate normal distribution N_{np} with parameters defined in Introduction, L^*B is a set of estimable linear combinations of parameters and the hypothesis $L^*B = \psi$ is true, then

$$(2.10) \quad S_H \sim W_p[r(L), \sigma^2\Sigma].$$

Theorem 2.4 (Oktaba and Kieloch [6]). *Subject to the assumption (2.9) we have*

$$(2.11) \quad S_e \sim W_p[r(\cdot; X) - r(X), \sigma^2\Sigma],$$

where S_e is defined in (2.4)

Theorem 2.5 (Oktaba and Kieloch [6]). *If the assumptions of Theorems 2.3 and 2.4 concerning the matrices S_H and S_e are fulfilled, then S_H and S_e are stochastically independent.*

As a consequence of the above five theorems we almost immediately obtain the following three theorems.

Theorem 2.6. *In the model (1.1) let the $p \times p$ matrix S_e satisfy*

$$(2.12) \quad S_e = U' C_1 U^{-1} W_p(\nu_e, \sigma^2 \Sigma),$$

where the $p \times p$ matrix Σ and the scalar σ^2 are positive,

$$(2.13) \quad \nu_e = r(V: X) - r(X),$$

C_1 is given as in (2.6). Then we have

$$(2.14) \quad \frac{|S_e|}{|\sigma^2 \Sigma|} = \prod_{i=1}^p \xi_i,$$

where ξ_i are independently distributed as $\chi_{\nu_e - i + 1}^2$, $i = 1, \dots, p$; with $\nu_e - i + 1$ degrees of freedom.

Proof. We observe (2.11) in model (1.1), where S_e and ν_e are as in (2.12) and (2.13). Replacing in Theorem 2.1 V^* , Σ , n and z_i by S_e , $\sigma^2 \Sigma$, ν_e and ξ_i , respectively, we get (2.14). \square

Theorem 2.7. *In the model (1.1) let the $p \times p$ matrix S_H (Oktaba and Kieloch [6]) in (2.3) satisfy*

$$(2.15) \quad S_H^{-1} W_p(\nu_H, \sigma^2 \Sigma)$$

under the true hypothesis

$$(2.16) \quad H_0: L^* B = \psi,$$

where the $p \times p$ matrix Σ and the scalar σ^2 are positive,

$$(2.17) \quad C_4 = (X', T^{-1} X)^{-1} - M, \quad \nu_H = r(L),$$

L , \tilde{B} and C_3 are as in (2.7), (2.5) and (2.6), respectively; the matrices T and M are defined in Introduction.

Then

$$(2.18) \quad \frac{|S_H|}{|\sigma^2 \Sigma|} \prod_{i=1}^p w_i,$$

where w_i are independently distributed as $\chi_{\nu_H - i + 1}^2$ with $\nu_H - i + 1$ degrees of freedom, $i = 1, \dots, p$.

Proof is analogous to that of Theorem 2.6. \square

Theorem 2.8. Under the true hypothesis (2.16) in the model (1.1) let the $p \times p$ matrix S_y satisfy

$$(2.19) \quad S_y = S_H + S_e \sim W_p(\nu_H + \nu_e, \sigma^2 \Sigma),$$

where the $p \times p$ matrix Σ and scalar σ^2 are positive. Then

$$(2.20) \quad \frac{|S_y|}{|\sigma^2 \Sigma|} \prod_{i=1}^p m_i,$$

where m_i are independently distributed as $\chi_{\nu_y - i + 1}^2$ with $\nu_y - i + 1 = \nu_h + \nu_e - i + 1$ degrees of freedom, $i = 1, \dots, p$.

Proof. By virtue of (2.10), (2.12) (Oktaba and Kieloch [6]) and the additive property of Wishart distributions given in Theorem 2.2 we conclude that (2.19) holds. Applying Theorem 2.1 where we put $V^* = S_e + S_H = S_y$ and replace n by $\nu_y = \nu_e + \nu_H = r(V^*:X) - r(X) + r(L)$, we obtain (2.20). \square

A particular case of the model (1.1) is the multivariate standard model

$$(2.21) \quad (U, 1 \cdot \mu', \Sigma \otimes I_n),$$

where U is a $n \times p$ matrix of observations, 1 —the vector $n \times 1$ involving unities, $\mu' = (\mu_1, \dots, \mu_p)$, the $p \times p$ matrix Σ is positive,

$$(2.22) \quad S_e = U' \left(I - \frac{1}{n} 1 1' \right) U = V^* \sim W_p(n-1, \Sigma)$$

and

$$(2.23) \quad |W| = \frac{|S_e|}{|\Sigma|} = \chi_{n-p}^2 \cdots \chi_{n-2}^2 \cdot \chi_{n-1}^2,$$

where $W = S_e \Sigma^{-1}$ (Anderson [1], (17), p. 171).

To prove (2.23) it is sufficient to set in (1.1): $X = 1$, $B = \mu'$, $V = I_n$, $\sigma^2 = 1$, $\nu_e = n - 1$. From (2.14) we get (2.23).

3. MOMENTS OF DETERMINANT RATIOS $|S_e|/(|\sigma^2\Sigma|)$ AND $|S_H|/(|\sigma^2\Sigma|)$
IN THE MGM MODEL WITH SINGULAR COVARIANCE MATRIX

For a general p a closed form density of the product of p independent central chi-squares is not available. It is possible to give the h -th moment of the determinant $|W| = |V^*||\Sigma|^{-1}$, where

$$(3.1) \quad V^* \sim W_p(n, \Sigma).$$

Its formula is as follows (Anderson [1]; Srivastava, Khatri [9], p. 83:

$$(3.2) \quad E(|W|^h) = E\left(\frac{|V^*|}{|\Sigma|}\right)^h = 2^{ph} \frac{\Gamma_p\left(\frac{n}{2} + h\right)}{\Gamma_p\left(\frac{n}{2}\right)},$$

where n denotes the number of degrees of freedom of V^* in (3.1),

$$(3.3) \quad \Gamma_p\left(\frac{n}{2}\right) = \Pi^{\frac{1}{2}p(p-1)} \prod_{i=1}^p \Gamma\left(\frac{n-i+1}{2}\right)$$

is the multivariate gamma function (James, [3], p. 483),

$$(3.4) \quad \Gamma(r) = \int_0^\infty y^{r-1} e^{-y} dy$$

is the complete gamma function.

Theorem 3.1. *The h -th moments of $|S_e|/(|\sigma^2\Sigma|)^{-1}$, $|S_H|/(|\sigma^2\Sigma|)^{-1}$ and $|S_y| \times (|\sigma^2\Sigma|)^{-1}$ in the MGM model of the form (1.1) with a singular covariance matrix are*

$$(3.5) \quad E\left(\frac{|S_e|}{|\sigma^2\Sigma|}\right)^h = 2^{ph} \cdot \frac{\Gamma_p\left(\frac{\nu_e}{2} + h\right)}{\Gamma_p\left(\frac{\nu_e}{2}\right)},$$

where S_e and ν_e are given as in (2.12) and (2.13).

$$(3.6) \quad E\left(\frac{|S_H|}{|\sigma^2\Sigma|}\right)^h = 2^{ph} \cdot \frac{\Gamma_p\left(\frac{\nu_H}{2} + h\right)}{\Gamma_p\left(\frac{\nu_H}{2}\right)},$$

where S_H and ν_H are given as in (2.3) and (2.17).

Moreover

$$(3.7) \quad E\left(\frac{|S_y|}{|\sigma^2\Sigma|}\right)^h = 2^{ph} \cdot \frac{\Gamma_p\left(\frac{\nu_y}{2} + h\right)}{\Gamma_p\left(\frac{\nu_y}{2}\right)},$$

where

$$(3.8) \quad \nu_y = \nu_e + \nu_H, \quad S_y = S_e + S_H.$$

Proof. Results of Oktaba and Kieloch [6] yield that S_e and S_H are distributed as $W_p(\nu_e, \sigma^2\Sigma)$ and $W_p(\nu_H, \sigma^2\Sigma)$, respectively. Formula (3.2), in which we replace V^* , n and Σ by S_e , $\nu_e = r(V^*X) - r(X)$ and $\sigma^2\Sigma$ respectively implies (3.5). Similarly, replacing V^* , n and Σ by S_H , $\nu_H = r(L)$ and $\sigma^2\Sigma$, we get (3.6). As a consequence of additivity given in Theorem 2.2 we state that $S_y = S_e + S_H \sim W_p(\nu_e + \nu_H, \sigma^2\Sigma)$, so by formula (3.2) we have (3.7). \square

4. SIMULTANEOUS CONFIDENCE INTERVAL FOR L^*B AND $|\sigma^2\Sigma|$ IN MULTIVARIATE MODEL (1.1) WITH SINGULAR COVARIANCE MATRIX

Let us consider the problem of constructing a simultaneous confidence region in model (1.1) for a set of estimable parametric functions L^*B and a determinant $|\sigma^2\Sigma|$. Applying Oktaba's and Kieloch's [6] results we state that in (1.1) under the assumption

$$(4.1) \quad U \sim N_{np}(XB, \sigma^2\Sigma \otimes V)$$

and the estimable true hypothesis

$$(4.2) \quad H_0: L^*B = \psi,$$

the matrices S_H and S_e in (2.3) and (2.4), respectively, are independently distributed.

From (2.20) we know that the density of the random variable

$$(4.3) \quad |W_y| = \frac{|S_y|}{|\sigma^2\Sigma|}$$

is the same as the density of p chi-squares independently distributed (cf. Wani and Kabe [10], p. 18).

The density of the random variable $|W_y|$ can be used to construct simultaneous confidence regions for the pair $(L^*B, |\sigma^2\Sigma|)$ under the assumption of estimability of L^*B in (4.2) (Wani and Kabe, [10], p. 18).

Remark 4.1. The moments of $|W_e| = \frac{|S_e|}{|\sigma^2 \Sigma|}$, $|W_H| = \frac{|S_H|}{|\sigma^2 \Sigma|}$ and $|W_y| = \frac{|S_y|}{|\sigma^2 \Sigma|}$ are given in (3.5), (3.6) and (3.7), so under the assumptions that S_e , S_H and S_y are distributed as $W_p(\nu_e, \sigma^2 \Sigma)$, $W_p(\nu_H, \sigma^2 \Sigma)$ and $W_p(\nu_y, \sigma^2 \Sigma)$ we can obtain an approximate density of variates $|W_e|$, $|W_H|$ and $|W_y|$ applying the method which is described by Srivastava and Khatri ([9], chapter 6.3). It is known that density is determined uniquely by the moments (loc. cit. p. 176).

References

- [1] *T.W. Anderson*: An Introduction to Multivariate Statistical Analysis. J. Wiley, New York, 1958.
- [2] *G.N. Browkovicz, J.W. Linnik*: A note to confidence estimation by least squares method. *Teoria verojatnostej i jejo primeneniya* 8 (1963), no. 2, 176. (In Russian.)
- [3] *A.T. James*: Distributions of matrix variates and latent roots derived from normal samples. *Ann. Math. Stat.* 35 (1964), no. 2, 475–501.
- [4] *R.J. Muirhead*: Aspects of Multivariate Statistical Theory. J. Wiley, New York, 1982.
- [5] *W. Oktaba*: Estimation of the parametric functions in the multivariate general Gauss-Markoff model. XV Colloquium Metodol. Agrobiom., Pol. Acad. Sci., Warsaw 1985. pp. 178–182. (In Polish.)
- [6] *W. Oktaba, A. Kieloch*: Wishart distributions in the multivariate Gauss-Markoff model with singular covariance matrix. *Appl. Math.* 38 (1993), 61–66.
- [7] *C.R. Rao*: Linear Statistical Inference and Its Applications, Sec. Ed. J. Wiley, New York, 1973.
- [8] *S.N. Roy*: Some Aspects of Multivariate Analysis. New York, 1957.
- [9] *M.S. Srivastava, G.G. Khatri*: An Introduction to Multivariate Statistics. Elsevier North Holland, Inc., New York, 1979.
- [10] *J.K. Wani, D.G. Kabe*: Generalized inverse and confidence estimation by least squares method. *Trabajos De. Estadistica Y de Investigation Operativa* 21. Cuadernos 1 y 2 Afio, Iberica, Tarragona 1970, vol. 34, Madrid, 1968, pp. 115–120.

Author's address: *Wiktor Oktaba*, Institute of Applied Mathematics, Academy of Agriculture, Akademicka 13, P.O.Box 325, 20-950 Lublin 1, Poland.