

Applications of Mathematics

Qun Lin; Shu Hua Zhang

A direct global superconvergence analysis for Sobolev and viscoelasticity type equations

Applications of Mathematics, Vol. 42 (1997), No. 1, 23--34

Persistent URL: <http://dml.cz/dmlcz/134342>

Terms of use:

© Institute of Mathematics AS CR, 1997

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

A DIRECT GLOBAL SUPERCONVERGENCE ANALYSIS
FOR SOBOLEV AND VISCOELASTICITY TYPE EQUATIONS

QUN LIN, SHUHUA ZHANG, Beijing

(Received June 16, 1995)

Abstract. In this paper we study the finite element approximations to the Sobolev and viscoelasticity type equations and present a direct analysis for global superconvergence for these problems, without using Ritz projection or its modified forms.

Keywords: Sobolev and viscoelasticity type equations, global superconvergence, direct analysis

MSC 2000: 65B05, 65N30

1. INTRODUCTION

Let Ω be a rectangular domain. In order to explain our superconvergence analysis for FEMs succinctly, we only consider the simple Sobolev type equation

$$(1.1) \quad \begin{cases} -\Delta u_t - \Delta u = f & \text{in } \Omega \times (0, T], \\ u = 0 & \text{on } \partial\Omega \times (0, T], \\ u(x, 0) = v & \text{in } \Omega, \end{cases}$$

and a viscoelasticity type equation

$$(1.2) \quad \begin{cases} u_{tt} - \Delta u_t - \Delta u = f & \text{in } \Omega \times (0, T], \\ u = 0 & \text{on } \partial\Omega \times (0, T], \\ u(x, 0) = v, u_t(x, 0) = w & \text{in } \Omega. \end{cases}$$

In (1.1) and (1.2), f , v and w are sufficiently smooth functions.

The problems (1.1) and (1.2) can arise from many physical processes. The numerical approximations to the solutions of these problems have been investigated

by Ewing [2, 3], Ford [4], Ford and Ting [5, 6], and Wahlbin [14]. Also Arnold, Douglas and Thomée [1] and Nakao [13] have considered Galerkin approximations to the solution of the problem (1.1) in a single space dimension with periodic boundary conditions. L^2 -error estimates and the interior pointwise superconvergence results have been derived by these authors. In particular, some further investigations of the finite element methods for the problems (1.1) and (1.2) have been carried out by Lin, Thomée and Wahlbin [12].

According to the conventional error analysis for FEMs of the time-dependent problems, either the Ritz projection initiated by Wheeler [15] or its modified forms e.g. the so-called Ritz-Volterra projection introduced by Lin et al. [12], have to be used as transitional tools. However, here we will use a new analysis in [7], i.e. an analysis for the “short side” in the FE-right triangle plus the sharp integral estimates of the “hypotenuse”, instead of using Ritz projection or its modified forms, to gain the global superconvergence for the problems (1.1) and (1.2) by an interpolation postprocessing technique, rather than the interior pointwise superconvergence by means of the average technique with which numerical analysts are familiar. Our analysis sharpens the results and shortens the proofs of error estimates that appeared in the previous literature under the rectangular mesh assumption.

2. SOBOLEV TYPE EQUATIONS

First of all, we discuss the model problem (1.1). Here and below, assume that T^h is a rectangular partition over Ω with mesh size h . The weak form of (1.1) consists in finding $u(\cdot, t) \in H_0^1(\Omega)$ for any $t \in [0, T]$ (the Sobolev space) such that

$$(2.1) \quad \begin{cases} (\nabla u_t, \nabla \varphi) + (\nabla u, \nabla \varphi) = (f, \varphi) & \forall \varphi \in H_0^1, \\ u(0) = v. \end{cases}$$

Let $S_0^h \subset H_0^1$ consist of piecewise bilinear functions. Thus, a continuous Galerkin approximation $u^h(x, t): [0, T] \rightarrow S_0^h$ is defined so that

$$(2.2) \quad \begin{cases} (\nabla u_t^h, \nabla \varphi) + (\nabla u^h, \nabla \varphi) = (f, \varphi) & \forall \varphi \in S_0^h, \\ u^h(0) = i_h v, \end{cases}$$

where $i_h v \in S_0^h$ is the bilinear interpolation function of v . From (2.1) and (2.2) we get the error equation

$$(2.3) \quad (\nabla(u_t^h - u_t), \nabla \varphi) + (\nabla(u^h - u), \nabla \varphi) = 0 \quad \forall \varphi \in S_0^h.$$

We need the following (see [9])

2.1. Lemma. For $\varphi \in S_0^h$,

$$|(\nabla(u - i_h u), \nabla\varphi)| \leq \begin{cases} ch^2 \|u\|_3 |\varphi|_1, \\ ch^2 \|u\|_4 \|\varphi\|_0. \end{cases}$$

2.1. Theorem. For sufficiently smooth u and u_t , we have

$$\|u^h - i_h u\|_1 \leq ch^2 \left[\int_0^t (\|u_t\|_3 + \|u\|_3)^2 ds \right]^{1/2}.$$

Proof. Let

$$\theta(x, t) = u^h - i_h u.$$

Then, by virtue of (2.3) we have for $\varphi \in S_0^h$

$$(2.4) \quad (\nabla\theta_t, \nabla\varphi) + (\nabla\theta, \nabla\varphi) = (\nabla(u_t - i_h u_t), \nabla\varphi) + (\nabla(u - i_h u), \nabla\varphi).$$

Hence, with $\varphi = \theta_t$ (where θ_t is the classical derivative) and Lemma 2.1,

$$|\theta_t|_1^2 + \frac{1}{2} \frac{d}{dt} |\theta|_1^2 \leq ch^2 (\|u_t\|_3 + \|u\|_3) |\theta_t|_1$$

or

$$|\theta_t|_1^2 + \frac{1}{2} \frac{d}{dt} |\theta|_1^2 \leq ch^4 (\|u_t\|_3 + \|u\|_3)^2 + \frac{1}{2} |\theta_t|_1^2.$$

By integration with respect to t , it follows from $\theta(x, 0) = 0$ that

$$|\theta|_1 \leq ch^2 \left[\int_0^t (\|u_t\|_3 + \|u\|_3)^2 ds \right]^{1/2}.$$

□

In order to derive L^∞ estimates, we introduce the discrete Green function $G_z^h \in S_0^h$ at any point $z \in \bar{\Omega}$ such that, for $\varphi \in S_0^h$,

$$(\nabla G_z^h, \nabla\varphi) = \varphi(z), \quad (\nabla D_z G_z^h, \nabla\varphi) = D_z \varphi.$$

We need the following

2.2. Lemma. ([16]) $\|G_z^h\|_0 \leq c$, $\|D_z G_z^h\|_0 \leq c \left(\log \frac{1}{h} \right)^{1/2}$.

2.2. Theorem. For sufficiently smooth u and u_t , we have

$$\|u_t^h - u_t\|_{0,\infty} \leq ch^2(\|u_t\|_{2,\infty} + \|u_t\|_4 + \|u\|_4).$$

P r o o f. Taking $\varphi = G_z^h$ in (2.4), we have by Lemma 2.1 and Lemma 2.2

$$(2.5) \quad \theta_t(z, t) + \theta(z, t) \leq ch^2(\|u_t\|_4 + \|u\|_4),$$

and hence, with $\theta(0) = 0$,

$$\theta_t(z, t) + \int_0^t \theta_t(z, s) \, ds \leq ch^2(\|u_t\|_4 + \|u\|_4).$$

Therefore, it follows from Gronwall's Lemma that

$$|\theta_t(z, t)| \leq ch^2(\|u_t\|_4 + \|u\|_4).$$

Thus, Theorem 2.2 holds in virtue of the triangle inequality

$$\|u_t^h - u_t\|_{0,\infty} \leq \|u_t^h - i_h u_t\|_{0,\infty} + \|i_h u_t - u_t\|_{0,\infty}.$$

□

2.3. Theorem. For sufficiently smooth u and u_t , we have

$$\|u^h - u\|_{0,\infty} \leq ch^2(\|u\|_{2,\infty} + \|u\|_4 + \|u_t\|_4).$$

P r o o f. From (2.5) and Theorem 2.2, we drive

$$|\theta(z, t)| \leq ch^2(\|u_t\|_4 + \|u\|_4),$$

and hence, Theorem 2.3 holds in virtue of the triangle inequality

$$\|u^h - u\|_{0,\infty} \leq \|u^h - i_h u\|_{0,\infty} + \|i_h u - u\|_{0,\infty}.$$

□

2.4. Theorem. For sufficiently smooth u and u_t , we have

$$\|u_t^h - i_h u_t\|_{1,\infty} \leq ch^2 \left(\log \frac{1}{h} \right)^{1/2} (\|u_t\|_4 + \|u\|_4).$$

P r o o f. Setting $\varphi = D_z G_z^h$ in (2.4), we get by means of Lemma 2.1 and 2.2

$$(2.6) \quad D_z \theta_t(z, t) + D_z \theta(z, t) \leq ch^2 \left(\log \frac{1}{h} \right)^{1/2} (\|u_t\|_4 + \|u\|_4).$$

And thus, with $D_z \theta(0) = 0$,

$$D_z \theta_t(z, t) + \int_0^t D_z \theta_t(z, s) \, ds \leq ch^2 \left(\log \frac{1}{h} \right)^{1/2} (\|u_t\|_4 + \|u\|_4).$$

Then, Theorem 2.4 follows from Gronwall's Lemma. □

2.5. Theorem. For sufficiently smooth u and u_t , we have

$$\|u^h - i_h u\|_{1,\infty} \leq ch^2 \left(\log \frac{1}{h} \right)^{1/2} (\|u_t\|_4 + \|u\|_4).$$

P r o o f. From (2.6) and Theorem 2.4 we obtain Theorem 2.5. □

Theorems 2.1, 2.4 and 2.5 play key roles in the analysis of the global superconvergence for the problem (1.1). Now we use an interpolation postprocessing technique from [8] to get the desired results. We assume that T^h has been gained from T^{2h} with mesh size $2h$ by subdividing each element of T^{2h} into four congruent elements. Thus, we can define a nodal biquadratic interpolation operator I_{2h}^2 associated with T^{2h} . It is easy to check that

$$I_{2h}^2 i_h = I_{2h}^2, \quad \|I_{2h}^2 \varphi\|_{1,p} \leq c \|\varphi\|_{1,p} \quad \forall \varphi \in S_0^h \quad (p = 2, \infty),$$

$$\|I_{2h}^2 \varphi - \varphi\|_{1,p} \leq ch^2 \|\varphi\|_{3,p} \quad (p = 2, \infty).$$

And thus, we have the following main results.

2.6. Theorem. For sufficiently smooth u and u_t , we have

$$\|I_{2h}^2 u^h - u\|_1 \leq ch^2 \left\{ \|u\|_3 + \left[\int_0^t (\|u_t\|_3 + \|u\|_3)^2 \, ds \right]^{1/2} \right\}.$$

Proof. Due to the property of I_{2h}^2 , we have

$$I_{2h}^2 u^h - u = I_{2h}^2(u^h - i_h u) + (I_{2h}^2 u - u).$$

Therefore, it follows from Theorem 2.1 and the interpolation theorem that

$$\begin{aligned} \|I_{2h}^2 u^h - u\|_1 &\leq c\|u^h - i_h u\|_1 + ch^2\|u\|_3 \\ &\leq ch^2 \left\{ \|u\|_3 + \left[\int_0^t (\|u_t\|_3 + \|u\|_3)^2 ds \right]^{1/2} \right\}. \end{aligned}$$

□

Analogously, by Theorems 2.4 and 2.5 we have the next assertions.

2.7. Theorem. For sufficiently smooth u and u_t , we have

$$\|I_{2h}^2 u_t^h - u_t\|_{1,\infty} \leq ch^2 \left(\log \frac{1}{h} \right)^{1/2} (\|u_t\|_{3,\infty} + \|u_t\|_4 + \|u\|_4).$$

2.8. Theorem. For sufficiently smooth u and u_t , we have

$$\|I_{2h}^2 u^h - u\|_{1,\infty} \leq ch^2 \left(\log \frac{1}{h} \right)^{1/2} (\|u\|_{3,\infty} + \|u_t\|_4 + \|u\|_4).$$

3. VISCOELASTICITY TYPE EQUATIONS

In this section, we will consider the semidiscrete Galerkin approximation to the problem (1.2). The weak form of (1.2) reads as follows: Find $u(\cdot, t) \in H_0^1(\Omega)$ for any fixed $t \in [0, T]$ such that

$$(3.1) \quad \begin{cases} (u_{tt}, \varphi) + (\nabla u_t, \nabla \varphi) + (\nabla u, \nabla \varphi) = (f, \varphi) \quad \forall \varphi \in H_0^1(\Omega), \\ u(0) = v, \quad u_t(0) = w. \end{cases}$$

Thus, a continuous Galerkin approximation $u^h(x, t): [0, T] \rightarrow S_0^h$ is defined so that

$$(3.2) \quad \begin{cases} (u_{tt}^h, \varphi) + (\nabla u_t^h, \nabla \varphi) + (\nabla u^h, \nabla \varphi) = (f, \varphi) \quad \forall \varphi \in S_0^h, \\ u^h(0) = i_h v, \quad u_t^h(0) = i_h w, \end{cases}$$

where $i_h v, i_h w \in S_0^h$ stand for the bilinear interpolation functions of v and w , respectively. Then, we obtain the error equation from (3.1) and (3.2):

$$(3.3) \quad (u_{tt} - u_{tt}^h, \varphi) + (\nabla(u_t - u_t^h), \nabla \varphi) + (\nabla(u - u^h), \nabla \varphi) = 0 \quad \forall \varphi \in S_0^h.$$

3.1. Theorem. For sufficiently smooth u, u_t and u_{tt} , we have

$$\|u^h - i_h u\|_1 + \|u_t^h - u_t\|_0 \leq ch^2 \left\{ \|u_t\|_2 + \left[\int_0^t (\|u_{tt}\|_2 + \|u_t\|_4 + \|u\|_4)^2 ds \right]^{1/2} \right\}.$$

Proof. Let

$$\theta(x, t) = u^h(x, t) - i_h u(x, t).$$

According to (3.3), we have for $\varphi \in S_0^h$ that

$$(3.4) \quad (\theta_{tt}, \varphi) + (\nabla \theta_t, \nabla \varphi) + (\nabla \theta, \nabla \varphi) = (u_{tt} - i_h u_{tt}, \varphi) + (\nabla(u_t - i_h u_t), \nabla \varphi) + (\nabla(u - i_h u), \nabla \varphi),$$

and hence, with $\varphi = \theta_t$ and Lemma 2.1,

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\theta_t\|_0^2 + c \|\theta_t\|_1^2 + \frac{1}{2} \frac{d}{dt} |\theta|_1^2 &\leq \frac{1}{2} \frac{d}{dt} \|\theta_t\|_0^2 + |\theta_t|_1^2 + \frac{1}{2} \frac{d}{dt} |\theta|_1^2 \\ &\leq ch^4 (\|u_{tt}\|_2 + \|u_t\|_4 + \|u\|_4)^2 + \frac{c}{2} \|\theta_t\|_0^2, \end{aligned}$$

or

$$\frac{d}{dt} (\|\theta_t\|_0^2 + |\theta|_1^2) \leq ch^4 (\|u_{tt}\|_2 + \|u_t\|_4 + \|u\|_4)^2.$$

By integration with respect to t , it follows from $\theta(0) = \theta_t(0) = 0$ that

$$\|\theta_t\|_0^2 + |\theta|_1^2 \leq ch^4 \int_0^t (\|u_{tt}\|_2 + \|u_t\|_4 + \|u\|_4)^2 ds,$$

that is

$$\|\theta_t\|_0 + |\theta|_1 \leq ch^2 \left[\int_0^t (\|u_{tt}\|_2 + \|u_t\|_4 + \|u\|_4)^2 ds \right]^{1/2}.$$

And thus, Theorem 3.1 follows from the triangle inequality

$$\|u_t^h - u_t\|_0 \leq \|u_t^h - i_h u_t\|_0 + \|i_h u_t - u_t\|_0.$$

□

3.2. Theorem. For sufficiently smooth u, u_t, u_{tt} and u_{ttt} , we have

$$\begin{aligned} \|u_t^h - i_h u_t\|_1 + \|u_{tt}^h - u_{tt}\|_0 &\leq ch^2 \left\{ \|u_{tt}\|_2 + [(\|u_{tt}(0)\|_2 + \|u_t(0)\|_4 + \|u(0)\|_4)^2 \right. \\ &\quad \left. + \int_0^t (\|u_{ttt}\|_2 + \|u_{tt}\|_4 + \|u_t\|_4)^2 ds]^{1/2} \right\}. \end{aligned}$$

P r o o f. Differentiating (3.4) with respect to t , we get, for $\varphi \in S_0^h$,

$$\begin{aligned} (\theta_{ttt}, \varphi) + (\nabla \theta_{tt}, \nabla \varphi) + (\nabla \theta_t, \nabla \varphi) &= (u_{ttt} - i_h u_{ttt}, \varphi) + (\nabla(u_{tt} - i_h u_{tt}), \nabla \varphi) \\ &\quad + (\nabla(u_t - i_h u_t), \nabla \varphi), \end{aligned}$$

and hence, with $\varphi = \theta_{tt}$ and Lemma 2.1,

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\theta_{tt}\|_0^2 + c \|\theta_{tt}\|_1^2 + \frac{1}{2} \frac{d}{dt} |\theta_t|_1^2 &\leq \frac{1}{2} \frac{d}{dt} \|\theta_{tt}\|_0^2 + |\theta_{tt}|_1^2 + \frac{1}{2} \frac{d}{dt} |\theta_t|_1^2 \\ &\leq ch^4 (\|u_{ttt}\|_2 + \|u_{tt}\|_4 + \|u_t\|_4)^2 + \frac{c}{2} \|\theta_{tt}\|_0^2, \end{aligned}$$

or

$$\frac{d}{dt} (\|\theta_{tt}\|_0^2 + |\theta_t|_1^2) \leq ch^4 (\|u_{ttt}\|_2 + \|u_{tt}\|_4 + \|u_t\|_4)^2.$$

Therefore, by virtue of $\theta_t(0) = 0$, we get

$$(3.5) \quad \|\theta_{tt}\|_0^2 + |\theta_t|_1^2 \leq \|\theta_{tt}(0)\|_0^2 + ch^4 \int_0^t (\|u_{ttt}\|_2 + \|u_{tt}\|_4 + \|u_t\|_4)^2 ds.$$

Let $t = 0$ and $\varphi = \theta_{tt}(0)$ in (3.4). Then

$$\|\theta_{tt}(0)\|_0 \leq ch^2 (\|u_{tt}(0)\|_2 + \|u_t(0)\|_4 + \|u(0)\|_4)$$

which, together with (3.5), leads to

$$\begin{aligned} \|\theta_{tt}\|_0^2 + |\theta_t|_1^2 &\leq ch^4 \left[(\|u_{tt}(0)\|_2 + \|u_t(0)\|_4 + \|u(0)\|_4)^2 \right. \\ &\quad \left. + \int_0^t (\|u_{ttt}\|_2 + \|u_{tt}\|_4 + \|u_t\|_4)^2 ds \right], \end{aligned}$$

or

$$\begin{aligned} \|\theta_{tt}\|_0 + \|\theta_t\|_1 &\leq ch^2 \left[(\|u_{tt}(0)\|_2 + \|u_t(0)\|_4 + \|u(0)\|_4)^2 \right. \\ &\quad \left. + \int_0^t (\|u_{ttt}\|_2 + \|u_{tt}\|_4 + \|u_t\|_4)^2 ds \right]^{1/2}, \end{aligned}$$

and Theorem 3.2 follows from the triangle inequality

$$\|u_{tt}^h - u_{tt}\|_0 \leq \|u_{tt}^h - i_h u_{tt}\|_0 + \|i_h u_{tt} - u_{tt}\|_0.$$

□

3.3. Theorem. For sufficiently smooth u , u_t , u_{tt} and u_{ttt} , we have

$$\begin{aligned} \|u_t^h - u_t\|_{0,\infty} &\leq ch^2 \left\{ \|u_{tt}\|_2 + \|u_t\|_4 + \|u_t\|_{2,\infty} + \|u\|_4 \right. \\ &\quad + [(\|u_{tt}(0)\|_2 + \|u_t(0)\|_4 + \|u(0)\|_4)^2 \\ &\quad \left. + \int_0^t (\|u_{ttt}\|_2 + \|u_{tt}\|_4 + \|u_t\|_4)^2 ds]^{1/2} \right\}. \end{aligned}$$

P r o o f. Setting $\varphi = G_z^h$ in (3.4), we have according to Lemmas 2.1 and 2.2

$$(3.6) \quad (\theta_{tt}, G_z^h) + \theta_t(z, t) + \theta(z, t) \leq ch^2(\|u_{tt}\|_2 + \|u_t\|_4 + \|u\|_4),$$

or

$$|\theta_t(z, t)| \leq \int_0^t |\theta_t(z, s)| ds + c\|\theta_{tt}\|_0 + ch^2(\|u_{tt}\|_2 + \|u_t\|_4 + \|u\|_4).$$

And thus, it follows from Gronwall's Lemma and Theorem 3.2 that

$$\begin{aligned} |\theta_t(z, t)| &\leq ch^2 \left\{ \|u_{tt}\|_2 + \|u_t\|_4 + \|u\|_4 \right. \\ &\quad + [(\|u_{tt}(0)\|_2 + \|u_t(0)\|_4 + \|u(0)\|_4)^2 \\ &\quad \left. + \int_0^t (\|u_{ttt}\|_2 + \|u_{tt}\|_4 + \|u_t\|_4)^2 ds]^{1/2} \right\}. \end{aligned}$$

Then, Theorem 3.3 holds by virtue of the triangle inequality

$$\|u_t^h - u_t\|_{0,\infty} \leq \|u_t^h - i_h u_t\|_{0,\infty} + \|i_h u_t - u_t\|_{0,\infty}.$$

□

3.4. Theorem. For sufficiently smooth u , u_t , u_{tt} and u_{ttt} , we have

$$\begin{aligned} \|u^h - u\|_{0,\infty} &\leq ch^2 \left\{ \|u\|_{2,\infty} + \|u\|_4 + \|u_t\|_4 + \|u_{tt}\|_2 \right. \\ &\quad + [(\|u_{tt}(0)\|_2 + \|u_t(0)\|_4 + \|u(0)\|_4)^2 \\ &\quad \left. + \int_0^t (\|u_{ttt}\|_2 + \|u_{tt}\|_4 + \|u_t\|_4)^2 ds]^{1/2} \right\}. \end{aligned}$$

P r o o f. From (3.6), Lemma 2.2 and Theorems 3.2, 3.3 we derive

$$\begin{aligned} |\theta(z, t)| &\leq |\theta_t(z, t)| + c\|\theta_{tt}\|_0 + ch^2(\|u_{tt}\|_2 + \|u_t\|_4 + \|u\|_4) \\ &\leq ch^2 \left\{ \|u_{tt}\|_2 + \|u_t\|_4 + \|u\|_4 + [(\|u_{tt}(0)\|_2 + \|u_t(0)\|_4 \right. \\ &\quad \left. + \|u(0)\|_4)^2 + \int_0^t (\|u_{ttt}\|_2 + \|u_{tt}\|_4 + \|u_t\|_4)^2 ds]^{1/2} \right\}, \end{aligned}$$

and hence, Theorem 3.4 holds in virtue of the triangle inequality

$$\|u^h - u\|_{0,\infty} \leq \|u^h - i_h u\|_{0,\infty} + \|i_h u - u\|_{0,\infty}.$$

□

3.5. Theorem. *For sufficiently smooth u , u_t , u_{tt} and u_{ttt} , we have*

$$\begin{aligned} \|u_t^h - i_h u_t\|_{1,\infty} &\leq ch^2 \left(\log \frac{1}{h} \right)^{1/2} \left\{ \|u_{tt}\|_2 + \|u_t\|_4 + \|u\|_4 \right. \\ &\quad + [(\|u_{tt}(0)\|_2 + \|u_t(0)\|_4 + \|u(0)\|_4)^2 \\ &\quad \left. + \int_0^t (\|u_{ttt}\|_2 + \|u_{tt}\|_4 + \|u_t\|_4)^2 ds]^{1/2} \right\}. \end{aligned}$$

P r o o f. Taking $\varphi = D_z G_z^h$ in (3.4), we obtain according to Lemmas 2.1 and 2.2

$$(3.7) \quad (\theta_{tt}, D_z G_z^h) + D_z \theta_t(z, t) + D_z \theta(z, t) \leq ch^2 \left(\log \frac{1}{h} \right)^{1/2} (\|u_{tt}\|_2 + \|u_t\|_4 + \|u\|_4),$$

or

$$\begin{aligned} |D_z \theta_t(z, t)| &\leq \int_0^t |D_z \theta_t(z, s)| ds + c \left(\log \frac{1}{h} \right)^{1/2} \|\theta_{tt}\|_0 \\ &\quad + ch^2 \left(\log \frac{1}{h} \right)^{1/2} (\|u_{tt}\|_2 + \|u_t\|_4 + \|u\|_4). \end{aligned}$$

And thus, Theorem 3.5 follows from Gronwall's Lemma and Theorem 3.2. □

3.6. Theorem. *For sufficiently smooth u , u_t , u_{tt} and u_{ttt} , we have*

$$\begin{aligned} \|u^h - i_h u\|_{1,\infty} &\leq ch^2 \left(\log \frac{1}{h} \right)^{1/2} \left\{ \|u_{tt}\|_2 + \|u_t\|_4 + \|u\|_4 \right. \\ &\quad + [(\|u_{tt}(0)\|_2 + \|u_t(0)\|_4 + \|u(0)\|_4)^2 \\ &\quad \left. + \int_0^t (\|u_{ttt}\|_2 + \|u_{tt}\|_4 + \|u_t\|_4)^2 ds]^{1/2} \right\}. \end{aligned}$$

P r o o f. From (3.7) and Theorems 3.2, 3.5 we obtain Theorem 3.6. □

Theorems 3.1, 3.2, 3.5 and 3.6 are essential, by them we can get the global superconvergence for the problem (1.2) instead of the interior pointwise superconvergence. Identically to Section 2, we derive the following main theorems by means of the interpolation postprocessing technique initiated in [8].

3.7. Theorem. For sufficiently smooth u, u_t and u_{tt} , we have

$$\|I_{2h}^2 u^h - u\|_1 \leq ch^2 \left\{ \|u_t\|_2 + \|u\|_3 + \left[\int_0^t (\|u_{tt}\|_2 + \|u_t\|_4 + \|u\|_4)^2 ds \right]^{1/2} \right\}.$$

3.8. Theorem. For sufficiently smooth u, u_t, u_{tt} and u_{ttt} , we have

$$\begin{aligned} \|I_{2h}^2 u_t^h - u_t\|_1 \leq ch^2 \left\{ \|u_{tt}\|_2 + \|u_t\|_3 \right. \\ \left. + [(\|u_{tt}(0)\|_2 + \|u_t(0)\|_4 + \|u(0)\|_4)^2 \right. \\ \left. + \int_0^t (\|u_{ttt}\|_2 + \|u_{tt}\|_4 + \|u_t\|_4)^2 ds]^{1/2} \right\}. \end{aligned}$$

3.9. Theorem. For sufficiently smooth u, u_t, u_{tt} and u_{ttt} , we have

$$\begin{aligned} \|I_{2h}^2 u_t^h - u_t\|_{1,\infty} \leq ch^2 \left(\log \frac{1}{h} \right)^{1/2} \left\{ \|u_{tt}\|_2 + \|u_t\|_4 + \|u_t\|_{3,\infty} \right. \\ \left. + \|u\|_4 + [(\|u_{tt}(0)\|_2 + \|u_t(0)\|_4 + \|u(0)\|_4)^2 \right. \\ \left. + \int_0^t (\|u_{ttt}\|_2 + \|u_{tt}\|_4 + \|u_t\|_4)^2 ds]^{1/2} \right\}. \end{aligned}$$

3.10. Theorem. For sufficiently smooth u, u_t, u_{tt} and u_{ttt} , we have

$$\begin{aligned} \|I_{2h}^2 u^h - u\|_{1,\infty} \leq ch^2 \left(\log \frac{1}{h} \right)^{1/2} \left\{ \|u_{tt}\|_2 + \|u_t\|_4 + \|u\|_{3,\infty} \right. \\ \left. + \|u\|_4 + [(\|u_{tt}(0)\|_2 + \|u_t(0)\|_4 + \|u(0)\|_4)^2 \right. \\ \left. + \int_0^t (\|u_{ttt}\|_2 + \|u_{tt}\|_4 + \|u_t\|_4)^2 ds]^{1/2} \right\}. \end{aligned}$$

Remark 1. In another paper, we will discuss the case of k ($k \geq 2$) which is the order of finite elements for the problems above.

Remark 2. When Ω is a convex quadrilateral domain, the corresponding superconvergent results hold for such problems as above if the quadrilateral meshes are almost uniform and are constructed by connecting the equi-proportional points of two opposite boundaries.

Acknowledgement. Prof. Y.P. Lin generously gave us his papers about Sobolev and viscoelasticity equations when the first author visited Canada in 1994. His work in this field has aroused our interest in such problems. The authors would like to thank Prof. M. Křížek whose comments improved this version of the paper.

References

- [1] *D. Arnold, J. Douglas, V. Thomée*: Superconvergence of a finite element approximation to the solution of a Sobolev equation in a single space variable. *Math. Comp.* *36* (1981), 53–63.
- [2] *R. Ewing*: The approximation of certain parabolic equations backward in time by Sobolev equations. *SIAM J. Math. Anal.* *6* (1975), 283–294.
- [3] *R. Ewing*: Numerical solution of Sobolev partial differential equations. *SIAM J. Numer. Anal.* *12* (1975), 345–363.
- [4] *W. Ford*: Galerkin approximation to nonlinear pseudoparabolic partial differential equation. *Aequationes Math.* *14* (1976), 271–291.
- [5] *W. Ford, T. Ting*: Stability and convergence of difference approximations to pseudoparabolic partial equations. *Math. Comp.* *27* (1973), 737–743.
- [6] *W. Ford, T. Ting*: Uniform error estimates for difference approximations to nonlinear pseudoparabolic partial differential equations. *SIAM J. Numer. Anal.* *11* (1974), 155–169.
- [7] *Q. Lin*: A new observation in FEM. *Proc. Syst. Sci. & Syst. Eng.* (1991), 389–391. Great Wall (H.K.) Culture Publish Co..
- [8] *Q. Lin, N. Yan, A. Zhou*: A rectangle test for interpolated finite elements, *ibid.*
- [9] *Q. Lin, S. Zhang*: An immediate analysis for global superconvergence for integrodifferential equations. *Appl. Math.* *42* (1997), 1–21.
- [10] *Y. Lin*: Galerkin methods for nonlinear Sobolev equations. *Aequationes Math.* *40* (1990), 54–56.
- [11] *Y. Lin, T. Zhang*: Finite element methods for nonlinear Sobolev equations with nonlinear boundary conditions. *J. Math. Anal. & Appl.* *165* (1992), 180–191.
- [12] *Y. Lin, V. Thomée, L. Wahlbin*: Ritz-Volterra projection on finite element spaces and applications to integrodifferential and related equations. *SIAM J. Numer. Anal.* *28* (1991), 1047–1070.
- [13] *M. Nakao*: Error estimates of a Galerkin method for some nonlinear Sobolev equations in one space dimension. *Numer. Math.* *47* (1985), 139–157.
- [14] *L. Wahlbin*: Error estimates for a Galerkin method for a class of model equations for long waves. *Numer. Math.* *23* (1975), 289–303.
- [15] *M. Wheeler*: A priori L_2 error estimates for Galerkin approximations to parabolic partial differential equations. *SIAM J. Numer. Anal.* *10* (1973), 723–759.
- [16] *Q. Zhu, Q. Lin*: Superconvergence Theory of the Finite Element Methods. Hunan Science Press, 1990.

Author's address: Qun Lin, Shuhua Zhang, Institute of Systems Science, Academia Sinica, Beijing 100080, P. R. China.