

# Applications of Mathematics

---

Jiří Anděl

An estimator for parameters of a nonlinear nonnegative multidimensional AR(1) process

*Applications of Mathematics*, Vol. 43 (1998), No. 5, 389--398

Persistent URL: <http://dml.cz/dmlcz/134395>

## Terms of use:

© Institute of Mathematics AS CR, 1998

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

AN ESTIMATOR FOR PARAMETERS OF A NONLINEAR  
NONNEGATIVE MULTIDIMENSIONAL AR(1) PROCESS

JIŘÍ ANDĚL, Praha

(Received February 16, 1998)

*Abstract.* Let  $e_t = (e_{t1}, \dots, e_{tp})'$  be a  $p$ -dimensional nonnegative strict white noise with finite second moments. Let  $h_{ij}(x)$  be nondecreasing functions from  $[0, \infty)$  onto  $[0, \infty)$  such that  $h_{ij}(x) \leq x$  for  $i, j = 1, \dots, p$ . Let  $U = (u_{ij})$  be a  $p \times p$  matrix with nonnegative elements having all its roots inside the unit circle. Define a process  $X_t = (X_{t1}, \dots, X_{tp})'$  by

$$X_{tj} = u_{j1}h_{1j}(X_{t-1,1}) + \dots + u_{jp}h_{pj}(X_{t-1,p}) + e_{tj}$$

for  $j = 1, \dots, p$ . A method for estimating  $U$  from a realization  $X_1, \dots, X_n$  is proposed. It is proved that the estimators are strongly consistent.

*Keywords:* autoregressive process, estimating parameters, multidimensional process, nonlinear process, nonnegative process

*MSC 2000:* 62M10

## 1. INTRODUCTION

Consider a one-dimensional AR(1) process  $X_t$  given by  $X_t = bX_{t-1} + e_t$  where  $b \in [0, 1)$  and  $e_t$  is a nonnegative strict white noise such that  $Ee_t^2 < \infty$ . Define  $F(x) = P(e_t < x)$ . Bell and Smith [9] proved that

$$(1.1) \quad b^* = \min \left( \frac{X_2}{X_1}, \dots, \frac{X_n}{X_{n-1}} \right)$$

is a strongly consistent estimator for  $b$  if and only if  $F(d) - F(c) < 1$  for all  $0 < c < d < \infty$ . Anděl [3] derived the distribution of  $b^*$  under the assumption that  $F$  is the distribution function of an exponential distribution. Davis and McCormick [10] obtained the asymptotic distribution of  $b^*$  when  $F$  varies regularly at 0 and satisfies a

---

Supported by grant 201/97/1176 from the Grant Agency of the Czech Republic.

suitable moment condition. For the case of AR( $p$ ) model with  $p > 1$  a straightforward generalization of (1.1) does not perform well. Anděl [5] suggested another estimator based on a maximum likelihood argument. His method was modified and generalized by An [1] and by An and Huang [2]. Asymptotic theory is developed in the paper by Feigin and Resnick [11]. The method was applied to nonlinear one-dimensional AR models by Anděl [4] and [6].

## 2. THE MODEL

Non-linear models of time series are important tools in statistical analysis of data. Statistical tests show that many series introduced in textbooks and papers are non-linear (see Tong [13]).

One of the simplest non-linear models is the one-dimensional non-linear AR(1) process  $X_t$  defined by

$$(2.1) \quad X_t = \lambda(X_{t-1}) + e_t$$

where  $\lambda$  is a non-linear function and  $e_t$  is a white noise. Even in this case it is quite hard task to estimate the function  $\lambda$  from a realization  $X_1, \dots, X_n$  (see Auestad and Tjøstheim [8]). It was proved by Jones [12] under general assumptions about the distribution function of  $e_t$  that the process  $\{X_t\}$  cannot be stationary because it has “a drift to infinity” with a positive probability if there exist constants  $\gamma > 0$  and  $q > 1$  such that  $|\lambda(x)| \geq \gamma|x|^q$  for large  $|x|$ .

It is a problem how to generalize (2.1) to  $p$ -dimensional case. The linear  $p$ -dimensional AR(1) process  $\mathbf{X}_t$  is given by  $\mathbf{X}_t = \mathbf{U}\mathbf{X}_{t-1} + \mathbf{e}_t$  where  $\mathbf{U}$  is a  $p \times p$  matrix and  $\mathbf{e}_t$  is a  $p$ -dimensional white noise. The model  $\mathbf{X}_t = \boldsymbol{\lambda}(\mathbf{X}_{t-1}) + \mathbf{e}_t$  with a non-linear function  $\boldsymbol{\lambda}: \mathbb{R}_p \rightarrow \mathbb{R}_p$  is too general. In this case the function  $\boldsymbol{\lambda}$  should be estimated in a whole, not only its parameters. In this paper we introduce a special non-linear model where  $\mathbf{X}_t$  is created as a linear combination of non-linear functions of components of  $\mathbf{X}_{t-1}$  plus a white noise (see (2.2) below). For simplicity we assume that the non-linear functions are known and only the parameters of their linear combinations must be estimated.

Remembering that a rapidly growing function  $\lambda$  in the one-dimensional model (2.1) leads to serious problems we assume that our non-linear functions in  $p$ -dimensional model do not grow faster than  $x$  (assumption A4).

The resulting model (2.2) seems to be quite flexible and its form enables to use procedures for estimating parameters similar to those in linear models.

Now, we introduce some assumptions valid throughout the paper.

A1. Let  $\mathbf{U} = (u_{ij})$  be a  $p \times p$  matrix with nonnegative elements such that all its roots are inside the unit circle.

A2. Let  $\mathbf{e}_t = (e_{t1}, \dots, e_{tp})'$  be independent identically distributed random vectors with nonnegative components and with finite second moments.

A3. Let  $\mathbf{Z}_t = (Z_{t1}, \dots, Z_{tp})'$  be the  $p$ -dimensional AR(1) process defined by  $\mathbf{Z}_t = \mathbf{U}\mathbf{Z}_{t-1} + \mathbf{e}_t$ .

A4. Let  $h_{ij}(x)$  be measurable nondecreasing functions mapping  $[0, \infty)$  onto  $[0, \infty)$  such that  $h_{ij}(x) \leq x$  for  $i, j = 1, \dots, p$ .

A5. Let  $P(e_{t1} < c, \dots, e_{tp} < c) > 0$  for every  $c > 0$ .

A6. There exists a constant  $K_{ij} > 0$  for each pair  $(i, j)$ ,  $i, j \in \{1, \dots, p\}$ , such that

$$P\{e_{t1} < c, \dots, e_{t,j-1} < c, h_{ji}(e_{tj}) > K_{ji}, e_{t,j+1} < c, \dots, e_{tp} < c\} > 0$$

for every  $c > 0$ .

Now, introduce a process  $\mathbf{X}_t = (X_{t1}, \dots, X_{tp})'$  in the following way. Define  $\mathbf{X}_t = \mathbf{Z}_t$  for  $t \leq 1$  and

$$(2.2) \quad X_{ti} = \sum_{j=1}^p u_{ij} h_{ji}(X_{t-1,j}) + e_{ti}, \quad i = 1, \dots, p,$$

for  $t \geq 2$ . Then we say that  $\{X_t, t \geq 1\}$  is a nonlinear  $p$ -dimensional AR(1) process. It follows from our assumptions that all the variables  $X_{ti}$  are nonnegative.

### 3. AUXILIARY RESULTS

**Lemma 3.1.** We have  $X_{ti} \leq Z_{ti}$  for all  $t$  and  $i$ .

*Proof.* The assertion can easily be proved by complete induction. □

Assume that a realization  $\mathbf{X}_1, \dots, \mathbf{X}_n$  is given. Define

$$u_{ij}^+ = \min_{2 \leq t \leq n} \frac{X_{ti}}{h_{ji}(X_{t-1,j})}, \quad i, j = 1, \dots, p.$$

**Theorem 3.2.** We have  $u_{ij}^+ \rightarrow u_{ij}$  a.s. for each  $i, j = 1, \dots, p$  as  $n \rightarrow \infty$ .

*Proof.* Let  $\mathbf{U}^k = (u_{ij}^{(k)})$  for  $k = 1, 2, \dots$ . Since

$$\mathbf{Z}_t = \sum_{k=0}^{\infty} \mathbf{U}^k \mathbf{e}_{t-k},$$

we can write

$$(3.1) \quad Z_{ti} = e_{ti} + \sum_{k=1}^{\infty} \sum_{j=1}^p u_{ij}^{(k)} e_{t-k,j}, \quad j = 1, \dots, p.$$

Since the elements  $u_{ij}$  are nonnegative, we have also  $u_{ij}^{(k)} \geq 0$  for all  $i, j, k$ . Assumption A1 also implies that  $\sum_k \sum_i \sum_j u_{ij}^{(k)} < \infty$  and thus there exists a constant  $L$  such that  $u_{ij}^{(k)} \leq L$  for all  $i, j, k$ .

To simplify the notation, we prove the assertion only for  $i = j = 1$ . The proof for any other pair  $(i, j)$  is the same. In our case we have

$$u_{11}^+ = u_{11} + \min_{2 \leq t \leq n} \frac{\sum_{j=2}^p u_{1j} h_{j1}(X_{t-1,j}) + e_{t1}}{h_{11}(X_{t-1,1})}.$$

Let  $\varepsilon > 0$  be a given number. Introduce the events

$$Q_t = \left\{ \omega: \frac{\sum_{j=2}^p u_{1j} Z_{t-1,j} + e_{t1}}{h_{11}(e_{t-1,1})} < \varepsilon \right\}.$$

From the inequalities

$$h_{j1}(X_{t-1,j}) \leq X_{t-1,j} \leq Z_{t-1,j}, \quad h_{11}(X_{t-1,1}) \geq h_{11}(e_{t-1,1})$$

it follows that  $u_{11}^+ < u_{11} + \varepsilon$  if at least one of the events  $Q_2, \dots, Q_n$  occurs. But  $u_{11}^+ \geq u_{11}$  trivially holds. We prove that for every  $\varepsilon > 0$  with probability one there exist infinitely many indices  $t \geq 2$  such that the events  $Q_t$  occur. This implies, of course, that  $u_{11}^+ \rightarrow u_{11}$  a.s. Using (3.1) we get

$$Q_t = \left\{ \omega: e_{t1} + \sum_{j=2}^p u_{1j} \left[ e_{t-1,j} + \sum_{k=1}^{\infty} \sum_{i=1}^p u_{ji}^{(k)} e_{t-1-k,i} \right] < \varepsilon h_{11}(e_{t-1,1}) \right\}.$$

Denote  $K = K_{11}$ . For every integer  $M \geq 1$  define the events

$$Q_{tM1} = \left\{ \omega: h_{11}(e_{t-1,1}) > K, \quad e_{t1} < \frac{\varepsilon K}{2Mp^2}, \right. \\ \left. u_{1j} e_{t-1,j} < \frac{\varepsilon K}{2Mp^2} \quad \text{for } j = 2, \dots, p, \right. \\ \left. u_{1j} u_{ji}^{(k)} e_{t-1-k,i} < \frac{\varepsilon K}{2Mp^2} \quad \text{for } j = 2, \dots, p; i = 1, \dots, p; \right. \\ \left. k = 1, \dots, M \right\},$$

$$Q_{tM2} = \left\{ \omega: \sum_{k=M+1}^{\infty} \sum_{j=2}^p \sum_{i=1}^p u_{1j} u_{ji}^{(k)} e_{t-1-k,i} < \frac{\varepsilon K}{2} \right\}.$$

It is clear that  $Q_t \supset Q_{tM1} \cap Q_{tM2}$ . Let

$$Q_{tM1}^* = \left\{ \omega: h_{11}(e_{t-1,1}) > K, e_{t1} < \frac{\varepsilon K}{2Mp^2}, \right. \\ \left. e_{t-1,j} < \frac{\varepsilon K}{2Mp^2L} \text{ for } j = 2, \dots, p, \right. \\ \left. e_{t-1-k,i} < \frac{\varepsilon K}{2Mp^2L^2} \text{ for } i = 1, \dots, p; k = 1, \dots, M \right\}.$$

Then  $Q_{tM1} \supset Q_{tM1}^*$  and our assumptions yield that neither  $P(Q_{tM1})$  nor  $P(Q_{tM1}^*)$  depend on  $t$ . Since

$$P(Q_{tM1}^*) = P\left(e_{t1} < \frac{\varepsilon K}{2Mp^2}\right) \\ \times P\left\{h_{11}(e_{t-1,1}) > K, e_{t-1,j} < \frac{\varepsilon K}{2Mp^2L} \text{ for } j = 2, \dots, p\right\} \\ \times \prod_{k=1}^M P\left\{e_{t-1-k,i} < \frac{\varepsilon K}{2Mp^2L^2} \text{ for } i = 1, \dots, p\right\} > 0,$$

we have also  $P(Q_{tM1}) > 0$ . Further,  $P(Q_{tM2}) \rightarrow 1$  as  $M \rightarrow \infty$  and the probability  $P(Q_{tM2})$  does not depend on  $t$ . Denote  $\pi_M = P(Q_{tM1})$ . Let  $w_M$  be the smallest integer such that  $w_M \pi_M \geq 1$  holds. Introduce the sets  $S_1, S_2, \dots$  in the following way. Let  $S_1$  contain the elements of  $w_1$  triples

$$(1, 2, 3), (4, 5, 6), \dots, (3w_1 - 2, 3w_1 - 1, 3w_1),$$

let  $S_2$  contain the elements of  $w_2$  four-tuples starting from  $3w_1 + 1, 3w_1 + 2, 3w_1 + 3, 3w_1 + 4$ ) and so on. The last elements of the triples, four-tuples etc. denote  $t_1, t_2, \dots$ . If  $t_r \in S_M$  we have  $Q_{t_r} \supset Q_{t_rM1} \cap Q_{t_rM2}$ . The events  $Q_{t_1M1}, Q_{t_2M1}, \dots$  are independent,

$$\sum_{r=1}^{\infty} P(Q_{t_rM1}) \geq \sum_{M=1}^{\infty} w_M \pi_M = \infty,$$

$P(Q_{t_rM2}) \rightarrow 1$  as  $r \rightarrow \infty$  and the events  $Q_{t_rM1}$  and  $Q_{t_rM2}$  are independent for every  $r = 1, 2, \dots$ . It follows from the generalized Borel lemma (see Anděl and Dupač [7]) that then with probability one infinitely many events  $Q_{t_rM1} \cap Q_{t_rM2}$  occur and thus also infinitely many events  $Q_t$  occur with probability one.  $\square$

#### 4. ESTIMATING PARAMETERS

To simplify the notation, we consider in this section the case  $p = 2$  only. The procedure for  $p > 2$  is analogous. First of all, we introduce a motivation of our method of estimation.

It is known from simulations that  $u_{ij}^+$  cannot be used as estimator for  $u_{ij}$  because the convergence  $u_{ij}^+ \rightarrow u_{ij}$  is very slow. Assume for a while that  $e_{t1}$  and  $e_{t2}$  are independent random variables having the exponential distribution  $Ex(\lambda_1)$  and  $Ex(\lambda_2)$ , respectively. Then the conditional likelihood of  $\mathbf{X}_2, \dots, \mathbf{X}_n$ , given  $\mathbf{X}_1$ , is

$$\begin{aligned} & \exp \left\{ - \sum_{t=2}^n [X_{t1} - u_{11}h_{11}(X_{t-1,1}) - u_{12}h_{21}(X_{t-1,2})] / \lambda_1 \right\} \\ & \times \exp \left\{ - \sum_{t=2}^n [X_{t2} - u_{21}h_{12}(X_{t-1,1}) - u_{22}h_{22}(X_{t-1,2})] / \lambda_2 \right\} \end{aligned}$$

under the conditions that

$$X_{t1} - u_{11}h_{11}(X_{t-1,1}) - u_{12}h_{21}(X_{t-1,2}) \geq 0, \quad t = 2, \dots, n,$$

and

$$X_{t2} - u_{21}h_{12}(X_{t-1,1}) - u_{22}h_{22}(X_{t-1,2}) \geq 0, \quad t = 2, \dots, n.$$

If these conditions are not satisfied then the conditional likelihood is zero. The maximum likelihood method leads to the conclusion to estimate  $u_{11}$ ,  $u_{12}$  by the variables which maximize

$$(4.1) \quad a \sum_{t=2}^n h_{11}(X_{t-1,1}) + b \sum_{t=2}^n h_{21}(X_{t-1,2})$$

for  $a \geq 0$ ,  $b \geq 0$  under the conditions

$$(4.2) \quad X_{t1} - ah_{11}(X_{t-1,1}) - bh_{21}(X_{t-1,2}) \geq 0, \quad t = 2, \dots, n,$$

and to estimate  $u_{21}$ ,  $u_{22}$  by the variables which maximize

$$(4.3) \quad c \sum_{t=2}^n h_{12}(X_{t-1,1}) + d \sum_{t=2}^n h_{22}(X_{t-1,2})$$

for  $c \geq 0$ ,  $d \geq 0$  under the conditions

$$(4.4) \quad X_{t2} - ch_{12}(X_{t-1,1}) - dh_{22}(X_{t-1,2}) \geq 0, \quad t = 2, \dots, n.$$

One can expect that the results will be approximately the same as in the case that we sum up to  $t = n + 1$  in (4.1) and (4.3). We prove that such estimators are strongly consistent. Note that this assertion will be proved under our general assumptions A1–A6 and that we do not use any specific distribution of  $e_t$ .

**Theorem 4.1.** *Let  $u_{11}^*, u_{12}^*$  be a solution of the linear program*

$$(4.5) \quad \max_{a,b} \left\{ a \sum_{t=1}^n h_{11}(X_{t1}) + b \sum_{t=1}^n h_{21}(X_{t2}) \right\}$$

for  $a \geq 0, b \geq 0$  under the conditions (4.2). Let  $u_{21}^*, u_{22}^*$  be a solution of the linear program

$$\max_{c,d} \left\{ c \sum_{t=1}^n h_{12}(X_{t1}) + d \sum_{t=1}^n h_{22}(X_{t2}) \right\}$$

for  $c \geq 0, d \geq 0$  under the conditions (4.4). Then  $u_{11}^* \rightarrow u_{11}, u_{12}^* \rightarrow u_{12}, u_{21}^* \rightarrow u_{21}, u_{22}^* \rightarrow u_{22}$  a.s. as  $n \rightarrow \infty$ .

*P r o o f.* Instead of (4.5) we consider an equivalent formula, namely to solve

$$(4.6) \quad \max_{a,b} \left\{ a \frac{1}{n} \sum_{t=1}^n h_{11}(X_{t1}) + b \frac{1}{n} \sum_{t=1}^n h_{21}(X_{t2}) \right\}$$

for  $a \geq 0, b \geq 0$  under (4.2). Assume first that  $u_{11} > 0, u_{12} > 0$ . Define

$$\begin{aligned} M_n &= \{(a, b) : a \geq 0, b \geq 0, X_{t1} - ah_{11}(X_{t-1,1}) - bh_{21}(X_{t-1,2}) \geq 0, \\ &\quad t = 2, \dots, n\}, \\ M &= \{(a, b) : 0 \leq a \leq u_{11}, 0 \leq b \leq u_{12}\}. \end{aligned}$$

It is clear that  $M_2 \supset M_3 \supset \dots$ . We prove that  $M_n \rightarrow M$  a.s. Theorem 2.2 implies that there exists a sequence of indices  $t_i$  such that

$$\frac{X_{t_i 1}}{h_{11}(X_{t_i-1,1})} \rightarrow u_{11} \quad \text{a.s.}$$

Since

$$\frac{X_{t1}}{h_{11}(X_{t-1,1})} = u_{11} + \frac{u_{12}h_{21}(X_{t-1,2}) + e_{t1}}{h_{11}(X_{t-1,1})}$$

we can see that

$$(4.7) \quad \frac{h_{21}(X_{t_i-1,2})}{h_{11}(X_{t_i-1,1})} \rightarrow 0 \quad \text{a.s.}$$



Because

$$\frac{X_{t1}}{h_{21}(X_{t-1,2})} = u_{12} + \frac{u_{11}h_{11}(X_{t-1,1}) + e_{t1}}{h_{21}(X_{t-1,2})}$$

using (4.7) we get

$$\frac{X_{t1}}{h_{21}(X_{t-1,2})} \rightarrow \infty \quad \text{a.s.}$$




Fig. 1

In this case the straight line  $p$  in Fig. 1 approaches the straight line  $q_1$ . Quite similarly it can be proved that with probability one there exists a sequence of straight lines  $p$  converging to  $q_2$ . It is easy to calculate that  $p$  intersects  $q_1$  at the point

$$\left( u_{11}, u_{12} + \frac{e_{t1}}{h_{21}(X_{t-1,2})} \right)$$

and thus no straight line  $p$  intersects  $M$ . Consider the sequence of linear programs (4.6) for  $n \rightarrow \infty$ . We know that

$$h_{11}(X_{t1}) \geq h_{11}(e_{t1}), \quad h_{21}(X_{t2}) \geq h_{21}(e_{t2}).$$

Denote

$$H_{11} = Eh_{11}(e_{t1}), \quad H_{21} = Eh_{21}(e_{t2}).$$

Since  $h_{11}(e_{t1})$  as well as  $h_{21}(e_{t2})$  are i.i.d. random variables with finite second moments, the law of large numbers yields

$$\frac{1}{n} \sum_{t=1}^n h_{11}(e_{t1}) \rightarrow H_{11}, \quad \frac{1}{n} \sum_{t=1}^n h_{21}(e_{t2}) \rightarrow H_{21} \quad \text{a.s.}$$

On the other hand,

$$h_{11}(X_{t1}) \leq X_{t1} \leq Z_{t1}, \quad h_{21}(X_{t2}) \leq X_{t2} \leq Z_{t2}.$$

Our assumptions imply that the process  $\mathbf{Z}_t = (Z_{t1}, Z_{t2})'$  is ergodic. If we denote  $L_1 = EZ_{t1}$ ,  $L_2 = EZ_{t2}$ , then the ergodic theorem gives

$$\frac{1}{n} \sum_{t=1}^n Z_{t1} \rightarrow L_1, \quad \frac{1}{n} \sum_{t=1}^n Z_{t2} \rightarrow L_2.$$

Thus for every fixed  $\varepsilon > 0$  there exists  $n_0$  such that with probability one we have for all  $n \geq n_0$  that

$$H_{11} - \varepsilon \leq \frac{1}{n} \sum_{t=1}^n h_{11}(X_{t1}) \leq L_1 + \varepsilon, \quad H_{21} - \varepsilon \leq \frac{1}{n} \sum_{t=1}^n h_{21}(X_{t2}) \leq L_2 + \varepsilon.$$

We choose  $\varepsilon$  so small that  $H_{11} - \varepsilon > 0$ ,  $H_{21} - \varepsilon > 0$ . Therefore, the coefficients standing by  $a$  and  $b$  in the expression (4.6) are positive and bounded for all sufficiently large  $n$  and  $M_n \rightarrow M$ . This implies that the sequence of any solutions of the linear programs (4.6) converges to the point  $(u_{11}, u_{12})$  in which (4.6) is maximized on  $M$ .

If  $u_{11} = 0$  and/or  $u_{12} = 0$ , the arguments are analogous. The proof for  $u_{21}^*$ ,  $u_{22}^*$  is similar.  $\square$

## 5. A NUMERICAL EXAMPLE

It is known from simulations as well as from theoretical results that the estimators like  $u_{ij}^*$  behave quite well in linear autoregressive processes. To illustrate our method in non-linear models we considered the model

$$\begin{aligned} X_{t1} &= u_{11}h_{11}(X_{t-1,1}) + u_{12}h_{21}(X_{t-1,2}) + e_{t1}, \\ X_{t2} &= u_{21}h_{12}(X_{t-1,1}) + u_{22}h_{22}(X_{t-1,2}) + e_{t2} \end{aligned}$$

where

$$\begin{aligned} h_{11}(x) &= \begin{cases} x^2 & \text{for } x \in [0, 1), \\ \sqrt{x} & \text{for } x \geq 1, \end{cases} & h_{21}(x) &= \begin{cases} 0 & \text{for } x \in [0, 1), \\ 1 & \text{for } x \geq 1, \end{cases} \\ h_{12}(x) &= \begin{cases} x & \text{for } x \in [0, 1), \\ 1 & \text{for } x \geq 1, \end{cases} & h_{22}(x) &= \begin{cases} \frac{x}{2} & \text{for } x \in [0, 1), \\ 0.5 + \ln x & \text{for } x \geq 1 \end{cases} \end{aligned}$$

and  $e_{t1}$ ,  $e_{t2}$  are independent random variables with exponential distribution  $Ex(1)$ . A realization of  $X_{t1}$ ,  $X_{t2}$  of the length 500 was simulated with

$$u_{11} = 0.7, \quad u_{12} = 0.3, \quad u_{21} = 0.1, \quad u_{22} = 0.5.$$

The estimates  $u_{ij}^*$  were calculated from the first  $n$  variables of this realization ( $n = 50, 100, 200, 500$ ). The results are given in Tab. 1.

## Results of a simulation

$n$	$u_{11}^*$	$u_{12}^*$	$u_{21}^*$	$u_{22}^*$
50	0.782	0.180	0.000	0.634
100	0.763	0.217	0.146	0.452
200	0.707	0.288	0.134	0.462
500	0.701	0.300	0.100	0.502

Tab. 1

The estimates seem to be satisfactory for  $n = 200$  and the agreement with theoretical values is quite good for  $n = 500$ . Of course, more extensive simulations are needed to see the small sample properties of our estimators in detail.

### References

- [1] *H.-Z. An*: Non-negative autoregressive models. *J. Time Ser. Anal.* *13* (1992), 283–295.
- [2] *H.-Z. An, F. Huang*: Estimation for regressive and autoregressive models with non-negative residual errors. *J. Time Ser. Anal.* *14* (1993), 179–191.
- [3] *J. Anděl*: On AR(1) processes with exponential white noise. *Commun. Statist. – Theory Methods* *17* (1988), 1481–1495.
- [4] *J. Anděl*: Nonlinear nonnegative AR(1) processes. *Commun. Statist. – Theory Methods* *18* (1989), 4029–4037.
- [5] *J. Anděl*: Non-negative autoregressive processes. *J. Time Ser. Anal.* *10* (1989), 1–11.
- [6] *J. Anděl*: Nonlinear positive AR(2) processes. *Statistics* *21* (1990), 591–600.
- [7] *J. Anděl, V. Dupač*: An extension of the Borel lemma. *Comment. Math. Univ. Carolinae* *30* (1989), 403–404.
- [8] *B. Auestad, D. Tjøstheim*: Identification of nonlinear time series: First order characterization and order determination. *Biometrika* *77* (1990), 669–687.
- [9] *C. B. Bell, E. P. Smith*: Inference for non-negative autoregressive schemes. *Commun. Statist. – Theory Methods* *15* (1986), 2267–2293.
- [10] *R. Davis, W. McCormick*: Estimation for first-order autoregressive process with positive or bounded innovations. *Stochastic Processes Appl.* *31* (1989), 237–250.
- [11] *P. D. Feigin, S. I. Resnick*: Estimation for autoregressive processes with positive innovations. *Commun. Statist. – Stochastic Models* *8* (1992), 479–498.
- [12] *D. A. Jones*: Stationarity of non-linear autoregressive processes. *Tech. Rep.*, Institute of Hydrology, Wallingford, Oxon, U.K., 1977.
- [13] *H. Tong*: *Non-linear Time Series*. Clarendon Press, Oxford, 1990.

*Author's address*: Prof. RNDr. *Jiří Anděl*, DrSc., Matematicko-fyzikální fakulta University Karlovy (Faculty of Mathematics and Physics, Charles University), Sokolovská 83, 186 00 Praha 8, Czech Republic, E-mail: [andel@karlin.mff.cuni.cz](mailto:andel@karlin.mff.cuni.cz).