

Applications of Mathematics

Stephen Montgomery-Smith

Conditions implying regularity of the three dimensional Navier-Stokes equation

Applications of Mathematics, Vol. 50 (2005), No. 5, 451--464

Persistent URL: <http://dml.cz/dmlcz/134617>

Terms of use:

© Institute of Mathematics AS CR, 2005

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

CONDITIONS IMPLYING REGULARITY OF
THE THREE DIMENSIONAL NAVIER-STOKES EQUATION*

STEPHEN MONTGOMERY-SMITH, Columbia

(Received July 14, 2003, in revised version January 12, 2004)

Abstract. We obtain logarithmic improvements for conditions for regularity of the Navier-Stokes equation, similar to those of Prodi-Serrin or Beale-Kato-Majda. Some of the proofs make use of a stochastic approach involving Feynman-Kac-like inequalities. As part of our methods, we give a different approach to a priori estimates of Foias, Guillopé and Temam.

Keywords: Navier-Stokes equation, vorticity, Prodi-Serrin condition, Beale-Kato-Majda condition, Orlicz norm, stochastic method

MSC 2000: 35Q30, 76D05, 60H30, 46E30

1. INTRODUCTION

The version of the three dimensional Navier-Stokes equation we study is the differential equation in $u = u(t) = u(x, t)$, where $t \geq 0$ and $x \in \mathbb{R}^3$:

$$\frac{\partial u}{\partial t} = \Delta u - L \operatorname{div}(u \otimes u), \quad u(0) = u_0.$$

Here L denotes the Leray projection. We will not usually be working with classical solutions. We define $u(t)$, $0 \leq t \leq T$, to be a solution of the Navier-Stokes equation if, whenever $u(t_0)$ is sufficiently regular for a mild solution

$$u(t) = e^{(t-t_0)\Delta} u(t_0) - \int_{t_0}^t e^{(t-s)\Delta} L \operatorname{div}(u(s) \otimes u(s)) ds$$

to exist for $t \in [t_0, t_0 + \tau)$ for some $\tau > 0$, then $u(t)$ is equal to that mild solution in $[t_0, t_0 + \tau)$.

* The author was partially supported by an NSF grant.

We also use other ways to describe the three dimensional Navier-Stokes equation. First, let us denote the vorticity by $w = w(t) = w(x, t) = \text{curl } u$. If w is sufficiently smooth then

$$\frac{\partial w}{\partial t} = \Delta w - u \cdot \nabla w + w \cdot \nabla u, \quad w(0) = \text{curl } u_0.$$

Another description is given by the so called magnetization variable [4], [16]. Let $m = m(t) = m(x, t)$ be a vector field satisfying an equation

$$\frac{\partial m}{\partial t} = \Delta m - u \cdot \nabla m - m \cdot (\nabla u)^T, \quad m(0) = u_0 + \nabla q_0$$

for some scalar field $q_0 = q_0(x)$. (Here the superscript T denotes the transpose.) Then under sufficient smoothness assumptions we have that u is the Leray projection of m .

A famous open problem is to prove regularity of the Navier-Stokes equation, that is, if the initial data u_0 is in L_2 and is regular (which in this paper we define to mean that it is in the Sobolev spaces $W^{n,q}$ for some $2 \leq q < \infty$ and all positive integers n), then the solution $u(t)$ is regular for all $t \geq 0$. Such regularity would also imply uniqueness of the solution $u(t)$. Currently only the existence of weak solutions is known. Also, it is known that for each regular u_0 there exists $t_0 > 0$ such that $u(t)$ is regular for $0 \leq t \leq t_0$. We refer the reader to [3], [6], [7], [14], [21].

In studying this problem, various conditions that imply regularity have been obtained. For example, the Prodi-Serrin conditions ([17], [19]) state that for some $2 \leq p < \infty$, $3 < q \leq \infty$ with $2/p + 3/q \leq 1$,

$$\int_0^T \|u(t)\|_q^p dt < \infty$$

for all $T > 0$. If u is a weak solution to the Navier-Stokes equation satisfying a Prodi-Serrin condition with regular initial data u_0 , then u is regular (see [20]). (Recently Escoriaza, Seregin and Šverák [8] showed that the condition when $q = 3$ and $p = \infty$ is also sufficient.) This is a long way from what is currently known for the so called Leray-Hopf weak solutions:

$$\int_0^T \|u(t)\|_q^p dt < \infty$$

for $2/p + 3/q \geq 3/2$, $2 \leq q \leq 6$.

Another condition is that of Beale, Kato and Majda [1]. They show that regularity follows from the condition

$$\int_0^T \|w(t)\|_\infty dt < \infty$$

for all $T > 0$. (In fact they proved this for the Euler equation, but the proof works also for the Navier-Stokes equation with only small modifications.) This was

strengthened by Kozono and Taniuchi [12] to show that regularity follows from the condition

$$\int_0^T \|\nabla u(t)\|_{\text{BMO}} dt \approx \int_0^T \|w(t)\|_{\text{BMO}} dt < \infty$$

for all $T > 0$, where BMO denotes here the space of functions with bounded mean oscillation.

The purpose of this paper is threefold. First, we would like to provide some logarithmic improvements to these conditions. Secondly, we would like to present a stochastic approach to the Navier-Stokes equation, obtaining our conditions using Feynman-Kac-like inequalities. Thirdly, we would like to present a different process for creating estimates of Foias, Guillopé and Temam.

To this end, the first result of this paper is the logarithmic improvement to the Prodi-Serrin conditions.

Theorem 1.1. *Let $2 < p < \infty$, $3 < q < \infty$ with $2/p + 3/q = 1$. If u is a solution to the Navier-Stokes equation satisfying*

$$\int_0^T \frac{\|u(t)\|_q^p}{1 + \log^+ \|u(t)\|_q} dt < \infty$$

for some $T > 0$, then $u(t)$ is regular for $0 < t \leq T$.

We first present a proof of this result (and indeed of a slightly stronger result) that uses a standard approach. Then we present a stochastic approach to the Navier-Stokes equation. This is a kind of Lagrangian coordinates approach to the Navier-Stokes equation, but with a probabilistic twist in that we follow the path of each particle with a stochastic perturbation. A similar approach was adopted by Busnello, Flandoli and Romito [2].

From this we obtain the following Beale-Kato-Majda type condition. For $1 \leq q < \infty$, define the function on $[0, \infty)$

$$\Phi_q(\lambda) = \left(\frac{e^\lambda - 1}{e - 1} \right)^q.$$

Define the Φ_q -Orlicz norm on any space of measurable functions by the formula

$$\|f\|_{\Phi_q} = \inf \left\{ \lambda > 0: \int \Phi_q(|f(x)|/\lambda) dx \leq 1 \right\}.$$

(Thus the triangle inequality is a consequence of the fact that Φ_q is convex, see [13].)

Theorem 1.2. *Let $1 < q < \infty$, $3 < r < \infty$, and $T > 0$. Suppose that u is a solution to the Navier-Stokes equation satisfying*

(1) *for all $T_0 \in (0, T)$,*

$$\int_{T_0}^T \|\nabla u(t)\|_{\Phi_q} dt < \infty,$$

and

(2) *either $q < 3$, or $\|u(t)\|_r < \infty$ for almost every $t \in [0, T]$.*

Then $u(t)$ is regular for $0 < t \leq T$.

Note that since $\|\cdot\|_{\Phi_{q_1}} \leq c\|\cdot\|_{\Phi_{q_2}}$ for $q_1 > q_2$, we may assume without loss of generality that $q > 3/2$. Next, if $3/2 < q < 3$, since $\|\cdot\|_q \leq (e-1)\|\cdot\|_{\Phi_q}$, by the Sobolev inequality we see that the second hypothesis is automatically satisfied with $r = 3q/(3-q)$. Also, this hypothesis is always satisfied for Leray-Hopf weak solutions with $r = 6$.

Next we demonstrate how to obtain Theorem 1.1 from Theorem 1.2 using the following result. If u is a solution to the Navier-Stokes equation, we define sets

$$A_{T_0, T_1}^{n, q}(\lambda) = \{t \in [T_0, T_1] : \|\nabla^n u(t)\|_q \geq \lambda\}.$$

Theorem 1.3. *Given $3 < q_1 \leq q_2 \leq \infty$ and a non-negative integer n , there exists constants $c_1, c_2, c_3 > 0$ such that if $u(t)$, $0 \leq t \leq T_2$, is a solution to the Navier-Stokes equation and if $0 \leq T_1 \leq T_2$, then for all $r \in (0, \sqrt{T_2 - T_1})$ we have*

$$|A_{T_1+r^2, T_2}^{n, q_2}(c_1 r^{3/q_2-n-1})| \leq c_2 |A_{T_1, T_2}^{0, q_1}(c_3 r^{3/q_1-1})|.$$

A similar result that one can obtain (but we do not prove here) is that for positive integers n we have $|A_{T_1+r^2, T_2}^{n, 2}(c_1 r^{1/2-n})| \leq c_2 |A_{T_1, T_2}^{1, 2}(c_3 r^{-1/2})|$.

Corollary 1.4. *Under the hypotheses of Theorem 1.3, there exists a constant $c > 0$ with the following properties. If $\Theta(\lambda)$ is a positive increasing function of $\lambda \geq 0$, define*

$$\kappa = \int_0^\infty \min\{(c\lambda^{-2} - T_0)^+, T_1\} d\Theta(\lambda).$$

Then

$$\int_{T_0}^{T_1} \Theta(\|\nabla^n u(s)\|_{q_2}^{1/(1+n-3/q_2)}) ds \leq c\kappa + c \int_0^{T_1} \Theta(c\|u(s)\|_{q_1}^{1/(1-3/q_1)}) ds.$$

Similarly,

$$\int_{T_0}^{T_1} \Theta(\|\nabla^n u(s)\|_2^{1/(n-1/2)}) ds \leq c\kappa + c \int_0^{T_1} \Theta(c\|\nabla u(s)\|_2^2) ds.$$

Since the Leray-Hopf weak solution to the Navier-Stokes equation satisfies $\int_0^T \|\nabla u(t)\|_2^2 dt < \infty$, one can quickly recover the results of Foias, Guillopé and Temam [9] that say that $\int_0^T \|\nabla^n u(t)\|_2^{1/(n-1/2)} dt < \infty$.

2. THEOREM 1.1

The hypothesis of Theorem 1.1 implies that, given $\varepsilon \in (0, T)$, there exists $T_0 \in (0, \varepsilon)$ with $u(T_0) \in L_q$. Let $T^* > T_0$ be the first point of non-regularity for $u(t)$. It is well known that in order to show that $T^* > T$, it is sufficient to show an *a priori* estimate, that is $\sup_{T_0 \leq t < \min\{T^*, T\}} \|u(t)\|_q < \infty$. This is because it is then possible to extend the regularity beyond T^* if $T^* \leq T$. Without loss of generality, it is sufficient to consider the case $T = T^*$ (so as to obtain a contradiction).

Proof of Theorem 1.1. We allow all constants to implicitly depend upon p and q . Let us define quantities

$$\begin{aligned} v &= u|u|^{q/2-1}, \\ A &= \sum_{i,j=1}^3 \left(|u|^{q/2-1} \frac{\partial u_i}{\partial x_j} \right)^2, \\ B &= \sum_{i,j=1}^3 \left(|u|^{q/2-3} u_i \sum_{k=1}^3 u_k \frac{\partial u_k}{\partial x_j} \right)^2. \end{aligned}$$

Note that

$$\begin{aligned} |\nabla v|^2 &:= \sum_{i,j=1}^3 \left(\frac{\partial v_i}{\partial x_j} \right)^2 \approx A + B, \\ \sum_{i,j=1}^3 \frac{\partial}{\partial x_j} (|u|^{q-2} u_i) \frac{\partial u_i}{\partial x_j} &\approx A + B, \\ \sum_{i,j=1}^3 \left(\frac{\partial}{\partial x_j} (|u|^{q-2} u_i) \right)^2 &\leq c |u|^{q-2} |\nabla v|^2. \end{aligned}$$

We start with the Navier-Stokes equation, take the inner product with $u|u|^{q-2}$, and integrate over \mathbb{R}^3 to obtain

$$|u|_q^{q-1} \frac{\partial}{\partial t} \|u\|_q = \int |u|^{q-2} u \cdot \Delta u \, dx - \int |u|^{q-2} u \cdot L \operatorname{div}(u \otimes u) \, dx.$$

Integrating by parts, we see that

$$\int |u|^{q-2} u \cdot \Delta u \, dx = - \int \sum_{i,j=1}^3 \frac{\partial}{\partial x_j} (|u|^{q-2} u_i) \frac{\partial u_i}{\partial x_j} \, dx \approx - \|\nabla v\|_2^2$$

and

$$\begin{aligned} \int |u|^{q-2} u \cdot L \operatorname{div}(u \otimes u) \, dx &= \int \sum_{i,j=1}^3 \frac{\partial}{\partial x_j} (|u|^{q-2} u_i) [L(u_j u)]_i \, dx \\ &\leq c \| |u|^{q/2-1} \|_s \| \nabla v \|_2 \| L(u \otimes u) \|_r \end{aligned}$$

where $r = 1 + q/2$ and $s = (2q + 4)/(q - 2)$. Now the Leray projection is a bounded operator on L_r , and hence $\|L(u \otimes u)\|_r \approx \|u\|_{2+q}^2$. Also $\| |u|^{q/2-1} \|_s \approx \|u\|_{2+q}^{q/2-1}$. Hence

$$\int |u|^{q-2} u \cdot L \operatorname{div}(u \otimes u) \, dx \leq c \|u\|_{2+q}^{1+q/2} \| \nabla v \|_2 = c \|v\|_{2+4/q}^{1+2/q} \| \nabla v \|_2.$$

From the Sobolev and interpolation inequalities we obtain

$$\|v\|_{2+4/q} \leq c \| |\nabla|^{3/(q+2)} v \|_2 \leq c \|v\|_2^{(q-1)/(q+2)} \| \nabla v \|_2^{3/(q+2)},$$

and hence

$$\int |u|^{q-2} u \cdot L \operatorname{div}(u \otimes u) \, dx \leq c \|v\|_2^{1-1/q} \| \nabla v \|_2^{1+3/q}.$$

Now apply Young's inequality $ab \leq ((q-3)a^{2q/(q-3)} + (q+3)b^{2q/(q+3)})/2q$ for $a, b \geq 0$ to obtain

$$\int |u|^{q-2} u \cdot L \operatorname{div}(u \otimes u) \, dx \leq c_1 \| \nabla v \|_2^2 + c_2 \|v\|_2^{2(q-1)/(q-3)},$$

where c_1 may be made as small as required by making c_2 larger. Hence

$$\|u\|_q^{q-1} \frac{\partial}{\partial t} \|u\|_q \leq c \|v\|_2^{2(q-1)/(q-3)},$$

that is,

$$\frac{\partial}{\partial t} \|u\|_q \leq c \|u\|_q^{p+1},$$

and so

$$\frac{\partial}{\partial t} \log(1 + \log^+ \|u\|_q) \leq \frac{c \|u\|_q^p}{1 + \log^+ \|u\|_q}.$$

Integrating, we see that for $T_0 \leq t < T$

$$\begin{aligned} \log(1 + \log^+ \|u(t)\|_q) &\leq \log(1 + \log^+ \|u(T_0)\|_q) \\ &\quad + c \int_{T_0}^T \frac{\|u(s)\|_q^p}{1 + \log^+ \|u(s)\|_q} \, ds, \end{aligned}$$

which provides a uniform bound for $\|u(t)\|_q$. □

Remark 2.1. Note that this proof can easily be adapted to show that a sufficient condition for regularity is that

$$\int_0^T \frac{\|u(s)\|_q^p}{\Theta(\|u(s)\|_q)} ds < \infty,$$

where Θ is any increasing function for which

$$\int_1^\infty \frac{1}{x\Theta(x)} dx = \infty.$$

3. A PRIORI ESTIMATES

This section is devoted to the proof of Theorem 1.3 and Corollary 1.4. The proof is very similar to the proof of Scheffer's Theorem [18] that states that the Hausdorff dimension of the set of t for which the solution $u(t)$ is not regular is $1/2$. The main tool is the following result due to Grujić and Kukavica [10] (see also [15]).

Theorem 3.1. *There exist constants $a, c > 0$ and a function $T: (0, \infty) \rightarrow (0, \infty)$ with $T(\lambda) \rightarrow \infty$ as $\lambda \rightarrow 0$, with the following properties. If $u_0 \in L_q(\mathbb{R}^3)$, then there is a solution $u(t)$ ($0 \leq t \leq T(\|u_0\|_q)$) to the Navier-Stokes equation with $u(0) = u_0$, and $u(x, t)$ is the restriction of an analytic function $u(x + iy, t) + iv(x + iy, t)$ in the region $\{x + iy \in \mathbb{C}^3: |y| \leq a\sqrt{t}\}$, and $\|u(\cdot + iy, t) + iv(\cdot + iy, t)\|_q \leq c\|u_0\|_q$ for $|y| \leq a\sqrt{t}$.*

Proof of Theorem 1.3. First let us show that there exist constants $c_1, c_3, c_4 > 0$ such that if $u(t)$, $t_0 - r^2 \leq t \leq t_0$, is a solution to the Navier-Stokes equation and $|A_{t_0-r^2, t_0}^{0, q_1}(c_3 r^{3/q_1-1})| < c_4 r^2$, then $\|\nabla^n u(t_0)\|_{q_2} < c_1 r^{3/q_2-n-1}$.

To see this, let us first consider the case when $t_0 = 0$ and $r = 1$. By hypothesis, we see that there exists $t \in [-1, -1 + c_4]$ with $\|u(t)\|_{q_1} < c_3$. By Theorem 3.1 and the appropriate Cauchy integrals, if c_4 is small enough, then there exists a constant $c_1 > 0$ such that $\|\nabla^n u(0)\|_{q_2} < c_1$.

Now, by replacing $u(x, t)$ by $r^{-1}u(r^{-1}x, r^{-2}(t - t_0))$, we can relax the restriction $r = 1$ and $t_0 = 0$, and we obtain the statement we asserted.

Next, given $\varepsilon > 0$, it is trivial to find a finite collection t_1, \dots, t_N in $A = A_{T_1+r^2, T_2}^{n, q_2}(c_1 r^{3/q_2-n-1})$ such that the sets $[t_n - r^2, t_n]$ are disjoint, but the sets $[t_n - r^2 - \varepsilon, t_n + \varepsilon]$ cover A . By the above observation, $|A_{t_0-r^2, t_0}^{0, q_1}(c_3 r^{3/q_1-1})| \geq c_4 r^2$.

Hence

$$\begin{aligned} \frac{r^2}{r^2 + 2\varepsilon} |A| &\leq N r^2 < c_4^{-1} \sum_{n=1}^N |A_{t_n-r^2, t_n}^{0, q_1}(c_3 r^{3/q_1-1})| \\ &\leq c_4^{-1} |A_{T_1, T_2}^{0, q_1}(c_3 r^{3/q_1-1})|. \end{aligned}$$

Since ε is arbitrary, the result follows. □

Proof of Corollary 1.4. We only prove the first inequality. By Theorem 1.3, there exist constants $c_1, c_2, c_3 > 0$ such that

$$\begin{aligned}
 & \int_{T_0}^{T_1} \Theta(\|\nabla^n u(s)\|_{q_2}^{1/(1+n-3/q_2)}) \, ds \\
 &= \int_0^\infty |\{s \in [T_0, T_1]: \|\nabla^n u(s)\|_{q_2}^{1/(1+n-3/q_2)} > \lambda\}| \, d\Theta(\lambda) \\
 &\leq c_1 \kappa + \int_0^\infty |\{s \in [c_2 \lambda^{-2}, T_1]: \|\nabla^n u(s)\|_{q_2}^{1/(1+n-3/q_2)} > \lambda\}| \, d\Theta(\lambda) \\
 &\leq c_1 \kappa + c_1 \int_0^\infty |\{s \in [0, T_1]: \|u(s)\|_{q_1}^{1/(1-3/q_1)} > c_3 \lambda\}| \, d\Theta(\lambda) \\
 &= c_1 \kappa + c_1 \int_0^{T_1} \Theta(c_3^{-1} \|u(s)\|_{q_1}^{1/(1-3/q_1)}) \, ds.
 \end{aligned}$$

□

4. A STOCHASTIC DESCRIPTION

Let us give a little motivation. Suppose that we defined $\varphi_{t_0, t_1}(x)$ to be $X(t_0)$, where X satisfies the equation

$$dX(t) = u(X(t), t) \, dt, \quad X(t_1) = x,$$

then φ_{t_0, t_1} would be the “back to coordinates map” that takes a point at $t = t_1$ to where it was carried from by the flow of the fluid at time $t = t_0$. For the Euler equation, this provides a very effective way to describe the solution, for example, the equation for vorticity can be rewritten in a Lagrangian form:

$$w(x, t) = w(\varphi_{0, t}(x), 0) + \int_0^t w(\varphi_{s, t}(x), s) \cdot \nabla u(\varphi_{s, t}(x), s) \, ds.$$

Similarly, for the magnetization variable we have

$$m(x, t) = m(\varphi_{0, t}(x), 0) - \int_0^t m(\varphi_{s, t}(x), s) \cdot (\nabla u(\varphi_{s, t}(x), s))^T \, ds.$$

For the Navier-Stokes equation this formula is not true, and the Laplacian term can make things complicated. One approach to dealing with this is described in the paper by Constantin [5]. However, we take a different approach using Brownian motion, using a kind of “randomly perturbed back to coordinates map.” Such a method was already discussed in the paper [16], here we make the discussion more rigorous. The

author recently found out that a similar approach was followed by Busnello, Flandoli and Romito in [2].

The hypotheses of Theorem 1.2 imply that, given $\varepsilon \in (0, T)$, there exists $t' \in (0, \varepsilon)$ with $u(t') \in L_r$. Then by known results (for example Theorem 3.1), it follows that there exists $0 < T_0 < \varepsilon$ such that $u(T_0) \in W^{n, r'}$ for all $r' \in [r, \infty]$ and all positive integers n . Furthermore, arguing as in Section 2, we only need to prove $\sup_{T_0 \leq t < \min\{T^*, T\}} \|u(t)\|_r < \infty$ under the *a priori* assumption that the solution is regular for $t \in [T_0, T]$.

If $f: \mathbb{R}^3 \rightarrow \mathbb{R}$ is regular and $T_0 \leq t_0 \leq t_1 < T$, define $A_{t_0, t_1} f(x) = \alpha(x, t_1)$, where α satisfies the transport equation

$$\frac{\partial \alpha}{\partial t} = \Delta \alpha - u \cdot \nabla \alpha, \quad \alpha(x, t_0) = f(x).$$

Since $\operatorname{div}(u) = 0$, an easy integration by parts argument shows that

$$\frac{\partial}{\partial t} \int \alpha(x, t) \, dx = 0,$$

and hence if f is also in L_1 , then

$$\int A_{t_0, t_1} f(x) \, dx = \int f(x) \, dx.$$

Since stochastic differential equations traditionally move forwards in time, it will be convenient to consider a time reversed equation. Let $b(t)$ be a three dimensional Brownian motion. For $T_0 \leq t_0 \leq t_1 < T_1$, define the random function $\varphi_{t_0, t_1}: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ by $\varphi_{t_0, t_1}(x) = X(-t_0)$, where X satisfies the stochastic differential equation

$$dX(t) = -u(X(t), t) \, dt + \sqrt{2} \, db(t), \quad X(-t_1) = x.$$

It follows by the Itô Calculus [11] that if $T_0 \leq t_0 \leq t_1 < T$, then

$$A_{t_0, t_1} f(x) = \mathbb{E}f(\varphi_{t_0, t_1}(x)).$$

(Here as in the rest of the paper, \mathbb{E} denotes the expected value.) Note that if f is also in L_1 , then

$$\int \mathbb{E}f(\varphi_{t_0, t_1}(x)) \, dx = \int f(x) \, dx.$$

Applying the usual dominated and monotone convergence theorems, it quickly follows that the last equality is also true if f is any function in L_1 , or if f is any positive function.

Now let us develop the equations for the magnetization variable. (The same approach will also work for the vorticity.) If we set $m(T_0) = u(T_0)$, then we note that m is the unique solution to the integral equation

$$m(t) = A_{T_0,t}u(T_0) - \int_{T_0}^t A_{s,t}(m(s) \cdot (\nabla u(s))^T) ds \quad (T_0 \leq t < T).$$

Uniqueness follows quickly by the usual fixed point argument over short intervals, remembering that $u(t)$ is regular for $T_0 \leq t < T$.

Consider also the random quantity $\tilde{m} = \tilde{m}(x, t)$ as the solution to the integral equation for $T_0 \leq t < T$

$$\tilde{m}(x, t) = u(\varphi_{T_0,t}(x), T_0) - \int_{T_0}^t \tilde{m}(\varphi_{s,t}(x), s) \cdot (\nabla u(\varphi_{s,t}(x), s))^T ds.$$

Again, it is very easy to show that a solution exists by using a fixed point argument over short time intervals. It is seen that $\mathbb{E}\tilde{m}$ satisfies the same equation as m , and hence $\mathbb{E}\tilde{m} = m$.

Next, $\varphi_{t_0,t_1}(\varphi_{t_1,t_2}(x)) = \varphi_{t_0,t_2}(x)$, since both are $Y(t_0)$ where $Y(t)$ is the solution to the integral equation

$$Y(t) = \varphi_{t_1,t_2}(x) + \int_{t_1}^t u(Y(s), s) ds + \sqrt{2}(b_{-t} - b_{-t_1}).$$

Hence

$$\tilde{m}(\varphi_{s_1,t}(x), s_1) - \tilde{m}(\varphi_{s_2,t}(x), s_2) = \int_{s_1}^{s_2} \tilde{m}(\varphi_{s,t}(x), s) \cdot (\nabla u(\varphi_{s,t}(x), s))^T ds.$$

Thus, by Gronwall's inequality, if $T_0 \leq t < T$ then

$$|\tilde{m}(x, t)| \leq \exp\left(\int_{T_0}^t |\nabla u(\varphi_{s,t}(x), s)| ds\right) |u(\varphi_{T_0,t}(x), T_0)|.$$

(This is essentially the Feynman-Kac formula.) The goal, then, is to find uniform estimates on the quantity

$$\exp\left(\int_{T_0}^t |\nabla u(\varphi_{s,t}(x), s)| ds\right).$$

This we proceed to do in the next section.

5. THEOREM 1.2

Let us fix q and r satisfying the hypotheses of Theorem 1.2, and allow all constants to implicitly depend upon q and r . We retain the notation from the previous section, in particular the definitions of T_0 , T^* and T .

P r o o f of Theorem 1.2. Since $\|u(t)\|_r < \infty$ for almost every $t \in [0, T]$, by Theorem 1.3, we see that $\|\nabla u(t)\|_\infty < \infty$ for almost every $t \in [0, T]$. Hence, there exists $\lambda > T_0^{-1}$ such that

$$\int_B \|\nabla u(t)\|_{\Phi_q} dt \leq \frac{1}{q},$$

where $B = \{t \in [T_0, T] : \|\nabla u(t)\|_\infty \geq c_2 \lambda\}$. Thus for $T_0 \leq t < T$, we have that $|\tilde{m}(x, t)|$ is bounded by

$$e^{c_2 \lambda(t-T_0)} \exp\left(\int_{B \cap [T_0, t]} |\nabla u(\varphi_{s,t}(x), s)| ds\right) |u(\varphi_{T_0,t}(x), T_0)|.$$

Hence by Jensen's and Hölder's inequalities,

$$\|m(t)\|_r^r \leq \int \mathbb{E}|\tilde{m}(t)|^r dx \leq e^{c_2 q \lambda(t-T_0)} (N_r^r + N_{rq'}^r \tilde{N}^r),$$

where $q' = q/(q-1)$,

$$N_s = \left(\int \mathbb{E}|u(\varphi_{T_0,t}(x), T_0)|^s dx\right)^{1/s} = \|u(T_0)\|_s,$$

and

$$\tilde{N}^q = \int \mathbb{E}\left(\exp\left(q \int_{B \cap [T_0, t]} \|\nabla u(\varphi_{s,t}(x), s)\| ds\right) - 1\right)^q dx.$$

Since the Orlicz norm satisfies the triangle inequality, we have

$$\left\| \int_{B \cap [T_0, t]} |\nabla u(\varphi_{s,t}(\cdot), s)| ds \right\|_{\Phi_q} \leq \frac{1}{q},$$

that is, $\tilde{N} \leq e - 1$. Since $a^r + b^r \leq (a+b)^r$ for $a, b \geq 0$, we conclude that

$$\|m(t)\|_r \leq \|u(T_0)\|_r + (e-1)e^{c_2 \lambda(t-T_0)} \|u(T_0)\|_{rq'}.$$

As the Leray projection is a bounded operator on L_r for $1 < r < \infty$, it follows that $\|u(t)\|_r$ is also uniformly bounded, and the result follows. \square

A second proof of Theorem 1.1 now follows from the next result.

Lemma 5.1. *There is a constant $c > 0$ such that if f is a measurable function, then*

$$\|f\|_{\Phi_q} \leq c \left(\|f\|_q + \frac{\|f\|_\infty}{1 + \Phi_q^{-1}((\|f\|_\infty/\|f\|_q)^q)} \right).$$

Proof. Let us assume that $\|f\|_\infty = 1$, and set $a = \|f\|_q$, $b = \Phi_q^{-1}(a^{-q})$ and $n = a + 1/(1 + b)$. Let $f^*: [0, \infty] \rightarrow [0, \infty]$ be the non-increasing rearrangement of $|f|$, that is,

$$f^*(t) = \sup\{\lambda > 0: |\{x: |f(x)| > \lambda\}| > t\},$$

so $\int F(|f(x)|) dx = \int_0^\infty F(f^*(t)) dt$ for any Borel measurable function F . Notice that $f^*(t) \leq \min\{1, at^{-1/q}\}$.

Let us first consider the case $a \leq 1$, so that $b \geq 1$, $2n \geq 1/b$ and $n \geq a$. Then

$$\int \Phi_q(|f(x)|/2n) dx \leq \int_0^\infty \Phi_q(f^*(t)/2n) dt.$$

We split this integral up into three pieces. First,

$$\int_0^{a^q} \Phi_q(f^*(t)/2n) dt \leq \int_0^{a^q} \Phi_q(b) dt = 1.$$

Next, since $(\Phi_q(\lambda))^{1/2q}$ is convex for $\lambda \geq 1$,

$$\begin{aligned} \int_{a^q}^{a^q b^q} \Phi_q(f^*(t)/2n) dt &\leq \int_{a^q}^{a^q b^q} \Phi_q(abt^{-1/q}) dt \\ &\leq \int_{a^q}^{a^q b^q} \frac{a^{2q}\Phi_q(b)}{t^2} dt \leq 1. \end{aligned}$$

Next, for $t \geq a^q b^q$, $f^*(t) \leq 1/b \leq 2n$, and $\Phi_q(\lambda) \leq \lambda^q$ for $0 \leq \lambda \leq 1$, we have

$$\int_{a^q b^q}^\infty \Phi_q(f^*(t)/2n) dt \leq \int_{a^q b^q}^\infty (f^*(t)/2n)^q dt \leq 1.$$

Since $\Phi_q(\lambda/3) \leq \Phi_q(\lambda)/3$ for $\lambda \geq 0$, we arrive at

$$\int \Phi_q(|f(x)|/6n) dx \leq 1,$$

that is, $\|f\|_{\Phi_q} \leq 6n$.

The case $a \geq 1$ (so $b \leq 1$ and $2n \geq 1 + 2a$) is simpler, as it is easy to estimate

$$\int_0^\infty \Phi_q(f^*(t)/2n) dt \leq \int_0^1 \Phi_q(1) dt + \int_1^\infty (f^*(t)/2n)^q dt \leq 2.$$

□

Second proof of Theorem 1.1. Applying Corollary 1.4 using the function

$$\Theta(\lambda) = \frac{\lambda^2}{1 + \log^+ \lambda},$$

we obtain for all $T_0 \in (0, T)$

$$\int_{T_0}^T \frac{\|\nabla u(s)\|_\infty}{1 + \log^+ \|\nabla u(s)\|_\infty} ds < \infty$$

and

$$\int_{T_0}^T \frac{\|\nabla u(s)\|_q^{2q/(2q-3)}}{1 + \log^+ \|\nabla u(s)\|_q} ds < \infty.$$

Hence if $1 < \alpha < 2q/(2q - 3)$ we have that

$$\int_{T_0}^T \|\nabla u(s)\|_q^\alpha ds < \infty.$$

Next, considering the cases $\|f\|_\infty > \|f\|_q^\alpha$ and $\|f\|_\infty \leq \|f\|_q^\alpha$, we see that

$$\frac{\|f\|_\infty}{1 + \Phi_q^{-1}((\|f\|_\infty/\|f\|_q)^q)} \leq c \left(\|f\|_q^\alpha + \frac{\|f\|_\infty}{1 + \log^+ \|f\|_\infty} \right).$$

Applying Lemma 5.1, we see that the hypothesis of Theorem 1.1 implies the hypotheses of Theorem 1.2 with $q = r$. \square

Acknowledgments.

The author wishes to extend his sincere gratitude to Michael Taksar for his help with understanding stochastic processes, and also to Pierre-Gilles Lemarié-Rieusset for very helpful e-mail discussions.

References

- [1] *J. T. Beale, T. Kato, and A. Majda*: Remarks on the breakdown of smooth solutions for the 3-D Euler equations. *Comm. Math. Phys.* 94 (1984), 61–66.
- [2] *B. Busnello, F. Flandoli, and M. Romito*: A probabilistic representation for the vorticity of a 3D viscous fluid and for general systems of parabolic equations. Preprint, <http://arxiv.org/abs/math/0306075>.
- [3] *M. Cannone*: Wavelets, paraproducts and Navier-Stokes. Diderot Editeur, Paris, 1995. (In French.)
- [4] *A. Chorin*: Vorticity and Turbulence. *Appl. Math. Sci.*, Vol. 103. Springer-Verlag, New York, 1994.

- [5] *P. Constantin*: An Eulerian-Lagrangian approach to the Navier-Stokes equations. *Commun. Math. Phys.* 216 (2001), 663–686.
- [6] *P. Constantin, C. Foias*: Navier-Stokes Equations. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, 1988.
- [7] *C. R. Doering, J. D. Gibbon*: Applied Analysis of the Navier-Stokes Equations. Cambridge Texts in Applied Mathematics. Cambridge University Press, Cambridge, 1995.
- [8] *L. Escauriaza, G. Seregin, and V. Šverák*: On $L_{3,\infty}$ -solutions to the Navier-Stokes equations and backward uniqueness. <http://www.ima.umn.edu/preprints/dec2002/dec2002.html>.
- [9] *C. Foias, C. Guillopé, and R. Temam*: New a priori estimates for Navier-Stokes equations in dimension 3. *Commun. Partial Differ. Equations* 6 (1981), 329–359.
- [10] *Z. Grujić, I. Kukavica*: Space analyticity for the Navier-Stokes and related equations with initial data in L^p . *J. Funct. Anal.* 152 (1998), 447–466.
- [11] *I. Karatzas, S. E. Shreve*: Brownian Motion and Stochastic Calculus, second edition. Graduate Texts in Mathematics Vol. 113. Springer-Verlag, New York, 1991.
- [12] *H. Kozono, Y. Taniuchi*: Bilinear estimates in BMO and the Navier-Stokes equations. *Math. Z.* 235 (2000), 173–194.
- [13] *M. A. Krasnosel'skiĭ, Ya. B. Rutitskiĭ*: Convex Functions and Orlicz Spaces. Translated from the first Russian edition. P. Noordhoff, Groningen, 1961.
- [14] *P. G. Lemarié-Rieusset*: Recent Developments in the Navier-Stokes Problem. Chapman and Hall/CRC, Boca Raton, 2002.
- [15] *P. G. Lemarié-Rieusset*: Further remarks on the analyticity of mild solutions for the Navier-Stokes equations in \mathbb{R}^3 . *C. R. Math. Acad. Sci. Paris* 338 (2004), 443–446. (In French.)
- [16] *S. J. Montgomery-Smith, M. Pokorný*: A counterexample to the smoothness of the solution to an equation arising in fluid mechanics. *Comment. Math. Univ. Carolin.* 43 (2002), 61–75.
- [17] *G. Prodi*: Un teorema di unicità per le equazioni di Navier-Stokes. *Ann. Mat. Pura Appl.* 48 (1959), 173–182. (In Italian.)
- [18] *V. Scheffer*: Turbulence and Hausdorff Dimension. Turbulence and Navier-Stokes Equations (Proc. Conf., Univ. Paris-Sud, Orsay, 1975). *Lect. Notes Math.* Vol. 565. Springer-Verlag, Berlin, 1976, pp. 174–183.
- [19] *J. Serrin*: On the interior regularity of weak solutions of the Navier-Stokes equations. *Arch. Ration. Mech. Anal.* 9 (1962), 187–195.
- [20] *H. Sohr*: Zur Regularitätstheorie der instationären Gleichungen von Navier-Stokes. *Math. Z.* 184 (1983), 359–375.
- [21] *R. Temam*: Infinite-Dimensional Dynamical Systems in Mechanics and Physics, second edition. Applied Mathematical Sciences Vol. 68. Springer-Verlag, New York, 1997.

Author's address: S. Montgomery-Smith, Department of Mathematics, University of Missouri, Columbia, MO 65211, U.S.A., e-mail: stephen@math.missouri.edu.