

# Applications of Mathematics

---

Albo Carlos Cavalheiro

The Neumann problem for some degenerate elliptic equations

*Applications of Mathematics*, Vol. 51 (2006), No. 6, 619--628

Persistent URL: <http://dml.cz/dmlcz/134657>

## Terms of use:

© Institute of Mathematics AS CR, 2006

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

THE NEUMANN PROBLEM FOR SOME DEGENERATE  
ELLIPTIC EQUATIONS

ALBO CARLOS CAVALHEIRO, Londrina

(Received April 21, 2005, in revised version August 22, 2005)

*Abstract.* In the paper we study the equation  $Lu = f$ , where  $L$  is a degenerate elliptic operator, with Neumann boundary condition in a bounded open set  $\Omega$ . We prove existence and uniqueness of solutions in the space  $H(\Omega)$  for the Neumann problem.

*Keywords:* Neumann problem, degenerate elliptic equations

*MSC 2000:* 35J70, 35J25

1. INTRODUCTION

In this paper we prove existence and uniqueness of solutions in the space  $H(\Omega)$  (see Definition 2.2) for the Neumann problem

$$(P) \quad \begin{cases} Lu(x) = f(x) & \text{on } \Omega, \\ \langle \mathcal{A}(x)\nabla u(x), \vec{\eta}(x) \rangle = 0 & \text{on } \partial\Omega, \end{cases}$$

where  $L$  is a degenerate elliptic operator

$$(1.1) \quad Lu(x) = - \sum_{i,j=1}^n D_j(a_{ij}(x)D_i u(x)) + \sum_{i=1}^n b_i(x)D_i u(x) + g(x)u(x) + \theta u(x)v(x)$$

with  $D_j = \partial/\partial x_j$  ( $j = 1, \dots, n$ ),  $\theta$  is a constant, the coefficients  $a_{ij}$ ,  $b_i$  and  $g$  are measurable, real-valued functions, the coefficient matrix  $\mathcal{A}(x) = (a_{ij}(x))$  is symmetric and satisfies the degenerate ellipticity condition

$$(1.2) \quad |\xi|^2\omega(x) \leq \langle \mathcal{A}(x)\xi, \xi \rangle \leq |\xi|^2v(x)$$

for all  $\xi \in \mathbb{R}^n$  and almost every  $x \in \Omega \subset \mathbb{R}^n$  where  $\Omega$  is a bounded open set with piecewise smooth boundary (i.e.,  $\partial\Omega \in C^{0,1}$ ),  $\omega$  and  $v$  are weight functions (that is, locally integrable and nonnegative functions on  $\mathbb{R}^n$ ),  $\vec{\eta}(x) = (\eta_1(x), \dots, \eta_m(x))$  is the unit outward normal to  $\partial\Omega$  at  $x$ ,  $\langle \cdot, \cdot \rangle$  denotes the usual inner product in  $\mathbb{R}^n$  and the symbol  $\nabla$  indicates the gradient.

In general, the Sobolev spaces  $W^{k,p}(\Omega)$  without weights occur as spaces of solutions for elliptic and parabolic partial differential equations. For degenerate partial differential equations, i.e., equations with various types of singularities in the coefficients, it is natural to look for solutions in weighted Sobolev spaces (see [1], [2] and [3]).

A class of weights which is particularly well understood, is the class of  $A_p$ -weights (or Muckenhoupt class) that was introduced by B. Muckenhoupt (see [4]). These weights have found many useful applications in harmonic analysis (see [5]). Another reason for studying  $A_p$ -weights is the fact that powers of distance to submanifolds of  $\mathbb{R}^n$  often belong to  $A_p$  (see [6] or [7]). There are, in fact, many interesting examples of weights (see [8] for  $p$ -admissible weights). In this paper we will consider only  $A_p$ -weights.

The following theorem will be proved in Section 3.

**Theorem 1.1.** *Let  $\Omega \subset \mathbb{R}^n$  be a bounded open set with boundary  $\partial\Omega \in C^{0,1}$ . Suppose that*

- (H1)  $\omega \in A_2$ ,  $v \in A_2$  and (1.2) holds;
- (H2)  $f/v \in L^2(\Omega, v)$ ;
- (H3)  $b_i/\omega \in L^\infty(\Omega)$  ( $i = 1, \dots, n$ ) and  $g/v \in L^\infty(\Omega)$ .

*Then there exists a constant  $\mathbf{C} > 0$  such that for all  $\theta \geq \mathbf{C}$  the Neumann problem (P) has a unique solution  $u \in H(\Omega)$ . Moreover, we have*

$$\|u\|_{H(\Omega)} \leq 2 \left\| \frac{f}{v} \right\|_{L^2(\Omega, v)}.$$

**Example 1.2.** Consider the domain  $\Omega = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 < 1\}$ , the weight functions

$$\omega(x, y) = (x^2 + y^2)^{-1/3} \quad \text{and} \quad v(x, y) = (x^2 + y^2)^{-1/2}$$

and the coefficient matrix

$$\mathcal{A}(x, y) = \begin{pmatrix} (x^2 + y^2)^{-1/3} & 0 \\ 0 & (x^2 + y^2)^{-1/2} \end{pmatrix}.$$

For all  $\xi \in \mathbb{R}^2$  and almost every  $(x, y) \in \Omega$  we have

$$\frac{1}{(x^2 + y^2)^{1/3}} |\xi|^2 \leq \langle \mathcal{A}(x, y) \xi, \xi \rangle \leq \frac{1}{(x^2 + y^2)^{1/2}} |\xi|^2.$$

If  $(x, y) \in \partial\Omega = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1\}$ , then  $\vec{\eta}(x, y) = (x, y)$  is the unit outward normal to  $\partial\Omega$ . By Theorem 1.1 the Neumann problem

$$\begin{cases} Lu(x, y) = (x^2 + y^2)^{-3/8} \cos(xy) & \text{on } \Omega, \\ \langle \mathcal{A}(x, y) \cdot \nabla u, \vec{\eta} \rangle = 0 & \text{on } \partial\Omega \end{cases}$$

where

$$\begin{aligned} Lu(x, y) = & - \left[ \frac{\partial}{\partial x} \left( \frac{1}{(x^2 + y^2)^{1/3}} \frac{\partial u}{\partial x} \right) + \frac{\partial}{\partial y} \left( \frac{1}{(x^2 + y^2)^{1/2}} \frac{\partial u}{\partial y} \right) \right] \\ & + \frac{\sin(xy)}{(x^2 + y^2)^{1/3}} \frac{\partial u}{\partial x} + \frac{\cos(xy)}{(x^2 + y^2)^{1/4}} \frac{\partial u}{\partial y} \\ & + \frac{u(x, y) \sin(xy)}{(x^2 + y^2)^{1/3}} + \theta \frac{u(x, y)}{(x^2 + y^2)^{1/2}} \end{aligned}$$

has a unique solution  $u \in H(\Omega)$  (if  $\theta \geq 2$ ).

## 2. DEFINITIONS AND BASIC RESULTS

Let  $\omega$  be a locally integrable nonnegative function in  $\mathbb{R}^n$  and assume that  $0 < \omega < \infty$  almost everywhere. We say that  $\omega$  belongs to the Muckenhoupt class  $A_p$ ,  $1 < p < \infty$ , or that  $\omega$  is an  $A_p$ -weight, if there is a constant  $C = C_{p,\omega}$  such that

$$\left( \frac{1}{|B|} \int_B \omega(x) \, dx \right) \left( \frac{1}{|B|} \int_B \omega^{1/(1-p)}(x) \, dx \right)^{p-1} \leq C$$

for all balls  $B \subset \mathbb{R}^n$ , where  $|\cdot|$  denotes the  $n$ -dimensional Lebesgue measure in  $\mathbb{R}^n$ . If  $1 < q \leq p$ , then  $A_q \subset A_p$  (see [5], [8] or [9] for more information about  $A_p$ -weights). The weight  $\omega$  satisfies the doubling condition if  $\omega(2B) \leq C\omega(B)$  for all balls  $B \subset \mathbb{R}^n$ , where  $\omega(B) = \int_B \omega(x) \, dx$  and  $2B$  denotes the ball with the same center as  $B$  which is twice as large. If  $\omega \in A_p$ , then  $\omega$  is doubling (see Corollary 15.7 in [8, p. 299]).

As an example of an  $A_p$ -weight, the function  $\omega(x) = |x|^\alpha$ ,  $x \in \mathbb{R}^n$ , is in  $A_p$  if and only if  $-n < \alpha < n(p-1)$  (see Corollary 4.4, Chapter IX in [10, p. 236]).

Given an open subset  $\Omega$  of  $\mathbb{R}^n$ , we will denote by  $L^p(\Omega, \omega)$  ( $1 \leq p < \infty$ ) the Banach space of all measurable functions  $f$  defined on  $\Omega$  for which

$$\|f\|_{L^p(\Omega, \omega)} = \left( \int_\Omega |f(x)|^p \omega(x) \, dx \right)^{1/p} < \infty.$$

If  $\omega \in A_p$ ,  $1 < p < \infty$ , then  $\omega^{-1/(p-1)}$  is locally integrable and we have  $L^p(\Omega, \omega) \subset L^1_{\text{loc}}(\Omega)$  for every open set  $\Omega$  (see Remark 1.2.4 in [10, p. 4]). It thus makes sense to talk about weak derivatives of functions in  $L^p(\Omega, \omega)$ .

**Definition 2.1.** Let  $\Omega \subset \mathbb{R}^n$  be open,  $1 < p < \infty$ ,  $k$  a nonnegative integer and  $\omega \in A_p$ . We define the weighted Sobolev space  $W^{k,p}(\Omega, \omega)$  as the set of functions  $u \in L^p(\Omega, \omega)$  with weak derivatives  $D^\alpha u \in L^p(\Omega, \omega)$ ,  $1 \leq |\alpha| \leq k$ . The norm of  $u$  in  $W^{k,p}(\Omega, \omega)$  is defined by

$$(2.1) \quad \|u\|_{W^{k,p}(\Omega, \omega)} = \left( \int_{\Omega} |u(x)|^p \omega(x) dx + \sum_{1 \leq |\alpha| \leq k} \int_{\Omega} |D^\alpha u(x)|^p \omega(x) dx \right)^{1/p}.$$

If  $\omega \in A_p$ , then  $W^{k,p}(\Omega, \omega)$  is the closure of  $C^\infty(\Omega)$  with respect to the norm (2.1) (see Proposition 3.5 in [11, p. 416] or Corollary 2.1.6 in [10, p. 18]). The space  $W_0^{k,p}(\Omega, \omega)$  is the closure of  $C_0^\infty(\Omega)$  with respect to the norm

$$\|u\|_{W_0^{k,p}(\Omega, \omega)} = \left( \sum_{1 \leq |\alpha| \leq k} \int_{\Omega} |D^\alpha u(x)|^p \omega(x) dx \right)^{1/p}.$$

The spaces  $W^{k,p}(\Omega, \omega)$  and  $W_0^{k,p}(\Omega, \omega)$  are Banach spaces and for  $k = 1$  and  $p = 2$  the spaces  $W^{1,2}(\Omega, \omega)$  and  $W_0^{1,2}(\Omega, \omega)$  are Hilbert spaces.

It is evident that the weight functions  $\omega$  which satisfy  $0 < c_1 \leq \omega(x) \leq c_2$  for  $x \in \Omega$  give nothing new (the space  $W^{k,p}(\Omega, \omega)$  is then identical with the classical Sobolev space  $W^{k,p}(\Omega)$ ). Consequently, we shall be interested above all in such weight functions  $\omega$  which either vanish somewhere in  $\bar{\Omega}$  or increase to infinity (or both).

**Definition 2.2.** Let  $\Omega \subset \mathbb{R}^n$  be a bounded and open set. We define the space  $H(\Omega)$  as the closure of  $C^\infty(\bar{\Omega})$  with respect to the norm

$$\|u\|_{H(\Omega)} = \left( \int_{\Omega} |u|^2 v dx + \int_{\Omega} \langle \mathcal{A} \nabla u, \nabla u \rangle dx \right)^{1/2},$$

where  $\mathcal{A} = (a_{ij})$  is the coefficient matrix of the operator  $L$  defined in (1.1),  $\langle \cdot, \cdot \rangle$  denotes the usual inner product in  $\mathbb{R}^n$  and the symbol  $\nabla$  indicates the gradient.

The space  $H(\Omega)$  is a Hilbert space with the inner product

$$a(u, \varphi) = \left( \int_{\Omega} u \varphi v dx + \int_{\Omega} \langle \mathcal{A} \nabla u, \nabla \varphi \rangle dx \right)^{1/2}.$$

Facts about  $H(\Omega)$  are given in [1, p. 1115].

**Remark 2.3.** By the degeneracy condition (1.2) we have

$$\int_{\Omega} |\nabla u|^2 \omega dx \leq \int_{\Omega} \langle \mathcal{A} \nabla u, \nabla u \rangle dx \leq \int_{\Omega} |\nabla u|^2 v dx.$$

Therefore,  $W^{1,2}(\Omega, v) \subset H(\Omega) \subset W^{1,2}(\Omega, \omega)$ .

Note also that since  $\mathcal{A}$  is symmetric,  $|\langle \mathcal{A}x, y \rangle| \leq \langle \mathcal{A}x, x \rangle^{1/2} \langle \mathcal{A}y, y \rangle^{1/2}$ .

**Remark 2.4.** Let  $\Omega \subset \mathbb{R}^n$  be a bounded open set with boundary  $\partial\Omega \in C^{0,1}$ . Using integration by parts with  $u, \varphi \in H(\Omega)$ , if  $u$  satisfies the boundary condition in problem (P), we have

$$\begin{aligned} \int_{\Omega} \varphi Lu \, dx &= \sum_{i,j=1}^n \int_{\Omega} a_{ij} D_i u D_j \varphi \, dx + \sum_{i=1}^n \int_{\Omega} b_i \varphi D_i u \, dx + \int_{\Omega} g u \varphi \, dx \\ &\quad + \theta \int_{\Omega} u \varphi v \, dx + \underbrace{\sum_{i,j=1}^n \int_{\partial\Omega} a_{ij} \frac{\partial u}{\partial x_j} \eta_i \varphi \, dx}_{=0} \\ &= B(u, \varphi) + \theta \int_{\Omega} u \varphi v \, dx, \end{aligned}$$

where

$$B(u, \varphi) = \sum_{i,j=1}^n \int_{\Omega} a_{ij} D_i u D_j \varphi \, dx + \sum_{i=1}^n \int_{\Omega} b_i \varphi D_i u \, dx + \int_{\Omega} g u \varphi \, dx$$

is a bilinear form.

We introduce the following definition of solutions for the Neumann problem (P).

**Definition 2.5.** Let  $\Omega \subset \mathbb{R}^n$  be a bounded open set with  $\partial\Omega \in C^{0,1}$  and suppose that  $f/v \in L^2(\Omega, v)$ . A function  $u \in H(\Omega)$  is a solution of the Neumann problem (P) if

$$\int_{\Omega} \sum_{i,j=1}^n a_{ij} D_i u D_j \varphi \, dx + \int_{\Omega} \left[ \sum_{i=1}^n b_i D_i u + g u \right] \varphi \, dx + \theta \int_{\Omega} u \varphi v \, dx = \int_{\Omega} f \varphi \, dx$$

for all  $\varphi \in H(\Omega)$ .

**Lemma 2.6.** Suppose that  $\omega \in A_2$ ,  $v \in A_2$ ,  $b_i/\omega \in L^\infty(\Omega)$  ( $i = 1, \dots, n$ ) and  $g/v \in L^\infty(\Omega)$ . Then there exists a constant  $\mathbf{C} > 0$  such that

$$B(u, u) + \mathbf{C} \|u\|_{L^2(\Omega, \omega)}^2 \geq \frac{1}{2} \|u\|_{H(\Omega)}^2$$

for all  $u \in H(\Omega)$ .

Proof. For all  $u \in H(\Omega)$  we have

$$\begin{aligned}
 (2.2) \quad B(u, u) &= \sum_{i,j=1}^n \int_{\Omega} a_{ij} D_i u D_j u \, dx + \sum_{i=1}^n \int_{\Omega} b_i u D_i u \, dx + \int_{\Omega} g u^2 \, dx \\
 &= \int_{\Omega} \langle \mathcal{A} \nabla u, \nabla u \rangle \, dx + \sum_{i=1}^n \int_{\Omega} \frac{b_i}{\omega} u D_i u \, dx + \int_{\Omega} \frac{g}{v} u^2 v \, dx \\
 &\geq \int_{\Omega} \langle \mathcal{A} \nabla u, \nabla u \rangle \, dx - \left( \max_{1 \leq i \leq n} \left\| \frac{b_i}{\omega} \right\|_{L^\infty(\Omega)} \right) \sum_{i=1}^n \int_{\Omega} |u| |D_i u| \omega \, dx \\
 &\quad - \left\| \frac{g}{v} \right\|_{L^\infty(\Omega)} \int_{\Omega} u^2 v \, dx \\
 &\geq \int_{\Omega} \langle \mathcal{A} \nabla u, \nabla u \rangle \, dx - C_1 \sum_{i=1}^n \left( \int_{\Omega} u^2 \omega \, dx \right)^{1/2} \left( \int_{\Omega} |D_i u|^2 \omega \, dx \right)^{1/2} \\
 &\quad - C_2 \int_{\Omega} u^2 v \, dx \\
 &\geq \int_{\Omega} \langle \mathcal{A} \nabla u, \nabla u \rangle \, dx - C_1 \left( \int_{\Omega} u^2 v \, dx \right)^{1/2} \left( \int_{\Omega} \langle \mathcal{A} \nabla u, \nabla u \rangle \, dx \right)^{1/2} \\
 &\quad - C_2 \|u\|_{L^2(\Omega, v)}^2
 \end{aligned}$$

where

$$C_1 = \max_{1 \leq i \leq n} \left\| \frac{b_i}{\omega} \right\|_{L^\infty(\Omega)} \quad \text{and} \quad C_2 = \left\| \frac{g}{v} \right\|_{L^\infty(\Omega)}.$$

Using the elementary inequality

$$ab \leq \varepsilon a^2 + \frac{1}{4\varepsilon} b^2 \quad \text{for all } \varepsilon > 0,$$

we obtain from (2.2)

$$\begin{aligned}
 (2.3) \quad B(u, u) &\geq \int_{\Omega} \langle \mathcal{A} \nabla u, \nabla u \rangle \, dx - C_1 \left( \varepsilon \|u\|_{L^2(\Omega, v)}^2 + \frac{1}{4\varepsilon} \int_{\Omega} \langle \mathcal{A} \nabla u, \nabla u \rangle \, dx \right) \\
 &\quad - C_2 \|u\|_{L^2(\Omega, v)}^2 \\
 &= \left( 1 - \frac{C_1}{4\varepsilon} \right) \int_{\Omega} \langle \mathcal{A} \nabla u, \nabla u \rangle \, dx - (C_1 \varepsilon + C_2) \|u\|_{L^2(\Omega, v)}^2.
 \end{aligned}$$

If  $C_1 > 0$ , we can choose  $\varepsilon > 0$  such that

$$1 - \frac{C_1}{4\varepsilon} = \frac{1}{2}, \quad \text{that is,} \quad \varepsilon = \frac{C_1}{2}.$$

Thus, (2.3) transforms to

$$\begin{aligned} B(u, u) &\geq \frac{1}{2} \int_{\Omega} \langle \mathcal{A} \nabla u, \nabla u \rangle \, dx - \left( \frac{C_1^2}{2} + C_2 \right) \|u\|_{L^2(\Omega)}^2 \\ &= \frac{1}{2} \left( \int_{\Omega} u^2 v \, dx + \int_{\Omega} \langle \mathcal{A} \nabla u, \nabla u \rangle \, dx \right) - \left( \frac{C_1^2}{2} + C_2 + \frac{1}{2} \right) \|u\|_{L^2(\Omega, v)}^2 \\ &= \frac{1}{2} \|u\|_{H(\Omega)}^2 - \mathbf{C} \|u\|_{L^2(\Omega, v)}^2, \end{aligned}$$

where  $\mathbf{C} = \frac{1}{2}C_1^2 + C_2 + \frac{1}{2} > 0$ . Therefore,

$$B(u, u) + \mathbf{C} \|u\|_{L^2(\Omega, v)}^2 \geq \frac{1}{2} \|u\|_{H(\Omega)}^2.$$

If  $C_1 = 0$  (that is,  $b_i(x) \equiv 0$ ,  $i = 1, \dots, n$ ) then (2.2) reduces to

$$\begin{aligned} B(u, u) &\geq \int_{\Omega} \langle \mathcal{A} \nabla u, \nabla u \rangle \, dx - C_2 \|u\|_{L^2(\Omega, v)}^2 \\ &\geq \frac{1}{2} \left( \int_{\Omega} |u|^2 v \, dx + \int_{\Omega} \langle \mathcal{A} \nabla u, \nabla u \rangle \, dx \right) - \left( C_2 + \frac{1}{2} \right) \|u\|_{L^2(\Omega, v)}^2 \\ &= \frac{1}{2} \|u\|_{H(\Omega)}^2 - \mathbf{C} \|u\|_{L^2(\Omega, v)}^2. \end{aligned}$$

Therefore, we also have

$$B(u, u) + \mathbf{C} \|u\|_{L^2(\Omega, v)}^2 \geq \frac{1}{2} \|u\|_{H(\Omega)}^2$$

for all  $u \in H(\Omega)$ , where  $\mathbf{C} = \frac{1}{2}C_1^2 + C_2 + \frac{1}{2}$ . □

### 3. PROOF OF THEOREM 1.1

We define a bilinear form

$$\tilde{B}: H(\Omega) \times H(\Omega) \longrightarrow \mathbb{R}, \quad \tilde{B}(u, \varphi) = B(u, \varphi) + \theta \int_{\Omega} u \varphi v \, dx$$

and a linear mapping

$$T: H(\Omega) \longrightarrow \mathbb{R}, \quad T(\varphi) = \int_{\Omega} f \varphi \, dx.$$

Then  $u \in H(\Omega)$  is a solution of the Neumann problem (P) if

$$\tilde{B}(u, \varphi) = T(\varphi)$$

for all  $\varphi \in H(\Omega)$ .



*Step 1.* If  $\theta \geq \mathbf{C}$  then  $\tilde{B}$  is coercive, that is, there exists a constant  $c > 0$  such that  $\tilde{B}(u, u) \geq c\|u\|_{H(\Omega)}^2$  for all  $u \in H(\Omega)$ . In fact, by Lemma 2.6 there exists a constant  $\mathbf{C} > 0$  such that

$$B(u, u) + \mathbf{C}\|u\|_{L^2(\Omega, v)}^2 \geq \frac{1}{2}\|u\|_{H(\Omega)}^2.$$

Hence, if  $\theta \geq \mathbf{C}$ , we have

$$\begin{aligned} \tilde{B}(u, u) &= B(u, u) + \theta \int_{\Omega} u^2 v \, dx = B(u, u) + \theta\|u\|_{L^2(\Omega, v)}^2 \\ &\geq B(u, u) + \mathbf{C}\|u\|_{L^2(\Omega, v)}^2 \geq \frac{1}{2}\|u\|_{H(\Omega)}^2. \end{aligned}$$

Therefore, for  $\theta \geq \mathbf{C}$  we have that

$$(3.1) \quad \tilde{B}(u, u) \geq \frac{1}{2}\|u\|_{H(\Omega)}^2$$

for all  $u \in H(\Omega)$ .

*Step 2.*  $\tilde{B}$  is bounded. In fact, using the fact that the coefficient matrix  $\mathcal{A} = (a_{ij})$  is symmetric, (H2) and (H3), we obtain

$$\begin{aligned} |\tilde{B}(u, \varphi)| &\leq |B(u, \varphi)| + \theta \left| \int_{\Omega} u\varphi v \, dx \right| \\ &\leq \int_{\Omega} |\langle \mathcal{A}\nabla u, \nabla\varphi \rangle| \, dx + \sum_{i=1}^n \int_{\Omega} |b_i||\varphi| |D_i u| \, dx + \int_{\Omega} |g||\varphi| |u| \, dx + \theta \int_{\Omega} |u||\varphi|v \, dx \\ &\leq \int_{\Omega} \langle \mathcal{A}\nabla u, \nabla u \rangle^{1/2} \langle \mathcal{A}\nabla\varphi, \nabla\varphi \rangle^{1/2} \, dx + \sum_{i=1}^n \int_{\Omega} \frac{|b_i|}{\omega} |\varphi| |D_i u| \omega \, dx \\ &\quad + \int_{\Omega} \frac{|g|}{v} |\varphi| |u|v \, dx + \theta \int_{\Omega} |u||\varphi|v \, dx \\ &\leq \left( \int_{\Omega} \langle \mathcal{A}\nabla u, \nabla u \rangle \, dx \right)^{1/2} \left( \int_{\Omega} \langle \mathcal{A}\nabla\varphi, \nabla\varphi \rangle \, dx \right)^{1/2} \\ &\quad + \left( \max_{1 \leq i \leq n} \left\| \frac{b_i}{\omega} \right\|_{L^\infty(\Omega)} \right) \sum_{i=1}^n \left( \int_{\Omega} |\varphi|^2 \omega \, dx \right)^{1/2} \left( \int_{\Omega} |D_i u|^2 \omega \, dx \right)^{1/2} \\ &\quad + \left\| \frac{g}{v} \right\|_{L^\infty(\Omega)} \left( \int_{\Omega} |u|^2 v \, dx \right)^{1/2} \left( \int_{\Omega} |\varphi|^2 v \, dx \right)^{1/2} \\ &\quad + \theta \left( \int_{\Omega} |u|^2 v \, dx \right)^{1/2} \left( \int_{\Omega} |\varphi|^2 v \, dx \right)^{1/2} \\ &\leq \left( 1 + \max_{1 \leq i \leq n} \left\| \frac{b_i}{\omega} \right\|_{L^\infty(\Omega)} + \left\| \frac{g}{v} \right\|_{L^\infty(\Omega)} + \theta \right) \|u\|_{H(\Omega)} \|\varphi\|_{H(\Omega)} \\ &= \tilde{C} \|u\|_{H(\Omega)} \|\varphi\|_{H(\Omega)} \end{aligned}$$

for all  $u, \varphi \in H(\Omega)$ .

*Step 3.* The linear mapping  $T$  is bounded (that is,  $T \in [H(\Omega)]^*$ ). In fact,

$$\begin{aligned} |T(\varphi)| &\leq \int_{\Omega} |f| |\varphi| \, dx = \int_{\Omega} \frac{|f|}{v} |\varphi| v \, dx \\ &\leq \left[ \int_{\Omega} \left( \frac{|f|}{v} \right)^2 v \, dx \right]^{1/2} \left[ \int_{\Omega} |\varphi|^2 v \, dx \right]^{1/2} \\ &\leq \left\| \frac{f}{v} \right\|_{L^2(\Omega, v)} \|\varphi\|_{H(\Omega)} \end{aligned}$$

for all  $\varphi \in H(\Omega)$ .

Therefore the bilinear form  $\tilde{B}$  and the linear functional  $T$  satisfy the hypotheses of the Lax-Milgram theorem. Thus, for every  $f$  with  $f/v \in L^2(\Omega, v)$ , there is a unique solution  $u \in H(\Omega)$  such that

$$\tilde{B}(u, \varphi) = T(\varphi)$$

for all  $\varphi \in H(\Omega)$ , that is,  $u$  is a unique solution of the Neumann problem (P).

In particular, by setting  $\varphi = u$ , we have

$$\tilde{B}(u, u) = \int_{\Omega} f u \, dx.$$

Using the definition of  $\tilde{B}$ , we obtain

$$\begin{aligned} \tilde{B}(u, u) &= B(u, u) + \theta \int_{\Omega} u^2 v \, dx = \int_{\Omega} \frac{f}{v} u v \, dx \\ &\leq \|u\|_{L^2(\Omega, v)} \left\| \frac{f}{v} \right\|_{L^2(\Omega, v)} \\ &\leq \|u\|_{H(\Omega)} \left\| \frac{f}{v} \right\|_{L^2(\Omega, v)}. \end{aligned}$$

Using (3.1), we obtain

$$\frac{1}{2} \|u\|_{H(\Omega)}^2 \leq \tilde{B}(u, u) \leq \|u\|_{H(\Omega)} \left\| \frac{f}{v} \right\|_{L^2(\Omega, v)}.$$

Therefore,

$$\|u\|_{H(\Omega)} \leq 2 \left\| \frac{f}{v} \right\|_{L^2(\Omega, v)}.$$

□

### References

- [1] *S. Chanillo, R. L. Wheeden*: Harnack's inequalities and mean-value inequalities for solutions of degenerate elliptic equations. *Commun. PDE* 11 (1986), 1111–1134. [Zbl 0634.3503](#)
- [2] *S. Chanillo, R. L. Wheeden*: Existence and estimates for Green's function for degenerate elliptic equations. *Ann. Sc. Norm. Super. Pisa IV ser.* 15, Fasc. II (1988), 309–340. [Zbl 0688.35002](#)
- [3] *B. Franchi, R. Serapioni*: Pointwise estimates for a class of strongly degenerate elliptic operators: A geometrical approach. *Ann. Sc. Norm. Sup. Pisa C1 ser.* 14 (1987), 527–568. [Zbl 0685.35046](#)
- [4] *B. Muckenhoupt*: Weighted norm inequalities for the Hardy maximal function. *Trans. Am. Math. Soc.* 165 (1972), 207–226. [Zbl 0236.26015](#)
- [5] *A. Torchinsky*: *Real-Variable Methods in Harmonic Analysis*. Academic Press, Orlando, 1986. [Zbl 0621.42001](#)
- [6] *A. Kufner, O. John, and S. Fučík*: *Function Spaces*. Noordhoff International Publishing, Leyden, 1977. [Zbl 0364.46022](#)
- [7] *A. Kufner*: *Weighted Sobolev Spaces*. John Wiley & Sons, New York, 1985. [Zbl 0567.46009](#)
- [8] *J. Heinonen, T. Kilpeläinen, and O. Martio*: *Nonlinear Potential Theory of Degenerate Elliptic Equations*. Oxford Math. Monographs. Clarendon Press, Oxford, 1993. [Zbl 0780.31001](#)
- [9] *J. Garcia-Cuerva, J. L. Rubio de Francia*: *Weighted Norm Inequalities and Related Topics*. North-Holland Mathematics Studies 116. North Holland, Amsterdam-New York-Oxford, 1985. [Zbl 0578.46046](#)
- [10] *B. O. Turesson*: *Nonlinear Potential Theory and Weighted Sobolev Spaces*. *Lectures Notes in Mathematics* Vol. 1736. Springer-Verlag, Berlin, 2000. [Zbl 0949.31006](#)
- [11] *V. Chiadò Piat, F. Serra Cassano*: Relaxation of degenerate variational integrals. *Nonlinear Anal., Theory Methods Appl.* 22 (1994), 409–424. [Zbl 0799.49012](#)

*Author's address*: A. C. Cavalheiro, Department of Mathematics, State University of Londrina, Londrina-PR, 86051-990, Brazil, e-mail: [accava@gmail.com](mailto:accava@gmail.com).