

Jiří Anděl; Karel Hrach

On calculation of stationary density of autoregressive processes

Kybernetika, Vol. 36 (2000), No. 3, [311]--319

Persistent URL: <http://dml.cz/dmlcz/135352>

Terms of use:

© Institute of Information Theory and Automation AS CR, 2000

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these

Terms of use.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library*
<http://project.dml.cz>

ON CALCULATION OF STATIONARY DENSITY OF AUTOREGRESSIVE PROCESSES

JIRÍ ANDĚL AND KAREL HRACH

An iterative procedure for computation of stationary density of autoregressive processes is proposed. On an example with exponentially distributed white noise it is demonstrated that the procedure converges geometrically fast. The AR(1) and AR(2) models are analyzed in detail.

1. INTRODUCTION AND PRELIMINARIES

Let $\{X_t\}$ be a stationary autoregressive process of the first order defined by

$$X_t = bX_{t-1} + e_t, \quad 0 \neq b \in (-1, 1) \quad (1.1)$$

where $\{e_t\}$ are i.i.d. random variables with a finite second moment. From (1.1) we have

$$X_t = e_t + be_{t-1} + b^2e_{t-2} + \dots \quad (1.2)$$

and the series converges in the quadratic mean. It is clear from (1.2) that $\{X_t\}$ is strictly stationary. We are interested in the stationary distribution of the process $\{X_t\}$. The following assertions show that this distribution is continuous.

Theorem 1.1. Let $\{\xi_t\}$ be i.i.d. random variables. If the series $X = \sum_{t=-\infty}^{\infty} k_t \xi_t$ converges almost surely and infinite many k_t are different from 0 then X has a continuous distribution.

Proof. See [10]. □

Theorem 1.2. Let $\{\eta_t\}$ be independent random variables such that $E\eta_t^2 < \infty$ and $\sum \text{var} \eta_t < \infty$. Then the series $\sum (\eta_t - E\eta_t)$ converges almost surely.

Proof. See [15], § 16.2 and § 17.2, or [18], Theorem IV.1.4, p. 241. □

Theorem 1.3. The stationary distribution of the process $\{X_t\}$ defined by (1.1) is continuous.

Proof. The assertion is a direct consequence of Theorems 1.1 and 1.2. \square

For example, if $b = 0.5$ and e_t is a discrete random variable such that $P(e_t = 0.5) = P(e_t = -0.5) = 0.5$ then X_t has the continuous rectangular distribution $R(-1, 1)$. A review of some results of this kind can be found in [2].

However, in many cases stronger assumptions about e_t are made.

Theorem 1.4. If e_t has a density, then X_t has also a density.

Proof. Using (1.2) we can write $X_t = e_t + Z_t$ where $Z_t = be_{t-1} + b^2e_{t-2} + \dots$. Since e_t and Z_t are independent and e_t has a density, their sum $e_t + Z_t = X_t$ has an absolutely continuous distribution (see [16], p. 196). \square

Now, we introduce a known equation for the stationary distribution of X_t and a formula for the characteristic function of X_t .

Theorem 1.5. Let e_t have a density f . Then the density h of X_t satisfies the equation

$$h(x) = \int f(x - bu)h(u) du. \quad (1.3)$$

Proof. The equation follows from (1.1), since X_{t-1} has also the density h and X_{t-1} and e_t are independent. \square

Theorem 1.6. Let $\psi(t)$ be the characteristic function of e_t and let $\rho(t)$ be the characteristic function of X_t . Then

$$\rho(t) = \prod_{n=0}^{\infty} \psi(b^n t).$$

Proof. Define

$$Y_{t,n} = e_t + be_{t-1} + \dots + b^n e_{t-n}. \quad (1.4)$$

Then $Y_{t,n} \rightarrow X_t$ in the quadratic mean as $n \rightarrow \infty$. Thus $Y_{t,n} \rightarrow X_t$ also in the distribution. It implies that the characteristic functions of the variables $Y_{t,n}$ converge pointwise to $\rho(t)$. But the characteristic function of $Y_{t,n}$ is $\psi(t)\psi(bt)\dots\psi(b^n t)$. \square

If e_t has normal distribution then it is well known that X_t is also normally distributed. One of the first attempts to find a connection between the distributions of X_t and e_t in a non-normal case was published in [9]. For some special non-normal distributions of e_t (continuous and discrete rectangular distributions, Laplace distribution) the stationary distribution of X_t is calculated in [2]. It is also known

that if e_t has a stable distribution of exponent θ , ($0 < \theta \leq 2$) then X_t also has a stable distribution of the same exponent (e. g., see [19], p. 208, Ex. 11).

By the way, the opposite problem was more popular, viz. to find a distribution of e_t for a given stationary distribution of X_t . The famous paper [12] contains the cases when X_t has exponential or gamma distributions. A review of such results can be found in [1] and [2].

The methods mentioned above were applicable only to some special distributions of X_t and e_t . Another approach was proposed in [3]. The goal of this paper was to find a distribution of e_t such that the stationary distribution of X_t has given moments. This method was derived for general linear processes so that autoregressive models form only a special class. Detailed results for AR(1) models are published in [4]. A different method based on Hermite polynomials can be found in [17].

The problem how to calculate stationary distribution of a process from the distribution of a white noise seems to be even more popular in non-linear models. The method of moments was applied in [6]. Exact explicit results are quite rare. One of them can be found in [7] but formulas for stationary density of the absolute autoregression published in [8] and [5] became more familiar (cf. [19], pp. 140–142 and p. 205). Numerical procedures suggested for computation of stationary density in non-linear models (see [19], p. 152) can be, of course, also used in the model (1.1). It corresponds to numerical solution of the equation (1.3).

Quite recently (see [13]) some theoretic results for the stationary density of X_t in the model (1.1) were derived in the case that e_t has the rectangular distribution on $(0, 1)$. It was proved that the stationary density belongs to the class C^∞ and some bounds for tails of this stationary distribution were derived. The derivation was based on investigation of asymptotic properties of $e_t + be_{t-1} + \dots + b^n e_{t-n}$ as $n \rightarrow \infty$.

2. AN ITERATIVE METHOD FOR AR(1) PROCESSES

An explicit formula for h satisfying (1.3) given f is known only in a few cases. Here we propose an iterative method for its computation. Let h_0 be an arbitrary density. For $n \geq 1$ define

$$h_n(x) = \int f(x - bu) h_{n-1}(u) du. \tag{2.1}$$

It is obvious that every function h_n defined by (2.1) is a density.

Theorem 2.1. Let h_0 be a density. Define h_n by (2.1). Assume that there exists an integer $m \geq 0$ such that

$$\int_{-\infty}^{\infty} |\psi(t) \psi(bt) \dots \psi(b^m t)| dt < \infty. \tag{2.2}$$

Then $h_n(x) \rightarrow h(x)$ for all x as $n \rightarrow \infty$.

Proof. Let λ_n be the characteristic function corresponding to h_n . Using (2.1) we obtain

$$\begin{aligned}\lambda_n(t) &= \int e^{itx} h_n(x) dx = \int e^{itx} \left[\int f(x-bu) h_{n-1}(u) du \right] dx \\ &= \int h_{n-1}(u) \left[\int e^{itx} f(x-bu) dx \right] du = \int h_{n-1}(u) \left[\int e^{itbu+ity} f(y) dy \right] du \\ &= \psi(t) \int e^{itbu} h_{n-1}(u) du = \psi(t) \lambda_{n-1}(bt).\end{aligned}$$

Thus

$$\lambda_n(t) = \psi(t) \psi(bt) \psi(b^2t) \dots \psi(b^{n-1}t) \lambda_0(b^nt).$$

From the continuity of the characteristic function we have

$$\lambda_0(b^nt) \rightarrow \lambda_0(0) = 1 \quad \text{as } n \rightarrow \infty.$$

Using Theorem 1.6 we get $\lambda_n(t) \rightarrow \rho(t)$ as $n \rightarrow \infty$. Assume that $n > m$. Because

$$h_n(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-itx} \lambda_n(t) dt \quad (2.3)$$

and $|e^{-itx} \lambda_n(t)| \leq |\psi(t) \psi(bt) \dots \psi(b^mt)|$, Lebesgue theorem gives

$$\lim_{n \rightarrow \infty} h_n(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-itx} \left[\lim_{n \rightarrow \infty} \lambda_n(t) \right] dt = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-itx} \rho(t) dt = h(x). \quad \square$$

Of course, speed of the convergence and complexity of formulas for h_n depend heavily on the choice of h_0 .

3. AN EXAMPLE

Sometimes it is easy to derive speed of convergence $h_n(x) \rightarrow h(x)$. From (2.3) we have

$$|h_{n+1}(x) - h_n(x)| \leq \frac{1}{2\pi} \int_{-\infty}^{\infty} |\lambda_{n+1}(t) - \lambda_n(t)| dt.$$

Define

$$\Delta_{n+1} = \sup_x |h_{n+1}(x) - h_n(x)|.$$

Then we get

$$\Delta_{n+1} \leq \frac{1}{2\pi} \int_{-\infty}^{\infty} |\psi(t) \psi(bt) \dots \psi(b^{n-1}t) [\psi(b^nt) \lambda_0(b^{n+1}t) - \lambda_0(b^nt)]| dt. \quad (3.1)$$

Consider the process $\{X_t\}$ given by (1.1) such that $b \in (0, 1)$ and $e_t \sim Ex(1)$, i. e., $f(x) = e^{-x}$ for $x > 0$. For simplicity, choose $h_0(x) = f(x)$. Then

$$\psi(t) = \lambda_0(t) = \frac{1}{1-it}.$$

Since

$$|\psi(t)| = \frac{1}{\sqrt{1+t^2}}, \tag{3.2}$$

the assumption (2.2) is fulfilled for $m = 1$. The inequality (3.1) can be written in the form

$$\Delta_{n+1} \leq \int_{-\infty}^{\infty} \left| \left[\prod_{s=0}^n \psi(b^s t) \right] [\psi(b^{n+1}t) - 1] \right| dt. \tag{3.3}$$

Assume that $n \geq 3$. Then (3.3) and (3.2) yield

$$\begin{aligned} \Delta_{n+1} &\leq \frac{1}{\pi} \int_{-\infty}^{\infty} |\psi(t) \psi(bt)| \cdot |\psi(b^2t)\psi(b^3t)| \cdot |\psi(b^{n+1}t) - 1| dt \\ &\leq \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{1}{1+b^2t^2} \frac{1}{1+b^6t^2} \frac{b^{n+1}t}{\sqrt{1+b^{2n+2}t^2}} dt \\ &\leq b^{n+1} \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{1}{1+b^2t^2} \frac{1}{1+b^6t^2} dt. \end{aligned}$$

Since

$$\frac{t}{1+b^6t^2} \leq \frac{1}{2b^3}, \quad \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{dt}{1+b^2t^2} = \frac{1}{2b},$$

we have

$$\Delta_{n+1} \leq \frac{1}{4} b^{n-3}.$$

In this case the iterative procedure converges to the limit density geometrically fast.

For small values of n we write (3.3) in the form

$$\Delta_{n+1} \leq \frac{b^{n+1}}{\pi} \int_0^{\infty} t \prod_{s=0}^{n+1} (1+b^{2s}t^2)^{-1/2} dt.$$

Especially,

$$\begin{aligned} \Delta_2 &\leq \frac{b^2}{\pi} \int_0^{\infty} t \frac{1}{\sqrt{1+t^2}} \frac{1}{\sqrt{1+b^2t^2}} \frac{1}{\sqrt{1+b^4t^2}} dt \\ &\leq \frac{b^2}{\pi} \int_0^{\infty} \frac{t}{(1+b^4t^2)\sqrt{1+t^2}} dt = \frac{b^2}{2\pi} \int_0^{\infty} \frac{dx}{(1+b^4x)\sqrt{1+x}} \\ &= \frac{\pi - 2 \operatorname{arctg} \frac{b^2}{\sqrt{1-b^4}}}{2\pi\sqrt{1-b^4}} = D_2(b). \end{aligned}$$

Similarly,

$$\begin{aligned} \Delta_3 &\leq \frac{b^3}{\pi} \int_0^{\infty} \frac{t}{\sqrt{1+t^2}} \frac{1}{\sqrt{1+b^2t^2}} \frac{1}{\sqrt{1+b^4t^2}} \frac{1}{\sqrt{1+b^6t^2}} dt \\ &\leq \frac{b^3}{\pi} \int_0^{\infty} \frac{1}{1+b^2t^2} \frac{1}{1+b^6t^2} dt = -\frac{2b \ln b}{\pi(1-b^4)} = D_3(b). \end{aligned}$$

Some values of $D_2(b)$ and $D_3(b)$ are introduced in Table 3.1.




Fig. 3.1.

Table 3.1.

b	0.01	0.1	0.5	0.9	0.99
$D_2(b)$	0.50	0.50	0.43	0.34	0.32
$D_3(b)$	0.03	0.15	0.24	0.18	0.16

Explicit formulas for $h_n(x)$ can be written down if n is small. For example,

$$h_1(x) = \frac{e^{-x}}{1-b} \left(1 - e^{x-x/b} \right),$$

$$h_2(x) = \frac{e^{-x}}{(1-b)^2} \left[1 - e^{x-x/b} - \frac{b - e^{x(b^2+b-1)/b-1/b}}{1+b} \right]$$

and so on. However, it is easier to use a program package for calculating and processing the functions $h_n(x)$. We used Mathematica. Figure 3.1 shows functions $h_0(x), \dots, h_5(x)$ in the case $b = 0.5$.

4. A GENERALIZATION TO AR PROCESSES OF HIGHER ORDER

The iterative method for calculating stationary density can be generalized to autoregressive processes of higher order. Here we present a derivation for AR(2) model.

Let X_t be a stationary AR(2) process defined by

$$X_t = b_1 X_{t-1} + b_2 X_{t-2} + e_t \tag{4.1}$$

where e_t is a strict white noise with a density f and a finite second moment. Let ψ be the characteristic function of e_t and

$$F = \begin{pmatrix} b_1 & b_2 \\ 1 & 0 \end{pmatrix}.$$

It is known that

$$X_t = \sum_{j=0}^{\infty} a_j e_{t-j}$$

where a_j is the (1,1)-element of the matrix F^j (see [14], p. 57). It follows from the assumption of stationarity that all roots of F lie inside the unit circle and thus the series (4.1) converges in the quadratic mean. If we define $c = (1, 0)'$ then $a_j = c' F^j c = c' F'^j c$ and the characteristic function λ of X_t is given by

$$\lambda(v) = \prod_{j=0}^{\infty} \psi(va_j) = \prod_{j=0}^{\infty} \psi(v c' F'^j c). \tag{4.2}$$

Since we assume that e_t has a density, it follows from (4.1) that the random vector $(X_t, X_{t-1})'$ has a joint density, say $q(x, y)$. The stationary density of $\{X_t\}$ is $\int q(x, y) dy$. Because $\{X_t\}$ is stationary, the vector $(X_{t-1}, X_{t-2})'$ has also the density q . The joint density of $(X_t, X_{t-1}, X_{t-2})'$ is $q(x_{t-1}, x_{t-2}) f(x_t - b_1 x_{t-1} - b_2 x_{t-2})$ and so we have an integral equation

$$q(x_t, x_{t-1}) = \int q(x_{t-1}, x_{t-2}) f(x_t - b_1 x_{t-1} - b_2 x_{t-2}) dx_{t-2}. \tag{4.3}$$

Let $q_0(y, z)$ be a density. Formula (4.3) suggests that a method for calculating q can be based on the recurrent relation

$$q_n(x, y) = \int q_{n-1}(y, z) f(x - b_1 y - b_2 z) dz. \tag{4.4}$$

We prove that under some conditions concerning q_0 the functions q_n converge pointwise to q .

Theorem 4.1. Let λ_n be the characteristic function corresponding to q_n . Then for arbitrary $t = (t_1, t_2)'$ we have $\lambda_n(t) \rightarrow \lambda(t)$.

Proof. Using (4.4) we get

$$\begin{aligned} \lambda_n(t_1, t_2) &= \iint e^{it_1 x + it_2 y} q_n(x, y) dx dy \\ &= \iiint e^{it_1 x + it_2 y} q_{n-1}(y, z) f(x - b_1 y - b_2 z) dz dx dy \end{aligned}$$

$$\begin{aligned}
&= \iiint e^{it_1(w+b_1y+b_2z)+it_2y} q_{n-1}(y, z) f(w) dz dw dy \\
&= \iint e^{i(t_1b_1+t_2)y+it_1b_2z} q_{n-1}(y, z) dy dz \int e^{it_1w} f(w) dw \\
&= \lambda_{n-1}(t_1b_1 + t_2, t_1b_2) \psi(t_1) \\
&= \lambda_{n-1}(\mathbf{F}'\mathbf{t}) \psi(\mathbf{c}'\mathbf{t}).
\end{aligned}$$

This gives

$$\lambda_n(\mathbf{t}) = \psi(\mathbf{c}'\mathbf{t}) \psi(\mathbf{c}'\mathbf{F}'\mathbf{t}) \dots \psi(\mathbf{c}'\mathbf{F}'^{n-1}\mathbf{t}) \lambda_0(\mathbf{F}'^n\mathbf{t}).$$

Since $\mathbf{F}'^n \rightarrow \mathbf{0}$ as $n \rightarrow \infty$, we have $\lambda_0(\mathbf{F}'^n\mathbf{t}) \rightarrow 1$ and in view of (4.2) it follows that $\lambda_n(\mathbf{t}) \rightarrow \lambda(\mathbf{t})$. \square

Theorem 4.2. Let q_0 be a density. Assume that there exists an integer $m \geq 0$ such that

$$\iint |\psi(\mathbf{c}'\mathbf{t}) \psi(\mathbf{c}'\mathbf{F}'\mathbf{t}) \dots \psi(\mathbf{c}'\mathbf{F}'^m\mathbf{t})| dt_1 dt_2 < \infty.$$

Then $q_n(x, y) \rightarrow q(x, y)$ for all (x, y) as $n \rightarrow \infty$.

Proof. For $n \geq m$ we have

$$|\lambda_n(t_1, t_2)| \leq |\psi(\mathbf{c}'\mathbf{t}) \psi(\mathbf{c}'\mathbf{F}'\mathbf{t}) \dots \psi(\mathbf{c}'\mathbf{F}'^m\mathbf{t})|$$

and thus $\iint |\lambda_n(t_1, t_2)| dt_1 dt_2 < \infty$. Then $q_n(x, y)$ is bounded, continuous, and

$$q_n(x, y) = \frac{1}{(2\pi)^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-i(xt_1+yt_2)} \lambda_n(t_1, t_2) dt_1 dt_2$$

(see [11], formula 7.12). Theorem 4.1 and Lebesgue theorem imply

$$\lim_{n \rightarrow \infty} \frac{1}{(2\pi)^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-i(xt_1+yt_2)} \lambda_n(t_1, t_2) dt_1 dt_2 = q(x, y). \quad \square$$

ACKNOWLEDGEMENT

The research was supported by Grant 201/97/1176 of the Grant Agency of the Czech Republic and by Grant CEZ:J13/98:113200008.

(Received April 13, 1999.)

REFERENCES

-
- [1] J. Anděl: Dependent random variables with a given marginal distribution. *Acta Univ. Carolin. – Math. Phys.* 24 (1983), 3–11.
 - [2] J. Anděl: Marginal distributions of autoregressive processes. In: *Trans. 9th Prague Conf. Inform. Theory, Statist. Decision Functions, Random Processes*, Academia, Prague 1983, pp. 127–135.

- [3] J. Anděl: On linear processes with given moments. *J. Time Ser. Anal.* 8 (1987), 373–378.
- [4] J. Anděl: AR(1) processes with given moments of marginal distribution. *Kybernetika* 22 (1989), 337–347.
- [5] J. Anděl and T. Bartoň: A note on the threshold AR(1) model with Cauchy innovations. *J. Time Ser. Anal.* 7 (1986), 1–5.
- [6] J. Anděl and M. Garrido: On stationary distributions of some time series models. In: *Trans. 10th Prague Conf. Inform. Theory, Statist. Decision Functions, Random Processes*, Academia, Prague 1988, pp. 193–202.
- [7] J. Anděl, M. Gómez and C. Vega: Stationary distribution of some nonlinear AR(1) processes. *Kybernetika* 25 (1989), 453–460.
- [8] J. Anděl, I. Netuka and K. Zvára: On threshold autoregressive processes. *Kybernetika* 20 (1984), 89–106.
- [9] J. Bernier: Inventaire des modèles et processus stochastique applicables de la description des débits journaliers des rivières. *Rev. Inst. Internat. Statist.* 38 (1970), 50–71.
- [10] R. A. Davis and M. Rosenblatt: Parameter estimation for some time series models without contiguity. *Statist. Probab. Lett.* 11 (1991), 515–521.
- [11] W. Feller: *An Introduction to Probability Theory and its Applications II*. Wiley, New York 1966.
- [12] D. P. Gaver and P. A. W. Lewis: First-order autoregressive gamma sequences and point processes. *Adv. in Appl. Probab.* 12 (1980), 727–745.
- [13] G. Haiman: Upper and lower bounds for the tail of the invariant distribution of some AR(1) processes. In: *Asymptotic Methods in Probability and Statistics* (B. Szyszkowicz, ed.), North-Holland/Elsevier, Amsterdam 1998, pp. 723–730.
- [14] J. D. Hamilton: *Time Series Analysis*. Princeton University Press, Princeton 1994.
- [15] M. Loève: *Probability Theory*. Second edition. Van Nostrand, Princeton 1955.
- [16] A. Rényi: *Probability Theory*. Akadémiai Kiadó, Budapest 1970.
- [17] M. M. Sondhi: Random processes with specified spectral density and first-order probability density. *Bell System Technical J.* 62 (1983), 679–701.
- [18] J. Štěpán: *Teorie pravděpodobnosti*. Academia, Praha 1987.
- [19] H. Tong: *Non-linear Time Series*. Clarendon Press, Oxford 1990.
- [20] Wolfram Research, Inc.: *Mathematica, Version 2.2*. Wolfram Research, Inc., Champaign, Illinois 1994.

*Prof. RNDr. Jiří Anděl, DrSc., Charles University – Faculty of Mathematics and Physics, Department of Statistics, Sokolovská 83, 186 00 Praha 8. Czech Republic.
e-mail: andel@karlin.mff.cuni.cz*

*Mgr. Karel Hrach, University of Jan Evangelista Purkyně, Faculty of Social Sciences and Economy, Department of Mathematics and Informatics, Moskevská 54, 400 01 Ústí n. L. Czech Republic.
e-mail: hrach@fse.ujep.cz*