

Gary Chartrand; Ronald J. Gould; S. F. Kapoor  
The min-max supergraph

*Mathematica Slovaca*, Vol. 30 (1980), No. 2, 175--179

Persistent URL: <http://dml.cz/dmlcz/136238>

## Terms of use:

© Mathematical Institute of the Slovak Academy of Sciences, 1980

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## THE MIN-MAX SUPERGRAPH

GARY CHARTRAND—RONALD J. GOULD—S. F. KAPOOR\*

Let the degree set (the set of degrees of the vertices) of a graph  $G$  be denoted by  $\vartheta_G$  in which  $\Delta$  and  $\delta$  represent the maximum and minimum elements respectively. If  $S$  is a finite set of positive integers with  $\Delta \in S \subseteq \vartheta_G$ , then there exists a graph  $H$  with degree set  $S$  containing  $G$  as an induced subgraph. In the case where  $S = \{\delta, \Delta\}$ , necessary and sufficient conditions are presented for the order of  $H$  to be minimum.

It is well known (see [1], Chap. 1, for example) that for any graph  $G$  with maximum degree  $\Delta$  there exists a  $\Delta$ -regular graph  $H$  containing  $G$  as an induced subgraph. (The graph  $H$  is called a supergraph of  $G$ .) Furthermore, Erdős and Kelly [3] have found a necessary and sufficient set of conditions which determine the minimum order of such a graph  $H$ . In this article we generalize the first of these results and extend the second.

The degree set  $\vartheta_G$  of a graph  $G$  is the set of degrees of the vertices of  $G$ . If  $\vartheta_G = \{a_1, a_2, \dots, a_n\}$ , where  $a_1 < a_2 < \dots < a_n$ , then  $\delta(G) = \delta = a_1$  is the minimum degree of  $G$  and  $\Delta(G) = \Delta = a_n$  is the maximum degree of  $G$ . As mentioned above, there exists a graph  $H$  with degree set  $\{\Delta\}$  containing  $G$  as an induced subgraph. We first present a generalization of this result.

**Theorem 1.** *Let  $G$  be a graph with degree set  $\vartheta_G$  and maximum degree  $\Delta$  and let  $S$  be a finite set of positive integers such that  $\Delta \in S \subseteq \vartheta_G$ . Then there exists a graph  $H$  with degree set  $S$  such that  $G$  is an induced subgraph of  $H$ .*

*Proof.* First, observe that if  $S = \vartheta_G$ , then we may take  $H = G$ . We have already noted that the result is true if  $S = \{\Delta\}$ , so we henceforth assume that  $2 \leq |S| < |\vartheta_G|$ .

Let  $\vartheta_G = \{a_1, a_2, \dots, a_n\}$  with  $\delta = a_1 < a_2 < \dots < a_n = \Delta$ , where  $n \geq 3$ . Define  $G_0 = G$ . For  $i \geq 1$ , define  $G_i$  to be that graph consisting of two disjoint copies of  $G_{i-1}$  together with those edges joining corresponding vertices, say with the same label  $v$  if  $\deg_{G_{i-1}} v \notin S$ . For each  $i = 1, 2, \dots, n$ , define

$$k_i = \min \{m \mid m \geq i, a_m \in S\}$$

and

$$k = \max \{a_{k_i} - a_i \mid i = 1, 2, \dots, n\}.$$

---

\* Research partially supported by a Faculty Research Fellowship from Western Michigan University.

Then  $H = G_k$  has degree set  $S$  and contains  $G$  as an induced subgraph.

In the case where  $S = \{\Delta\}$  Erdős and Kelly described a method for determining the minimum order of the graph  $H$  mentioned in the statement of Theorem 1. We do the same thing if  $S = \{\delta, \Delta\}$ . Prior to presenting a set of conditions which give the minimum order of  $H$  in this case, we find it necessary to introduce some terminology.

Let  $G$  be a graph of order  $p$  and degree set  $\vartheta_G = \{a_1, a_2, \dots, a_n\}$ . Let the vertex set  $V(G)$  of  $G$  be expressed as  $V(G) = V_1 \cup V_2 \cup \dots \cup V_n$ , where for  $1 \leq i \leq n$ ,  $|V_i| = m_i \geq 1$  such that  $v \in V_i$  implies that  $\deg_G v = a_i$ . Let  $V^* = \bigcup_{i=2}^{n-1} V_i$  and let

$\sigma = \sum_{v \in V^*} (\Delta - \deg_G v)$  denote the regular deficiency of  $G$ . Further, let  $H$  be a graph having degree set  $\{\delta(G), \Delta(G)\}$  and containing  $G$  as an induced subgraph, and let  $I = V(H) - V(G)$  be the set of vertices that need to be added to  $G$  in order to obtain  $H$ . From the  $s = |I|$  vertices in  $H$  which are not in  $G$ , let  $s_1$  have degree  $\delta$  and  $s_2 = s - s_1$  have degree  $\Delta$  in  $H$ . Let  $j$  represent the number of vertices in  $V_1$  that have degree  $\Delta$  in  $H$ . Then  $0 \leq j \leq m_1$ , and  $j = m_1$  forces  $s_1$  to be at least one. Let  $F = \langle I \rangle$  denote the subgraph of  $H$  induced by the set  $I$ . With  $k = |E(F)|$  denoting the size of  $F$ , we observe that  $0 \leq k \leq s(s-1)/2$ .

The graph  $H$  contains  $(m_1 - j + s_1)$  vertices of degree  $\delta$  and  $(p - m_1 + j + s_2)$  vertices of degree  $\Delta$ . Hence

$$\delta(m_1 - j + s_1) + \Delta(p - m_1 + j + s_2) \text{ is even.} \tag{1}$$

We may observe that  $H = G$  in case  $V^* = \emptyset$ . Otherwise, if  $u \in V_1$  and  $\deg_H u = \Delta$ , then  $s \geq \Delta - \delta$ ; and, if  $v \in V_2$  so that  $\deg_G v = a_2$  and  $\deg_H v = \Delta$ , then  $s \geq \Delta - a_2$ . Thus,

$$s = s_1 + s_2 \geq \begin{cases} \Delta - a_2 & \text{if } j = 0 \\ \Delta - \delta & \text{if } j \geq 1. \end{cases} \tag{2}$$

We may count the number  $e$  of edges in  $H$  between the sets  $V(G)$  and  $I$  in two ways. The set  $V_1$  has  $j$  vertices of degree  $\Delta$  in  $H$  and the vertices in  $V^*$  result in the regular deficiency  $\sigma$ . Then  $e = \sigma + j(\Delta - \delta)$ . Moreover the graph  $F$  has size  $k$ , and the set  $I$  contains  $s_1$  vertices of degree  $\delta$  and  $s_2$  vertices of degree  $\Delta$  in  $H$ . So  $e = \delta s_1 + \Delta s_2 - 2k$ . Thus

$$\delta s_1 + \Delta s_2 - 2k = \sigma + j(\Delta - \delta). \tag{3}$$

We also observe that these  $e$  edges induce a bipartite graph on the set  $V(G) \cup I$ . It is possible to describe this more precisely, which we now do. Let  $\mathcal{F}: f_1, f_2, \dots, f_s$  denote the sequence of degrees of the vertices of  $F$ , where  $\sum_{i=1}^s f_i = 2k$ . For each permutation  $\pi$  on  $\{1, 2, \dots, s\}$ , consider the sequence  $\mathcal{S}_\pi: b_1, b_2, \dots, b_s$ , where

$$b_i = \begin{cases} \Delta - f_{\pi(i)} & \text{if } 1 \leq i \leq s_2 \\ \delta - f_{\pi(i)} & \text{if } s_2 < i \leq s_1 + s_2 \end{cases}$$

is nonnegative. Also consider the sequence  $\mathcal{S}_2$  whose terms are  $(\Delta - \deg_G v)$ , where  $v \in V^*$  if  $j=0$ , and  $1 \leq j \leq m_1$  implies that  $v \in V^* \cup V_1(j)$ , where  $V_1(j)$  denotes a  $j$ -element subset of  $V_1$ . Then  $\mathcal{S}_2$  has  $n = p - (m_1 + m_n) + j$  terms. Let us write this sequence as  $\mathcal{S}_2: c_1, c_2, \dots, c_n$ .

The pair of sequences  $\mathcal{S}_1: b_1, b_2, \dots, b_s$  and  $\mathcal{S}_2: c_1, c_2, \dots, c_n$  is called **bigraphical** (see [2]) if there exists a bipartite graph  $B$  with partite sets  $U_1 = \{u_1, u_2, \dots, u_s\}$  and  $U_2 = \{w_1, w_2, \dots, w_n\}$  such that  $\deg_B u_i = b_i$ ,  $1 \leq i \leq s$ , and  $\deg_B w_j = c_j$ ,  $1 \leq j \leq n$ . Necessary and sufficient conditions were obtained in [2] for a pair of sequences of nonnegative integers to be bigraphical. We state one such condition for later use.

**Theorem 2.** *Let  $\mathcal{S}_1: b_1, b_2, \dots, b_s$  and  $\mathcal{S}_2: c_1, c_2, \dots, c_n$  be a pair of sequences of nonnegative integers with*

$$b_1 \geq b_2 \geq \dots \geq b_s,$$

$$c_1 \geq c_2 \geq \dots \geq c_n,$$

and

$$\sum_{i=1}^s b_i = \sum_{j=1}^n c_j.$$

Then the pair of sequences  $(\mathcal{S}_1; \mathcal{S}_2)$  is bigraphical if and only if the pair of sequences  $(\mathcal{S}'_1; \mathcal{S}'_2)$  is bigraphical where

$$\mathcal{S}'_1: b_1 - 1, b_2 - 1, \dots, b_{c_1} - 1, b_{c_1+1}, \dots, b_s$$

and

$$\mathcal{S}'_2: c_2, c_3, \dots, c_n.$$

We can now state the following condition:

there exists a graphical sequence  $\mathcal{F}$  for which some pair of  
sequences  $(\mathcal{S}_1; \mathcal{S}_2)$  is bigraphical. (4)

We have now shown that the conditions (1)—(4) are necessary for a graph  $H$  of minimum order  $p + s$  (where  $s = s_1 + s_2$ ) to exist. These conditions also prove to be sufficient. In order to see this let  $G$  be a given graph with degree set  $\vartheta_G = \{a_1, a_2, \dots, a_n\}$ , where  $\delta = a_1 < a_2 < \dots < a_n = \Delta$  and  $n \geq 2$ , and let  $s = s_1 + s_2$  (where  $s_1, s_2$  are nonnegative integers) be the least positive integer for which there exist integers  $j$  and  $k$ ,  $0 \leq j \leq m_1$  and  $0 \leq k \leq s(s-1)/2$  such that (1)—(4) are satisfied. By (4) there exists a graphical sequence  $\mathcal{F}: f_1, f_2, \dots, f_s$ . Let  $F$  be a graph having degree sequence  $\mathcal{F}$  where, then, the size of  $F$  is  $k$ . Also by (4) some pair of sequences  $(\mathcal{S}_1; \mathcal{S}_2)$  is bigraphical, so there exists a bipartite graph  $B$  with partite sets  $U_1 = \{u_1, u_2, \dots, u_s\}$  and  $U_2 = \{w_1, w_2, \dots, w_n\}$  such that  $\deg_B u_i = b_i$ ,  $1 \leq i \leq s$ , and

$\deg_B w_j = c_j$ ,  $1 \leq j \leq n$ . We now define a graph  $H$  by  $V(H) = U_1 \cup V(G)$ , where  $U_1 = V(F)$ ,  $U_2 = V^* \cup V_1(j)$  and  $E(H) = E(B) \cup E(F) \cup E(G)$ . Clearly  $G$  is an induced subgraph of  $H$  and  $\vartheta_H = \{\delta, \Delta\}$ . Thus, the following result has been verified.

**Theorem 3.** *Let  $G$  be a graph with degree set  $\vartheta_G = \{a_1, a_2, \dots, a_n\}$ , where  $\delta = a_1 < a_2 < \dots < a_n = \Delta$  and  $n \geq 2$ . Let  $H$  be a graph with degree set  $\vartheta_H = \{\delta, \Delta\}$  containing  $G$  as an induced subgraph. A necessary and sufficient condition that  $p + s$  be the least possible order for  $H$  is that  $s = s_1 + s_2$  is the least integer satisfying:*

- (1)  $\delta(m_1 - j + s_1) + \Delta(p - m_1 + j + s_2)$  is even,
- (2)  $s = s_1 + s_2 \geq \begin{cases} \Delta - a_2 & \text{if } j = 0 \\ \Delta - \delta & \text{if } j \geq 1, \end{cases}$
- (3)  $\delta s_1 + \Delta s_2 - 2k = \sigma + j(\Delta - \delta)$ , and
- (4) there exists a graphical sequence  $\mathcal{F}$  for which some pair of sequences  $(\mathcal{S}_1; \mathcal{S}_2)$  is bigraphical.


We illustrate the procedure by an example. Let  $G$  be a graph with degree sequence

$$9, 9, 9, 8, 8, 6, 6, 6, 6, 6, 5, 5, 5.$$

Here  $\delta = 5$ ,  $\Delta = 9$ ,  $\sigma = 17$ ,  $m_1 = 3$ ,  $p = 13$  and  $a_2 = 6$ . By (1),  $5(3 - j + s_1) + 9(13 - 3 + j + s_2)$  is even, and this implies that  $s_1$  and  $s_2$  have opposite parity and  $s$  is odd. Condition (2) implies that  $s_1 + s_2 \geq 3$  if  $j = 0$ ; and  $j = 1, 2$  or  $3$  implies that  $s_1 + s_2 \geq 5$ , since  $s = s_1 + s_2$  is odd and at least 4. Also, (3) states that  $5s_1 + 9s_2 - 2k = 17 + 4j$ .

Consider  $s = s_1 + s_2 = 3$ . Then  $j$  must be zero, and  $0 \leq k \leq \binom{3}{2} = 3$ . (i) If  $s_1 = 0$  and  $s_2 = 3$ , then  $k = 5$ . (ii) If  $s_1 = 1$  and  $s_2 = 2$ , then  $k = 3$  and  $F \cong K_3$ . This implies that  $\mathcal{F}: 2, 2, 2; \mathcal{S}_1: 7, 7, 3$ ; and  $\mathcal{S}_2: 3, 3, 3, 3, 3, 1, 1$ . Here the conditions (1), (2) and (3) hold. Moreover the sequence  $\mathcal{F}$  is graphical. But a repeated application of Theorem 2 shows that the pair of sequences  $(\mathcal{S}_1; \mathcal{S}_2)$  is not bigraphical. So (4) fails to hold. (iii) If  $s_1 = 2$  and  $s_2 = 1$ , then  $k = 1$  and  $F \cong K_1 \cup K_2$ . Hence  $\mathcal{F}: 1, 1, 0; \mathcal{S}_1: 9, 4, 4$  or  $\mathcal{S}_1: 8, 5, 4$ ; and  $\mathcal{S}_2: 3, 3, 3, 3, 3, 1, 1$ . Once again we use Theorem 2 to observe that  $(\mathcal{S}_1; \mathcal{S}_2)$  is not bigraphical. (iv) If  $s_1 = 3$  and  $s_2 = 0$ , then  $k < 0$ . Thus,  $s \geq 5$ .

We consider  $s_1 = 0$ ,  $s_2 = 5$ ,  $j = 2$ . Then  $k = 10$  and  $F \cong K_5$ . Now  $\mathcal{F}: 4, 4, 4, 4, 4; \mathcal{S}_1: 5, 5, 5, 5, 5$ ; and  $\mathcal{S}_2: 4, 4, 3, 3, 3, 3, 3, 1, 1$ . The pair  $(\mathcal{S}_1; \mathcal{S}_2)$  is easily seen to be bigraphical (by Theorem 2). In the figure below we have shown the essential sequences and the graph  $B$ . ( $E(G)$  and  $E(F)$  are not shown.) Clearly  $\vartheta_H = \{5, 9\}$ .



#### REFERENCES

- [1] BEHZAD, M.—CHARTRAND, G.—LESNIAK-FOSTER, L.: Graphs and Digraphs. Prindle, Weber and Schmidt, Boston 1979.
- [2] CHARTRAND, G.—GOULD, R. J.—KAPOOR, S. F.: Bigraphical sequences, submitted for publication.
- [3] ERDŐS, P.—KELLY, P.: The minimal regular graph containing a given graph. A Seminar on Graph Theory (F. Harary Ed.) Holt, Rinehart and Winston, New York 1967, 65—69.

Received June 14, 1978

*Gary Chartrand*  
*San José State University*  
*Ronald J. Gould*  
*Western Michigan University*  
*S. F. Kapoor*  
*Western Michigan University*

#### МИНИМАКСНЫЙ НАДГРАФ

Гары Чартрэнд—Роналд Дж. Гулд—Ц. Ф. Капур

#### Резюме

Пусть  $\vartheta_G$  обозначает множество всех степеней вершин графа  $G$ ,  $\max \vartheta_G = \Delta$ ,  $\min \vartheta_G = \delta$ . Если  $S$  — множество такое, что  $\Delta \in S \subseteq \vartheta_G$ , то существует граф  $H$  с множеством  $\vartheta_H = S$ , для которого  $G$  является порожденным подграфом. В случае  $S = \{\delta, \Delta\}$  находится необходимое и достаточное условие для того, чтобы число вершин графа  $H$  было минимальным.