

Zbigniew Walczak

On a class of Szász-Mirakyan type operators

Czechoslovak Mathematical Journal, Vol. 58 (2008), No. 3, 705–716

Persistent URL: <http://dml.cz/dmlcz/140415>

Terms of use:

© Institute of Mathematics AS CR, 2008

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON A CLASS OF SZÁSZ-MIRAKYAN TYPE OPERATORS

ZBIGNIEW WALCZAK, Poznań

(Received June 27, 2006)

Abstract. The actual construction of the Szász-Mirakyan operators and its various modifications require estimations of infinite series which in a certain sense restrict their usefulness from the computational point of view. Thus the question arises whether the Szász-Mirakyan operators and their generalizations cannot be replaced by a finite sum. In connection with this question we propose a new family of linear positive operators.

Keywords: linear positive operator, polynomial weighted space

MSC 2010: 41A36

1. INTRODUCTION

Approximation properties of the Szász-Mirakyan operators

$$(1) \quad S_n(f; x) := e^{-nx} \sum_{k=0}^{\infty} \frac{(nx)^k}{k!} f\left(\frac{k}{n}\right), \quad x \in \mathbb{R}_0 = [0, +\infty), \quad n \in \mathbb{N} := \{1, 2, \dots\},$$

in polynomial weighted spaces C_p were examined in [2]. The space C_p , $p \in \mathbb{N}_0 := \{0, 1, 2, \dots\}$, considered in [2] is associated with the weight function

$$(2) \quad w_0(x) := 1, \quad w_p(x) := (1 + x^p)^{-1} \quad \text{if } p \geq 1,$$

and consists of all real-valued functions f continuous on \mathbb{R}_0 and such that $w_p f$ is uniformly continuous and bounded on \mathbb{R}_0 . The norm on C_p is defined by

$$(3) \quad \|f\|_p \equiv \|f(\cdot)\|_p := \sup_{x \in \mathbb{R}_0} w_p(x) |f(x)|.$$

These operators are very interesting approximation processes and have many nice properties.

For f the results on the degree of approximation were studied in [2] for the usual Szász-Mirakyan operators (1). From these theorems it was deduced that

$$(4) \quad \lim_{n \rightarrow \infty} S_n(f; x) = f(x)$$

for every $f \in C_p$, $p \in \mathbb{N}_0$ and $x \in \mathbb{R}_0$. Moreover, the above convergence is uniform on every interval $[x_1, x_2]$, $x_1 \geq 0$.

Recently, in many papers various modifications of S_n were introduced and examined. They have been studied intensively in connection with different branches of analysis such as convex and numerical analysis. We refer the reader to P. Gupta and V. Gupta [9], V. Gupta [10], N. Ispir and C. Atakut [1], [18], V. Gupta, V. Vasishtha and M. K. Gupta [13], G. Feng [7], [8], A. Ciupa [5], N. Ispir [17], S. Li [21], X. Linsen and Z. Xiaoping [22]. Their results improve other related results in literature.

The actual construction of the Szász-Mirakyan operators (the Baskakov operators, the Favard operators, the Meyer-König and Zeller operators) and its many various modifications (see, for example, the works cited above) requires estimations of infinite series which in a certain sense restrict their usefulness from the computational point of view. Thus the question arises whether the Szász-Mirakyan operators and their generalizations cannot be replaced by a finite sum. In connection with this question, in [33] certain positive linear operators were considered, namely

$$(5) \quad L_n(f; x) := \frac{1}{(1 + (x + n^{-1})^2)^n} \sum_{k=0}^n \binom{n}{k} (x + n^{-1})^{2k} f\left(\frac{k}{n} \cdot \frac{1 + (x + n^{-1})^2}{x + n^{-1}}\right),$$

$x \in \mathbb{R}_0$, $n \in \mathbb{N}$, for a function of one variable.

In [33] it was proved that if $f \in C_p$, $p \in \mathbb{N}_0$, then

$$(6) \quad \lim_{n \rightarrow \infty} \|L_n(f; \cdot) - f(\cdot)\|_p = 0.$$

The operators L_n are defined in terms of a sample of the given function f on the points $(k/n) \cdot (1 + (x + n^{-1})^2)/(x + n^{-1})$ for $k \in \mathbb{N}_0$, $n \in \mathbb{N}$ and $x \in \mathbb{R}_0$.

Thus new questions arise whether the knots $(k/n) \cdot (1 + (x + n^{-1})^2)/(x + n^{-1})$ cannot be replaced by a given subset of points which are independent of x , provided this would not change essentially the approximation properties.

In connection to these questions we propose a new family of linear positive operators. This together with the simple form of the operator makes the results given in the present paper more helpful from the computational point of view.

This note was inspired by the results obtained in our previous papers. In the papers [34], [29] and [36], for each function f of polynomial type defined on \mathbb{R}_0 , the

following operators were considered:

$$(7) \quad A_n^{\{1\}}(f; r; x) = \frac{1}{g(nx; r)} \sum_{k=0}^{\infty} \frac{(nx)^k}{(k+r)!} f\left(\frac{k+r}{n}\right), \quad x \in \mathbb{R}_0,$$

$$(8) \quad A_n^{\{2\}}(f; r; x) = \frac{1}{g((nx+1)^2; r)} \sum_{k=0}^{\infty} \frac{(nx+1)^{2k}}{(k+r)!} f\left(\frac{k+r}{n(nx+1)}\right),$$

and

$$(9) \quad A_n^{\{3\}}(f; p, r, s; x) = \frac{1}{g(n^s x; r)} \sum_{k=0}^{\infty} \frac{(n^s x)^k}{(k+r)!} \sum_{j=0}^p \frac{f^{(j)}\left(\frac{k+r}{n^s}\right)}{j!} \left(x - \frac{k+r}{n^s}\right)^j,$$

where $n, r, p \in \mathbb{N}$, $s > 0$ and

$$(10) \quad g(t; r) = \sum_{k=0}^{\infty} \frac{t^k}{(k+r)!}, \quad t \in \mathbb{R}_0,$$

i.e.

$$g(0; r) = \frac{1}{r!}, \quad g(t, r) = \frac{1}{t^r} \left(e^t - \sum_{j=0}^{r-1} \frac{t^j}{j!} \right) \quad \text{if } t > 0.$$

Similar results in exponential weighted spaces can be found in [28], [30].

In this paper we will use the same method to obtain a new operator.

We introduce the following class of operators in C_p , $p \in \mathbb{N}$.

Definition 1. We introduce a class of new operators defined by

$$(11) \quad B_n(f; r; a_n; x) := \frac{1}{g(nx; r)} \sum_{k=0}^{[n(x+a_n)]} \frac{(nx)^k}{(k+r)!} f\left(\frac{k+r}{n}\right), \quad x \in \mathbb{R}_0, \quad n \in \mathbb{N},$$

where r is a fixed natural number and $(a_n)_1^\infty$ is a sequence of positive numbers such that $\lim_{n \rightarrow \infty} n^{1/2} a_n = \infty$ and $[n(x+a_n)]$ denotes the integral part of $n(x+a_n)$.

Observe that the operator B_n is linear and positive.

Moreover, we will introduce certain linear positive operators in polynomial weighted spaces of functions of two variables.

Let $p, q \in \mathbb{N}_0$ and let

$$(12) \quad w_{p,q}(x, y) := w_p(x)w_q(y), \quad (x, y) \in \mathbb{R}_0^2 := \mathbb{R}_0 \times \mathbb{R}_0,$$

where $w_p(\cdot)$ is defined by (2). Denote by $C_{p,q}$ the weighted space of all real-valued functions f continuous on \mathbb{R}_0^2 for which $w_{p,q}f$ is uniformly continuous and bounded on \mathbb{R}_0^2 . The norm on $C_{p,q}$ is defined by

$$(13) \quad \|f\|_{p,q} \equiv \|f(\cdot, \cdot)\|_{p,q} := \sup_{(x,y) \in \mathbb{R}_0^2} w_{p,q}(x,y)|f(x,y)|.$$

Approximation properties of linear positive operators

$$(14) \quad L_{m,n}(f; x, y) = \frac{1}{(1 + (x + m^{-1})^2)^m (1 + (y + n^{-1})^2)^n} \sum_{j=0}^m \sum_{k=0}^n \binom{m}{j} \binom{n}{k} \\ \times (x + m^{-1})^{2j} (y + n^{-1})^{2k} f\left(\frac{j(1 + (x + m^{-1})^2)}{m(x + m^{-1})}, \frac{k(1 + (y + n^{-1})^2)}{n(y + n^{-1})}\right), \\ A_{m,n}^{\{1\}}(f; r, s; x, y) = \frac{1}{g(mx; r)g(ny; s)} \\ \times \sum_{j=0}^{\infty} \sum_{k=0}^{\infty} \frac{(mx)^j}{(j+r)!} \frac{(ny)^k}{(k+s)!} f\left(\frac{j+r}{m}, \frac{k+s}{n}\right), \quad r, s \in \mathbb{N},$$

in polynomial weighted spaces of functions of two variables were examined in [31] and [34].

In this paper we will give some properties of the following operators.

Definition 2. Fix $r, s \in \mathbb{N}$ and $p, q \in \mathbb{N}$. We define the class of operators $B_{m,n}$ by

$$(15) \quad B_{m,n}(f; r, s; a_m, b_n; x, y) := \frac{1}{g(mx; r)g(ny; s)} \\ \times \sum_{j=0}^{[m(x+a_m)]} \sum_{k=0}^{[n(y+b_n)]} \frac{(mx)^j}{(j+r)!} \frac{(ny)^k}{(k+s)!} f\left(\frac{j+r}{m}, \frac{k+s}{n}\right), \quad f \in C_{p,q}, \quad (x, y) \in \mathbb{R}_0^2,$$

where $(a_m)_1^\infty$ and $(b_n)_1^\infty$ are given sequences of positive numbers such that

$$\lim_{m \rightarrow \infty} m^{1/2} a_m = \infty \quad \text{and} \quad \lim_{n \rightarrow \infty} n^{1/2} b_n = \infty.$$

Observe that the operator $B_{m,n}$ is linear and positive.

In this paper, by $K_i(\alpha, \beta)$, $i = 1, 2, \dots$, we denote suitable positive constants depending only on the parameters α and β .

2. PRELIMINARIES

In this section we will give some properties of the above operators which we will apply to the proofs of the main theorems.

It is known ([2]) that

$$(16) \quad S_n(1; x) = 1, \quad S_n(t - x; x) = 0,$$

$$(17) \quad S_n((t - x)^{q+1}; x) = \frac{x}{n} \{S'_n((t - x)^q; x) + qS_n((t - x)^{q-1}; x)\},$$

for $x \in \mathbb{R}_0$, $n \in \mathbb{N}$ and $q \in \mathbb{N}$.

Using (16), (17) and mathematical induction on $q \in \mathbb{N}$ we can prove the following lemma.

Lemma 1 ([25]). *For every $2 \leq q \in \mathbb{N}$ we have*

$$S_n((t - x)^q; x) = \sum_{j=1}^{[q/2]} c_{j,q} \frac{x^j}{n^{q-j}}, \quad x \in \mathbb{R}_0, \quad n \in \mathbb{N},$$

where $c_{j,q}$ are positive numerical coefficients depending only on j and q ($[y]$ denotes the integral part of $y \in \mathbb{R}_0$).

In the paper [34] the following results were proved for $A_n^{\{1\}}(f)$ defined by (7).

Lemma 2. $A_n^{\{1\}}$ defines a positive linear operator $C_p \rightarrow C_p$.

Theorem 1. For every fixed $r \in \mathbb{N}$ and $f \in C_p$, $p \in \mathbb{N}_0$, we have

$$(18) \quad \lim_{n \rightarrow \infty} \{A_n^{\{1\}}(f; r; x) - f(x)\} = 0, \quad x \in \mathbb{R}_0.$$

Moreover, (18) holds uniformly on every interval $[x_1, x_2]$, $x_2 > x_1 \geq 0$.

In [34] it was proved that if $f \in C_{p,q}$, $p, q \in \mathbb{N}_0$, then $A_{m,n}^{\{1\}}$ is a positive linear operator $C_{p,q} \rightarrow C_{p,q}$. Moreover, we derived

Theorem 2. Suppose that $f \in C_{p,q}$, $p, q \in \mathbb{N}_0$. Then there exists a positive constant $K_3(p, q, r, s)$ such that for all $(x, y) \in \mathbb{R}_0^2$

$$(19) \quad w_{p,q}(x, y) |A_{m,n}^{\{1\}}(f; r, s; x, y) - f(x, y)| \\ \leq K_3(p, q, r, s) \omega \left(f, C_{p,q}; \sqrt{\frac{x+1}{m}}, \sqrt{\frac{y+1}{n}} \right),$$

$m, n \in \mathbb{N}$, $r, s \in \mathbb{N}$, where

$$(20) \quad \omega(f, C_{p,q}; t, s) := \sup_{0 \leq h \leq t, 0 \leq \delta \leq s} \|\Delta_{h,\delta} f(\cdot, \cdot)\|_{p,q}, \quad t, s \geq 0,$$

$\Delta_{h,\delta} f(x, y) := f(x+h, y+\delta) - f(x, y)$, $(x+h, y+\delta) \in \mathbb{R}_0^2$ is the modulus of continuity of $f \in C_{p,q}$.

From (20) it follows that

$$(21) \quad \lim_{t,s \rightarrow 0+} \omega(f, C_{p,q}; t, s) = 0$$

for every $f \in C_{p,q}$, $p, q \in \mathbb{N}_0$. This implies that

$$(22) \quad \lim_{m,n \rightarrow \infty} A_{m,n}^{\{1\}}(f; r, s; x, y) = f(x, y), \quad (x, y) \in \mathbb{R}_0^2$$

uniformly on every rectangle $0 \leq x \leq x_0$, $0 \leq y \leq y_0$.

3. MAIN RESULTS

Now we give an approximation theorem for B_n .

Theorem 3. Fix $p, r \in \mathbb{N}$. Then for B_n defined by (11) we have

$$(23) \quad \lim_{n \rightarrow \infty} \{B_n(f; r; a_n; x) - f(x)\} = 0, \quad f \in C_p$$

uniformly on every interval $[x_1, x_2]$, $x_2 > x_1 \geq 0$.

Proof. We first suppose that $f \in C_p$, $p \in \mathbb{N}$. From (11) and (7) we obtain

$$\begin{aligned} B_n(f; r; a_n; x) - f(x) &= \frac{1}{g(nx; r)} \sum_{k=0}^{[n(x+a_n)]} \frac{(nx)^k}{(k+r)!} f\left(\frac{k+r}{n}\right) - f(x) \\ &= \frac{1}{g(nx; r)} \sum_{k=0}^{\infty} \frac{(nx)^k}{(k+r)!} f\left(\frac{k+r}{n}\right) - f(x) \\ &\quad - \frac{1}{g(nx; r)} \sum_{k=[n(x+a_n)]+1}^{\infty} \frac{(nx)^k}{(k+r)!} f\left(\frac{k+r}{n}\right) \\ &= A_n^{\{1\}}(f; r; x) - f(x) - M_n(f; r; x), \quad x \in \mathbb{R}_0, \quad n, r \in \mathbb{N}. \end{aligned}$$

By our assumption, using the elementary inequality $(a+b)^k \leq 2^{k-1}(a^k + b^k)$, $a, b > 0$, $k \in \mathbb{N}_0$, we get

$$(24) \quad |f(t)| \leq K_1(1+t^p) \leq K_1(1+(|t-x|+x)^p) \leq K_1(1+2^{p-1}(|t-x|^p+x^p)).$$

Observe that

$$\begin{aligned}
 |M_n(f; r; x)| &\leq \frac{1}{g(nx; r)} \sum_{k=[n(x+a_n)]+1}^{\infty} \frac{(nx)^k}{(k+r)!} \left| f\left(\frac{k+r}{n}\right) \right| \\
 &\leq \frac{1}{g(nx; r)} \sum_{k=[n(x+a_n)]+r+1}^{\infty} \frac{(nx)^{k-r}}{k!} \left| f\left(\frac{k}{n}\right) \right| \\
 &\leq \frac{1}{(nx)^r g(nx; r)} \sum_{k=[n(x+a_n)]+1}^{\infty} \frac{(nx)^k}{k!} \left| f\left(\frac{k}{n}\right) \right|.
 \end{aligned}$$

This together with (10), (24) and (1) yields

$$\begin{aligned}
 |M_n(f; r; x)| &\leq \frac{1}{(nx)^r g(nx; r)} \sum_{k=[n(x+a_n)]+1}^{\infty} \frac{(nx)^k}{k!} K_1 \left(1 + 2^{p-1} \left(\left| \frac{k}{n} - x \right|^p + x^p \right) \right) \\
 &\leq \frac{e^{nx}}{e^{nx} - \sum_{j=0}^{r-1} (nx)^j / j!} K_1 \\
 &\quad \times \left((1 + 2^{p-1} x^p) e^{-nx} \sum_{k=[n(x+a_n)]+1}^{\infty} \frac{(nx)^k}{k!} + 2^{p-1} e^{-nx} \sum_{k=0}^{\infty} \frac{(nx)^k}{k!} \left| \frac{k}{n} - x \right|^p \right) \\
 &= \frac{e^{nx}}{e^{nx} - \sum_{j=0}^{r-1} (nx)^j / j!} K_1 \\
 &\quad \times \left((1 + 2^{p-1} x^p) e^{-nx} \sum_{k=[n(x+a_n)]+1}^{\infty} \frac{(nx)^k}{k!} + 2^{p-1} S_n(|t-x|^p; x) \right).
 \end{aligned}$$

We remark that

$$\frac{e^t}{e^t - \sum_{j=0}^{r-1} t^j / j!} = O(1)$$

and

$$\begin{aligned}
 e^{-nx} \sum_{k=[n(x+a_n)]+1}^{\infty} \frac{(nx)^k}{k!} &\leq e^{-nx} \sum_{a_n < |k/n - x|}^{\infty} \frac{(nx)^k}{k!} \\
 &\leq e^{-nx} \sum_{a_n < |k/n - x|}^{\infty} \frac{(nx)^k}{k!} \left| \frac{k}{n} - x \right|^p \frac{1}{a_n^p} \\
 &\leq \frac{1}{a_n^p} e^{-nx} \sum_{k=0}^{\infty} \frac{(nx)^k}{k!} \left| \frac{k}{n} - x \right|^p = \frac{1}{a_n^p} S_n(|t-x|^p; x).
 \end{aligned}$$

This implies that

$$|M_n(f; r; x)| \leq K_2 \left(\frac{(1 + 2^{p-1}x^p)}{a_n^p} + 2^{p-1} \right) S_n(|t - x|^p; x).$$

Consequently, in view of Lemma 1, the Hölder inequality and (16), we further have

$$\begin{aligned} |M_n(f; r; x)| &\leq K_2 \left(\frac{(1 + 2^{p-1}x^p)}{a_n^p} + 2^{p-1} \right) \{S_n((t - x)^{2p}; x)S_n(1; x)\}^{1/2} \\ &= K_2 \left(\frac{(1 + 2^{p-1}x^p)}{a_n^p} + 2^{p-1} \right) \left\{ \sum_{j=1}^p c_{j,2p} \frac{x^j}{n^{2p-j}} \right\}^{1/2} \\ &\leq \frac{K_2}{n^{p/2}} \left(\frac{(1 + 2^{p-1}x^p)}{a_n^p} + 2^{p-1} \right) \left\{ \sum_{j=1}^p c_{j,2p} x^j \right\}^{1/2}. \end{aligned}$$

The relation

$$\lim_{n \rightarrow \infty} n^{1/2} a_n = \infty$$

implies that

$$\lim_{n \rightarrow \infty} M_n(f; r; x) = 0$$

uniformly on every interval $[x_1, x_2]$, $x_2 > x_1 \geq 0$. In view of Theorem 1 this yields

$$\lim_{n \rightarrow \infty} \{B_n(f; r; a_n; x) - f(x)\} = 0,$$

uniformly on every interval $[x_1, x_2]$, $x_2 > x_1 \geq 0$. This completes the proof of (23). \square

Applying Theorem 2, we can prove the basic property of $B_{m,n}$.

Theorem 4. Fix $p, q, r, s \in \mathbb{N}$. Then for $B_{m,n}$ defined by (15) we have

$$(25) \quad \lim_{m,n \rightarrow \infty} B_{m,n}(f; r, s; a_m, b_n; x, y) = f(x, y), \quad f \in C_{p,q}.$$

Moreover, (25) holds uniformly on every rectangle $0 \leq x \leq x_0$, $0 \leq y \leq y_0$.

Proof. Suppose that $f \in C_{p,q}$, $p, q \in \mathbb{N}$ and $r, s \in \mathbb{N}$. This implies that

$$|f(t, z)| \leq K_4(1 + t^p)(1 + z^q) \leq K_4(1 + 2^{p-1}(|t - x|^p + x^p))(1 + 2^{q-1}(|z - y|^q + y^q)).$$

From (15) and (14) we have

$$B_{m,n}(f; r, s; a_m, b_n; x) - f(x, y) = A_{m,n}^{\{1\}}(f; r, s; x, y) - f(x, y) - M_{m,n}(f; r, s; x, y)$$

where

$$M_{m,n}(f; r, s; x, y) = \frac{1}{g(mx; r)g(ny; s)} \times \sum_{j=[m(x+a_m)]+1}^{\infty} \sum_{k=[n(y+b_n)]+1}^{\infty} \frac{(mx)^j}{(j+r)!} \frac{(ny)^k}{(k+s)!} f\left(\frac{j+r}{m}, \frac{k+s}{n}\right),$$

$$(x, y) \in \mathbb{R}_0^2.$$

Observe that

$$|M_{m,n}(f; r, s; x, y)| \leq \frac{1}{g(mx; r)g(ny; s)} \times \sum_{j=[m(x+a_m)]+1}^{\infty} \sum_{k=[n(y+b_n)]+1}^{\infty} \frac{(mx)^j}{(j+r)!} \frac{(ny)^k}{(k+s)!} \left| f\left(\frac{j+r}{m}, \frac{k+s}{n}\right) \right|$$

$$\leq K_4 \frac{1}{g(mx; r)} \sum_{j=[m(x+a_m)]+r+1}^{\infty} \frac{(mx)^{j-r}}{j!} \left(1 + 2^{p-1} \left(\left|\frac{j}{m} - x\right|^p + x^p\right)\right)$$

$$\times \frac{1}{g(ny; s)} \sum_{k=[n(y+b_n)]+s+1}^{\infty} \frac{(ny)^{k-s}}{k!} \left(1 + 2^{q-1} \left(\left|\frac{k}{n} - y\right|^q + y^q\right)\right).$$

Arguing as in the second part of Theorem 3 we derive

$$\frac{1}{g(mx; r)} \sum_{j=[m(x+a_m)]+r+1}^{\infty} \frac{(mx)^{j-r}}{j!} \left(1 + 2^{p-1} \left(\left|\frac{j}{m} - x\right|^p + x^p\right)\right)$$

$$\leq \frac{K_5}{m^{p/2}} \left(\frac{(1 + 2^{p-1}x^p)}{a_m^p} + 2^{p-1}\right) \left\{ \sum_{j=1}^p c_{j,2p} x^j \right\}^{1/2},$$

$$\frac{1}{g(ny; s)} \sum_{k=[n(y+b_n)]+s+1}^{\infty} \frac{(ny)^{k-s}}{k!} \left(1 + 2^{q-1} \left(\left|\frac{k}{n} - y\right|^q + y^q\right)\right)$$

$$\leq \frac{K_5}{n^{q/2}} \left(\frac{(1 + 2^{q-1}y^q)}{b_n^q} + 2^{q-1}\right) \left\{ \sum_{j=1}^q c_{j,2q} y^j \right\}^{1/2}.$$

This yields in view of Definition 2

$$\lim_{m,n \rightarrow \infty} M_{m,n}(f; r, s; x, y) = 0$$

uniformly on every rectangle $0 \leq x \leq x_0$, $0 \leq y \leq y_0$. Applying Theorem 2 and (26) we immediately obtain (25). \square

It is similarly verified that analogous approximation properties hold for the two operators

$$T_n(f; r; x) := \frac{1}{g(nx; r)} \sum_{k=0}^{nr} \frac{(nx)^k}{(k+r)!} f\left(\frac{k+r}{n}\right),$$

$f \in C_{[0,2r]}$, $x \in [0, r]$, $n, r \in \mathbb{N}$,

$$T_{m,n}(f; r, s; x, y) := \frac{1}{g(mx; r)g(ny; s)} \\ \times \sum_{j=0}^{mr} \sum_{k=0}^{ns} \frac{(mx)^j}{(j+r)!} \frac{(ny)^k}{(k+s)!} f\left(\frac{j+r}{m}, \frac{k+s}{n}\right),$$

$f \in C_{[0,2r],[0,2s]}$, $(x, y) \in [0, r] \times [0, s]$, $m, n, r, s \in \mathbb{N}$.

Observe that the operators T_n , $n \in \mathbb{N}$, are obtained from (11) for $a_n = r - x$, $x \in [0, r]$, $r \in \mathbb{N}$.

Analogously we obtain

$$B_{m,n}(f; r, s; r - x, s - y; x, y) = T_{m,n}(f; r, s; x, y), \\ (x, y) \in [0, r] \times [0, s], \quad m, n, r, s \in \mathbb{N}.$$

The methods used to prove the Theorems are similar to those used in the construction of the modified Szász-Mirakjan operators [19], [31], [34], [38].

Acknowledgment. The author would like to thank the referee for his valuable comments and proposals.

References

- [1] *C. Atakut, N. Ispir*: The order of approximation by certain linear positive operators. *Math. Balk., New Ser.* 15 (2001), 25–33.
- [2] *M. Becker*: Global approximation theorems for Szasz-Mirakjan and Baskakov operators in polynomial weight spaces. *Indiana Univ. Math. J.* 27 (1978), 127–142.
- [3] *M. Becker, D. Kucharski, R. J. Nessel*: Global Approximation theorems for the Szasz-Mirakjan operators in exponential weight spaces. *Linear Spaces and Approximation (Proc. Conf. Oberwolfach, 1977)*. ISNM, Int. Ser. Numer. Math. 40 (1978), 319–333.
- [4] *A. Ciupa*: On the approximation by Favard-Szasz type operators. *Rev. Anal. Numér. Théor. Approx.* 25 (1996), 57–61.
- [5] *A. Ciupa*: Approximation by a generalized Szasz type operators. *J. Comput. Anal. Appl.* 5 (2003), 413–424.
- [6] *R. A. De Vore, G. G. Lorentz*: *Constructive Approximation*. Springer, Berlin, 1993.
- [7] *G. Feng*: Weighted approximation on Szasz-type operators. *Anal. Theory Appl.* 19 (2003), 47–54.
- [8] *G. Feng*: A characterization of pointwise approximation for linear combinations of Szász-type operators. *Chinese Quart. J. Math.* 19 (2004), 379–384.

- [9] *P. Gupta, V. Gupta*: Rate of convergence on Baskakov-Szasz type operators. *Fasc. Math.* 31 (2001), 37–44.
- [10] *V. Gupta*: Error estimation for mixed summation integral type operators. *J. Math. Anal. Appl.* 313 (2006), 632–641.
- [11] *V. Gupta, P. Maheshwari*: On Baskakov-Szasz type operators. *Kyungpook Math. J.* 43 (2003), 315–325.
- [12] *V. Gupta, R. P. Pant*: Rate of convergence for the modified Szasz-Mirakjan operators on functions of bounded variation. *J. Math. Anal. Appl.* 233 (1999), 476–483.
- [13] *V. Gupta, V. Vasishtha, M. K. Gupta*: Rate of convergence of the Szasz-Kantorovitch-Bezier operators for bounded variation functions. *Publ. Inst. Math., Nouv. Sér.* 72 (2002), 137–143.
- [14] *S. Guo*: On the rate of convergence of the integrated Meyer-König and Zeller operators for functions of bounded variation. *J. Approximation Theory* 56 (1989), 245–255.
- [15] *S. Guo, C. Li, Y. Sun, G. Yand, S. Yue*: Pointwise estimate for Szasz-type operators. *J. Approximation Theory* 94 (1998), 160–171.
- [16] *M. Herzog*: Approximation theorems for modified Szasz-Mirakjan operators in polynomial weight spaces. *Matematyczne* 54 (1999), 77–90.
- [17] *N. Ispir*: Weighted approximation by modified Favard-Szász operators. *Int. Math. J.* 3 (2003), 1053–1060.
- [18] *N. Ispir, C. Atakut*: Approximation by modified Szasz-Mirakjan operators on weighted spaces. *Proc. Indian Acad. Sci., Math. Sci.* 112 (2002), 571–578.
- [19] *H. G. Lehnhoff*: On a modified Szasz-Mirakjan operator. *J. Approximation Theory* 42 (1984), 278–282.
- [20] *M. Lesniewicz, L. Rempulska*: Approximation by some operators of the Szasz-Mirakjan type in exponential weight spaces. *Glas. Mat., III. Ser.* 32 (1997), 57–69.
- [21] *S. Li*: Local smoothness of functions and Baskakov-Durrmeyer operators. *J. Approximation Theory* 88 (1997), 139–153.
- [22] *X. Linsen, Z. Xiaoping*: Pointwise characterization for combinations of Baskakov operators. *Approximation Theory Appl.* 18 (2002), 76–89.
- [23] *L. Rempulska, Z. Walczak*: On modified Baskakov operators. *Proc. A. Razmadze Math. Inst.* 133((2003), 109–117.
- [24] *L. Rempulska, Z. Walczak*: Approximation by some operators of Szasz-Mirakjan type. *Anal. Theory Appl.* 20 (2004), 1–15.
- [25] *L. Rempulska, Z. Walczak*: Modified Szasz-Mirakjan operators. *Math. Balk., New Ser.* 18 (2004), 53–63.
- [26] *A. Sahai, G. Prasad*: On simultaneous approximation by modified Lupas operators. *J. Approximation Theory* 45 (1985), 122–128.
- [27] *V. Totik*: Uniform approximation by Szász-Mirakjan type operators. *Acta Math.* 41 (1983), 291–307.
- [28] *Z. Walczak*: On certain linear positive operators in exponential weighted spaces. *Math. J. Toyama Univ.* 25 (2002), 109–118.
- [29] *Z. Walczak*: On certain positive linear operators in weighted polynomial spaces. *Acta Math.* 101 (2003), 179–191.
- [30] *Z. Walczak*: Approximation properties of certain linear positive operators in exponential weighted spaces. *Acta Univ. Palacki. Olomuc., Fac. Rerum Nat., Math.* 42 (2003), 123–130.
- [31] *Z. Walczak*: Approximation by some linear positive operators of functions of two variables. *Saitama Math. J.* 21 (2003), 23–31.
- [32] *Z. Walczak*: On the rate of convergence for modified Baskakov operators. *Liet. matem. rink* 44 (2004), 124–130.

- [33] *Z. Walczak*: Approximation by some linear positive operators in polynomial weighted spaces. *Publ. Math. Debrecen* *64* (2004), 353–367.
- [34] *Z. Walczak*: Approximation properties of certain linear positive operators in polynomial weighted spaces of functions of one and two variables. *Publ. Elektrotehn. Fak. Univ. Beograd* *15* (2004), 52–65.
- [35] *Z. Walczak*: On the convergence of the modified Szasz-Mirakyan operators. *Yokohama Math. J.* *51* (2004), 11–18.
- [36] *Z. Walczak*: On the rate of convergence for some linear operators. *Hiroshima Math. J.* *35* (2005), 115–124.
- [37] *B. Wood*: Uniform approximation with positive linear operators generated by binomial expansions. *J. Approximation Theory* *56* (1989), 48–58.
- [38] *S. Xiehua*: On the convergence of the modified Szasz-Mirakjan operator. *Approximation Theory Appl.* *10* (1994), 20–25.

Author's address: Z. Walczak, Institute of Mathematics, Poznań University of Technology, Piotrowo 3A, 60-965 Poznań, Poland, e-mail: zwalczak@math.put.poznan.pl.