

Jianping Liu; Bo Lian Liu

The maximum clique and the signless Laplacian eigenvalues

Czechoslovak Mathematical Journal, Vol. 58 (2008), No. 4, 1233–1240

Persistent URL: <http://dml.cz/dmlcz/140453>

Terms of use:

© Institute of Mathematics AS CR, 2008

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

THE MAXIMUM CLIQUE AND THE SIGNLESS
LAPLACIAN EIGENVALUES

JIANPING LIU, Fujian, BOLIAN LIU, Guangzhou

(Received June 27, 2007)

Abstract. Lower and upper bounds are obtained for the clique number $\omega(G)$ and the independence number $\alpha(G)$, in terms of the eigenvalues of the signless Laplacian matrix of a graph G .

Keywords: bound, clique number, independence number, signless Laplacian eigenvalues

MSC 2010: 05C50

1. INTRODUCTION

Let $G = (V, E)$ be a simple graph with n vertices. Denote $V(G) = \{v_1, v_2, \dots, v_n\}$. For any two vertices $v_i, v_j \in V(G)$ with $i < j$ we will use the symbol $i \sim j$ to denote the edge $v_i v_j$. Let A be the adjacency matrix of G and $D = \text{diag}(d_1, d_2, \dots, d_n)$ the diagonal matrix of vertex degrees. The matrix $L = D - A$ is known as the Laplacian of G and has been studied extensively in literature (see, e.g., [1]). The matrix $Q = D + A$ is called the signless Laplacian in [4] and appears very rarely in published papers (see [1]), the paper [2] being one of the very few research papers concerning this matrix. Since the signless Laplacian is a positive semi-definite matrix, all its eigenvalues are non-negative. Throughout the paper, we order the eigenvalues of the signless Laplacian matrix Q (Q -eigenvalues for short) of a graph G as $q_1 \geq q_2 \geq \dots \geq q_n \geq 0$, and let u_1, \dots, u_n be the corresponding normalized eigenvectors. We can suppose that $u_1 > 0$.

This work was supported by the National Natural Science Foundation of China (No. 10771080), SRFDP of China (No. 20070574006) and by the Foundation to the Educational Committee of Fujian (No. JB07020).

Given a graph G , define $\omega(G)$ and $\alpha(G)$ (ω and α for short), the clique number and the independence number of G to be the numbers of vertices of the largest clique and the largest independent set in G , respectively. Obviously, $\omega(G) = \alpha(G^c)$ where G^c is the complement of G .

In this paper we obtain lower and upper bounds for the independence number α and the clique number ω of a graph G . The bounds involve the signless Laplacian eigenvalues.

2. BOUNDS ON THE CLIQUE NUMBER AND INDEPENDENCE NUMBER OF ARBITRARY GRAPHS

We give bounds for the clique number ω and for the independence number α . They involve the largest Q -eigenvalue q_1 and the least Q -eigenvalue q_n of the graph G . The derivation of these bounds rests on a theorem of Motzkin and Straus. We recall the following important result.

Theorem 2.1 ([6]). *Let S be the simplex in E^n given by $x_i \geq 0$, $\sum_{i=1}^n x_i = 1$. Let ω be the order of the maximal clique in G and let A be the adjacency matrix of G . Then*

$$(1) \quad 1 - \frac{1}{\omega} = \max_{x \in S} \langle x, Ax \rangle.$$

Theorem 2.1 offers great possibilities for the investigation of the clique number by means of spectra. It will be used many times in the sequel. The dual theorem of Theorem 2.1 also holds, as shown by the following theorem.

Theorem 2.2. *Let α be the size of the largest independent set of G . Then*

$$(1') \quad \frac{1}{\alpha} = \min_{x \in S} \langle x, (I + A)x \rangle$$

where S and A are the same as in Theorem 2.1.

Lemma 2.1. *Let G be a graph with minimum degree δ . Then $q_n \leq \delta$.*

Proof. By the min-max characterization of the eigenvalues of a symmetric matrix, we have $q_n = \min_{x \neq 0} (x^t Q x / x^t x)$. Let $e_j = (0, \dots, 0, \overset{j}{1}, 0, \dots, 0)$, where j is a vertex with the minimum degree δ . Then $q_n \leq e_j^t Q e_j / e_j^t e_j = \delta$. □

Theorem 2.3. Let G be a graph with n vertices, m edges and minimum degree δ . Then

$$(2) \quad w \geq \frac{2m}{2m - (q_n - \delta)^2}.$$

Proof. Let $u_n = (y_1, y_2, \dots, y_n)$ be the normalized eigenvector corresponding to q_n . Then

$$\begin{aligned} q_n &= (u_n)^T Q u_n = \sum_{i \sim j} (y_i + y_j)^2 = \sum_{i=1}^n d_i y_i^2 + \sum_{i \sim j} 2y_i y_j \\ &\geq \delta \sum_{i=1}^n y_i^2 + \sum_{i \sim j} 2y_i y_j = \delta + \sum_{i \sim j} 2y_i y_j. \end{aligned}$$

By Lemma 2.1, $q_n - \delta \leq 0$. We have $(q_n - \delta)^2 \leq \left(\sum_{i \sim j} 2y_i y_j \right)^2$.

By the Cauchy inequality we obtain

$$\left(\sum_{i \sim j} 2y_i y_j \right)^2 \leq 4 \left(\sum_{i=1}^m 1^2 \right) \left(\sum_{i \sim j} (y_i y_j)^2 \right) = 2m \left(2 \sum_{i \sim j} y_i^2 y_j^2 \right).$$

Since $(y_1^2, y_2^2, \dots, y_n^2) \geq 0$ and $y_1^2 + y_2^2 + \dots + y_n^2 = 1$, by Theorem 2.1 we have

$$2 \sum_{i \sim j} y_i^2 y_j^2 \leq 1 - \frac{1}{w}.$$

Therefore

$$\frac{(q_n - \delta)^2}{2m} \leq 1 - \frac{1}{w},$$

that is

$$w \geq \frac{2m}{2m - (q_n - \delta)^2}.$$

□

Remark 2.1. In [5] it was proved that

$$(3) \quad w \geq \frac{2m}{2m - (\lambda_1 - \Delta)^2}$$

where Δ is the maximum degree and λ_1 is the largest Laplacian eigenvalue. We now show that the bounds (3) and (2) are incomparable.

Let G_1, G_2 be the graphs shown in Fig 1.




Fig. 1

By direct computation, we find that for G_1 , the bound (3) is 1.143 and better than the bound (2) 1.041 numerically, but the bounds are in practice equivalent (both bounds imply that the clique number is at least 2); and for G_2 , the bound (2) is 1.33 and better than the bound (3) 1.067 numerically, but the bounds are in practice equivalent (both bounds imply that the clique number is at least 2). And if $G = K_{1,n-1}$ (the star) or $G = K_n$ (the complete graph), the two bounds are equivalent numerically.

Note that the eigenvector u_1 is positive number. Let U be the sum of the entries of u_1 . Then the vector $x = u_1/U$ belongs to the simplex S of Eq. (1).

Theorem 2.4. *Let G be a graph with maximum degree Δ . Then*

$$(4) \quad w \geq \frac{n}{n - q_1 + \Delta}.$$

Proof. Let $u_1 = (r_1, r_2, \dots, r_n)$ be the normalized eigenvector corresponding to q_1 . Let $U = \sum_{i=1}^n r_i$ and $x = u_1/U$. Then

$$\langle x, Qx \rangle = \langle x, Dx \rangle + \langle x, Ax \rangle \leq \Delta \langle x, x \rangle + \left(1 - \frac{1}{w}\right) = \frac{\Delta}{U^2} + 1 - \frac{1}{w}.$$

On the other hand,

$$\langle x, Qx \rangle = \sum_{i \sim j} (x_i + x_j)^2 = \frac{1}{U^2} \sum_{i \sim j} (r_i + r_j)^2 = \frac{q_1}{U^2}.$$

Therefore

$$\frac{q_1}{U^2} \leq \frac{\Delta}{U^2} + \left(1 - \frac{1}{w}\right).$$

Since

$$U^2 = \left(\sum_{i=1}^n r_i\right)^2 \leq n \sum_{i=1}^n r_i^2 = n,$$

we obtain,

$$\frac{q_1 - \Delta}{n} \leq \frac{q_1 - \Delta}{U^2} \leq 1 - \frac{1}{w}.$$

That is,

$$w \geq \frac{n}{n - q_1 + \Delta}.$$

□

Remark 2.2. We now show that in many cases the bound (4) is better than both (2) and (3). For example, for $G = K_n$ the bound (4) is nicely equal to n and is better than (2) and (3), which gives $\omega(K_n) \geq (n^2 - n)/(n^2 - n - 1)$ (this implies that the clique number is at least 2, which is very trivial!); and for $n \leq 5$ the bound (4) is always better than (3), except for the P_5 (path of order 5). The bound (3) gives $\omega(P_5) \geq 1.486$, while the bound (4) gives $\omega(P_5) \geq 1.478$ (both bounds imply that the clique number is at least 2). And if G is a bipartite regular graph, the three bounds are equivalent.

By adapting a similar technique, we find the analogous result for α .

Theorem 2.5. *Let G be a graph with minimum degree δ . Then*

$$\alpha \geq \frac{U^2}{q_1^2 - \delta + 1},$$

where U is the same as in Theorem 2.4.

3. SOME RESULTS FOR REGULAR GRAPHS

We proceed now to the investigation of the clique number of regular graphs. For d -regular graphs we have $u_1 = e/\sqrt{n}$, where $e = (1, 1, \dots, 1)$ and $q_1 = 2d$. Naturally, all theorems of Section 2 hold also for regular graphs.

Theorem 3.1. *Let G be a d -regular graph. Then*

$$w \geq \frac{n^2}{n^2 - nd + (d - q_n)M^2},$$

where $M = \min_{y_i \neq 0} \{1/|y_i|\}$ and $u_n = (y_1, y_2, \dots, y_n)$ is the normalized eigenvector corresponding to q_n .

Proof. Let $\theta = M/n$ and $x = n^{-1}e + \theta u_n$. Then $\theta y_i \geq -1/n$ ($i = 1, 2, \dots, n$), and since $\langle e, u_n \rangle = 0$, it follows that x belongs to the simplex S of Eq. (1).

By Eq. (1) we have

$$\begin{aligned}\langle x, Qx \rangle &= \langle x, Dx \rangle + \langle x, Ax \rangle \\ &\leq d \langle x, x \rangle + \left(1 - \frac{1}{w}\right) = d\left(\frac{1}{n} + \theta^2\right) + \left(1 - \frac{1}{w}\right).\end{aligned}$$

On the other hand,

$$\begin{aligned}\langle x, Qx \rangle &= \left\langle \frac{e}{n} + \theta u_n, Q\left(\frac{e}{n} + \theta u_n\right) \right\rangle \\ &= \left\langle \frac{e}{n} + Q\frac{e}{n} \right\rangle + \left\langle \frac{e}{n}, Q\theta u_n \right\rangle + \left\langle \theta u_n, Q\frac{e}{n} \right\rangle + \left\langle \theta u_n, Q\theta u_n \right\rangle \\ &= \frac{1}{n^2} \langle e, Qe \rangle + 0 + 0 + \theta^2 \langle u_n, Qu_n \rangle = \frac{2nd}{n^2} + \theta^2 q_n.\end{aligned}$$

Therefore $2d/n + \theta^2 q_n \leq d(1/n + \theta^2) + 1 - 1/w$, that is

$$w \geq \frac{1}{1 - \frac{d}{n} + \theta^2(d - q_n)}.$$

Since $\theta = M/n$, $M = \min_{y_i \neq 0} \{1/|y_i|\}$, we find

$$w \geq \frac{n^2}{n^2 - nd + (d - q_n)M^2},$$

which gives the required inequality. \square

By adapting a similar technique, we find the following similar result for α .

Theorem 3.2. *Let G be a d -regular graph. Let $u_n = (y_1, y_2, \dots, y_n)$ be the normalized eigenvector corresponding to q_n , and let $M = \min_{y_i \neq 0} \{1/|y_i|\}$. Then*

$$\alpha \geq \frac{n^2}{nd + n + M^2(q_n - d + 1)}.$$

Given an $n \times n$ matrix A and an ordered partition (X_1, \dots, X_m) of the ordered set $\{1, 2, \dots, n\}$, $A_{n \times n}$ can be presented as a partitioned matrix

$$\begin{pmatrix} A_{1,1} & \dots & A_{1,m} \\ \vdots & \ddots & \vdots \\ A_{m,1} & \dots & A_{m,m} \end{pmatrix},$$

where $A_{i,j}$ has X_i as the set of its row numbers and X_j as the set of its column numbers. We always use Q_A hereafter to denote the quotient matrix of the partitioned

matrix A , which is defined to be the $m \times m$ matrix whose entries are the average row sums of the blocks of $A_{n \times n}$; namely, each entry $x_{i,j}$ is obtained by dividing the sum of all row sums of $A_{i,j}$ by $|X_i|$, where $1 \leq i, j \leq m$.

Consider two sequences of real numbers: $\alpha_1 \geq \alpha_2 \geq \dots \geq \alpha_n$ and $\beta_1 \geq \beta_2 \geq \dots \geq \beta_m$ with $m < n$. The latter sequence is said to interlace the former whenever

$$\alpha_i \geq \beta_i \geq \alpha_{n-m+i} \quad \text{for } i = 1, 2, \dots, m.$$

The interlacing is called tight if there exists an integer $k \in [0, m]$ such that

$$\lambda_i = \mu_i \quad \text{for } 1 \leq i \leq k \quad \text{and} \quad \lambda_{n-m+i} = \mu_i \quad \text{for } k+1 \leq i \leq m.$$

Lemma 3.1 ([3]). *For a symmetric partitioned matrix A , the eigenvalues of the quotient matrix Q_A interlace the eigenvalues of A .*

Theorem 3.3. *Suppose $\{v_1, v_2, \dots, v_\omega\}$ is the largest clique vertices set of the d -regular graph G . If $n > q_1 - q_n$, then*

$$\frac{n(q_n + 1 - d)}{n - 2d + q_n} \leq \omega \leq \frac{n(q_2 + 1 - d)}{n - 2d + q_2}.$$

The equalities $n(q_n + 1 - d)/(n - 2d + q_n) = \omega = n(q_2 + 1 - d)/(n - 2d + q_2)$ hold if and only if $G = O_n$ (the empty graph of order n).

Proof. The clique vertices set $\{v_1, v_2, \dots, v_\omega\}$ gives rise to a partition of $Q(G)$ with the quotient matrix

$$Q_{Q(G)} = \begin{pmatrix} d + (\omega - 1) & d - (\omega - 1) \\ \frac{[d - (\omega - 1)]\omega}{n - \omega} & 2d - \frac{[d - (\omega - 1)]\omega}{n - \omega} \end{pmatrix}.$$

$Q_{Q(G)}$ has eigenvalues $\lambda_1 = 2d$, $\lambda_2 = d + (\omega - 1) - ([d - (\omega - 1)]\omega)/(n - \omega)$. By Lemma 3.1 we have $q_n \leq \lambda_2 \leq q_2$, which gives the required inequality (note that $n > q_1 - q_n$). If the right equality holds, then $\lambda_2 = q_n$, and since $\lambda_1 = q_1 = 2d$ the interlacing is tight. Since $n(q_n + 1 - d)/(n - 2d + q_n) = \omega = n(q_2 + 1 - d)/(n - 2d + q_2)$ holds, if and only if $q_n = \lambda_2 = q_2$, and since $\sum_{i=1}^n q_i = nd$, we obtain $\omega = 1$, that is $G = O_n$. □

Note that the condition $n > q_1 - q_n$ is necessary. For example, if $G = K_n$ or G is a bipartite $n/2$ -regular graph then $n - 2d + q_n = 0$, the inequalities in Theorem 3.3 are invalid. However, in these two cases the clique number ω is known.

Now we will give some bounds for the independence number α by adapting a similar technique.

Theorem 3.4. Suppose $\{v_1, v_2, \dots, v_\alpha\}$ is a largest independent vertices set of a d -regular graph G , then

$$\frac{n(d - q_2)}{2d - q_2} \leq \alpha \leq \frac{n(d - q_n)}{2d - q_n}.$$

The equalities $n(d - q_2)/(2d - q_2) = \alpha = n(d - q_n)/(2d - q_n)$ hold, if and only if $G = K_n$ (the complete graph).

PROOF. The independent set $\{v_1, v_2, \dots, v_\alpha\}$ gives rise to a partition of $Q(G)$ with the quotient matrix

$$Q_{Q(G)} = \begin{pmatrix} d & d \\ \frac{d\alpha}{n - \alpha} & 2d - \frac{d\alpha}{n - \alpha} \end{pmatrix}.$$

$Q_{Q(G)}$ has eigenvalues $\lambda_1 = 2d$, $\lambda_2 = d - d\alpha/(n - \alpha)$ and so $q_n \leq \lambda_2 \leq q_2$ gives the required inequality. If the right equality holds, then $\lambda_2 = q_n$, and since $\lambda_1 = q_1 = 2d$, the interlacing is tight. Since $(n(d - q_2))/(2d - q_2) = \alpha = (n(d - q_n))/(2d - q_n)$ holds if and only if $q_n = \lambda_2 = d - d\alpha/(n - \alpha) = q_2$, and since $\sum_{i=1}^n q_i = 2m = nd$, we obtain $\alpha = 1$, that is $G = K_n$. \square

References

- [1] *D. Cvetković, M. Doob, H. Sachs*: Spectra of Graphs, third ed. Johann Ambrosius Barth Verlag, Heidelberg-Leipzig, 1995.
- [2] *M. Desai, V. Rao*: A characterization of the smallest eigenvalue of a graph. *J. Graph Theory* 18 (1994), 181–194.
- [3] *W. Haemers*: Interlacing eigenvalues and graphs. *Linear Algebra Appl.* 227–228 (1995), 593–616.
- [4] *W. Haemers, E. Spence*: Enumeration of cospectral graph. *Europ. J. Combin.* 25 (2004), 199–211.
- [5] *M. Lu, H. Liu, F. Tian*: Laplacian spectral bounds for clique and independence numbers of graphs. *J. Combin. Theory Ser. B* 97 (2007), 726–732.
- [6] *T. Motzkin, E. G. Straus*: Maxima for graphs and a new proof of a theorem of Turán. *Canad. J. Math.* 17 (1965), 533–540.

Authors' addresses: J. Liu, College of Mathematics and Computer Science, Fuzhou University, Fujian, 350 002, P. R. China, e-mail: Ljiping010@163.com, B. Liu (corresponding author), School of Mathematics Sciences, South China Normal University, Guangzhou, 510 631, P. R. China, e-mail: liubl@scnu.edu.cn.