

Kangkang Wang

A class of strong limit theorems for countable nonhomogeneous Markov chains on the generalized gambling system

Czechoslovak Mathematical Journal, Vol. 59 (2009), No. 1, 23–37

Persistent URL: <http://dml.cz/dmlcz/140461>

Terms of use:

© Institute of Mathematics AS CR, 2009

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A CLASS OF STRONG LIMIT THEOREMS FOR
COUNTABLE NONHOMOGENEOUS MARKOV CHAINS
ON THE GENERALIZED GAMBLING SYSTEM

KANGKANG WANG, Zhenjiang

(Received April 11, 2006)

Abstract. In this paper, we study the limit properties of countable nonhomogeneous Markov chains in the generalized gambling system by means of constructing compatible distributions and martingales. By allowing random selection functions to take values in arbitrary intervals, the concept of random selection is generalized. As corollaries, some strong limit theorems and the asymptotic equipartition property (AEP) theorems for countable nonhomogeneous Markov chains in the generalized gambling system are established. Some results obtained are extended.

Keywords: local convergence theorem, stochastic adapted sequence, martingale

MSC 2010: 60F15

1. INTRODUCTION

Consider a sequence of Bernoulli trials and suppose that at each trial the bettor has the free choice of whether or not to bet. A theorem on gambling system asserts that under any non-anticipative system the successive bets form a sequence of Bernoulli trials with unchanged probability for success. The importance of this statement was recognized by von Mises, who introduced the impossibility of a successful gambling system as a fundamental axiom (see [1], [2]). This topic was discussed still further by Kolmogorov (see [3]) and Liu and Wang (see [4] and [5]). Yang and Liu (see [14]) and Wang (see [15]) have studied the limit properties for Markov chains on the tree and on the random transform, respectively. Wang and Li (see [16-25]) have studied the strong limit theorems for nonhomogeneous Markov chains and Markov chains field on trees and gambling systems. On the basis of [3-7] and [15] we studied strong limit theorems for nonhomogeneous Markov chains on the generalized gambling system.

The purpose of this paper is to extend the discussion to the case of strong limit theorem for countable nonhomogeneous Markov chains by using the martingale method and constructing compatible distribution. By allowing random selection functions to take values in arbitrary intervals, the concept of random selection is generalized. As corollaries, the results of Liu and Yang (see [6]) are extended.

Let $\{X_n, n \geq 0\}$ be a stochastic sequence defined on a probability space (Ω, \mathcal{F}, P) which takes values in $S = \{s_1, s_2, \dots\}$. The joint distribution is

$$(1) \quad P(X_0 = x_0, \dots, X_n = x_n) = p(x_0, \dots, x_n) > 0, \quad x_i \in S, \quad 0 \leq i \leq n.$$

Let $\{X_n, n \geq 0\}$ be a nonhomogeneous Markov chain. The initial distribution and the transition matrix are respectively:

$$(2) \quad (p(s_1), p(s_2) \dots), \quad p(i) > 0, \quad i \in S$$

and

$$(3) \quad P_n = (p_n(i, j)), p_n(i, j) > 0, \quad i, j \in S, \quad n \geq 1,$$

where $p_n(i, j) = P(X_n = j | X_{n-1} = i)$ ($n \geq 1$). Then

$$(4) \quad p(x_0, \dots, x_n) = p(x_0) \prod_{k=1}^n p_k(x_{k-1}, x_k).$$

In order to extend the concept of random selection, which is the crucial part of the gambling system, we first give a set of real-valued functions $f_n(x_1, \dots, x_n)$ defined on S^n ($n = 1, 2, \dots$), which will be called the A -valued selection function if they take values in a set A of real numbers. Then let

$$(5) \quad \begin{aligned} Y_1 &= y \quad (y \text{ is an arbitrary real number}), \\ Y_{n+1} &= f_n(X_1, \dots, X_n), \quad n \geq 1, \end{aligned}$$

where $\{Y_n, n \geq 1\}$ will be called a generalized gambling system (the generalized random selection system). Let $\delta_i(j)$ be the Kronecker delta function on S , that is for $i, j \in S$

$$\delta_i(j) = \begin{cases} 0, & i \neq j, \\ 1, & i = j. \end{cases}$$

In order to explain the real meaning of the extended notion of the random selection, we consider the following gambling model. Let $\{X_n, n \geq 0\}$ be a nonhomogeneous Markov chain with the initial distribution (2) and the transition matrix

(3), and $\{g_n(x, y), n \geq 1\}$ be a real-valued function sequence defined on S^2 . Interpret X_n as the result of the n th trial, the type of which may change at each step. Let $\mu_n = Y_n g_n(X_{n-1}, X_n)$ denote the gain of the bettor at the n th trial, where Y_n represents the bet size, $g_n(X_{n-1}, X_n)$ is determined by the gambling rules, and $\{Y_n, n \geq 0\}$ is called a generalized gambling system or a generalized random selection system. The bettor's strategy is to determine $\{Y_n, n \geq 1\}$ by the results of the last trial. Let the entrance fee that the bettor pays at the n th trial be b_n . Also suppose that b_n depends on X_{n-1} as $n \geq 1$, and b_1 is a constant. Thus $\sum_{k=1}^n Y_k g_k(X_{k-1}, X_k)$ represents the total gain in the first n trials, $\sum_{k=1}^n b_k$ the accumulated entrance fees, and $\sum_{k=1}^n [Y_k g_k(X_{k-1}, X_k) - b_k]$ the accumulated net gain. Motivated by the classical definition of "fairness" of game of chance (see [3]), we introduce the following definition:

Definition. The game is said to be fair, if for almost all $\omega \in \left\{ \omega: \sum_{k=1}^{\infty} Y_k = \infty \right\}$, the accumulated net gain in the first n trials is of smaller order of magnitude than the accumulated stake $\sum_{k=1}^n Y_k$ as n tends to infinity, that is

$$\lim_{n \rightarrow \infty} \frac{1}{\sum_{k=1}^n Y_k} \sum_{k=1}^n [Y_k g_k(X_{k-1}, X_k) - b_k] = 0 \quad \text{a.s. on } \left\{ \omega: \sum_{k=1}^{\infty} Y_k = \infty \right\}$$

We can obtain the following conclusion.

2. MAIN RESULTS AND ITS PROOF

Theorem 1. Let $\{X_n, n \geq 0\}$ be a nonhomogeneous Markov chain with the initial distribution (2) and the transition matrix (3) and $\{Y_n, n \geq 1\}$ be defined as before. Let $\{\sigma_n, n \geq 1\}$ be an arbitrary nonnegative stochastic aequence. Let $\{g_n(x, y), n \geq 1\}$ be a real-valued function sequence defined on S^2 and let $\alpha > 0$ be a constant. Let

$$(6) \quad D = \left\{ \omega: \lim_{n \rightarrow \infty} \sigma_n = \infty, \right. \\ \left. \limsup_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n E[\exp\{\alpha |Y_k g_k(X_{k-1}, X_k)|\} | X_{k-1}] < \infty \right\}.$$

Then

$$(7) \quad \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n Y_k \{g_k(X_{k-1}, X_k) - E(g_k(X_{k-1}, X_k) | X_{k-1})\} = 0 \quad \text{a.s. } \omega \in D$$

Proof. Let

$$(8) \quad D_{x_0 \dots x_n} = \{\omega : X_k = x_k, 0 \leq k \leq n\}, \quad x_k \in S, \quad 1 \leq k \leq n.$$

Then

$$P(D_{x_0 \dots x_n}) = p(x_0, \dots, x_n)$$

and

$$(9) \quad P(D_{x_0 \dots x_n}) = p(x_0, \dots, x_n) = p(x_0) \prod_{k=1}^n p_k(x_{k-1}, x_k), \quad n \geq 1.$$

$D_{x_0 \dots x_n}$ is called an n th-order elementary cylinder. Let N_n be the collection of n th-order elementary cylinders, N the collection consisting of \emptyset , Ω and all cylinder sets and let $|\lambda| \leq \alpha$. Define a set function μ on N as follows:

$$(10) \quad \begin{aligned} \mu(\emptyset) &= 0, & \mu(\Omega) &= 1, \\ \mu(D_{x_0 \dots x_n}) &= p(x_0) \prod_{k=1}^n \frac{\exp\{\lambda y_k g(x_{k-1}, x_k)\} p_k(x_{k-1}, x_k)}{E[\exp\{\lambda y_k g(X_{k-1}, X_k)\} | X_{k-1} = x_{k-1}]}, \end{aligned}$$

where y_1 is an arbitrary real number, and

$$(11) \quad y_k = f_{k-1}(x_0, \dots, x_{k-1}), \quad k \geq 1.$$

We have by (10)

$$(12) \quad \begin{aligned} \sum_{x_n \in S} \mu(D_{x_0 \dots x_n}) &= \sum_{x_n \in S} \mu(D_{x_0 \dots x_{n-1}}) \frac{\exp\{\lambda y_n g(x_{n-1}, x_n)\} p_n(x_{n-1}, x_n)}{E[\exp\{\lambda y_n g(X_{n-1}, X_n)\} | X_{n-1} = x_{n-1}]} \\ &= \mu(D_{x_0 \dots x_{n-1}}) \frac{\sum_{x_n \in S} \exp\{\lambda y_n g(x_{n-1}, x_n)\} p_n(x_{n-1}, x_n)}{E[\exp\{\lambda y_n g(X_{n-1}, X_n)\} | X_{n-1} = x_{n-1}]} \\ &= \mu(D_{x_0 \dots x_{n-1}}) \frac{E[\exp\{\lambda y_n g(X_{n-1}, X_n)\} | X_{n-1} = x_{n-1}]}{E[\exp\{\lambda y_n g(X_{n-1}, X_n)\} | X_{n-1} = x_{n-1}]} \\ &= \mu(D_{x_0 \dots x_{n-1}}). \end{aligned}$$

It follows from (10)–(12) that μ is a measure on N . Since N is semialgebra, μ has a unique extension to the σ -field $\sigma(N)$. Let

$$(13) \quad T_n(\lambda, \omega) = \sum_{D \in N_n} \frac{\mu(D_{x_0 \dots x_n})}{P(D_{x_0 \dots x_n})} I_{D_{x_0 \dots x_n}}$$

where $I_{D_{x_0 \dots x_n}}$ denotes the indicator function of $D_{x_0 \dots x_n}$, that is

$$(14) \quad T_n(\lambda, \omega) = \frac{\mu(D_{X_0(\omega) \dots X_n(\omega)})}{P(D_{X_0(\omega) \dots X_n(\omega)})}.$$

It is easy to see that $\{N_n, n \geq 0\}$ is a net relative to (Ω, A, P) , where A denotes the σ -algebra of events on which P is defined. By the differentiation on a net of Hewitt and Stromberg (see [7], p. 373), there exists $A(\lambda) \in \sigma(N)$ with $P(A(\lambda)) = 1$ such that

$$\lim_n T_n(\lambda, \omega) = T_\infty(\lambda, \omega) < \infty \quad \omega \in A(\lambda)$$

that is

$$(15) \quad \lim_n T_n(\lambda, \omega) = T_\infty(\lambda, \omega) < \infty \quad \text{a.s.}$$

By (6) and (15) we have

$$(16) \quad \limsup_{n \rightarrow \infty} \frac{1}{\sigma_n} \log T_n(\lambda, \omega) \leq 0 \quad \text{a.s.} \quad \omega \in D.$$

By (9), (10), (13) and (14), we have

$$(17) \quad \log T_n(\lambda; \omega) = \sum_{k=1}^n \lambda Y_k g_k(X_{k-1}, X_k) - \sum_{k=1}^n \log E[\exp\{\lambda Y_k g_k(X_{k-1}, X_k) | X_{k-1}\}].$$

By (16) and (17) we have

$$(18) \quad \limsup_{n \rightarrow \infty} \frac{1}{\sigma_n} \left\{ \sum_{k=1}^n \lambda Y_k g_k(X_{k-1}, X_k) - \sum_{k=1}^n \log E[\exp\{\lambda Y_k g_k(X_{k-1}, X_k) | X_{k-1}\}] \right\} \leq 0 \quad \text{a.s.} \quad \omega \in D.$$

By virtue of the property of limes superior:

$$\limsup_{n \rightarrow \infty} (a_n - b_n) \leq d \Rightarrow \limsup_{n \rightarrow \infty} (a_n - c_n) \leq \limsup_{n \rightarrow \infty} (b_n - c_n) + d,$$

and the inequalities $\log x \leq x - 1$ ($x > 0$), $e^x - 1 - x \leq \frac{1}{2}x^2 e^{|x|}$, noticing

$$\max\{x^2 e^{-hx}, x \geq 0\} = 4e^{-2}/h^2 \quad (h > 0),$$

and letting $0 \leq |\lambda| < \alpha$, we have by (18)

$$\begin{aligned}
(19) \quad & \limsup_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n \lambda Y_k \{g_k(X_{k-1}, X_k) - E(g_k(X_{k-1}, X_k) | X_{k-1})\} \\
& \leq \limsup_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n \{\log E[\exp\{\lambda Y_k g_k(X_{k-1}, X_k)\} | X_{k-1}] \\
& \qquad \qquad \qquad - \lambda Y_k E(g_k(X_{k-1}, X_k) | X_{k-1})\} \\
& \leq \limsup_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n \{E[\exp\{\lambda Y_k g_k(X_{k-1}, X_k)\} | X_{k-1}] - 1 \\
& \qquad \qquad \qquad - \lambda Y_k E(g_k(X_{k-1}, X_k) | X_{k-1})\} \\
& \leq \frac{1}{2} \limsup_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n (\lambda Y_k)^2 E\{[g_k(X_{k-1}, X_k)]^2 e^{|\lambda Y_k g_k(X_{k-1}, X_k)|} | X_{k-1}\} \\
& \leq \frac{1}{2} (\lambda Y_k)^2 \limsup_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n E\{[g_k(X_{k-1}, X_k)]^2 e^{|\lambda Y_k| \cdot |g_k(X_{k-1}, X_k)|} | X_{k-1}\} \\
& = \frac{1}{2} (\lambda Y_k)^2 \limsup_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n E\{g_k(X_{k-1}, X_k)^2 e^{(|\lambda| - \alpha) |Y_k g_k(X_{k-1}, X_k)|} \\
& \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \times e^{\alpha |Y_k g_k(X_{k-1}, X_k)|} | X_{k-1}\} \\
& \leq \frac{1}{2} (\lambda Y_k)^2 \limsup_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n E\{e^{\alpha |Y_k g_k(X_{k-1}, X_k)|} \\
& \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \times 4e^{-2} / (Y_k \cdot (|\lambda| - \alpha))^2 | X_{k-1}\} \text{ a.s. } \omega \in D.
\end{aligned}$$

When $0 < \lambda < \alpha$, we have by (6) and (19)

$$\begin{aligned}
(20) \quad & \limsup_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n Y_k \{g_k(X_{k-1}, X_k) - E(g_k(X_{k-1}, X_k) | X_{k-1})\} \\
& \leq \frac{1}{2} \lambda \cdot \limsup_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n E\{e^{\alpha |Y_k g_k(X_{k-1}, X_k)|} 4e^{-2} / (\lambda - \alpha)^2 | X_{k-1}\} \text{ a.s. } \omega \in D.
\end{aligned}$$

Choose $0 < \lambda_i < \alpha$, $i = 1, 2, \dots$ such that $\lambda_i \rightarrow 0$ (as $i \rightarrow \infty$). Therefore, for all i , we have by (20)

$$(21) \quad \limsup_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n Y_k \{g_k(X_{k-1}, X_k) - E(g_k(X_{k-1}, X_k) | X_{k-1})\} \leq 0 \text{ a.s. } \omega \in D$$

When $-\alpha < \lambda < 0$, we have by (6) and (19)

$$(22) \quad \liminf_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n Y_k \{g_k(X_{k-1}, X_k) - E(g_k(X_{k-1}, X_k)|X_{k-1})\} \\ \geq \frac{1}{2} \lambda \cdot \limsup_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n E\{e^{\alpha|Y_k g(X_{k-1}, X_k)|} 4e^{-2}/(\lambda + \alpha)^2 |X_{k-1}\} \quad \text{a.s. } \omega \in D.$$

Choose $-\alpha < \lambda_i < 0$, $i = 1, 2, \dots$ such that $\lambda_i \rightarrow 0$ (as $i \rightarrow \infty$). Therefore, for all i , we have by (22)

$$(23) \quad \liminf_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n Y_k \{g_k(X_{k-1}, X_k) - E(g_k(X_{k-1}, X_k)|X_{k-1})\} \geq 0 \quad \text{a.s. } \omega \in D.$$

Therefore (7) follows from (21) and (23). \square

Corollary 1 (see [2]). *Let $\{X_n, n \geq 0\}$ be a nonhomogeneous Markov chain with the initial distribution (2) and the transition matrix (3) and let $\{Y_n, n \geq 1\}$ and $\{g_n(x, y), n \geq 1\}$ be defined as before. Let*

$$(24) \quad D_0 = \left\{ \omega : \lim_{n \rightarrow \infty} \sum_{k=1}^n E[\exp\{\alpha|Y_k g_k(X_{k-1}, X_k)|\} |X_{k-1}] = \infty \right\}.$$

where $\alpha > 0$ is a constant. Then

$$(25) \quad \lim_{n \rightarrow \infty} \frac{\sum_{k=1}^n Y_k \{g_k(X_{k-1}, X_k) - E(g_k(X_{k-1}, X_k)|X_{k-1})\}}{\sum_{k=1}^n E[\exp\{\alpha|Y_k g_k(X_{k-1}, X_k)|\} |X_{k-1}]} = 0 \quad \text{a.s. } \omega \in D_0.$$

Proof. Let

$$\sigma_n = \sum_{k=1}^n E[\exp\{\alpha|Y_k g_k(X_{k-1}, X_k)|\} |X_{k-1}], \quad n \geq 1.$$

Then

$$(26) \quad \limsup_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n E[\exp\{\alpha|Y_k g_k(X_{k-1}, X_k)|\} |X_{k-1}] \leq 1.$$

It can be known that $D = D_0$ from (24) and (26). Therefore (25) follows from (7).

Corollary 2. Let $\{X_n, n \geq 0\}$ be a nonhomogeneous Markov chain with the initial distribution (2) and the transition matrix (3) and let $\{\sigma_n, n \geq 1\}$ and $\{g_n(x, y), n \geq 1\}$ be defined as before, where $\alpha > 0$ is a constant. Let

$$(27) \quad D_2 = \left\{ \omega: \lim_{n \rightarrow \infty} \sigma_n = \infty, \right. \\ \left. \limsup_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n E[\exp\{\alpha |g_k(X_{k-1}, X_k)|\} | X_{k-1}] < \infty \right\}.$$

Then

$$(28) \quad \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n \{g_k(X_{k-1}, X_k) - E(g_k(X_{k-1}, X_k) | X_{k-1})\} = 0 \quad \text{a.s. } \omega \in D_2.$$

Proof. Let $Y_k \equiv 1$ in Theorem 1. It is easy to see that $D_2 = D$ then. Thus (28) can follow from (7). \square

Remark. It can be seen that the condition (27) weakens the condition of Theorem 1 in the paper of Liu and Yang (see [6]):

$$\limsup_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n E[g_k^2(X_{k-1}, X_k) e^{\alpha |g_k(X_{k-1}, X_k)|} | X_{k-1}] < \infty.$$

Correspondingly the conclusion is strengthened.

3. THE GENERALIZATION FOR AEP THEOREMS FOR NONHOMOGENEOUS MARKOV CHAINS IN THE GENERALIZED GAMBLING SYSTEM

Let $\{X_n, n \geq 0\}$ be an arbitrary stochastic sequence defined on the probability space (Ω, \mathcal{F}, P) which takes values in $S_0 = \{s_1, s_2, \dots, s_N\}$. The joint distribution is defined as in (1). Let

$$(29) \quad f_n(\omega) = -\frac{1}{n} \log p(X_0, \dots, X_n).$$

$f_n(\omega)$ is called as the relative entropy density of $\{X_n, n \geq 0\}$. If $\{X_n, n \geq 0\}$ is the nonhomogeneous Markov chain with the initial distribution (2) and the transition matrix (3), then by virtue of (4) and (29) we have

$$(30) \quad f_n(\omega) = -\frac{1}{n} \left[\log p(X_0) + \sum_{k=1}^n \log p_k(X_{k-1}, X_k) \right].$$

The limit property of the relative entropy density is an important problem in information theory. Shannon (see [8]) first showed that for stationary ergodic Markov chains $f_n(\omega)$ converges in probability to a constant. McMillan (see [9]) and Breiman (see [10]) proved, respectively, that if $\{X_n, n \geq 0\}$ is stationary and ergodic, then $f_n(\omega)$ converges in L_1 and almost everywhere to a constant. This is the famous Shannon-McMillan theorem. The extension of the Shannon-McMillan theorem to the general stochastic process can be found, for example, in Barron (see [11]), Chung (see [12]) and Feinstein (see [13]). In this paper we mainly study the some limit properties and asymptotic equipartition property (AEP) theorems for countable nonhomogeneous Markov chains in the generalized gambling system.

Corollary 3 (see [3]). *Let $\{X_n, n \geq 0\}$ be a nonhomogeneous Markov chain with the initial distribution (2) and the transition matrix (3), and let $f_n(\omega)$ be the relative entropy density defined as in (30). Then*

$$(31) \quad \lim_{n \rightarrow \infty} \left\{ f_n(\omega) - \frac{1}{n} \sum_{k=1}^n \sum_{j \in S_0} p_k(X_{k-1}, j) \log p_k(X_{k-1}, j) \right\} = 0 \quad \text{a.s.}$$

Proof. Let $g_k(i, j) = -\log p_k(i, j)$, $Y_k \equiv 1$, $\sigma_n = n$ and $\alpha = 1$ in Theorem 1. Then

$$(32) \quad \begin{aligned} & \limsup_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n E[\exp\{\alpha |Y_k g_k(X_{k-1}, X_k)|\} | X_{k-1}] \\ & \leq \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n E[\exp\{|-\log p_k(X_{k-1}, X_k)|\} | X_{k-1}] \\ & = \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \sum_{j \in S_0} \frac{p_k(X_{k-1}, j)}{p_k(X_{k-1}, j)} \leq N. \end{aligned}$$

It can be shown that $D = \Omega$ from $\sigma_n = n$ and (32). Therefore (31) follows from (7), (30) and the assumption conditions above.

Remark. The corollary is just the result of Theorem 2 in the paper of Liu and Yang (1996). We consider the problem of the generalized random selection for a arbitrary stochastic sequence. We choose a subsequence of $X_0, X_1, \dots, X_n, \dots$ and $(X_0, X_1), (X_1, X_2), \dots, (X_{n-1}, X_n), \dots$ (where $(X_0, X_1), (X_1, X_2), \dots, (X_{n-1}, X_n), \dots$ are the ordered couples of random variables $\{X_n, n \geq 0\}$) according to the value Y_n takes. We select X_n and (X_{n-1}, X_n) if and only if $Y_n \in [-m, 0) \cup (0, m]$. Therefore we obtain a subsequence of the above sequence. We let for $i, j \in S_0$, $S_n(i; \omega)$ be the number of the states i in the sequence X_1, X_2, \dots, X_n ,

which are selected by $\{Y_k, 1 \leq k \leq n\}$; $S_n(i, j; \omega)$ be the number of the ordered couples (i, j) in the ordered couples $(X_0, X_1), (X_1, X_2), \dots, (X_{n-1}, X_n)$, which are selected by $\{Y_k, 1 \leq k \leq n\}$. That is $S_n(i; \omega) = \sum_{k=1}^n Y_k \delta_i(X_k)$, $S_n(i, j; \omega) = \sum_{k=1}^n Y_k \delta_i(X_{k-1}) \delta_j(X_k)$, where $\delta_i(j)$ is defined as before. Then we can conclude the following results.

Corollary 4. *Let $\{X_n, n \geq 0\}$ be a nonhomogeneous Markov chain with the initial distribution (2) and the transition matrix (3) and let $\{\sigma_n, n \geq 1\}$, $S_n(i; \omega)$ and $S_n(i, j; \omega)$ be defined as before. $|Y_n| \leq m, n \geq 1$. assume that*

$$(33) \quad \limsup_{n \rightarrow \infty} \frac{n}{\sigma_n} \leq M \quad \text{a.s.}$$

Then

$$(34) \quad \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \left\{ S_n(i; \omega) - \sum_{k=1}^n Y_k p_k(X_{k-1}, i) \right\} = 0 \quad \text{a.s.},$$

$$(35) \quad \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \left\{ S_n(i, j; \omega) - \sum_{k=1}^n Y_k \delta_i(X_{k-1}) p_k(i, j) \right\} = 0 \quad \text{a.s.}$$

Proof. Letting $g_k(X_{k-1}, X_k) = \delta_i(X_k)$, $k \geq 1$, then

$$(36) \quad \begin{aligned} & \sum_{k=1}^n Y_k \{g_k(X_{k-1}, X_k) - E(g_k(X_{k-1}, X_k) | X_{k-1})\} \\ &= \sum_{k=1}^n Y_k \{\delta_i(X_k) - E(\delta_i(X_k) | X_{k-1})\} = S_n(i; \omega) - \sum_{k=1}^n Y_k p_k(X_{k-1}, i). \end{aligned}$$

Noticing that $|Y_n| \leq m, n \geq 1$. We have by (33) that

$$(37) \quad \begin{aligned} & \limsup_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n E[\exp\{\alpha |Y_k g_k(X_{k-1}, X_k)|\} | X_{k-1}] \\ & \leq \limsup_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n E[e^{m\alpha} | X_{k-1}] \leq \limsup_{n \rightarrow \infty} e^{m\alpha} \frac{n}{\sigma_n} \leq M e^{\alpha m} \quad \text{a.s.} \end{aligned}$$

It is also easy to see from (33) that $\lim_{n \rightarrow \infty} \sigma_n = \infty$. Therefore $D = \Omega$. It is easy to see from (7) and (36) that (34) holds. Similarly, let $g_k(X_{k-1}, X_k) = \delta_i(X_{k-1}) \delta_j(X_k)$,

$k \geq 1$. Noticing that

$$\begin{aligned}
(38) \quad & \sum_{k=1}^n Y_k \{g_k(X_{k-1}, X_k) - E(g_k(X_{k-1}, X_k)|X_{k-1})\} \\
&= \sum_{k=1}^n Y_k \{\delta_i(X_{k-1})\delta_j(X_k) - E(\delta_i(X_{k-1})\delta_j(X_k)|X_{k-1})\} \\
&= S_n(i, j; \omega) - \sum_{k=1}^n Y_k \delta_i(X_{k-1})p_k(X_{k-1}, j),
\end{aligned}$$

it can also be seen that (37) holds. Therefore, $D = \Omega$ holds too. (35) follows from (7) and (38). \square

Theorem 2. *Let $\{X_n, n \geq 0\}$ be a nonhomogeneous Markov chain with the initial distribution (41) and the transition matrix (42). Let $S_n(i; \omega)$ and $S_n(i, j; \omega)$ be defined as before. Let $\sigma_n = \sum_{k=1}^n Y_k$, where Y_k is defined as in Corollary 4. Let $P = (p(i, j))$ be another transition matrix and be ergodic. $\{g(x, y)\}$ be a real-valued function defined on S_0^2 . If*

$$(39) \quad \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n |Y_k p_k(i, j) - Y_{k-1} p(i, j)| = 0 \quad \text{for all } i, j \in S_0,$$

Then

$$(40) \quad \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} S_n(i; \omega) = \pi_i \quad \text{a.s.},$$

$$(41) \quad \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} S_n(i, j; \omega) = \pi_i p(i, j) \quad \text{a.s.},$$

$$(42) \quad \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n Y_k g(X_{k-1}, X_k) = \sum_{i \in S_0} \sum_{j \in S_0} \pi_i g(i, j) p(i, j) \quad \text{a.s.},$$

where $(\pi_{s_1}, \pi_{s_2}, \dots, \pi_{s_N})$ is the unique stationary distribution determined by P .

Proof. (1) We have by (34) in Corollary 4,

$$(43) \quad \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \left\{ S_n(j; \omega) - \sum_{k=1}^n Y_k p_k(X_{k-1}, j) \right\} = 0 \quad \text{a.s.}$$

Since

$$(44) \quad \sum_{k=1}^n Y_k p_k(X_{k-1}, j) = \sum_{k=1}^n \sum_{i \in S_0} Y_k \delta_i(X_{k-1}) p_k(i, j),$$

by (39) and the definition of σ_n in Corollary 4, we have

$$\begin{aligned}
(45) \quad & \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \left| \sum_{k=1}^n \sum_{i \in S_0} \delta_i(X_{k-1})(Y_k p_k(i, j) - Y_{k-1} p(i, j)) \right| \\
& \leq \sum_{i \in S_0} \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \left| \sum_{k=1}^n (Y_k p_k(i, j) - Y_{k-1} p(i, j)) \right| \\
& \leq \sum_{i \in S_0} \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n |Y_k p_k(i, j) - Y_{k-1} p(i, j)|.
\end{aligned}$$

By (43), (44), (45) and the definition of $S_n(i; \omega)$ we have

$$\begin{aligned}
(46) \quad & \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \left\{ S_n(j; \omega) - \sum_{i \in S_0} S_n(i; \omega) p(i, j) \right\} \\
& = \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n \sum_{i \in S_0} \delta_i(X_{k-1})(Y_k p_k(i, j) - Y_{k-1} p(i, j)) = 0 \quad \text{a.s.}
\end{aligned}$$

Multiplying (46) by $p(j, l)$, adding them together for $j \in S_0$ and using (46) once again, we have

$$\begin{aligned}
(47) \quad & 0 = \sum_{j \in S_0} p(j, l) \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \left\{ S_n(j; \omega) - \sum_{i \in S_0} S_n(i; \omega) p(i, j) \right\} \\
& = \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \left\{ \sum_{j \in S_0} S_n(j; \omega) p(j, l) - S_n(l; \omega) \right\} \\
& \quad + \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \left\{ S_n(l; \omega) - \sum_{i \in S_0} \sum_{j \in S_0} S_n(i; \omega) p(i, j) p(j, l) \right\} \\
& = \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \left\{ S_n(l; \omega) - \sum_{i \in S_0} S_n(i; \omega) p^{(2)}(i, l) \right\} \quad \text{a.s.},
\end{aligned}$$

where $p^{(k)}(i, l)$ (k is an integer) is the k -step transition probability of P . By induction, we have

$$(48) \quad \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \left\{ S_n(l; \omega) - \sum_{i \in S_0} S_n(i; \omega) p^{(k)}(i, l) \right\} = 0 \quad \text{a.s.}$$

By (48) we have

$$(49) \quad \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \left\{ S_n(l; \omega) - \sum_{i \in S_0} S_n(i; \omega) \frac{1}{N} \sum_{k=1}^N p^{(k)}(i, l) \right\} = 0 \quad \text{a.s.}$$

Since

$$(50) \quad \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=1}^N p^{(k)}(i, l) = \pi_l$$

and

$$(51) \quad \frac{1}{\sigma_n} \sum_{i \in S_0} S_n(i; \omega) = \sum_{k=1}^n \sum_{i \in S_0} Y_k \delta_i(X_k) / \sum_{k=1}^n Y_k = 1,$$

it is easy to see from (49)–(51) that (40) holds .

(2) By (35) in Corollary 4, we have

$$(52) \quad \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \left\{ S_n(i, j; \omega) - \sum_{k=1}^n Y_k \delta_i(X_{k-1}) p_k(i, j) \right\} = 0 \quad \text{a.s.}$$

It is easy to show by (39) and (45)

$$(53) \quad \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n \delta_i(X_{k-1}) (Y_k p_k(i, j) - Y_{k-1} p(i, j)) = 0.$$

By (52), (53) and the definition of $S_n(i; \omega)$ we have

$$(54) \quad \begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \{ S_n(i, j; \omega) - S_n(i; \omega) p(i, j) \} \\ = \lim_{n \rightarrow \infty} \frac{1}{\sigma_n} \sum_{k=1}^n \delta_i(X_{k-1}) (Y_k p_k(i, j) - Y_{k-1} p(i, j)) = 0 \quad \text{a.s.} \end{aligned}$$

It can be seen from (40) and (54) that (41) holds.

(3) Since

$$(55) \quad \begin{aligned} \frac{1}{\sigma_n} \sum_{k=1}^n Y_k g(X_{k-1}, X_k) &= \frac{1}{\sigma_n} \sum_{k=1}^n \sum_{i \in S_0} \sum_{j \in S_0} Y_k \delta_i(X_{k-1}) \delta_j(X_k) g(i, j) \\ &= \sum_{i \in S_0} \sum_{j \in S_0} g(i, j) \frac{1}{\sigma_n} S_n(i, j; \omega), \end{aligned}$$

(42) follows from (41) and (55). □

Corollary 5. Let $\{X_n, n \geq 0\}$ be a nonhomogeneous Markov chain with the initial distribution (2) and the transition matrix (3) and $f_n(\omega)$ be the relative entropy density defined as in (30). Let $P = (p(i, j))$ be defined as in Theorem 2. If (39) holds, then

$$(56) \quad \lim_{n \rightarrow \infty} f_n(\omega) = - \sum_{i \in S_0} \pi_i \sum_{j \in S_0} p(i, j) \log p(i, j) \quad \text{a.s.}$$

Proof. Let $Y_n \equiv 1$, $g(x, y) = -\log p(x, y)$ in Theorem 2, then $\sigma_n = n$. Moreover,

$$(57) \quad \limsup_{n \rightarrow \infty} \left(\frac{n}{\sigma_n} \right) = \limsup_{n \rightarrow \infty} \frac{n}{n} \leq 1 \quad \text{a.s.}$$

From (42) in Theorem 2 it follows immediately that (56) holds.

References

- [1] *P. Billingsley*: Probability and Measure. Wiley, New York, 1986.
- [2] *R. V. Mises*: Mathematical Theory of Probability and Statistics. Academic Press. New York, 1964.
- [3] *A. N. Kolmogorov*: On the logical foundation of probability theory. Lecture Notes in Mathematics. Springer-Verlag, New York, vol. 1021, 1982, pp. 1–2.
- [4] *W. Liu and Z. Wang*: An extension of a theorem on gambling systems to arbitrary binary random variables. Statistics and Probability Letters, vol. 28, 1996, pp. 51–58.
- [5] *Z. Wang*: A strong limit theorem on random selection for the N-valued random variables. Pure and Applied Mathematics 15 (1999), 56–61.
- [6] *W. Liu and W. Yang*: An extension of Shannon-McMillan theorem and some limit properties for nonhomogeneous Markov chains. Stochastic Process. Appl. 61 (1996), 279–292.
- [7] *K. R. Stromberg and E. Hewitt*: Real and abstract analysis—a modern treatment of the theory of functions of real variable. Springer, New York, 1994.
- [8] *C. Shannon*: A mathematical theory of communication. Bell System Tech J. 27 (1948), 379–423.
- [9] *B. Mcmillan*: The Basic Theorem of information theory. Ann. Math. Statist. 24 (1953), 196–219.
- [10] *L. Breiman*: The individual ergodic theorem of information theory. Ann. Math. Statist. 28 (1957), 809–811.
- [11] *A. R. Barron*: The strong ergodic theorem of densities; Generalized Shannon-McMillan-Breiman theorem. Ann. Probab. 13 (1985), 1292–1303.
- [12] *K. L. Chung*: The ergodic theorem of information theory. Ann. Math. Statist 32 (1961), 612–614.
- [13] *A. Feinstein*: A new basic theory of information. IRE Trans. P.G.I.T. (1954), 2–22.
- [14] *W. Yang and W. Liu*: Strong law of large numbers and Shannon-McMillan theorem for Markov fields on trees. IEEE Trans. Inform. Theory 48 (2002), 313–318.
- [15] *Z. Wang and W. Yang*: Some strong limit theorems for both nonhomogeneous Markov chains of order two and their random transforms. J. Sys. Sci. and Math. Sci 24 (2004), 451–462.

- [16] *K. Wang and W. Yang*: Research on strong limit theorem for Cantor-like stochastic sequence of Science and Technology (in Chinese). *J. Jiangsu Univ. Sci-tech. Nat. Sci.* *20* (2006), 26–29.
- [17] *K. Wang*: Strong large number law for Markov chains field on arbitrary Cayley tree (in Chinese). *J. Jiangsu Univ. Sci-tech. Nat. Sci.* *20* (2006), 28–32.
- [18] *K. Wang*: Some research on strong limit theorems for Cantor-like nonhomogeneous Markov chains (in Chinese). *J. Jiangsu Univ. Sci-tech. Nat. Sci.* *20* (2006), 19–23.
- [19] *K. Wang and Z. Qin*: A class of strong limit theorems for arbitrary stochastic sequence in random selection system (in Chinese). *J. Jiangsu Univ. Sci-tech. Nat. Sci.* *20* (2006), 40–44.
- [20] *K. Wang*: A class of strong limit theorems for stochastic sequence on product distribution in gambling system (in Chinese). *J. Jiangsu Univ. Sci-tech. Nat. Sci.* *21* (2007), 33–36.
- [21] *K. Wang and H. Ye*: A class of strong limit theorems for Markov chains field on arbitrary Bethe tree (in Chinese). *J. Jiangsu Univ. Sci-tech. Nat. Sci.* *21* (2007), 37–40.
- [22] *K. Wang*: A class of strong limit theorems for random sum of Three-order countable nonhomogeneous Markov chains (in Chinese). *J. Jiangsu Univ. Sci-tech. Nat. Sci.* *21* (2007), 42–45.
- [23] *K. Wang and H. Ye*: A class of local strong limit theorems for random sum of Cantor-like random function sequences (in Chinese). *J. Jiangsu Univ. Sci-tech. Nat. Sci.* *22* (2008), 87–90.
- [24] *K. Wang*: A class of strong limit theorems on generalized gambling system for arbitrary continuous random variable sequence (in Chinese). *J. Jiangsu Univ. Sci-tech. Nat. Sci.* *22* (2008), 86–90.
- [25] *M. Li*: Some limit properties for the sequence of arbitrary random variables on their generalized random selection system (in Chinese). *J. Jiangsu Univ. Sci-tech. Nat. Sci.* *22* (2008), 90–94.

Author's address: Kang kang Wang, School of Mathematics and physics, Jiangsu University of Science and Technology, Zhenjiang 212003, China, e-mail: wkk.cn@126.com.