

Lili Hu; Chunhui Lai; Ping Wang
On potentially $K_5 - H$ -graphic sequences

Czechoslovak Mathematical Journal, Vol. 59 (2009), No. 1, 173–182

Persistent URL: <http://dml.cz/dmlcz/140471>

Terms of use:

© Institute of Mathematics AS CR, 2009

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON POTENTIALLY $K_5 - H$ -GRAPHIC SEQUENCESLILI HU, Zhangzhou, CHUNHUI LAI, Zhangzhou,
PING WANG, Antigonish

(Received March 16, 2007)

Abstract. Let $K_m - H$ be the graph obtained from K_m by removing the edges set $E(H)$ of H where H is a subgraph of K_m . In this paper, we characterize the potentially $K_5 - P_4$ and $K_5 - Y_4$ -graphic sequences where Y_4 is a tree on 5 vertices and 3 leaves.

Keywords: graph, degree sequence, potentially $K_5 - H$ -graphic sequence

MSC 2010: 05C07, 05C35

1. INTRODUCTION

We consider finite simple graphs. Any undefined notation follows that of Bondy and Murty [1]. An n -term non-increasing nonnegative integer sequence $\pi = (d_1, d_2, \dots, d_n)$ is said to be graphic if it is the degree sequence of a simple graph G of order n ; such a graph G is referred as a realization of π . Let C_k and P_k denote a cycle on k vertices and a path on $k + 1$ vertices, respectively. We use the symbol E_4 to denote graphs on 5 vertices and 4 edges. Let $\sigma(\pi)$ be the sum of all the terms of π , and let $[x]$ be the largest integer less than or equal to x . Let Y_4 denote a tree on 5 vertices and 3 leaves. Z_4 is $K_4 - P_2$. A graphic sequence π is said to be potentially H -graphic if it has a realization G containing H as a subgraph. Let $G - H$ denote the graph obtained from G by removing the edges set $E(H)$ where H is a subgraph of G . In the degree sequence, r^t means r repeats t times, that is, in the realization of the sequence there are t vertices of degree r .

In 1907, Mantel first proposed the problem of determining the maximum number of edges in a graph without containing 3-cycles. In general, this problem can be

Research was supported by NNSF of China (10271105) and by NSF of Fujian (Z0511034), Fujian Provincial Training Foundation for “Bai-Quan-Wan Talents Engineering”, Project of Fujian Education Department, Project of Zhangzhou Teachers College and by NSERC.

phrased as determining the maximum number of edges, denoted $\text{ex}(n, H)$, of a graph with n vertices not containing H as a subgraph. This area of research is called extremal graph theory. In terms of graphic sequences, the number $2 \text{ex}(n, H) + 2$ is the minimum even integer l such that every n -term graphic sequence π with $\sigma(\pi) \geq l$ is forcibly H -graphic. In 1991, Erdős, Jacobson and Lehel [2] showed $\sigma(K_k, n) \geq (k-2)(2n-k+1)+2$ and conjectured that the equality holds. In the same paper, they proved that the conjecture is true for the case $k = 3$ and $n \geq 6$. The cases $k = 4$ and 5 were proved separately in [3], [16] and [17]. Based on linear algebraic techniques, Li, Song and Luo [18] proved the conjecture true for $k \geq 6$ and $n \geq \binom{k}{2} + 3$. Recently, Ferrara, Gould and Schmitt proved the conjecture [5] and they also determined in [6] $\sigma(F_k, n)$ where F_k denotes the graph of k triangles intersecting at exactly one common vertex.

In 1999, Gould, Jacobson and Lehel [3] considered the following generalized problem: determine the smallest even integer $\sigma(H, n)$ such that every n -term positive graphic sequence $\pi = (d_1, d_2, \dots, d_n)$ with $\sigma(\pi) \geq \sigma(H, n)$ has a realization G containing H as a subgraph. They proved $\sigma(pK_2, n) = (p-1)(2n-p) + 2$ for $p \geq 2$ and $\sigma(C_4, n) = 2\lfloor \frac{1}{2}(3n-1) \rfloor$ for $n \geq 4$. Lai [10] determined $\sigma(K_4 - e, n)$ for $n \geq 4$. Yin, Li, and Mao [24] determined $\sigma(K_{r+1} - e, n)$ for $r \geq 3$ and $r+1 \leq n \leq 2r$ and $\sigma(K_5 - e, n)$ for $n \geq 5$, and Yin and Li [23] further determined $\sigma(K_{r+1} - e, n)$ for $r \geq 2$ and $n \geq 3r^2 - r - 1$. Moreover, Yin and Li in [23] also gave two sufficient conditions for a sequence $\pi \in GS_n$ to be potentially $(K_{r+1} - e)$ -graphic. Yin [26] determined $\sigma(K_{r+1} - K_3, n)$ for $r \geq 3$ and $n \geq 3r + 5$. Lai [11]–[13] determined $\sigma(K_5 - P_3, n)$, $\sigma(K_5 - P_4, n)$, $\sigma(K_5 - C_4, n)$ and $\sigma(K_5 - K_3, n)$ for $n \geq 5$. Lai and Hu [14] determined $\sigma(K_{r+1} - H, n)$ for $n \geq 4r + 10$, $r \geq 3$, $r+1 \geq k \geq 4$ and H be a graph on k vertices which containing a tree on 4 vertices but not containing a cycle on 3 vertices and $\sigma(K_{r+1} - P_2, n)$ for $n \geq 4r + 8$, $r \geq 3$. Lai [15] determined $\sigma(K_{r+1} - Z_4, n)$, $\sigma(K_{r+1} - (K_4 - e), n)$, $\sigma(K_{r+1} - K_4, n)$ for $n \geq 5r + 16$, $r \geq 4$ and $\sigma(K_{r+1} - Z, n)$ for $n \geq 5r + 19$, $r+1 \geq k \geq 5$, $j \geq 5$ where Z is a graph on k vertices and j edges which contains a graph Z_4 but does not contain a cycle on 4 vertices.

A harder question is to characterize the potentially H -graphic sequences without zero terms. That is, finding necessary and sufficient conditions for a sequence to be a potentially H -graphic sequence. Luo [20] characterized the potentially C_k -graphic sequences for each $k = 3, 4$ and 5 . Recently, in [21], Luo and Warner also characterized the potentially K_4 -graphic sequences. Eschen and Niu [22] characterized the potentially $K_4 - e$ -graphic sequences. Hu and Lai [7]–[8] characterized the potentially $K_5 - C_4$ and $K_5 - Z_4$ -graphic sequences. Yin and Chen [25] characterized the potentially $K_{r,s}$ -graphic sequences for $r = 2, s = 3$ and $r = 2, s = 4$, where $K_{r,s}$ is an $r \times s$ complete bipartite graph. Gupta, Joshi and Tripathi [4] gave a necessary and sufficient condition for the existence of a tree of order n with a given degree set.

In attempt to completely characterize the potentially $K_5 - E_4$ -graphic sequences, we will characterize the potentially $K_5 - P_4$ and $K_5 - Y_4$ -graphic sequences in this paper. The problem of characterizing the potentially $K_5 - E_4$ -graphic sequences has not been solved so far.

Let $\pi = (d_1, d_2, \dots, d_n)$ be a nonincreasing positive integer sequence. We write $m(\pi)$ and $h(\pi)$ to denote the largest positive terms of π and the smallest positive terms of π , respectively. $\pi'' = (d_1 - 1, d_2 - 1, \dots, d_{d_n} - 1, d_{d_n+1}, \dots, d_{n-1})$ is the residual sequence obtained by laying off d_n from π . We denote $\pi' = (d'_1, d'_2, \dots, d'_{n-1})$ where $d'_1 \geq d'_2 \geq \dots \geq d'_{n-1}$ is a rearrangement of the $n - 1$ terms in π'' . We denote by π' the residual sequence obtained by laying off d_n from π and all the graphic sequences have no zero terms. We need the following results.

Theorem 1.1 ([3]). *If $\pi = (d_1, d_2, \dots, d_n)$ is a graphic sequence with a realization G containing H as a subgraph, then there exists a realization G' of π containing H as a subgraph so that the vertices of H have the largest degrees of π .*

Theorem 1.2 ([19]). *If $\pi = (d_1, d_2, \dots, d_n)$ is a sequence of nonnegative integers with $1 \leq m(\pi) \leq 2$, $h(\pi) = 1$ and $\sigma(\pi)$ even, then π is graphic.*

Theorem 1.3 ([9]). *π is graphic if and only if π' is graphic.*

The following corollary is obvious.

Corollary 1.4. *Let H be a simple graph. If π' is potentially H -graphic, then π is potentially H -graphic.*

2. MAIN THEOREMS

Theorem 2.1. *Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence with $n \geq 5$. Then π is potentially $K_5 - P_4$ -graphic if and only if the following conditions hold:*

- (1) $d_2 \geq 3$.
- (2) $d_5 \geq 2$.
- (3) $\pi \neq (n - 1, k, 2^t, 1^{n-2-t})$ where $n \geq 5$, $k, t = 3, 4, \dots, n - 2$, and k and t have different parities.
- (4) For $n \geq 5$, $\pi \neq (n - k, k + i, 2^i, 1^{n-i-2})$ where $i = 3, 4, \dots, n - 2k$ and $k = 1, 2, \dots, \lfloor \frac{1}{2}(n - 1) \rfloor - 1$.
- (5) If $n = 6, 7$, then $\pi \neq (3^2, 2^{n-2})$.

Proof. First we show that the conditions (1)–(5) are necessary conditions for π to be potentially $K_5 - P_4$ -graphic. Assume that π is potentially $K_5 - P_4$ -graphic. (1), (2) and (5) are obvious. If $\pi = (n - 1, k, 2^t, 1^{n-2-t})$ is potentially

$K_5 - P_4$ -graphic, then according to Theorem 1.1, there exists a realization G of π containing $K_5 - P_4$ as a subgraph so that the vertices of $K_5 - P_4$ have the largest degrees of π . Therefore, the sequence $\pi^* = (n-4, k-3, 2^{t-3}, 1^{n-2-t})$ obtained from $G - (K_5 - P_4)$ must be graphic. Since the edge between two vertices with degree $n-4$ and $k-3$ has been removed from the realization of π^* , thus, $\Delta(G - (K_5 - P_4)) \leq n-5$, a contradiction. Hence, (3) holds. If $\pi = (n-k, k+i, 2^i, 1^{n-i-2})$ is potentially $K_5 - P_4$ -graphic, then according to Theorem 1.1, there exists a realization G of π containing $K_5 - P_4$ as a subgraph so that the vertices of $K_5 - P_4$ have the largest degrees of π . Therefore, the sequence $\pi^* = (n-k-3, k+i-3, 2^{i-3}, 1^{n-i-2})$ obtained from $G - (K_5 - P_4)$ must be graphic and there is no edge between two vertices with degree $n-k-3$ and $k+i-3$ in the realization of π^* . Let G^* be a realization of π^* , and, $d_{G^*}(x) = n-k-3$ and $d_{G^*}(y) = k+i-3$. Consider a partition of G^* where $X = \{x, y\}$ and $Y = V(G^*) - \{x, y\}$. It follows that the number of edges between X and Y equals $(n-k-3) + (k+i-3) \leq 2(i-3) + (n-i-2)$, that is, $[(n-k-3) + (k+i-3)] - [2(i-3) + (n-i-2)] = 2 \leq 0$, a contradiction. Hence, (4) holds.

Now we show that the conditions (1)–(5) are sufficient conditions for π to be potentially $K_5 - P_4$ -graphic. Suppose the graphic sequence π satisfies the conditions (1) to (5). Our proof is by induction on n . We first prove the base case where $n = 5$. Since $\pi \neq (4^2, 2^3)$, π must be one of the following sequences: (4^5) , $(4^3, 3^2)$, $(4^2, 3^2, 2)$, $(4, 3^4)$, $(4, 3^2, 2^2)$, $(3^4, 2)$, $(3^2, 2^3)$. It is easy to check that all of these are potentially $K_5 - P_4$ -graphic. Now we assume that the sufficiency holds for $n-1$ ($n \geq 6$). We will prove that π is potentially $K_5 - P_4$ -graphic.

Case 1: $\pi' = (3^2, 2^4)$. Clearly, $n = 7$ and π must be one of the following sequences $(4^2, 2^5)$, $(4, 3^2, 2^4)$, $(3^4, 2^3)$, $(4, 3, 2^4, 1)$ or $(3^3, 2^3, 1)$. It is easy to check that all of these are potentially $K_5 - P_4$ -graphic.

Case 2: $\pi' = (3^2, 2^5)$. Clearly, $n = 8$ and π must be one of the following sequences $(4^2, 2^6)$, $(4, 3^2, 2^5)$, $(3^4, 2^4)$, $(4, 3, 2^5, 1)$ or $(3^3, 2^4, 1)$. It is easy to check that all of these are potentially $K_5 - P_4$ -graphic.

Case 3: $d_n \geq 3$. Clearly, π' satisfies the assumption, and thus, by the induction hypothesis, π' is potentially $K_5 - P_4$ -graphic, and hence so is π . In the following, we only consider the cases $d_n = 1$ or $d_n = 2$.

Case 4: $\pi' = (n-2, k, 2^t, 1^{n-3-t})$ where $n-1 \geq 5$, $k, t = 3, 4, \dots, n-3$, and, k and t have different parities.

If $d_n = 2$, then $\pi' = (n-2, k, 2^{n-3})$. If $k \geq 4$, then $\pi = (n-1, k+1, 2^{n-2})$ which contradicts condition (3). If $k = 3$, that is $\pi' = (n-2, 3, 2^{n-3})$, then $\pi = (n-1, 4, 2^{n-2})$ or $\pi = (n-1, 3^2, 2^{n-3})$. But $\pi = (n-1, 4, 2^{n-2})$ contradicts condition (3), thus $\pi = (n-1, 3^2, 2^{n-3})$ where n is odd. We will show that $\pi = (n-1, 3^2, 2^{n-3})$

is potentially $K_5 - P_4$ -graphic. In other words, we would like to show that $\pi_1 = (n-4, 2^{n-5}, 1)$ is graphic. It suffices to show that $\pi_2 = (1^{n-5})$ where $n \geq 7$ is graphic. By $\sigma(\pi_2)$ being even and Theorem 1.2, π_2 is graphic. Thus, $\pi = (n-1, 3^2, 2^{n-3})$ is potentially $K_5 - P_4$ -graphic.

If $d_n = 1$, then $\pi = (n-1, k, 2^t, 1^{n-2-t})$ which contradicts condition (3).

Case 5: $\pi' = (n-1-k, k+i, 2^i, 1^{n-i-3})$ where $i = 3, 4, \dots, n-1-2k$ and $k = 1, 2, \dots, \lfloor n/2 \rfloor - 2$.

If $d_n = 2$, then $n-i-3 = 0$ and $\pi = (n-k, k+i+1, 2^{i+1})$ which contradicts condition (4).

If $d_n = 1$ and $n-1-k = k+i+1$, then $\pi = (n-k, k+i, 2^i, 1^{n-i-2})$ or $\pi = ((n-1-k)^2, 2^i, 1^{n-i-2})$, both of which contradict condition (4). If $d_n = 1$ and $n-1-k = k+i$ or $n-1-k \geq k+i+2$, then $\pi = (n-k, k+i, 2^i, 1^{n-i-2})$ which also contradicts condition (4).

Case 6: $d_n = 2$, $\pi' \neq (n-2, k, 2^{n-3})$, $\pi' \neq (n-1-k, n+k-3, 2^{n-3})$, $\pi' \neq (3^2, 2^4)$, and $\pi' \neq (3^2, 2^5)$.

Consider $\pi' = (d'_1, d'_2, \dots, d'_{n-1})$. Since $d_2 \geq 3$, we have $d'_{n-1} \geq 2$. If $d'_2 \geq 3$, then π' satisfies the assumption. Thus, π' is potentially $K_5 - P_4$ -graphic. Hence, we may assume $d'_2 = 2$, that is, $d_2 = 3$ and $d_3 = d_4 = \dots = d_n = 2$. It follows that $\pi = (d_1, 3, 2^{n-2})$. Since $\sigma(\pi)$ is even, d_1 must be odd. If $d_1 = 3$, then $\pi = (3^2, 2^{n-2})$. Since $\pi \neq (3^2, 2^4)$ and $\pi \neq (3^2, 2^5)$, we have $n \geq 8$. We will show that π is potentially $K_5 - P_4$ -graphic. It suffices to show $\pi_1 = (2^{n-5})$ is graphic. Clearly, C_{n-5} is a realization of π_1 . If $d_1 \geq 5$, since $\pi \neq (n-1, 3, 2^{n-2})$, we have $d_1 \leq n-2$. We will prove that $\pi = (d_1, 3, 2^{n-2})$, where $d_1 \geq 5$ and $n \geq d_1 + 2$, is potentially $K_5 - P_4$ -graphic. We would like to show that $\pi_1 = (d_1 - 3, 2^{n-5})$ is graphic. It suffices to show that $\pi_2 = (2^{n-d_1-2}, 1^{d_1-3})$ is graphic. Since $\sigma(\pi_2)$ is even, π_2 is graphic by Theorem 1.2. Thus, $\pi = (d_1, 3, 2^{n-2})$ is potentially $K_5 - P_4$ -graphic.

Case 7: $d_n = 1$, $\pi' \neq (n-2, k, 2^t, 1^{n-3-t})$, $\pi' \neq (n-1-k, k+i, 2^i, 1^{n-i-3})$, $\pi' \neq (3^2, 2^4)$, and $\pi' \neq (3^2, 2^5)$.

Consider $\pi' = (d'_1, d'_2, \dots, d'_{n-1})$. Since $d_2 \geq 3$ and $d_5 \geq 2$, we have $d'_1 \geq 3$ and $d'_5 \geq 2$. If $d'_2 \geq 3$, then π' satisfies the assumption. Thus, π' is potentially $(K_5 - P_4)$ -graphic. Hence, we may assume $d'_2 = 2$, that is, $d_1 = d_2 = 3$ and $d_3 = d_4 = d_5 = 2$. Thus $\pi = (3^2, 2^t, 1^{n-2-t})$ where $t \geq 3$ and $n-2-t \geq 1$. Since $\sigma(\pi)$ is even, $n-2-t$ must be even. We will prove π is potentially $K_5 - P_4$ -graphic. It suffices to show that $\pi_1 = (2^{t-3}, 1^{n-2-t})$ is graphic. Since $\sigma(\pi_1)$ is even, π_1 is graphic by Theorem 1.2 and, in turn, π is potentially $K_5 - P_4$ -graphic.

This completes the proof. □

Theorem 2.2. Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence with $n \geq 5$. Then π is potentially $K_5 - Y_4$ -graphic if and only if the following conditions hold:

- (1) $d_3 \geq 3$.
- (2) $d_4 \geq 2$.
- (3) $\pi \neq (3^6)$.

Proof. Assume that π is potentially $K_5 - Y_4$ -graphic. In this case the necessary conditions (1) to (3) are obvious.

Now we prove the sufficient conditions. Suppose the graphic sequence π satisfies the conditions (1) to (3). Our proof is by induction on n . We first prove the base case where $n = 5$. In this case, π is one of the following sequences: (4^5) , $(4^3, 3^2)$, $(4^2, 3^2, 2)$, $(4, 3^4)$, $(4, 3^3, 1)$, $(4, 3^2, 2^2)$, $(3^4, 2)$, or $(3^3, 2, 1)$. It is easy to check that all of these are potentially $K_5 - Y_4$ -graphic. Now suppose the sufficiency holds for $n - 1$ ($n \geq 6$), and let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence which satisfies (1) to (3). We will prove π is potentially $K_5 - Y_4$ -graphic.

Case 1: $\pi' = (3^6)$. We have $n = 7$ and π is one of the following sequences $(4^3, 3^4)$, $(4^2, 3^4, 2)$ or $(4, 3^5, 1)$. It is easy to check that all of these are potentially $K_5 - Y_4$ -graphic.

Case 2: $d_n \geq 3$ and $\pi' \neq (3^6)$. Clearly, $d'_4 \geq 2$. If $d_3 \geq 4$, then $d'_3 \geq 3$. If $d_3 = \dots = d_n = 3$ and $n \geq 6$, $d'_3 \geq 3$. It follows conditions (1) and (2) hold. Thus, by the induction hypothesis, π' is potentially $K_5 - Y_4$ -graphic. Therefore, π is potentially $K_5 - Y_4$ -graphic by Corollary 1.4. In the following, we only consider the cases where $d_n = 2$ or $d_n = 1$.

Case 3: $d_n = 2$ and $\pi' \neq (3^6)$. Consider $\pi' = (d'_1, d'_2, \dots, d'_{n-1})$. Since $d_3 \geq 3$ and $d_n = 2$, we have $d'_1 \geq 3$ and $d'_{n-1} \geq 2$. If $d'_3 \geq 3$, then π' satisfies the assumption and it follows π' is potentially $K_5 - Y_4$ -graphic. Therefore, π is potentially $K_5 - Y_4$ -graphic by Corollary 1.4. Hence, we may assume $d'_3 = 2$. We will proceed with the following two cases $d_1 \geq 4$ and $d_1 = 3$.

Subcase 1: $d_1 \geq 4$. It suffices to consider the case where $d_2 = d_3 = 3$ and $d_4 = d_5 = \dots = d_n = 2$. That is, $\pi = (d_1, 3^2, 2^{n-3})$. Since $\sigma(\pi)$ is even, d_1 must be even. We will prove π is potentially $K_5 - Y_4$ -graphic. It is enough to show that $\pi_1 = (d_1 - 3, 2^{n-5}, 1)$ is graphic. If $d_1 = n - 1$, then $\pi_1 = (n - 4, 2^{n-5}, 1)$. It suffices to show that $\pi_2 = (1^{n-5})$ is graphic. Since $\sigma(\pi_2)$ is even, π_2 is graphic by Theorem 1.2. If $d_1 \leq n - 2$, it suffices to show that $\pi_2 = (2^{n-2-d_1}, 1^{d_1-2})$ (or $\pi_2 = (2^{n-1-d_1}, 1^{d_1-4})$) is graphic. Similarly, one can show π_2 is graphic. Thus, $\pi_1 = (d_1 - 3, 2^{n-5}, 1)$ is graphic and, in turn, π is potentially $K_5 - Y_4$ -graphic.

Subcase 2: $d_1 = 3$. It suffices to consider the case where $d_1 = d_2 = d_3 = d_4 = 3$ and $d_5 = \dots = d_n = 2$. That is, $\pi = (3^4, 2^{n-4})$. We will prove π is potentially $(K_5 - Y_4)$ -

graphic. It is enough to show that $\pi_1 = (2^{n-5}, 1^2)$ is graphic. Since $\sigma(\pi_1)$ is even, π_1 is graphic by Theorem 1.2 and, in turn, π is potentially $K_5 - Y_4$ -graphic.

Case 4: $d_n = 1$ and $\pi' \neq (3^6)$. Consider $\pi' = (d'_1, d'_2, \dots, d'_{n-1})$. Since $d_3 \geq 3$ and $d_4 \geq 2$, we have $d'_2 \geq 3$ and $d'_4 \geq 2$. If $d'_3 \geq 3$, then π' satisfies the assumptions and it follows π' is potentially $K_5 - Y_4$ -graphic. Therefore, π is potentially $K_5 - Y_4$ -graphic by Corollary 1.4. Hence, we may assume $d'_3 = 2$. It suffices to consider the case where $d_1 = d_2 = d_3 = 3$ and $d_4 = 2$. That is, $\pi = (3^3, 2^t, 1^{n-3-t})$ where $t \geq 1$ and $n - 3 - t \geq 1$. Since $\sigma(\pi)$ is even, $n - t$ must be even. We will prove π is potentially $K_5 - Y_4$ -graphic. It is enough to show that $\pi_1 = (2^{t-2}, 1^{n-2-t})$ is graphic when $t \geq 2$. Since $\sigma(\pi_1)$ is even, π_1 is graphic by Theorem 1.2. If $t = 1$, then $\pi = (3^3, 2, 1^{n-4})$. Similarly we can show that $\pi_2 = (1^{n-5})$ is graphic and, in turn, π is potentially $K_5 - Y_4$ -graphic.

This completes the proof. □

In the remainder of this section, we will use the above two theorems to find exact values of $\sigma(K_5 - P_4, n)$, $\sigma(K_5 - C_5, n)$, $\sigma(K_5 - Y_4, n)$, $\sigma(K_5 - (Y_4 + e), n)$ and $\sigma(K_5 - K_{2,3}, n)$. Note that the value of $\sigma(K_5 - P_4, n)$ was determined by Lai in [11] so a much simpler proof is given here.

Corollary 2.3 ([11]). *For $n \geq 5$, $\sigma(K_5 - P_4, n) = 4n - 4$.*

Proof. First we claim that $\sigma(K_5 - P_4, n) \geq 4n - 4$ for $n \geq 5$. We would like to show there exists π_1 with $\sigma(\pi_1) = 4n - 6$ such that π_1 is not potentially $K_5 - P_4$ -graphic. Let $\pi_1 = ((n - 1)^2, 2^{n-2})$. It is easy to see that $\sigma(\pi_1) = 4n - 6$ and π_1 is not potentially $K_5 - P_4$ -graphic by Theorem 2.1.

Now we show if π is an n -term ($n \geq 5$) graphical sequence with $\sigma(\pi) \geq 4n - 4$, then there exists a realization of π containing a $K_5 - P_4$. If $d_5 = 1$, then $\sigma(\pi) = d_1 + d_2 + d_3 + d_4 + (n - 4)$. Let X be the four vertices of the largest degrees of G and $Y = V(G) - X$. Since there are at most six edges in X , $d_1 + d_2 + d_3 + d_4 \leq 12 + |E(X, Y)| \leq 12 + (n - 4) = n + 8$. This leads to $\sigma(\pi) \leq 2n + 4 < 4n - 4$, a contradiction. Thus, $d_5 \geq 2$. If $d_2 \leq 2$, then $\sigma(\pi) \leq d_1 + 2(n - 1) \leq 3n - 3 < 4n - 4$, a contradiction. Thus, $d_2 \geq 3$. Since $\sigma(\pi) \geq 4n - 4$, then π is not one of the following: $(3^2, 2^4)$, $(3^2, 2^5)$, and $(n - 1, k, 2^t, 1^{n-2-t})$ where $n \geq 6$ and $k, t = 3, 4, \dots, n - 2$, $(n - k, k + i, 2^i, 1^{n-i-2})$ where $i = 3, 4, \dots, n - 2k$ and $k = 1, 2, \dots, \lfloor \frac{1}{2}(n - 1) \rfloor - 1$. Thus, π satisfies the conditions (1) to (5) in Theorem 2.1. Therefore, π is potentially $K_5 - P_4$ -graphic by Theorem 2.1. □

Corollary 2.4 ([14]). *For $n \geq 5$, $\sigma(K_5 - C_5, n) = 4n - 4$.*

Proof. Obviously, for $n \geq 5$, $\sigma(K_5 - C_5, n) \leq \sigma(K_5 - P_4, n) = 4n - 4$. Now we claim that $\sigma(K_5 - C_5, n) \geq 4n - 4$ for $n \geq 5$. We would like to show there

exists π_1 with $\sigma(\pi_1) = 4n - 6$, such that π_1 is not potentially $K_5 - C_5$ -graphic. Let $\pi_1 = ((n - 1)^2, 2^{n-2})$. It is easy to see that $\sigma(\pi_1) = 4n - 6$ and the only realization of π_1 does not contain $K_5 - C_5$. Thus, $\sigma(K_5 - C_5, n) = 4n - 4$. \square

Corollary 2.5 ([14]). *For $n \geq 5$, $\sigma(K_5 - Y_4, n) = 4n - 4$.*

Proof. First we claim that $\sigma(K_5 - Y_4, n) \geq 4n - 4$ if $n \geq 5$. We would like to show there exists π_1 with $\sigma(\pi_1) = 4n - 6$, such that π_1 is not potentially $K_5 - Y_4$ -graphic. Let $\pi_1 = ((n - 1)^2, 2^{n-2})$. It is easy to see that $\sigma(\pi_1) = 4n - 6$ and π_1 is not potentially $K_5 - Y_4$ -graphic by Theorem 2.2.

Now we show if π is an n -term ($n \geq 5$) graphical sequence with $\sigma(\pi) \geq 4n - 4$, then there exists a realization of π containing a $K_5 - Y_4$. If $d_4 = 1$, then $\sigma(\pi) = d_1 + d_2 + d_3 + (n - 3)$. Using a similar argument as in the above corollary, we have $d_1 + d_2 + d_3 \leq 6 + (n - 3) = n + 3$. This leads to $\sigma(\pi) \leq 2n < 4n - 4$, a contradiction. Thus, $d_4 \geq 2$. Similarly, if $d_3 \leq 2$, then $\sigma(\pi) \leq d_1 + d_2 + 2(n - 2) \leq 2(n - 1) + 2(n - 2) = 4n - 6 < 4n - 4$, a contradiction. Thus, $d_3 \geq 3$. Since $\sigma(\pi) \geq 4n - 4$, necessarily $\pi \neq (3^6)$. Thus, π satisfies the conditions (1) to (3) in Theorem 2.2. Therefore, π is potentially $K_5 - Y_4$ -graphic by Theorem 2.2. \square

Corollary 2.6 ([14]). *For $n \geq 5$, $\sigma(K_5 - (Y_4 + e), n) = 4n - 4$ where the two vertices of e are the leaves of Y_4 whose distance is 3.*

Proof. Obviously, for $n \geq 5$, $\sigma(K_5 - (Y_4 + e), n) \leq \sigma(K_5 - Y_4, n) = 4n - 4$. Now we claim that $\sigma(K_5 - (Y_4 + e), n) \geq 4n - 4$ for $n \geq 5$. We would like to show there exists π_1 with $\sigma(\pi_1) = 4n - 6$, such that π_1 is not potentially $K_5 - (Y_4 + e)$ -graphic. Let $\pi_1 = ((n - 1)^2, 2^{n-2})$. It is easy to see that $\sigma(\pi_1) = 4n - 6$ and the only realization of π_1 does not contain $K_5 - (Y_4 + e)$. Thus, $\sigma(K_5 - (Y_4 + e), n) = 4n - 4$. \square

Corollary 2.7 ([14]). *For $n \geq 5$, $\sigma(K_5 - K_{2,3}, n) = 4n - 4$.*

Proof. Obviously, for $n \geq 5$, $\sigma(K_5 - K_{2,3}, n) \leq \sigma(K_5 - Y_4, n) = 4n - 4$. Now we claim $\sigma(K_5 - K_{2,3}, n) \geq 4n - 4$ for $n \geq 5$. We would like to show there exists π_1 with $\sigma(\pi_1) = 4n - 6$, such that π_1 is not potentially $K_5 - K_{2,3}$ -graphic. Let $\pi_1 = ((n - 1)^2, 2^{n-2})$. It is easy to see that $\sigma(\pi_1) = 4n - 6$ and the only realization of π_1 does not contain $K_5 - K_{2,3}$. Thus, $\sigma(K_5 - K_{2,3}, n) = 4n - 4$. \square

Acknowledgements. The authors wish to thank R. J. Gould, Jiongsheng Li, Rong Luo, J. Schmitt, Amitabha Tripathi, Jianhua Yin and Mengxiao Yin for sending their papers to us.

References

- [1] *J. A. Bondy, U. S. R. Murty*: Graph Theory with Applications. Macmillan Press, 1976.
- [2] *P. Erdős, M. S. Jacobson, J. Lehel*: Graphs realizing the same degree sequences and their respective clique numbers. In: Graph Theory, Combinatorics and Application, Vol. 1 (Y. Alavi et al., eds.). John Wiley and Sons, New York, 1991, pp. 439–449.
- [3] *R. J. Gould, M. S. Jacobson, J. Lehel*: Potentially G -graphic degree sequences. Combinatorics, Graph Theory and Algorithms, Vol. 2 (Y. Alavi et al., eds.). New Issues Press, Kalamazoo, 1999, pp. 451–460.
- [4] *G. Gupta, P. Joshi, A. Tripathi*: Graphic sequences of trees and a problem of Frobenius. Czech. Math. J. *57* (2007), 49–52.
- [5] *M. Ferrara, R. Gould, J. Schmitt*: Potentially K_s^t -graphic degree sequences. Submitted.
- [6] *M. Ferrara, R. Gould, J. Schmitt*: Graphic sequences with a realization containing a friendship graph. Ars Comb. Accepted.
- [7] *Lili Hu, Chunhui Lai*: On potentially $K_5 - C_4$ -graphic sequences. Ars Comb. Accepted.
- [8] *Lili Hu, Chunhui Lai*: On potentially $K_5 - Z_4$ -graphic sequences. Submitted.
- [9] *D. J. Kleitman, D. L. Wang*: Algorithm for constructing graphs and digraphs with given valences and factors. Discrete Math. *6* (1973), 79–88.
- [10] *Chunhui Lai*: A note on potentially $K_4 - e$ graphical sequences. Australas J. Comb. *24* (2001), 123–127.
- [11] *Chunhui Lai*: An extremal problem on potentially $K_m - P_k$ -graphic sequences. Int. J. Pure Appl. Math. Accepted.
- [12] *Chunhui Lai*: An extremal problem on potentially $K_m - C_4$ -graphic sequences. J. Comb. Math. Comb. Comput. *61* (2007), 59–63.
- [13] *Chunhui Lai*: An extremal problem on potentially $K_{p,1,1}$ -graphic sequences. Discret. Math. Theor. Comput. Sci. *7* (2005), 75–80.
- [14] *Chunhui Lai, Lili Hu*: An extremal problem on potentially $K_{r+1} - H$ -graphic sequences. Ars Comb. Accepted.
- [15] *Chunhui Lai*: The smallest degree sum that yields potentially $K_{r+1} - Z$ -graphical sequences. Ars Comb. Accepted.
- [16] *Jiong-Sheng Li, Zi-Xia Song*: An extremal problem on the potentially P_k -graphic sequences. In: Proc. International Symposium on Combinatorics and Applications, June 28–30, 1996 (W. Y. C. Chen et al., eds.). Nankai University, Tianjin, 1996, pp. 269–276.
- [17] *Jiong-Sheng Li, Zi-Xia Song*: The smallest degree sum that yields potentially P_k -graphical sequences. J. Graph Theory *29* (1998), 63–72.
- [18] *Jiong-sheng Li, Zi-Xia Song, Rong Luo*: The Erdős-Jacobson-Lehel conjecture on potentially P_k -graphic sequence is true. Sci. China (Ser. A) *41* (1998), 510–520.
- [19] *Jiong-sheng Li, Jianhua Yin*: A variation of an extremal theorem due to Woodall. Southeast Asian Bull. Math. *25* (2001), 427–434.
- [20] *R. Luo*: On potentially C_k -graphic sequences. Ars Comb. *64* (2002), 301–318.
- [21] *R. Luo, M. Warner*: On potentially K_k -graphic sequences. Ars Combin. *75* (2005), 233–239.
- [22] *E. M. Eschen, J. Niu*: On potentially $K_4 - e$ -graphic sequences. Australas J. Comb. *29* (2004), 59–65.
- [23] *J.-H. Yin, J. S. Li*: Two sufficient conditions for a graphic sequence to have a realization with prescribed clique size. Discrete Math. *301* (2005), 218–227.
- [24] *J.-H. Yin, J.-S. Li, R. Mao*: An extremal problem on the potentially $K_{r+1} - e$ -graphic sequences. Ars Comb. *74* (2005), 151–159.
- [25] *J.-H. Yin, G. Chen*: On potentially K_{r_1, r_2, \dots, r_m} -graphic sequences. Util. Math. *72* (2007), 149–161.

- [26] *M. Yin*: The smallest degree sum that yields potentially $K_{r+1} - K_3$ -graphic sequences. *Acta Math. Appl. Sin., Engl. Ser.* 22 (2006), 451–456.

Authors' addresses: L. H u, Department of Mathematics, Zhangzhou Teachers College, Zhangzhou 363000, P. R. China, e-mail: jackey2591924@163.com; C h. L a i (corresponding author), Department of Mathematics, Zhangzhou Teachers College, Zhangzhou 363000, P. R. China, e-mail: zjlaichu@public.zzptt.fj.cn; P. W a n g, Dept. of Math., Stats. and Computer Science, St. Francis Xavier University, Antigonish, NS, Canada, B2G 2W5, e-mail: pwang@stfx.ca.