

Amir Mafi

Matlis reflexive and generalized local cohomology modules

Czechoslovak Mathematical Journal, Vol. 59 (2009), No. 4, 1095–1102

Persistent URL: <http://dml.cz/dmlcz/140539>

Terms of use:

© Institute of Mathematics AS CR, 2009

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

MATLIS REFLEXIVE AND GENERALIZED LOCAL
COHOMOLOGY MODULES

AMIR MAFI, Sanandaj

(Received June 5, 2008)

Abstract. Let (R, \mathfrak{m}) be a complete local ring, \mathfrak{a} an ideal of R and N and L two Matlis reflexive R -modules with $\text{Supp}(L) \subseteq V(\mathfrak{a})$. We prove that if M is a finitely generated R -module, then $\text{Ext}_R^i(L, H_{\mathfrak{a}}^j(M, N))$ is Matlis reflexive for all i and j in the following cases:

- (a) $\dim R/\mathfrak{a} = 1$;
- (b) $\text{cd}(\mathfrak{a}) = 1$; where cd is the cohomological dimension of \mathfrak{a} in R ;
- (c) $\dim R \leq 2$.

In these cases we also prove that the Bass numbers of $H_{\mathfrak{a}}^j(M, N)$ are finite.

Keywords: Bass numbers, generalized local cohomology modules, Matlis reflexive

MSC 2010: 13D45, 13E99, 13D07

1. INTRODUCTION

Let (R, \mathfrak{m}) be a commutative Noetherian local ring and \mathfrak{a} an ideal of R . For an integer $j \geq 0$, the j th generalized local cohomology module $H_{\mathfrak{a}}^j(M, N)$ of two R -modules M and N with respect to an ideal \mathfrak{a} was defined by Herzog [10] as follows:

$$H_{\mathfrak{a}}^j(M, N) = \varinjlim_n \text{Ext}_R^j(M/\mathfrak{a}^n M, N).$$

It is clear that $H_{\mathfrak{a}}^j(R, N)$ is just the ordinary local cohomology module $H_{\mathfrak{a}}^j(N)$ of N with respect to \mathfrak{a} (cf. [1]).

Hartshorne [9] defined a module T to be \mathfrak{a} -cofinite if $\text{Supp}(T) \subseteq V(\mathfrak{a})$ and $\text{Ext}_R^i(R/\mathfrak{a}, T)$ is finitely generated for all i . He proved that if R is a complete regular ring and \mathfrak{a} is either a principal ideal or a prime ideal with $\dim R/\mathfrak{a} = 1$, then the

This work was partially supported by the grant from IPM (No. 87130024).

local cohomology modules $H_{\mathfrak{a}}^j(N)$ are \mathfrak{a} -cofinite for all finitely generated modules N . Kawasaki [13] showed that, in general, the local cohomology modules $H_{\mathfrak{a}}^j(N)$ are \mathfrak{a} -cofinite for all finitely generated modules N , where the ideal \mathfrak{a} is principal. Delfino and Marley [6] and Yoshida [22], in general, proved that if \mathfrak{a} is an ideal of R with $\dim R/\mathfrak{a} = 1$ and L is finitely generated with $\text{Supp}(L) \subseteq V(\mathfrak{a})$, then $\text{Ext}_R^i(L, H_{\mathfrak{a}}^j(N))$ is finitely generated for all i and j and for all finitely generated modules N (see also [5] and [11]). Melkersson [17] proved that if R is a local ring with $\dim R \leq 2$, then $H_{\mathfrak{a}}^j(N)$ is \mathfrak{a} -cofinite for all j and all finitely generated modules N (see also [18]). Belshoff, Slattery and Wickham [3] and Belshoff and Slattery [2] extended the results of Hartshorne to larger class of modules. In fact, they showed that if R is a complete Gorenstein domain, \mathfrak{a} is either a principal ideal or an ideal with $\dim R/\mathfrak{a} = 1$, and M and N are Matlis reflexive modules with $\text{Supp}(M) \subseteq V(\mathfrak{a})$, then $\text{Ext}_R^i(M, H_{\mathfrak{a}}^j(N))$ is Matlis reflexive for all i and j . Recall that an R -module N is Matlis reflexive if $D(D(N)) = N$, where $D(-) = \text{Hom}_R(-, E(R/\mathfrak{m}))$ is the Matlis duality functor. Khashyarmansh and Khosh-Ahang [14] proved that if R is a complete ring and M and N are Matlis reflexive modules with $\text{Supp}(M) \subseteq V(\mathfrak{a})$, then $\text{Ext}_R^i(M, H_{\mathfrak{a}}^j(N))$ is Matlis reflexive for all i and j in the following cases:

- (a) $\dim R/\mathfrak{a} = 1$;
- (b) \mathfrak{a} is a principal ideal;
- (c) $\dim R \leq 2$.

The goal of the present paper is to extend the main results of Khashyarmansh and Khosh-Ahang [14] to generalized local cohomology modules.

2. PRELIMINARIES

Throughout this paper we assume that R is a commutative Noetherian ring, \mathfrak{a} an ideal of R , M a finitely generated R -module and N a Matlis reflexive R -module. We now briefly recall some known facts on generalized local cohomology modules.

Lemma 2.1 (see [15]). *Let X be an \mathfrak{a} -torsion R -module; that is, $\Gamma_{\mathfrak{a}}(X) = X$. Then $H_{\mathfrak{a}}^j(M, X) \cong \text{Ext}_R^j(M, X)$ for all $j \geq 0$.*

Lemma 2.2 (see [8]). *The following assertions hold:*

- (i) *If $0 \rightarrow N \rightarrow E^{\bullet}$ is an injective resolution of N , then $H_{\mathfrak{a}}^j(M, N) \cong H^j(\Gamma_{\mathfrak{a}}(\text{Hom}_R(M, E^{\bullet}))) \cong H^j(\text{Hom}_R(M, \Gamma_{\mathfrak{a}}(E^{\bullet})))$ for all $j \geq 0$. In particular, $H_{\mathfrak{a}}^j(M, N) \cong H_{\sqrt{\mathfrak{a}}}^j(M, N)$ for all $j \geq 0$.*
- (ii) *If $f: R \rightarrow S$ is a flat ring homomorphism, then $H_{\mathfrak{a}}^j(M, N) \otimes_R S \cong H_{\mathfrak{a}S}^j(M \otimes_R S, N \otimes_R S)$ for all $j \geq 0$.*

Lemma 2.3 (see [16]). *Let X be a finitely generated R -module. Then $H_{\mathfrak{a}}^j(M, X)$ is \mathfrak{a} -cofinite for all $j \geq 0$, whenever one of the following conditions holds:*

- (i) $\dim R \leq 2$;
- (ii) $\text{cd}(\mathfrak{a}) = 1$.

Lemma 2.4. *Let (R, \mathfrak{m}) be a local ring, \mathfrak{p} a prime ideal of R with $\dim R/\mathfrak{p} = 1$, and X a finitely generated module. Then $H_{\mathfrak{p}}^j(M, X)$ is \mathfrak{p} -cofinite for all $j \geq 0$.*

Proof. By [6, Proposition 2] and Lemma 2.2, we can assume that R is a complete regular local ring. Hence the result follows by [7, Theorem 2.9]. \square

Lemma 2.5 (see [6]). *Let X be an R -module. Then the following assertions are equivalent:*

- (i) $\text{Ext}_R^i(R/\mathfrak{a}, X)$ is finitely generated for all $i \geq 0$;
- (ii) $\text{Ext}_R^i(L, X)$ is finitely generated for all finitely generated modules L with $\text{Supp}(L) \subseteq V(\mathfrak{a})$ and all $i \geq 0$.

3. THE RESULTS

We begin this section by recalling some general facts about Matlis reflexive modules. First, any module of finite length and any Artinian module over a local ring are Matlis reflexive (see, for example, [19] and [20]).

Lemma 3.1 (see [4]). *Let (R, \mathfrak{m}) be a local ring. If N is a Matlis reflexive R -module, then there is a short exact sequence*

$$0 \longrightarrow S \longrightarrow N \longrightarrow A \longrightarrow 0$$

with S finitely generated and complete, and A Artinian.

Lemma 3.2 (see [20]). *Let $0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$ be a short exact sequence of R -modules and R -homomorphisms. Then B is Matlis reflexive if and only if A and C are Matlis reflexive.*

Theorem 3.3. Let (R, \mathfrak{m}) be a complete local ring and let L be a finitely generated R -module with $\text{Supp}(L) \subseteq V(\mathfrak{a})$. Suppose that one of the following cases hold:

(α) $\text{cd}(\mathfrak{a}) = 1$;

(β) $\dim R \leq 2$.

Then $\text{Ext}_R^i(L, H_{\mathfrak{a}}^j(M, N))$ is Matlis reflexive for all i and j .

Proof. Since N is Matlis reflexive, by Lemma 3.1 there is a short exact sequence

$$0 \longrightarrow S \longrightarrow N \longrightarrow A \longrightarrow 0$$

with S finitely generated and A Artinian. Hence we obtain the long exact sequence of generalized local cohomology

$$\dots \longrightarrow H_{\mathfrak{a}}^j(M, S) \xrightarrow{h^j} H_{\mathfrak{a}}^j(M, N) \xrightarrow{f^j} \text{Ext}_R^j(M, A) \xrightarrow{g^j} H_{\mathfrak{a}}^{j+1}(M, S) \longrightarrow \dots$$

Set $X^j = \text{Im } f^j$, $Y^j = \text{Im } h^j$ and $Z^j = \text{Im } g^j$. Now, consider the exact sequences

(\dagger) $0 \longrightarrow Z^{j-1} \longrightarrow H_{\mathfrak{a}}^j(M, S) \longrightarrow Y^j \longrightarrow 0,$

(\ddagger) $0 \longrightarrow Y^j \longrightarrow H_{\mathfrak{a}}^j(M, N) \longrightarrow X^j \longrightarrow 0.$

Hence we note that X^j and Z^j are Artinian for all $j \geq 0$, since $\text{Ext}_R^j(M, A)$ is Artinian for all $j \geq 0$. Let $j \geq 0$ be fixed arbitrary. By the exact sequence (\dagger), we have an exact sequence:

$$\begin{aligned} \dots \longrightarrow \text{Ext}_R^i(L, Z^{j-1}) &\longrightarrow \text{Ext}_R^i(L, H_{\mathfrak{a}}^j(M, S)) \longrightarrow \text{Ext}_R^i(L, Y^j) \\ &\longrightarrow \text{Ext}_R^{i+1}(L, Z^{j-1}) \longrightarrow \dots \end{aligned}$$

Now, $\text{Ext}_R^i(L, Z^{j-1})$ is Matlis reflexive for all $i \geq 0$ and by Lemmas 2.3, 2.5 $\text{Ext}_R^i(L, H_{\mathfrak{a}}^j(M, S))$ is Matlis reflexive for all $i \geq 0$. Hence $\text{Ext}_R^i(L, Y^j)$ is Matlis reflexive for all $i \geq 0$. Furthermore, we obtain an exact sequence by (\ddagger):

$$\dots \longrightarrow \text{Ext}_R^i(L, Y^j) \longrightarrow \text{Ext}_R^i(L, H_{\mathfrak{a}}^j(M, N)) \longrightarrow \text{Ext}_R^i(L, X^j) \longrightarrow \dots$$

Since $\text{Ext}_R^i(L, Y^j)$ and $\text{Ext}_R^i(L, X^j)$ are Matlis reflexive for all $i \geq 0$, $\text{Ext}_R^i(L, H_{\mathfrak{a}}^j(M, N))$ is Matlis reflexive for all $i \geq 0$. The proof is complete. \square

Lemma 3.4. *Let (R, \mathfrak{m}) be a local ring. Then $H_{\mathfrak{m}}^j(M, N)$ is Artinian for all $j \geq 0$.*

Proof. By Lemma 3.1, there is an exact sequence

$$0 \longrightarrow S \longrightarrow N \longrightarrow A \longrightarrow 0$$

with S finitely generated and A Artinian. This induces the long exact sequence

$$\dots \longrightarrow H_{\mathfrak{m}}^j(M, S) \longrightarrow H_{\mathfrak{m}}^j(M, N) \longrightarrow \text{Ext}_R^j(M, A) \longrightarrow \dots$$

Since $H_{\mathfrak{m}}^j(M, S)$ by [8, Theorem 2.2] and $\text{Ext}_R^j(M, A)$ are Artinian for all $i \geq 0$, we have that $H_{\mathfrak{m}}^j(M, N)$ is Artinian for all $i \geq 0$. \square

Theorem 3.5. *Let (R, \mathfrak{m}) be a complete local ring and \mathfrak{a} an ideal of R with $\dim R/\mathfrak{a} = 1$. Then for any finitely generated R -module L with $\text{Supp}(L) \subseteq V(\mathfrak{a})$, $\text{Ext}_R^i(L, H_{\mathfrak{a}}^j(M, N))$ is Matlis reflexive for all i and j .*

Proof. We may assume that $\mathfrak{a} = \sqrt{\mathfrak{a}}$ by Lemma 2.2. Let $\mathfrak{a} = \bigcap_{i=1}^n \mathfrak{p}_i$ be the irredundant primary decomposition. Now, we proceed by induction on the number n . Let $n = 1$. We consider the exact sequence

$$0 \longrightarrow S \longrightarrow N \longrightarrow A \longrightarrow 0$$

with S finitely generated and A Artinian. This induces a long exact sequence

$$\dots \longrightarrow H_{\mathfrak{a}}^j(M, S) \longrightarrow H_{\mathfrak{a}}^j(M, N) \longrightarrow \text{Ext}_R^j(M, A) \longrightarrow \dots$$

By Lemmas 2.4, 2.5 and using the same arguments as in the proof of Theorem 3.3 the result follows in this case. Now suppose that $n \geq 2$, and that the assertion holds for $n-1$. Put $\mathfrak{a}_1 = \mathfrak{p}_1$ and $\mathfrak{a}_2 = \bigcap_{i=2}^n \mathfrak{p}_i$. One can easily see that $V(\mathfrak{a}_1) \cup V(\mathfrak{a}_2) = V(\mathfrak{a})$ and $V(\mathfrak{a}_1) \cap V(\mathfrak{a}_2) = V(\mathfrak{m})$, since \mathfrak{a} is an ideal of dimension one. We have a Mayer-Vietoris exact sequence (cf. [21, Corollary 2.14])

$$\dots \longrightarrow H_{\mathfrak{m}}^j(M, N) \xrightarrow{f^j} H_{\mathfrak{a}_1}^j(M, N) \oplus H_{\mathfrak{a}_2}^j(M, N) \xrightarrow{h^j} H_{\mathfrak{a}}^j(M, N) \xrightarrow{g^j} \dots$$

Set $X^j = \text{Im } f^j$, $Y^j = \text{Im } h^j$ and $Z^j = \text{Im } g^j$. Hence there are exact sequences

$$\begin{aligned} (\dagger) \quad & 0 \longrightarrow Y^j \longrightarrow H_{\mathfrak{a}}^j(M, N) \longrightarrow Z^j \longrightarrow 0, \\ (\ddagger) \quad & 0 \longrightarrow X^j \longrightarrow H_{\mathfrak{a}_1}^j(M, N) \oplus H_{\mathfrak{a}_2}^j(M, N) \longrightarrow Y^j \longrightarrow 0. \end{aligned}$$

Here we note that X^j and Z^j are Artinian for all $j \geq 0$, since by Lemma 3.4 $H_m^j(M, N)$ is Artinian for all $j \geq 0$. Let $j \geq 0$ be fixed arbitrary. By the exact sequence (\ddagger) , we have an exact sequence

$$\begin{aligned} \dots \longrightarrow \text{Ext}_R^i(L, X^j) \longrightarrow \text{Ext}_R^i(L, H_{\mathfrak{a}_1}^j(M, N)) \oplus \text{Ext}_R^i(L, H_{\mathfrak{a}_2}^j(M, N)) \longrightarrow \\ \text{Ext}_R^i(L, Y^j) \longrightarrow \text{Ext}_R^{i+1}(L, X^j) \longrightarrow \dots \end{aligned}$$

Now $\text{Ext}_R^i(L, H_{\mathfrak{a}_1}^j(M, N))$ and $\text{Ext}_R^i(L, H_{\mathfrak{a}_2}^j(M, N))$ are Matlis reflexive for all $i \geq 0$, by induction hypothesis. Since $\text{Ext}_R^i(L, X^j)$ is Matlis reflexive, $\text{Ext}_R^i(L, Y^j)$ is Matlis reflexive for all $i \geq 0$. Moreover, we obtain an exact sequence by (\ddagger) :

$$\dots \longrightarrow \text{Ext}_R^i(L, Y^j) \longrightarrow \text{Ext}_R^i(L, H_{\mathfrak{a}}^j(M, N)) \longrightarrow \text{Ext}_R^i(L, Z^j) \longrightarrow \dots$$

Since Z^j is also Artinian, $\text{Ext}_R^i(L, H_{\mathfrak{a}}^j(M, N))$ is Matlis reflexive for all $i \geq 0$ by the above exact sequence, as required. \square

The following result extends [12, Theorem 1].

Corollary 3.6. *Let (R, \mathfrak{m}) be a complete local ring and suppose that one of the following cases occurs*

- (a) \mathfrak{a} is an ideal of R with $\dim R/\mathfrak{a} = 1$;
- (b) $\text{cd}(\mathfrak{a}) = 1$;
- (c) $\dim R \leq 2$.

Then the Bass numbers of generalized local cohomology modules $H_{\mathfrak{a}}^j(M, N)$ are finite for all $j \geq 0$.

Proof. Let k be the residue field of R . Then $\text{Ext}_R^i(k, H_{\mathfrak{a}}^j(M, N))$ is Matlis reflexive by Theorems 3.3, 3.5. Since $\text{Ext}_R^i(k, H_{\mathfrak{a}}^j(M, N))$ is also a k vector space, it must be finitely generated. If \mathfrak{p} is any non-maximal prime ideal, it follows from Lemma 3.1 that $N_{\mathfrak{p}}$ is finitely generated over $R_{\mathfrak{p}}$. We have $(H_{\mathfrak{a}}^j(M, N))_{\mathfrak{p}} \cong H_{\mathfrak{a}R_{\mathfrak{p}}}^j(M_{\mathfrak{p}}, N_{\mathfrak{p}})$ if $\mathfrak{p} \supseteq \mathfrak{a}$ or $(H_{\mathfrak{a}}^j(M, N))_{\mathfrak{p}} = 0$ if $\mathfrak{p} \not\supseteq \mathfrak{a}$. In either case, it follows that $\text{Ext}_R^i(R/\mathfrak{p}, H_{\mathfrak{a}}^j(M, N))_{\mathfrak{p}}$ is finitely generated over $R_{\mathfrak{p}}$. \square

Corollary 3.7. *Let (R, \mathfrak{m}) be a complete local ring and suppose that one of the following cases occurs:*

- (a) \mathfrak{a} is an ideal of R with $\dim R/\mathfrak{a} = 1$;
- (b) $\text{cd}(\mathfrak{a}) = 1$;
- (c) $\dim R \leq 2$.

If L is Artinian, then $\text{Ext}_R^i(L, H_{\mathfrak{a}}^j(M, N))$ is finitely generated (and thus Matlis reflexive) for all i and j .

Proof. By Corollary 3.6, $H_{\mathfrak{a}}^j(M, N)$ has finite Bass numbers. Fix j . Let $0 \rightarrow H_{\mathfrak{a}}^j(M, N) \rightarrow E^0 \rightarrow E^1 \rightarrow \dots$ be a minimal injective resolution of $H_{\mathfrak{a}}^j(M, N)$. Hence for each t , $\text{Hom}_R(L, E^t) = \bigoplus \text{Hom}_R(L, E(R/\mathfrak{m}))$ where the direct sum is a finite direct sum, since $H_{\mathfrak{a}}^j(M, N)$ has finite Bass numbers. By Matlis duality, $\text{Hom}_R(L, E(R/\mathfrak{m}))$ is finitely generated. Thus $\text{Hom}_R(L, E^t)$, and hence $\text{Ext}_R^i(L, H_{\mathfrak{a}}^j(M, N))$, is finitely generated. \square

The following corollary is a generalization of [14, Theorem 2.2].

Corollary 3.8. *Let (R, \mathfrak{m}) be a complete local ring and suppose that one of the following cases occurs*

- (a) \mathfrak{a} is an ideal of R with $\dim R/\mathfrak{a} = 1$;
- (b) $\text{cd}(\mathfrak{a}) = 1$;
- (c) $\dim R \leq 2$.

If L is Matlis reflexive with $\text{Supp}(L) \subseteq V(\mathfrak{a})$, then $\text{Ext}_R^i(L, H_{\mathfrak{a}}^j(M, N))$ is Matlis reflexive for all i and j .

Proof. Since L is Matlis reflexive, there is a short exact sequence

$$0 \rightarrow S \rightarrow L \rightarrow A \rightarrow 0$$

with S finitely generated and A Artinian. This induces a long exact sequence

$$\dots \rightarrow \text{Ext}_R^i(A, H_{\mathfrak{a}}^j(M, N)) \rightarrow \text{Ext}_R^i(L, H_{\mathfrak{a}}^j(M, N)) \rightarrow \text{Ext}_R^i(S, H_{\mathfrak{a}}^j(M, N)) \rightarrow \dots$$

By Theorems 3.3, 3.5 $\text{Ext}_R^i(S, H_{\mathfrak{a}}^j(M, N))$ is Matlis reflexive. By Corollary 3.7, $\text{Ext}_R^i(A, H_{\mathfrak{a}}^j(M, N))$ is Matlis reflexive. Thus $\text{Ext}_R^i(L, H_{\mathfrak{a}}^j(M, N))$ is Matlis reflexive for all i and j . \square

References

- [1] *M. P. Brodmann, R. Y. Sharp: Local Cohomology. An Algebraic Introduction with Geometric Applications.* Cambridge University Press, Cambridge, 1998.
- [2] *R. Belshoff, S. P. Slattery, C. Wickham: Finiteness properties for Matlis reflexive modules.* Commun. Algebra *24* (1996), 1371–1376.
- [3] *R. Belshoff, S. P. Slattery, C. Wickham: The local cohomology modules of Matlis reflexive modules are almost cofinite.* Proc. Am. Math. Soc. *124* (1996), 2649–2654.
- [4] *R. Belshoff, C. Wickham: A note on local duality.* Bull. Lond. Math. Soc. *29* (1997), 25–31.
- [5] *D. Delfino: On the cofiniteness of local cohomology modules.* Math. Proc. Camb. Philos. Soc. *115* (1994), 79–84.
- [6] *D. Delfino, T. Marley: Cofinite modules and local cohomology.* J. Pure Appl. Algebra *121* (1997), 45–52.

- [7] *K. Divaani-Aazar, R. Sazeedeh*: Cofiniteness of generalized local cohomology modules. *Colloq. Math.* *99* (2004), 283–290.
- [8] *K. Divaani-Aazar, R. Sazeedeh, M. Tousi*: On vanishing of generalized local cohomology modules. *Algebra Colloq.* *12* (2005), 213–218.
- [9] *R. Hartshorne*: Affine duality and cofiniteness. *Invent. Math.* *9* (1970), 145–164.
- [10] *J. Herzog*: Komplexe Auflösungen und Dualität in der lokalen Algebra. Habilitationsschrift. Universität Regensburg, Regensburg, 1970. (In German.)
- [11] *C. Huneke, J. Koh*: Cofiniteness and vanishing of local cohomology modules. *Math. Proc. Camb. Philos. Soc.* *110* (1991), 421–429.
- [12] *S. Kawakami, K.-I. Kawasaki*: On the finiteness of Bass numbers of generalized local cohomology modules. *Toyama Math. J.* *29* (2006), 59–64.
- [13] *K.-I. Kawasaki*: Cofiniteness of local cohomology modules for principal ideals. *Bull. Lond. Math. Soc.* *30* (1998), 241–246.
- [14] *K. Khashyarmanesh, F. Khosh-Ahang*: On the local cohomology of Matlis reflexive modules. *Commun. Algebra* *36* (2008), 665–669.
- [15] *A. Mafi*: A generalization of the finiteness problem in local cohomology modules. *Proc. Indian Acad. Sci. (Math. Sci.)* *119* (2009), 159–164.
- [16] *A. Mafi, H. Saremi*: Cofinite modules and generalized local cohomology. *Houston J. Math.* To appear.
- [17] *L. Melkersson*: Properties of cofinite modules and applications to local cohomology. *Math. Proc. Camb. Philos. Soc.* *125* (1999), 417–423.
- [18] *L. Melkersson*: Modules cofinite with respect to an ideal. *J. Algebra* *285* (2005), 649–668.
- [19] *A. Ooishi*: Matlis duality and width of a module. *Hiroshima Math. J.* *6* (1976), 573–587.
- [20] *J. Strooker*: *Homological Questions in Local Algebra*. Lecture Notes Series 145. Cambridge University Press, Cambridge, 1990.
- [21] *S. Yassemi*: Generalized section functors. *J. Pure Appl. Algebra* *95* (1994), 103–119.
- [22] *K. I. Yoshida*: Cofiniteness of local cohomology modules for ideals of dimension one. *Nagoya Math. J.* *147* (1997), 179–191.

Author's address: A. Mafi, Department of Mathematics, University of Kurdistan, P.O. Box 416, Sanandaj, Iran, and Institute for Studies in Theoretical Physics and Mathematics, P.O. Box 19395-5746, Tehran, Iran, e-mail: a_mafi@ipm.ir.