

Lingyu Jiang; Yidong Wang

On the blow up criterion for the 2-D compressible Navier-Stokes equations

Czechoslovak Mathematical Journal, Vol. 60 (2010), No. 1, 195–209

Persistent URL: <http://dml.cz/dmlcz/140562>

Terms of use:

© Institute of Mathematics AS CR, 2010

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON THE BLOW UP CRITERION FOR THE 2-D COMPRESSIBLE
NAVIER-STOKES EQUATIONS

LINGYU JIANG, YIDONG WANG, Beijing

(Received September 12, 2008)

Abstract. Motivated by [10], we prove that the upper bound of the density function ϱ controls the finite time blow up of the classical solutions to the 2-D compressible isentropic Navier-Stokes equations. This result generalizes the corresponding result in [3] concerning the regularities to the weak solutions of the 2-D compressible Navier-Stokes equations in the periodic domain.

Keywords: compressible Navier-Stokes equations, classical solutions, blow up criterion

MSC 2010: 35Q30, 76D03

1. INTRODUCTION

In this paper we consider the blow up criterion for the classical solutions to the following 2-D compressible isentropic Navier-Stokes equations in the periodic domain:

$$(1.1) \quad \begin{cases} \partial_t \varrho + \operatorname{div}(\varrho U) = 0, & (t, x) \in \mathbb{R}^+ \times \mathbb{T}^2, \\ \partial_t(\varrho U) + \operatorname{div}(\varrho U \otimes U) - \mu \Delta U - (\mu + \lambda) \nabla \operatorname{div} U + \nabla p(\varrho) = 0, \end{cases}$$

where $\varrho, U = (u, v)$ stand for the density and velocity of the viscous compressible fluid respectively, and μ, λ are the dynamical and volume viscosities such that $\mu > 0$ and $3\lambda + 2\mu \geq 0$. For simplicity, in what follows, we always take $\mu = 1$. We complement the above system with the initial data

$$(1.2) \quad \varrho|_{t=0} = \varrho_0, \quad U|_{t=0} = U_0.$$

Furthermore, we assume that there exist two positive constants m and M such that

$$(1.3) \quad m \leq \varrho_0(x) \leq M \quad \text{for } x \in \mathbb{T}^2.$$

The Navier-Stokes equations are the basic model describing the evolution of a viscous compressible gas. Before the celebrated works of P.L. Lions, very little was known about the global solutions to the multi-dimensional compressible Navier-Stokes equations. In particular, Lions [7] proved the global existence of weak solutions to (1.1) under the assumptions that

$$p(\varrho) = A\varrho^\gamma, \quad \varrho_0 \in L^1(\mathbb{T}^d) \cap L^\gamma(\mathbb{T}^d), \quad \varrho_0 \geq 0 \quad \text{and} \quad \frac{m_0^2}{\varrho_0} \in L^1(\mathbb{T}^d),$$

$$\gamma \geq \frac{3}{2} \quad \text{if } d = 2, \quad \gamma \geq \frac{9}{5} \quad \text{if } d = 3, \quad \gamma > \frac{d}{2} \quad \text{if } d \geq 4,$$

where we agree that $m_0^2/\varrho_0 = 0$ on the set $\{x \in \mathbb{T}^d \text{ such that } \varrho_0(x) = 0\}$. This result was improved later by Feireisl [4] et al to $\gamma > \frac{1}{2}d$.

On the other hand, as was emphasized in many papers related to the viscous compressible fluid dynamics, vacuum might be a major difficulty when one tries to prove the global classical solutions to (1.1), like in [9], where Xin proved the finite time blow up of classical solutions to the full compressible Navier-Stokes equations when the initial density has compact support. As a matter of fact, starting from initial densities with positive lower bound, local existence of smooth solutions to (1.1) can be proved by classical method (see [5], [8]), while in [3], Desjardins proved that $\sup_{t \in [0, T]} \|\varrho(t)\|_{L^\infty}$ controls the regularities of weak solutions to the 2-D isentropic compressible Navier-Stokes equations.

Motivated by [10], where the authors proved the global existence of classical solutions to (1.1) for $\lambda = \varrho^\beta$ with $\beta > 3$, we shall prove in this paper that $\sup_{t \in [0, T]} \|\varrho(t)\|_{L^\infty}$ controls the finite time blow up of classical solutions to (1.1). More precisely, we prove

Theorem 1.1. *Given $\varrho_0, U_0 \in H^3(\mathbb{T}^2)$ with ϱ_0 satisfying (1.3), and $p(\varrho) = A\varrho^\gamma$ for $A, \gamma > 0$, there exists a positive constant T such that (1.1)–(1.2) has a unique solution (ϱ, U) with $\varrho \in C([0, T]; H^3(\mathbb{T}^2))$, $U \in C([0, T]; H^3(\mathbb{T}^2)) \cap L^2((0, T); H^4(\mathbb{T}^2))$. We denote by T^* the maximal time of the existence of (ϱ, U) . Then if $T^* < \infty$, we have*

$$(1.4) \quad \lim_{T \rightarrow T^*} \sup_{t \in [0, T]} \|\varrho(t)\|_{L^\infty} = \infty.$$

Let us end this section with the notation we are going to use in this paper.

Notation. In the following, we shall denote by C and C_T uniform positive constants which may be different on different lines. We shall denote by (a, b) the $L^2(\mathbb{T}^2)$ inner product of a and b , and $\|a\|_{L^p}, \|a\|_{H^s}$ the standard $L^p(\mathbb{T}^2)$ and $H^s(\mathbb{T}^2)$ norms of a .

2. THE PROOF OF THE THEOREM

Under the assumptions of Theorem 1.1, in standard method can be applied to prove the local well posedness of (1.1)–(1.2), like the [5], [8], and we omit the details here. Now let (ϱ, U) be the local classical solution of (1.1)–(1.2) given by Theorem 1.1. Then one gets by the standard energy estimate that

$$(2.1) \quad \begin{aligned} & \frac{1}{2} \sup_{0 \leq s \leq t} \|U(s)\|_{L^2}^2 + \sup_{0 \leq s \leq t} \int_{\mathbb{T}^2} \varphi(\varrho) \, dx \, dy + \int_0^t [\|\nabla U\|_{L^2}^2 + (1 + \lambda) \|\operatorname{div} U\|_{L^2}^2] \, ds \\ & \leq \frac{1}{2} \|U_0\|_{L^2}^2 + \int_{\mathbb{T}^2} \varphi(\varrho_0) \, dx \, dy \quad \text{for } 0 \leq t < T, \end{aligned}$$

with $\varphi(\varrho) \stackrel{\text{def}}{=} \varrho \int_1^\varrho \tau^{-2} p(\tau) \, d\tau$.

To get an estimate of the derivatives of U , motivated by [10] we denote

$$(2.2) \quad \begin{aligned} A & \stackrel{\text{def}}{=} u_y - v_x, & B & \stackrel{\text{def}}{=} (2 + \lambda)(u_x + v_y) - p(\varrho), \\ L & \stackrel{\text{def}}{=} \frac{1}{\varrho}(A_y + B_x), & \text{and } H & \stackrel{\text{def}}{=} \frac{1}{\varrho}(-A_x + B_y). \end{aligned}$$

Then one can rewrite the momentum equations of (1.1) as

$$(2.3) \quad \begin{cases} u_t + u\partial_x u + v\partial_y u = \frac{1}{\varrho}(A_y + B_x), \\ v_t + u\partial_x v + v\partial_y v = \frac{1}{\varrho}(-A_x + B_y). \end{cases}$$

Lemma 2.1. *Let (ϱ, U) be the classical solution of (1.1)–(1.2) given by Theorem 1.1. We assume that $\sup_{t \in [0, T)} \|\varrho(t)\|_{L^\infty} < \infty$, then*

$$(2.4) \quad \sup_{0 \leq t < T} (\|A(t)\|_{L^2}^2 + \|B(t)\|_{L^2}^2) + \int_0^T (\|\nabla A(t)\|_{L^2}^2 + \|\nabla B(t)\|_{L^2}^2) \, dt \leq C_T,$$

where C_T depends only on $\|U_0\|_1$, $\|\varphi(\varrho_0)\|_{L^1}$ and $\sup_{t \in [0, T)} \|\varrho(t)\|_{L^\infty}$.

Proof. First, thanks to (2.2) and (2.3) we obtain

$$(2.5) \quad A_t + u\partial_x A + v\partial_y A + A \operatorname{div} U = L_y - H_x$$

and

$$(\operatorname{div} U)_t + u\partial_x(\operatorname{div} U) + v\partial_y(\operatorname{div} U) + u_x^2 + 2u_y v_x + v_y^2 = L_x + H_y.$$

On the other hand, thanks to the continuity equation of (1.1), one has

$$p(\varrho)_t + u\partial_x p(\varrho) + v\partial_y p(\varrho) = -\varrho p'(\varrho) \operatorname{div} U,$$

from which we deduce that

$$(2.6) \quad \frac{1}{2+\lambda} [B_t + u\partial_x B + v\partial_y B - \varrho p'(\varrho) \operatorname{div} U] + u_x^2 + 2u_y v_x + v_y^2 = L_x + H_y.$$

By virtue of (2.5) and (2.6) we get by the standard energy estimate that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\mathbb{T}^2} \left(A^2 + \frac{B^2}{2+\lambda} \right) dx dy + \int_{\mathbb{T}^2} [(LA_y - HA_x) + (LB_x + HB_y)] dx dy \\ & - \frac{1}{2+\lambda} \int_{\mathbb{T}^2} \varrho p'(\varrho) B \operatorname{div} U dx dy + \int_{\mathbb{T}^2} (uA_x + vA_y) A dx dy + \int_{\mathbb{T}^2} \operatorname{div} UA^2 dx dy \\ & + \frac{1}{2+\lambda} \int_{\mathbb{T}^2} (uB_x + vB_y) B dx dy + \int_{\mathbb{T}^2} B(u_x^2 + 2u_y v_x + v_y^2) dx dy = 0. \end{aligned}$$

However, notice that

$$(LA_y - HA_x) + (LB_x + HB_y) = \frac{1}{\varrho} (A_y + B_x)^2 + \frac{1}{\varrho} (-A_x + B_y)^2,$$

and

$$\begin{aligned} & \left[\sup_{t \in [0, T]} \|\varrho(t)\|_{L^\infty} \right]^{-1} \int_{\mathbb{T}^2} (|\nabla A|^2 + |\nabla B|^2) dx dy \\ & \leq \int_{\mathbb{T}^2} \frac{1}{\varrho} [(A_y + B_x)^2 + (-A_x + B_y)^2] dx dy \end{aligned}$$

for $0 \leq t < T$. Then one gets by using integration by parts

$$\begin{aligned} (2.7) \quad & \frac{1}{2} \frac{d}{dt} \int_{\mathbb{T}^2} \left(A^2 + \frac{B^2}{2+\lambda} \right) dx dy \\ & + \left[\sup_{t \in [0, T]} \|\varrho(t)\|_{L^\infty} \right]^{-1} \int_{\mathbb{T}^2} [(A_y + B_x)^2 + (-A_x + B_y)^2] dx dy \\ & \leq \frac{1}{2} \int_{\mathbb{T}^2} \left(-A^2 + \frac{B^2}{2+\lambda} \right) \operatorname{div} U dx dy + \frac{1}{2+\lambda} \int_{\mathbb{T}^2} \varrho p'(\varrho) B \operatorname{div} U dx dy \\ & - \int_{\mathbb{T}^2} B(u_x^2 + 2u_y v_x + v_y^2) dx dy, \quad \text{for } 0 \leq t < T. \end{aligned}$$

Note that for any $\varepsilon > 0$, Young's inequality applied gives

$$\begin{aligned} & \left| \int_{\mathbb{T}^2} \left(-A^2 + \frac{B^2}{2+\lambda} \right) \operatorname{div} U dx dy \right| \leq C(\|A\|_{L^4}^2 + \|B\|_{L^4}^2)(\|u_x\|_{L^2} + \|v_y\|_{L^2}) \\ & \leq \varepsilon(\|\nabla A\|_{L^2}^2 + \|\nabla B\|_{L^2}^2) + \frac{C}{\varepsilon}(\|u_x\|_{L^2}^2 + \|v_y\|_{L^2}^2)(\|A\|_{L^2}^2 + \|B\|_{L^2}^2), \end{aligned}$$

where we have used the trivial interpolation inequality in 2-D,

$$(2.8) \quad \|A\|_{L^4} \leq C \|A\|_{L^2}^{\frac{1}{2}} \|\nabla A\|_{L^2}^{\frac{1}{2}},$$

and it is easy to observe that

$$\left| \int_{\mathbb{T}^2} B(u_x^2 + 2u_y v_x + v_y^2) dx dy \right| \leq C \|B\|_{L^4} (\|\nabla u\|_{L^{\frac{8}{3}}}^2 + \|\nabla v\|_{L^{\frac{8}{3}}}^2).$$

On the other hand, thanks to (2.2) we have

$$\begin{aligned} \|\nabla u\|_{L^{\frac{8}{3}}}^2 + \|\nabla v\|_{L^{\frac{8}{3}}}^2 &\leq C (\|u_x + v_y\|_{L^{\frac{8}{3}}}^2 + \|u_y - v_x\|_{L^{\frac{8}{3}}}^2), \quad \text{and} \\ \|u_y - v_x\|_{L^{\frac{8}{3}}} &= \|A\|_{L^{\frac{8}{3}}} \leq C \|A\|_{L^2}^{\frac{3}{4}} \|\nabla A\|_{L^2}^{\frac{1}{4}}, \\ \|u_x + v_y\|_{L^{\frac{8}{3}}} &\leq \left\| \frac{B}{2 + \lambda} + p(\varrho) \right\|_{L^{\frac{8}{3}}} \leq C_T (1 + \|B\|_{L^2}^{\frac{3}{4}} \|\nabla B\|_{L^2}^{\frac{1}{4}}), \end{aligned}$$

with C_T depending only on $\sup_{t \in [0, T]} \|\varrho(t)\|_{L^\infty}$. As a consequence, we obtain

$$\left| \int_{\mathbb{T}^2} B(u_x^2 + 2u_y v_x + v_y^2) dx dy \right| \leq \varepsilon (\|\nabla A\|_{L^2}^2 + \|\nabla B\|_{L^2}^2) + \frac{C_T}{\varepsilon} (1 + \|A\|_{L^2}^4 + \|B\|_{L^2}^4)$$

for any $\varepsilon > 0$. Then thanks to (2.7), we get by taking $\varepsilon \leq \frac{1}{4} \left[\sup_{t \in [0, T]} \|\varrho(t)\|_{L^\infty} \right]^{-1}$ that

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{T}^2} \left(A^2 + \frac{B^2}{2 + \lambda} \right) dx dy + \left[\sup_{t \in [0, T]} \|\varrho(t)\|_{L^\infty} \right]^{-1} \int_{\mathbb{T}^2} (|\nabla A|^2 + |\nabla B|^2) dx dy \\ \leq C_T (1 + \|\nabla u\|_{L^2}^2 + \|\nabla v\|_{L^2}^2) \int_{\mathbb{T}^2} \left(A^2 + \frac{B^2}{2 + \lambda} \right) dx dy, \end{aligned}$$

and applying (2.1) and the Gronwall inequality gives (2.4). \square

Lemma 2.2. *Under the assumptions of Lemma 2.1, one can find a positive constant m_T which depends only on $\sup_{t \in [0, T]} \|\varrho(t)\|_{L^\infty}$, $\|U_0\|_{H^1}$ and m, M in (1.3) such that*

$$(2.9) \quad \varrho(t, x) \geq m_T \quad \text{for } 0 \leq t < T.$$

Proof. We first get by taking divergence to the momentum equation of (1.1) that

$$\Delta [(2 + \lambda) \operatorname{div} U - p(\varrho)] = \partial_t \operatorname{div}(\varrho U) + \operatorname{div}(\operatorname{div}(\varrho U \otimes U)),$$

which implies that

$$\operatorname{div} U = \frac{1}{2+\lambda} [\partial_t \Delta^{-1} \operatorname{div}(\varrho U) + \Delta^{-1} \operatorname{div}(\operatorname{div}(\varrho U \otimes U)) + p(\varrho)],$$

where $\Delta^{-1} f$ is defined as the unique solution of

$$\Delta N = f \quad \text{on } \mathbb{T}^2, \quad \int_{\mathbb{T}^2} N \, dx \, dy = 0.$$

Then thanks to the continuity equation of (1.1),

$$\partial_t \varrho + U \cdot \nabla \varrho + \varrho \operatorname{div} U = 0,$$

one has

$$\partial_t \log \varrho + u \cdot \nabla \log \varrho + \frac{1}{2+\lambda} [\partial_t \Delta^{-1} \operatorname{div}(\varrho U) + \Delta^{-1} \operatorname{div}(\operatorname{div}(\varrho U \otimes U)) + p(\varrho)] = 0,$$

which gives

$$(2.10) \quad \begin{aligned} & \partial_t \left[\log \varrho + \frac{1}{2+\lambda} \operatorname{div} \Delta^{-1}(\varrho U) \right] + u \cdot \nabla \left[\log \varrho + \frac{1}{2+\lambda} \operatorname{div} \Delta^{-1}(\varrho U) \right] \\ &= \frac{1}{2+\lambda} [u \cdot \nabla \operatorname{div} \Delta^{-1}(\varrho U) - \Delta^{-1} \operatorname{div} \operatorname{div}(\varrho U \otimes U) - p(\varrho)]. \end{aligned}$$

On the other hand, motivated by [2], we denote $D \stackrel{\text{def}}{=} \Delta^{-1} \operatorname{div} \operatorname{div}(\varrho U \otimes U) - u \cdot \nabla \operatorname{div} \Delta^{-1}(\varrho U)$ and $G \stackrel{\text{def}}{=} \operatorname{div} \Delta^{-1}(\varrho U)$. Then

$$\Delta D = \operatorname{div} [\varrho U \cdot \nabla U + \operatorname{div} U \nabla G - \nabla U \cdot \nabla G + \nabla G \cdot \nabla U],$$

from which and (2.2) one gets by the standard potential theory that

$$\begin{aligned} \|\nabla D\|_{L^3} &\leq C \|(\varrho U + |\nabla G|) |\nabla U|\|_{L^3} \\ &\leq C \|\varrho\|_{L^\infty} \|U\|_{L^6} \|\nabla U\|_{L^6} \leq C_T \|U\|_{L^2}^{\frac{1}{3}} \|\nabla U\|_{L^2}^{\frac{2}{3}} [\|A\|_{L^6} + \|B\|_{L^6} + 1]. \end{aligned}$$

However, thanks to (2.4) we have

$$\begin{aligned} & \int_0^T (\|A(t)\|_{L^6} + \|B(t)\|_{L^6}) \, dt \\ & \leq C \int_0^T [\|A(t)\|_{L^2}^{\frac{1}{3}} \|\nabla A(t)\|_{L^2}^{\frac{2}{3}} + \|B(t)\|_{L^2}^{\frac{1}{3}} \|\nabla B(t)\|_{L^2}^{\frac{2}{3}}] \, dt \leq C_T, \end{aligned}$$

which gives

$$(2.11) \quad \int_0^T \|\nabla D\|_{L^3} dt \leq C_T.$$

Similarly to the proof of (2.11) we have

$$\|D\|_{L^3} \leq C\|(\varrho U + |\nabla G|)|\nabla U\|_{L^{\frac{6}{5}}} \leq C\|\varrho\|_{L^\infty} \|U\|_{L^2}^{\frac{2}{3}} \|\nabla U\|_{L^2}^{\frac{4}{3}} \leq C_T,$$

which gives

$$(2.12) \quad \int_0^T \|D(t)\|_{L^\infty} dt \leq C_T.$$

Thanks to (1.3), (2.10) and (2.12), one gets by using the characteristic method that there exists a positive constant C_T such that

$$(2.13) \quad \log \varrho(t, x) + \frac{1}{2 + \lambda} G(t, x) \geq -C_T \quad \text{for } 0 \leq t < T.$$

Now we note that

$$\begin{aligned} \|G\|_{L^4} &\leq C\|\varrho\|_{L^\infty} \|U\|_{L^{\frac{4}{3}}} \leq C\|U\|_{L^2}, \\ \|\nabla G\|_{L^4} &\leq C\|\varrho\|_{L^\infty} \|U\|_{L^4} \leq C\|\varrho\|_{L^\infty} \|U\|_{L^2}^{\frac{1}{2}} \|\nabla U\|_{L^2}^{\frac{1}{2}}, \end{aligned}$$

which together with (2.1) and (2.4) implies that

$$\sup_{t \in [0, T)} \|G(t)\|_{L^\infty} \leq C_T,$$

from which and (2.13) one obtains (2.9). This completes the proof of Lemma 2.2. \square

An estimate of higher derivatives of U is obtained by carrying out calculations similar to but easier than (15) and (16) in [10]:

$$(2.14) \quad \begin{cases} \varrho(L_t + U \cdot \nabla L) - \varrho L \operatorname{div} U + u_y A_x + v_y A_y + u_x B_x + v_x B_y + (A \operatorname{div} U)_y \\ \quad - (p'(\varrho)\varrho \operatorname{div} U)_x + (2 + \lambda)(u_x^2 + v_y^2 + 2v_x u_y) \\ = (L_y - H_x)_y + (2 + \lambda)(L_x + H_y)_x, \\ \varrho(H_t + U \cdot \nabla H) - \varrho H \operatorname{div} U - u_x A_x - v_x A_y + u_y B_x + v_y B_y - (A \operatorname{div} U)_x \\ \quad - (p'(\varrho)\varrho \operatorname{div} U)_y + (2 + \lambda)(u_x^2 + v_y^2 + 2v_x u_y)_y \\ = -(L_y - H_x)_x + (2 + \lambda)(L_x + H_y)_y. \end{cases}$$

Now let us turn to the estimate of L, H .

Lemma 2.3. *Under the assumptions of Lemma 2.1, there exists a positive constant C_T which depends on $\sup_{t \in [0, T]} \|\varrho(t)\|_{L^\infty}$, $\|U_0\|_{H^2}$, $\|\varrho_0\|_{H^1}$ and m, M in (1.3) such that*

$$(2.15) \quad \sup_{0 \leq t < T} \int_{\mathbb{T}^2} (L^2 + H^2) dx dy + \int_0^T \int_{\mathbb{T}^2} [(L_y - H_x)^2 + (L_x + H_y)^2] dx dy dt \leq C_T.$$

Proof. Thanks to (2.14), one gets by using the standard energy estimate that

$$(2.16) \quad \begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\mathbb{T}^2} \varrho(L^2 + H^2) dx dy + \int_{\mathbb{T}^2} [(L_y - H_x)^2 + (2 + \lambda)(L_x + H_y)^2] dx dy \\ & - \int_{\mathbb{T}^2} \varrho(L^2 + H^2) \operatorname{div} U dx dy + \int_{\mathbb{T}^2} A \operatorname{div} U (H_x - L_y) dx dy \\ & + \int_{\mathbb{T}^2} \varrho(L_x + H_y) \operatorname{div} U p'(\varrho) dx dy \\ & + \int_{\mathbb{T}^2} L(u_y A_x + v_y A_y + u_x B_x + v_x B_y) dx dy \\ & + \int_{\mathbb{T}^2} H(-u_x A_x - v_x A_y + u_y B_x + v_y B_y) dx dy \\ & - (2 + \lambda) \int_{\mathbb{T}^2} (L_x + H_y)(u_x^2 + 2v_x u_y + v_y^2) dx dy = 0. \end{aligned}$$

Applying the Cauchy-Schwartz inequality, we obtain

$$\left| \int_{\mathbb{T}^2} \varrho(L^2 + H^2) \operatorname{div} U dx dy \right| \leq \|\varrho\|_{L^\infty} \|\nabla U\|_{L^2} (\|L\|_{L^4}^2 + \|H\|_{L^4}^2),$$

and using (2.8) we arrive at

$$\begin{aligned} \|L\|_{L^4}^2 + \|H\|_{L^4}^2 & \leq C(\|L\|_{L^2} + \|H\|_{L^2})(\|\nabla L\|_{L^2} + \|\nabla H\|_{L^2}) \\ & \leq C(\|L\|_{L^2} + \|H\|_{L^2})(\|L_y - H_x\|_{L^2} + \|L_x + H_y\|_{L^2}). \end{aligned}$$

As a consequence, for any $\varepsilon > 0$ one has

$$\begin{aligned} & \left| \int_{\mathbb{T}^2} \varrho(L^2 + H^2) \operatorname{div} U dx dy \right| \\ & \leq \varepsilon(\|L_y - H_x\|_{L^2}^2 + \|L_x + H_y\|_{L^2}^2) + \frac{C_T}{\varepsilon} \|\nabla U\|_{L^2}^2 (\|L\|_{L^2}^2 + \|H\|_{L^2}^2). \end{aligned}$$

A similar argument yields

$$\begin{aligned} & \left| \int_{\mathbb{T}^2} A \operatorname{div} U (H_x - L_y) dx dy \right| \leq \varepsilon \|H_x - L_y\|_{L^2}^2 + \frac{C}{\varepsilon} \|A\|_{L^4}^2 \|\operatorname{div} U\|_{L^4}^2 \\ & \leq \varepsilon \|H_x - L_y\|_{L^2}^2 + \frac{C_T}{\varepsilon} (1 + \|A\|_{L^2}^2 \|\nabla A\|_{L^2}^2 + \|B\|_{L^2}^2 \|\nabla B\|_{L^2}^2), \end{aligned}$$

where we used (2.2) and (2.8), so that

$$(2.17) \quad \begin{aligned} \|\nabla U\|_{L^4} &\leq C(\|\operatorname{div} U\|_{L^4} + \|u_y - v_x\|_{L^4}) \\ &\leq C_T(1 + \|A\|_{L^2}^{\frac{1}{2}}\|\nabla A\|_{L^2}^{\frac{1}{2}} + \|B\|_{L^2}^{\frac{1}{2}}\|\nabla B\|_{L^2}^{\frac{1}{2}}). \end{aligned}$$

It is easy to observe that

$$\left| \int_{\mathbb{T}^2} \varrho(L_x + H_y) \operatorname{div} U p'(\varrho) \, dx \, dy \right| \leq \varepsilon \|L_x + H_y\|_{L^2}^2 + \frac{C}{\varepsilon} \|\varrho p'(\varrho)\|_{L^\infty}^2 \|\nabla U\|_{L^2}^2$$

and

$$\left| \int_{\mathbb{T}^2} L(u_y A_x + v_y A_y + u_x B_x + v_x B_y) \, dx \, dy \right| \leq \|L\|_{L^2} \|\nabla U\|_{L^4} (\|\nabla A\|_{L^4} + \|\nabla B\|_{L^4}),$$

while thanks to (2.2) and (2.9) one has

$$\begin{aligned} \|\nabla A\|_{L^4} + \|\nabla B\|_{L^4} &\leq \frac{C}{m_T} (\|L\|_{L^4} + \|H\|_{L^4}) \\ &\leq C_T (\|L\|_{L^2} + \|H\|_{L^2})^{\frac{1}{2}} (\|L_x + H_y\|_{L^2} + \|L_y - H_x\|_{L^2})^{\frac{1}{2}}. \end{aligned}$$

Then one obtains

$$\begin{aligned} &\left| \int_{\mathbb{T}^2} L(u_y A_x + v_y A_y + u_x B_x + v_x B_y) \, dx \, dy \right| \\ &\leq \varepsilon (\|L_x + H_y\|_{L^2}^2 + \|L_y - H_x\|_{L^2}^2) \\ &\quad + \frac{C_T}{\varepsilon} (\|L\|_{L^2}^2 + \|H\|_{L^2}^2) (1 + \|A\|_{L^2}^{\frac{1}{2}}\|\nabla A\|_{L^2}^{\frac{1}{2}} + \|B\|_{L^2}^{\frac{1}{2}}\|\nabla B\|_{L^2}^{\frac{1}{2}})^{\frac{4}{3}}. \end{aligned}$$

A similar estimate holds for $\int_{\mathbb{T}^2} H(-u_x A_x - v_x A_y + u_y B_x + v_y B_y) \, dx \, dy$.

Finally, again thanks to (2.17) one has

$$\begin{aligned} &\left| \int_{\mathbb{T}^2} (L_x + H_y)(u_x^2 + 2v_x u_y + v_y^2) \, dx \, dy \right| \\ &\leq \varepsilon \|L_x + H_y\|_{L^2}^2 + \frac{C_T}{\varepsilon} (1 + \|A\|_{L^2}^2 \|\nabla A\|_{L^2}^2 + \|B\|_{L^2}^2 \|\nabla B\|_{L^2}^2). \end{aligned}$$

Combining the above estimates, we get by taking ε small enough

$$(2.18) \quad \begin{aligned} &\frac{1}{2} \frac{d}{dt} \int_{\mathbb{T}^2} \varrho(L^2 + H^2) \, dx \, dy + \int_{\mathbb{T}^2} [(L_y - H_x)^2 + (2 + \lambda)(L_x + H_y)^2] \, dx \, dy \\ &\leq \frac{C_T}{\varepsilon} (1 + \|L\|_{L^2}^2 + \|H\|_{L^2}^2) \\ &\quad \times [1 + \|\nabla U\|_{L^2}^2 + \|A\|_{L^2}^2 \|\nabla A\|_{L^2}^2 + \|B\|_{L^2}^2 \|\nabla B\|_{L^2}^2]. \end{aligned}$$

Then thanks to (2.1), (2.4) and (2.9) one obtains (2.15) by using the Gronwall inequality. This completes the proof of the lemma. □

With the above estimate for the first derivatives of U , we now turn to an estimate of the first derivatives of ϱ .

Proposition 2.1. *Under the assumptions of Theorem 1.1, there is a positive constant C_T , which depends on $\sup_{t \in [0, T]} \|\varrho(t)\|_{L^\infty}$, $\|\varrho_0\|_{H^2}$, $\|U_0\|_{H^2}$ and m, M in (1.3) such that*

$$(2.19) \quad \sup_{0 \leq t \leq T} \|\nabla \varrho(t)\|_{L^q} \leq C_T \quad \text{for any } q < \infty.$$

Proof. First, thanks to (2.15) we deduce that both L and H are bounded in $L^p([0, T]; L^q(\mathbb{T}^2))$ with p, q satisfying $1/q = \frac{1}{2} - 1/p$. As a consequence, we get using (2.2) and $\sup_{t \in [0, T]} \|\varrho(t)\|_{L^\infty} < \infty$ that

$$(2.20) \quad \nabla A, \nabla B \quad \text{are bounded in } L^p([0, T]; L^q(\mathbb{T}^2)),$$

in particular, $B \in L^p([0, T]; L^\infty(\mathbb{T}^2))$ for any $p < \infty$.

Now let X be the flow of U given by

$$\partial_t X(t, s, x) = U(t, X(t, s, x)), \quad X(t, s, x)|_{t=s} = x,$$

from which, (2.2) and (2.20) we deduce that

$$(2.21) \quad \det \left(\frac{\partial X(t, s, x)}{\partial x} \right) = \exp \left(\int_0^t \operatorname{div} U(t', X(t', s, x)) dt' \right) \\ = \exp \left(\frac{1}{2 + \lambda} \int_0^t (B + p(\varrho))(t', X(t', s, x)) dt' \right) \leq C_T.$$

Thanks to the continuity equation of (1.1) and (2.2) we have

$$\frac{d}{dt} \varrho(t, X(t, s, x)) = -\frac{\varrho}{2 + \lambda} (B + A\varrho^\gamma)(t, X(t, s, x)),$$

which gives

$$\varrho^{-\gamma}(t, x) = \varrho_0^{-\gamma}(X(0, t, x)) e^{\gamma(2+\lambda)^{-1} \int_0^t B(s, X(s, t, x)) ds} \\ + \frac{A\gamma}{2 + \lambda} \int_0^t e^{\gamma(2+\lambda)^{-1} \int_\tau^t B(s, X(s, t, x)) ds} d\tau \stackrel{\text{def}}{=} M_1(t, x) + M_2(t, x),$$

and

$$(2.22) \quad \varrho(t, x) = (M_1(t, x) + M_2(t, x))^{-1/\gamma} \quad \text{and} \\ \nabla \varrho(t, x) = -\frac{1}{\gamma} (M_1(t, x) + M_2(t, x))^{-1/\gamma - 1} (\nabla M_1(t, x) + \nabla M_2(t, x)).$$

Note that

$$\begin{aligned} \nabla M_1(t, x) &= \nabla \varrho_0^{-\gamma}(X(0, t, x)) \cdot \nabla X(0, t, x) e^{\gamma(2+\lambda)^{-1} \int_0^t B(s, X(s, t, x)) ds} \\ &+ \frac{\gamma}{2+\lambda} \varrho_0^{-\gamma}(X(0, t, x)) e^{\gamma(2+\lambda)^{-1} \int_0^t B(s, X(s, t, x)) ds} \int_0^t \nabla B(s, X(s, t, x)) \cdot \nabla X(s, t, x) ds \end{aligned}$$

and

$$\sup_{(s, t) \in [0, T]^2} |\nabla X(s, t, x)| \leq \exp \left(\int_0^T |\nabla U(s, x)| ds \right).$$

Then thanks to (2.2) and (2.15), we obtain that $\nabla U \in L^2([0, T]; BMO(\mathbb{T}^2))$, and

$$\begin{aligned} (2.23) \quad \|\nabla M_1(t, \cdot)\|_{L^q} &\leq C_T \left(\int_{\mathbb{T}^2} \exp \left(q' \int_0^T |\nabla U(s, x)| ds \right) dx \right)^{1/q'} \\ &\leq C_T \left(\frac{1}{T} \int_0^T \int_{\mathbb{T}^2} \exp(q' |\nabla U(s, x)|) dx ds \right)^{1/q'} \leq C_T. \end{aligned}$$

A similar estimate holds for ∇M_2 as well, which together with (2.20) and (2.22) completes the proof of the proposition. \square

Now we are in position to complete the proof of Theorem 1.1.

Proof of Theorem 1.1. Let (ϱ, U) be the local classical solution of (1.1)–(1.2) given by Theorem 1.1. Then if $T^* < \infty$, we have

$$\lim_{T \rightarrow T^*} \sup_{t \in [0, T]} [\|\varrho(t)\|_{H^3} + \|U(t)\|_{H^3}] = \infty.$$

Therefore to complete the proof, we need to prove that if $\sup_{t \in [0, T]} \|\varrho(t)\|_{L^\infty} < \infty$, then

$$(2.24) \quad \sup_{t \in [0, T]} [\|\varrho(t)\|_{H^3} + \|U(t)\|_{H^3}] < \infty.$$

In fact, thanks to (2.5) and (2.6), we get by the standard energy estimate that

$$\begin{aligned} (2.25) \quad &\frac{1}{2} \frac{d}{dt} \left(\|A\|_{H^2}^2 + \frac{1}{2+\lambda} \|B\|_{H^2}^2 \right) \\ &+ \sum_{|\alpha| \leq 2} \int_{\mathbb{T}^2} [\partial^\alpha H(\partial^\alpha B_y - \partial^\alpha A_x) + \partial^\alpha L(\partial^\alpha A_y + \partial^\alpha B_x)] dx dy \\ &= \sum_{|\alpha| \leq 2} \left[-(\partial^\alpha(U \cdot \nabla A), \partial^\alpha A) + \frac{1}{2+\lambda} (\partial^\alpha(U \cdot \nabla B), \partial^\alpha B) \right. \\ &\quad \left. - \frac{1}{2+\lambda} (\partial^\alpha[(B + p(\varrho))A], \partial^\alpha A) \right. \\ &\quad \left. + \left(\partial^\alpha \left[\frac{1}{2+\lambda} \varrho p'(\varrho) \operatorname{div} U - (u_x^2 + 2u_y v_x + v_y^2) \right], \partial^\alpha B \right) \right]. \end{aligned}$$

Taking into account (2.2) we get

$$\begin{aligned}
& \sum_{|\alpha| \leq 2} \int_{\mathbb{T}^2} [\partial^\alpha H(\partial^\alpha B_y - \partial^\alpha A_x) + \partial^\alpha L(\partial^\alpha A_y + \partial^\alpha B_x)] dx dy \\
&= \int_{\mathbb{T}^2} \frac{1}{\varrho} \sum_{|\alpha| \leq 2} [|\partial^\alpha B_y - \partial^\alpha A_x|^2 + |\partial^\alpha A_y + \partial^\alpha B_x|^2] dx dy \\
&\quad + \sum_{|\alpha| \leq 2} \int_{\mathbb{T}^2} \left\{ \left[\partial^\alpha; \frac{1}{\varrho} \right] (-A_x + B_y) \times (\partial^\alpha B_y - \partial^\alpha A_x) \right. \\
&\quad \left. + \left[\partial^\alpha; \frac{1}{\varrho} \right] (A_y + B_x) \times (\partial^\alpha A_y + \partial^\alpha B_x) \right\} dx dy \stackrel{\text{def}}{=} R_1 + R_2.
\end{aligned}$$

However, it is easy to observe that for any $\varepsilon > 0$, we have

$$|R_2| \leq \varepsilon (\|B_y - A_x\|_{H^2}^2 + \|A_y + B_x\|_{H^2}^2) + \frac{C_T}{\varepsilon} (\|\nabla A\|_{H^1}^2 + \|\nabla B\|_{H^1}^2) \|\varrho\|_{H^3}^2,$$

from which we deduce that

$$\begin{aligned}
& \sum_{|\alpha| \leq 2} \int_{\mathbb{T}^2} [\partial^\alpha H(\partial^\alpha B_y - \partial^\alpha A_x) + \partial^\alpha L(\partial^\alpha A_y + \partial^\alpha B_x)] dx dy \\
& \geq (c_T - \varepsilon) (\|B_y - A_x\|_{H^2}^2 + \|A_y + B_x\|_{H^2}^2) - \frac{C_T}{\varepsilon} (\|\nabla A\|_{H^1}^2 + \|\nabla B\|_{H^1}^2) \|\varrho\|_{H^3}^2.
\end{aligned}$$

Applying a Moser type inequality gives

$$\begin{aligned}
& \left| \sum_{|\alpha| \leq 2} (\partial^\alpha (U \cdot \nabla A), \partial^\alpha A) \right| \\
&= \left| \sum_{|\alpha| \leq 2} [(U \cdot \nabla \partial^\alpha A, \partial^\alpha A) + (\partial^\alpha (U \cdot \nabla A) - U \cdot \nabla \partial^\alpha A, \partial^\alpha A)] \right| \\
&\leq C [\|\nabla U\|_{L^\infty} \|A\|_{H^2} + \|\nabla A\|_{H^1} \|U\|_{H^s}] \|A\|_{H^2},
\end{aligned}$$

and thanks to (2.2), we have

$$(2.26) \quad \|U\|_{H^s} \leq \|U\|_{L^2} + C[\|\varrho\|_{H^{s-1}} + \|A\|_{H^{s-1}} + \|B\|_{H^{s-1}}],$$

which gives

$$\left| \sum_{|\alpha| \leq 2} (\partial^\alpha (U \cdot \nabla A), \partial^\alpha A) \right| \leq C(\|\nabla A\|_{H^1} + \|\nabla U\|_{L^\infty})(1 + \|\varrho\|_{H^2}^2 + \|A\|_{H^2}^2 + \|B\|_{H^2}^2).$$

A similar estimate holds for $\sum_{|\alpha| \leq 2} (\partial^\alpha (U \cdot \nabla B), \partial^\alpha B)$.

Applying a Moser type inequality again gives

$$\left| \sum_{|\alpha| \leq 2} (\partial^\alpha [(B+p(\varrho))A], \partial^\alpha A) \right| \leq C_T(1+\|A\|_{L^\infty}+\|B\|_{L^\infty})(1+\|\varrho\|_{H^2}^2+\|A\|_{H^2}^2+\|B\|_{H^2}^2)$$

and

$$\begin{aligned} & \left| \sum_{|\alpha| \leq 2} \left(\partial^\alpha \left[\frac{1}{2+\lambda} \varrho p'(\varrho) \operatorname{div} U - (u_x^2 + 2u_y v_x + v_y^2) \right], \partial^\alpha B \right) \right| \\ & \leq C_T(1 + \|\nabla U\|_{L^\infty})(1 + \|\varrho\|_{H^2}^2 + \|A\|_{H^2}^2 + \|B\|_{H^2}^2). \end{aligned}$$

Then thanks to (2.25) we obtain by taking ε small enough in the above inequalities that

$$(2.27) \quad \begin{aligned} & \frac{d}{dt} \left(\|A\|_{H^2}^2 + \frac{1}{2+\lambda} \|B\|_{H^2}^2 \right) + c_T(\|B_y - A_x\|_{H^2}^2 + \|A_y + B_x\|_{H^2}^2) \\ & \leq C_T(1 + \|\nabla U\|_{L^\infty} + \|A\|_{H^2}^2 + \|B\|_{H^2}^2)(1 + \|\varrho\|_{H^3}^2 + \|A\|_{H^2}^2 + \|B\|_{H^2}^2). \end{aligned}$$

On the other hand, rewriting the continuity equation of (1.1) as

$$\partial_t \varrho + U \cdot \nabla \varrho + \frac{A}{2+\lambda} \varrho^{1+\gamma} + \frac{1}{2+\lambda} B \varrho = 0,$$

we get by the standard energy estimate that

$$\frac{d}{dt} \|\varrho\|_{H^3}^2 \leq C[(1+\|B\|_{L^\infty}+\|\nabla U\|_{L^\infty})\|\varrho\|_{H^3}+\|U\|_{H^3}\|\nabla \varrho\|_{L^\infty}+\|\varrho\|_{L^\infty}\|B\|_{H^3}]\|\varrho\|_{H^3}.$$

However, trivial interpolation inequality gives

$$\|U\|_{H^3} \leq C\|U\|_{H^1}^{\frac{1}{3}}\|U\|_{H^4}^{\frac{2}{3}}, \quad \|\nabla \varrho\|_{L^\infty} \leq \|\nabla \varrho\|_{L^2}^{\frac{2}{3}}\|\varrho\|_{H^3}^{\frac{1}{3}},$$

which together with (2.26) implies that

$$\|U\|_{H^3}\|\nabla \varrho\|_{L^\infty}\|\varrho\|_{H^3} \leq \varepsilon(\|A\|_{H^3}^2 + \|B\|_{H^3}^2) + \frac{C}{\varepsilon}(\|U\|_{H^1} + \|\nabla \varrho\|_{L^2})(1 + \|\varrho\|_{H^3}^2).$$

Therefore, we get

$$(2.28) \quad \begin{aligned} \frac{d}{dt} \|\varrho\|_{H^3}^3 & \leq \varepsilon(\|A\|_{H^3}^2 + \|B\|_{H^3}^2) + \frac{C_T}{\varepsilon}(\|U\|_{H^1} + \|B\|_{L^\infty} \\ & \quad + \|\nabla \varrho\|_{L^2} + \|\nabla U\|_{L^\infty})(1 + \|\varrho\|_{H^3}^2). \end{aligned}$$

Finally, thanks to [6] and (2.26) we have

$$\begin{aligned}
 (2.29) \quad & \|\nabla U\|_{L^\infty} \\
 & \leq C(1 + \|\nabla U\|_{BMO} \times \log^+ \|U\|_{H^3}) \\
 & \leq C_T [1 + (\|L\|_{H^1} + \|H\|_{H^1}) \log^+(1 + \|\varrho\|_{H^2} + \|A\|_{H^2} + \|B\|_{H^2})].
 \end{aligned}$$

On the other hand, we deduce from (2.2), (2.15) and (2.19) that $\varrho L, \varrho H \in L^2((0, T); H^1(\mathbb{T}^2))$, which together with (2.2) yields that both ∇A and ∇B are bounded in $L^2((0, T); H^1(\mathbb{T}^2))$. Then combining (2.27)–(2.29), we get

$$\begin{aligned}
 & \sup_{0 \leq t < T} (\|A(t)\|_{H^2}^2 + \|B(t)\|_{H^2}^2 + \|\varrho(t)\|_{H^3}^2) + \int_0^T (\|B_y - A_x\|_{H^2}^2 + \|A_y + B_x\|_{H^2}^2) dt \\
 & \leq C(m, M, \|\varrho_0\|_{H^3}, \|U_0\|_{H^3}, T),
 \end{aligned}$$

which together with (2.1) implies that

$$\sup_{0 \leq t < T} (\|\varrho(t)\|_{H^3}^2 + \|U(t)\|_{H^3}^2) + \int_0^T \|\nabla U\|_{H^3}^2 dt \leq C(m, M, \|\varrho_0\|_{H^3}, \|U_0\|_{H^3}, T).$$

This implies (2.24), and we have completed the proof of Theorem 1.1. \square

Acknowledgments. The authors would like to thank Professors Ping Zhang and Zhifei Zhang for profitable discussions concerning this topic. We also thank the referee for valuable suggestions.

References

- [1] *J. T. Beale, T. Kato and A. Majda*: Remarks on the breakdown of smooth solutions for the 3-D Euler equations. *Comm. Math. Phys.* *94* (1984), 61–66.
- [2] *H. J. Choe and B. J. Jin*: Regularity of weak solutions of the compressible Navier-Stokes equations. *J. Korean Math. Soc.* *40* (2003), 1031–1050.
- [3] *B. Desjardins*: Regularity of weak solutions of the compressible isentropic Navier-Stokes equations. *Comm. P. D. E.* *22* (1997), 977–1008.
- [4] *E. Feireisl, A. Novotný and H. Petzeltová*: On the existence of globally defined weak solutions to the Navier-Stokes equations. *J. Math. Fluid Mech.* *3* (2001), 358–392.
- [5] *N. Itaya*: On the Cauchy problem for the system of fundamental equations describing movement of compressible viscous fluids. *Kōdai Math. Sem. Rep.* *23* (1971), 60–120.
- [6] *H. Kozono and Y. Taniuchi*: Limiting case of the Sobolev inequality in BMO, with application to the Euler equations. *Comm. Math. Phys.* *214* (2000), 191–200.
- [7] *P. L. Lions*: *Mathematical Topics in Fluid Mechanics, Vol 2. Compressible Models*. Oxford lecture series in Mathematics and its Applications, 10, Oxford Sciences Publications. The Clarendon Press, Oxford University Press, New York, 1998.
- [8] *A. Tani*: On the first initial-boundary value problem of compressible viscous fluid motion. *Publ. RIMS. Kyoto Univ.* *13* (1977), 193–253.

- [9] *Z. P. Xin*: Blow up of smooth solutions to the compressible Navier-Stokes equation with compact density. *Comm. Pure Appl. Math.* *51* (1998), 229–240.
- [10] *V. A. Vaigant and A. V. Kazhikhov*: On the existence of global solutions of two-dimensional Navier-Stokes equations of a compressible viscous fluid. *Sibirsk. Mat. Zh.* *36* (1995), 1283–1316 translation in it *Siberian Math. J.* *36* (1995). (In Russian.)

Authors' address: Lingyu Jiang, Yidong Wang, Department of Mathematics, Central University of Finance and Economics, Beijing 100081, China, e-mail: jiangly1221@yahoo.com.cn.