

Milan Matejdes

Upper quasi continuous maps and quasi continuous selections

Czechoslovak Mathematical Journal, Vol. 60 (2010), No. 2, 517–525

Persistent URL: <http://dml.cz/dmlcz/140586>

Terms of use:

© Institute of Mathematics AS CR, 2010

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

UPPER QUASI CONTINUOUS MAPS AND
QUASI CONTINUOUS SELECTIONS

MILAN MATEJDES, Zlín

(Received December 19, 2008)

Abstract. The paper deals with the existence of a quasi continuous selection of a multifunction for which upper inverse image of any open set with compact complement contains a set of the form $(G \setminus I) \cup J$, where G is open and I, J are from a given ideal. The methods are based on the properties of a minimal multifunction which is generated by a cluster process with respect to a system of subsets of the form $(G \setminus I) \cup J$.

Keywords: selection, quasi continuity, minimal usco multifunction, cluster point, generalized quasi continuity

MSC 2010: 54C60, 54C65, 26E25

1. INTRODUCTION

The upper and lower semi continuity, their generalizations and the problem of finding a desirable selection are intensively studied in the theory of multifunctions and they play crucial role in many applications. Recall that a multifunction F is upper semi continuous/lower semi continuous (briefly usc/lsc), if $F^+(V) = \{x: F(x) \subset V\}/F^-(V) = \{x \in X: F(x) \cap V \neq \emptyset\}$ is open for any open set V . In the paper we will be interested in searching a quasi continuous selection. For this goal the upper semi continuity is too strong and the existence of a quasi continuous selection has been studied in a series of papers [1], [9], [10] for more general continuity. Perhaps the most general technic was presented in [1] based on Zorn's lemma. Namely, a compact valued upper Baire continuous multifunction (definition is below) acting from an arbitrary topological space into a regular T_1 -space has a quasi continuous selection. Compactness of the values is necessary. A multifunction from \mathbb{R} to \mathbb{R} defined by $F(x) = \{1/x\}$ for $x \neq 0$ and $F(0) = \mathbb{R}$ is upper Baire continuous (even usc) without a quasi continuous selection. The multifunction F above has closed graph which

is closely connected to c -upper semi continuity. Namely, F is c -usc (c -upper semi continuous), if $F^+(V)$ is open for any open set V with compact complement (see [5], [7], [12]). The dual notion of c -lower semi continuity, briefly c -lsc, means that $F^-(V)$ is open for any open set V with compact complement. From the continuity point of view, c -lower semi continuity has very nice behavior. Under reasonable conditions, c -lower semi continuity of F guarantees lower semi continuity of F except for a nowhere dense set [5]. On the other hand, c -upper semi continuity is rather strange. Namely, a c -upper semi continuous multifunction need not be usc/lsc at any point. An example can be found in [5]. The question if a c -upper semi continuous multifunction has a selection (submultifunction) which is quasi continuous (minimal usco) except for a nowhere dense set is the main stimulation for our investigation (see Theorem 4). Moreover, c -upper semi continuity will be replaced by c - u - \mathcal{E} -continuity (see Definition 2) which seems to be suitable for finding a selection being quasi continuous except for a nowhere dense set. It is more general than the notion of c -upper semi continuity and closedness of graph, even than the upper Baire continuity, and on the other hand it still leads to reasonable results. The notion of u - \mathcal{E} -continuity (formally derived from the upper quasi continuity) is based on a family $\mathcal{E} \subset 2^X \setminus \{\emptyset\}$ and the results obtained flexibly depend on a specification of \mathcal{E} .

2. BASIC DEFINITIONS AND PRELIMINARY RESULTS

In the sequel X, Y are topological spaces. By \overline{A} , A° we denote the closure and the interior of A , respectively. A σ -compact space Y (i.e., $Y = \bigcup_{n=1}^{\infty} C_n$, where C_n are compact) is understood to be Hausdorff. By a multifunction F we understand a subset of the cartesian product $X \times Y$ with the values $\{y \in Y : [x, y] \in F\} =: F(x)$ (it can be empty valued at some points). For a multifunction F and a set $C \subset Y$, $F \cap C$ denotes the multifunction with the values $F(x) \cap C$. By $\text{Dom}(F)$, we denote the domain of F , i.e., the set of all arguments x at which $F(x)$ is non-empty. A function f is understood as a special multifunction with values $\{f(x)\}$, $x \in \text{Dom}(f)$. For a function f , we will prefer traditional notation of its values as $f(x)$.

A multifunction can be considered as a set-valued mapping from its domain to Y denoted as $F: A \rightarrow Y$, where $A = \text{Dom}(F)$. Then the set $\{[x, y] \in A \times Y : y \in F(x)\}$ is the graph of F . In the paper, we identify the mapping with its graph.

A multifunction F is bounded on a set A if $F(A) := \bigcup\{F(x) : x \in A\}$ is a subset of some compact set, and F is locally bounded at x if there is an open set U containing x and a compact set C such that $F(U) \subset C$. If $S \subset F$, then S is called a submultifunction of F . A function f is a selection of a multifunction F , if $f(x) \in F(x)$ for all $x \in \text{Dom}(f) = \text{Dom}(F)$. If $f(x) \in F(x)$ for all $x \in A \subset \text{Dom}(f)$,

then f is called a selection of F on a set A . A multifunction F is usco, if $F(x)$ is compact and F is usc at x for all $x \in \text{Dom}(F)$.

Any non-empty system $\mathcal{E} \subset 2^X \setminus \{\emptyset\}$ will be called a cluster system. For some special cluster systems we will use a special notation. For example, $\mathcal{O}, \mathcal{B}r$ is a cluster system containing all non-empty open sets or all sets being of second category with the Baire property, respectively. For an ideal \mathbb{I} on X , put $\mathcal{E}_{\mathbb{I}} = \{(G \setminus S) \cup T\}$ where $S, T \in \mathbb{I}$ and G is open such that none of its non-empty open subsets is from \mathbb{I} .

The next two definitions introduce the notion of an \mathcal{E} -cluster point and an upper \mathcal{E} -continuity (u - \mathcal{E} -continuity), as a basic tool for investigation of the properties of multifunctions. In this form it was studied for the first time in [9], later in [10] and for the functions in [3]. Formally, upper \mathcal{E} -continuity (see Definition 2 below) is motivated by the notion of the upper quasi continuity, which is a special case of our approach.

Definition 1. A point $y \in Y$ is an \mathcal{E} -cluster point of F at a point x , if for any open sets $V \ni y$ and $U \ni x$ there is a set $E \in \mathcal{E}$, $E \subset U$ such that $F(e) \cap V \neq \emptyset$ for any $e \in E$. The set of all \mathcal{E} -cluster points of F at x is denoted by $\mathcal{E}_F(x)$. A multifunction \mathcal{E}_F with the values $\mathcal{E}_F(x)$ is called \mathcal{E} -cluster multifunction of F .

Definition 2. A multifunction F is u - \mathcal{E} -continuous at $x \in \text{Dom}(F)$ (c - u - \mathcal{E} -continuous), if for any open sets V, U ($Y \setminus V$ is compact) such that $F(x) \subset V$ and $x \in U$ there is a set $E \in \mathcal{E}$, $E \subset U \cap \text{Dom}(F)$ such that $F(e) \subset V$ for any $e \in E$. The global definition is given by the local one at any point of $\text{Dom}(F)$. Notation “ c - u - \mathcal{E} -continuity” is derived from the notion of c -upper semi continuity (see [5], [7]).

Since a function is a special case of a multifunction when upper and lower inverse images coincide, we will say that f is \mathcal{E} -continuous, c - \mathcal{E} -continuous, respectively. It is evident that if f is \mathcal{E} -continuous at x , then $f(x) \in \mathcal{E}_f(x)$. For the system \mathcal{O} we have the notion of upper quasi continuity/ c -upper quasi continuity, which is intensively studied, see a survey [13]. A few new characterizations of quasi continuity have been studied in [11]. A u - $\mathcal{B}r$ -continuous multifunction is called upper Baire continuous and this is one of the most general notions of continuity which guarantees the existence of a quasi continuous selection, see [1], [9], [10].

It can happen that some open sets need not contain a set from a given cluster system \mathcal{E} . Avoiding such case we can enlarge \mathcal{E} by some reasonable sets, for example by open ones. That is the case of the cluster system $\mathcal{E}_{\mathbb{I}}$ above, which is of our main interest. So we will deal with a cluster system $\mathcal{O} \cup \mathcal{E}_{\mathbb{I}}$ and the continuity introduced in Definition 2 can be considered as the local definition of measurability, i.e., $F^+(V) \cap U$ contains a set of the form $(G \setminus S) \cup T$ (G is open, $S, T \in \mathbb{I}$) whenever $F^+(V) \cap U$ is non-empty. For example, a compact valued multifunction F acting from a Baire

space to a metric one has the Baire property if and only if F is $u\text{-}\mathcal{E}_{\mathbb{I}}$ -continuous except for a set of first category, where \mathbb{I} is the ideal of all sets of first category (see [10]).

Now we give a definition which is a natural generalization of a minimal multifunction ([2], [6], [8]) and in this form has been studied in [11].

Definition 3. A multifunction F is \mathcal{E} -minimal at a point x , if $F(x)$ is non-empty and for any open sets U, V such that $U \ni x$ and $V \cap F(x) \neq \emptyset$ there is a set $E \subset U \cap \text{Dom}(F)$. The global definition is given by the local one at any point from $\text{Dom}(F)$. It is evident that any selection of an \mathcal{E} -minimal multifunction is \mathcal{E} -continuous.

Lemma 1 (see also [4]). *For any net $\{x_t\}$ converging to x and $y_t \in \mathcal{E}_F(x_t)$, $\mathcal{E}_F(x)$ contains all accumulation points of the net $\{y_t\}$. Consequently, \mathcal{E}_F has a closed graph and closed values.*

Proof. Let y be an accumulation point of $\{y_t\}$. Then for any open sets $V \ni y$ and $U \ni x$ there are frequently given indexes t' such that $x_{t'} \in U$ and $y_{t'} \in V \cap \mathcal{E}_F(x_{t'})$. Hence there is $E \in \mathcal{E}, E \subset U$ such that $F(e) \cap V \neq \emptyset$ for any $e \in E$. That means $y \in \mathcal{E}_F(x)$. \square

Remark 1. Since \mathcal{E}_F has a closed graph, $\mathcal{E}_F^-(K)$ is closed for any compact set K or equivalently, $\mathcal{E}_F^+(G)$ is open for any open G with compact complement in Y . Hence, \mathcal{E}_F is c -upper semi continuous. Consequently, if $\mathcal{E}_F^-(K)$ is dense in an open set G , then $G \subset \mathcal{E}_F^-(K)$, so \mathcal{E}_F is non-empty valued on G .

3. MAIN RESULTS

Lemma 2. *Let Y be Hausdorff.*

- (1) *If $F(x)$ is closed, F is c - u - \mathcal{E} -continuous at x and S is usco at x , then $F \cap S$ is c - u - \mathcal{E} -continuous at x provided $F \cap S$ is non-empty on some neighborhood of x .*
- (2) *If F is locally bounded and c - u - \mathcal{E} -continuous at x , then F is u - \mathcal{E} -continuous at x .*

Proof. 1. Let $G \supset F(x) \cap S(x)$ be open with compact complement and let W be open containing x . Then $S(x)$ is disjoint with $F(x) \setminus G$ and since $S(x)$ is compact, there are two disjoint open sets $G_1 \supset S(x)$ and $G_2 \supset F(x) \setminus G$. The complement of $G \cup G_2$ is compact, $G \cup G_2 \supset F(x)$ and by virtue of usc of S and c - u - \mathcal{E} -continuity of F , there is an open set $U \subset W$ containing x and there is $E \in \mathcal{E}, E \subset U \cap \text{Dom}(F)$ such that $F(E) \subset G \cup G_2$ and $S(U) \subset G_1$. Then $F(E) \cap S(E) \subset (G \cup G_2) \cap G_1 \subset (G \cup G_2) \cap (Y \setminus G_2) \subset G$. So $F \cap S$ is c - u - \mathcal{E} -continuous.

2. F is a locally bounded multifunction, so there is an open set U containing x and a compact set K such that $F(U) \subset K$. Let $H \supset F(x)$, let H be open and $U_0 \subset U$ open containing x . Since the complement of $(Y \setminus K) \cup H$ is compact, there is a set $E \subset U_0 \cap \text{Dom}(F)$ such that $F(E) \subset (Y \setminus K) \cup H$. So $F(E) = F(E) \cap ((Y \setminus K) \cup H) = F(E) \cap H$, which means $F(E) \subset H$. \square

Theorem 1. *Let Y be Hausdorff and F compact valued (it can be empty valued at some points) c - u - \mathcal{E} -continuous. Then F has a c - \mathcal{E} -continuous selection.*

Proof. Let \mathcal{M} be the family of all c - u - \mathcal{E} -continuous non-empty compact valued submultifunctions of F which is partially ordered by inclusion. It is non-empty, since $F \in \mathcal{M}$. For any linearly ordered subfamily \mathcal{M}_0 , a multifunction $M_0(x) := \bigcap \{M(x) : M \in \mathcal{M}_0\}$ is a non-empty compact valued submultifunction of F , and for any open sets $V \supset M_0(x)$, $Y \setminus V$ compact, and U containing x there is $M \in \mathcal{M}_0$ such that $M(x) \subset V$. By the c - u - \mathcal{E} -continuity of M there is a set $E \in \mathcal{E}$, $E \subset \text{Dom}(M) \cap U \cap M^+(V)$, hence for any $e \in E$ we have $M_0(e) \subset M(e) \subset V$. That means M_0 is c - u - \mathcal{E} -continuous and \mathcal{M} has a minimal element M_m with respect to inclusion. Now we will prove that M_m is \mathcal{E} -minimal with respect to co-compact topology on Y given by all open sets with compact complement. If not at $a \in \text{Dom}(M_m)$, there is an open set V intersecting $M_m(a)$, $Y \setminus V$ compact and an open set U containing a such that for any $E \subset U \cap \text{Dom}(M_m)$ from \mathcal{E} there is a point $e \in E$ such that $M_m(e)$ is not a subset of V . Since M_m is c - u - \mathcal{E} -continuous, hence for all $u \in U \cap \text{Dom}(M_m)$, $M_m(u)$ is not a subset of V . Define a multifunction N as $N(x) := M_m(x)$ if $x \in \text{Dom}(M_m) \setminus U$ and $N(x) := M_m(x) \cap (Y \setminus V)$ if $x \in U \cap \text{Dom}(M_m)$. Then N is a non-empty compact valued submultifunction of F . We will show that N is c - u - \mathcal{E} -continuous. If $x \in \text{Dom}(M_m) \setminus U$ there is nothing to prove. Let $x \in U \cap \text{Dom}(M_m)$, $N(x) \subset W$, let $Y \setminus W$ be compact, $x \in H \subset U$ and H, W be open. Then $M_m(x) \subset V \cup W$ and by the c - u - \mathcal{E} -continuity of M_m there is a set $E \in \mathcal{E}$, $E \subset H \cap \text{Dom}(M_m)$ such that $M_m(e) \subset V \cup W$ for any $e \in E$. That means $N(e) \subset W$ for any $e \in E$. Hence $N \in \mathcal{M}$ and $N(a)$ is a proper subset of $M_m(a)$, a contradiction with the minimality of M_m . Finally, since M_m is \mathcal{E} -minimal with respect to the co-compact topology, any selection of M_m is c - \mathcal{E} -continuous. \square

Remark 2. In a similar way we can prove the next result: If Y is Hausdorff and F is compact valued u - \mathcal{E} -continuous, then F has an \mathcal{E} -continuous selection. For $\mathcal{E} = \mathcal{B}r$ it was proved in [1].

Definition 4. A multifunction is partially \mathcal{E} -bounded if for any non-empty open set G there is a set $E \in \mathcal{E}$, $E \subset G$ and a compact set C such that $F(e) \cap C \neq \emptyset$ for any $e \in E$. Hence, a multifunction $F \cap C$ is bounded on E .

Theorem 2. *Let Y be Hausdorff, $\mathcal{E} \subset \mathcal{O} \cup \mathcal{E}_1$ and let F defined on X (i.e., $X = \text{Dom}(F)$) be closed valued and c - u - \mathcal{E} -continuous. Then F is partially \mathcal{E} -bounded if and only if F has a selection which is both locally bounded and \mathcal{E} -continuous except for a nowhere dense set.*

Proof. \Rightarrow We will prove that for any non-empty open set G there is a non-empty open set $G_0 \subset G$ and a compact set C such that $F \cap C$ is non-empty valued on G_0 .

By assumption, there are a set $E \in \mathcal{E}$, $E \subset G$ and a compact set C such that

$$(*) \quad F(e) \cap C \neq \emptyset \text{ for any } e \in E.$$

There are two possibilities. Either the set E is open ($E \in \mathcal{O}$) or $E = (G_0 \setminus I) \cup J$ ($E \in \mathcal{E}_1$), where G is open and $I, J \in \mathbb{I}$ and no non-empty open subset of G_0 is from \mathbb{I} . First, if E is open, we can put $G_0 = E$. Secondly, if $E = (G_0 \setminus I) \cup J$, we will show that $F(x) \cap C \neq \emptyset$ for any $x \in G_0$. If $F(x) \cap C = \emptyset$ for some $x \in G_0$, then by the c - u - \mathcal{E} -continuity there is $E' \subset G_0 \cap \text{Dom}(F)$, $E' \in \mathcal{E}$, such that $F(E') \subset Y \setminus C$. If E' is open, then E' is not from \mathbb{I} , so $\emptyset \neq E' \setminus I \subset G_0 \setminus I \subset E$, a contradiction with (*). If $E' = (G' \setminus I') \cup J' \in \mathcal{E}_1$, then E' is not from \mathbb{I} either, and $G' \cap G_0$ is non-empty (otherwise, $E' = E' \cap G_0 \subset (G' \cap G_0) \cup (J' \cap G_0) = J' \cap G_0 \in \mathbb{I}$, a contradiction), so there is a point $a \in G' \cap G_0 \setminus (I' \cup I) \subset E' \cap E$. Hence $F(a) \subset Y \setminus C$ and $F(a) \cap C \neq \emptyset$ (see (*)), a contradiction. That means that in both cases $F \cap C$ is a multifunction which is non-empty valued on G_0 .

By Lemma 2 (1), $F \cap C$ is non-empty compact valued and c - u - \mathcal{E} -continuous on G_0 and by Theorem 1, $F \cap C$ has a c - \mathcal{E} -continuous selection f_{G_0} on G_0 . Again, f_{G_0} is bounded, so it is \mathcal{E} -continuous by Lemma 2 (2).

We have proved for any non-empty open set G there is a non-empty open set $G_0 \subset G$ such that F has a selection that is both bounded and \mathcal{E} -continuous on G_0 .

Using Zorn's lemma, we can prove the existence of an open set H and a function $f: H \rightarrow Y$ such that f is both locally bounded and \mathcal{E} -continuous and $X \setminus H$ is nowhere dense. So, a function $g: X \rightarrow Y$ such that $g = f$ on H and $g(x) \in F(x)$ for $x \in X \setminus H$ is a desirable selection.

The converse implication is obvious. □

4. APPLICATIONS

Global \mathcal{E} -continuity on an open set has a very interesting feature. For some cluster systems, global \mathcal{E} -continuity of the functions implies quasi continuity. It is the case when Y is regular and $\mathcal{E} \subset \mathcal{O} \cup \mathcal{E}_\perp$ (see the next theorem or Corollary 1). But in multifunction setting the two notions are different as we can see from the multifunction F defined as $F(x) = \langle 0, 1 \rangle$, if x is rational and $F(x) = \{0\}$ otherwise. It is $u\text{-}\mathcal{B}r$ -continuous but not upper quasi continuous. This is a nice methodological feature of the upper Baire continuity, when a more general continuity of a multifunction guarantees a stronger continuity of a selection, see [1], [9], [10].

Theorem 3. *Suppose that the interior of $\text{Dom}(\mathcal{E}_f)$ is non-empty, where f is an arbitrary function. If Y is a regular topological space, then \mathcal{E}_f is \mathcal{O} -minimal on the interior of $\text{Dom}(\mathcal{E}_f)$ provided $\mathcal{E} \subset \mathcal{O} \cup \mathcal{E}_\perp$.*

Proof. Recall that no $E \in \mathcal{E}_\perp$ is from \perp . If not at $x \in (\text{Dom}(\mathcal{E}_f))^\circ$, there are the open sets $U \ni x$, V and a set $A \subset U \subset (\text{Dom}(\mathcal{E}_f))^\circ$ dense in U such that $\mathcal{E}_f(x) \cap V \neq \emptyset$ and $\mathcal{E}_f(a) \cap (Y \setminus \overline{V}) \neq \emptyset$ for any $a \in A$. Let $y \in \mathcal{E}_f(x) \cap V$. Then there is a set $E \in \mathcal{E}$, $E \subset U$ such that $f(E) \subset V$.

First, suppose that the set E is of the form $E = (G \setminus S) \cup T \in \mathcal{E}_\perp$, where G is open and $S, T \in \perp$. Then the intersection $G \cap U \neq \emptyset$ (otherwise $E \subset (G \cap U) \cup (T \cap U) = T \cap U \in \perp$, a contradiction) so there is a point $a \in A \cap G \cap U$ such that $\mathcal{E}_f(a) \cap (Y \setminus \overline{V}) \neq \emptyset$. Pick up $z \in \mathcal{E}_f(a) \cap (Y \setminus \overline{V})$. Then there is a set $E_0 \in \mathcal{E}$, $E_0 \subset G \cap U$ such that $f(E_0) \subset Y \setminus \overline{V}$ and E_0 is of the form $E_0 = (G_0 \setminus S_0) \cup T_0 \in \mathcal{E}_\perp$, where G_0 is open and $S_0, T_0 \in \perp$ or $E_0 \in \mathcal{O}$. In the first case, the intersection $G \cap U \cap G_0 \neq \emptyset$ (otherwise $E_0 = G \cap U \cap ((G_0 \setminus S_0) \cup T_0) \subset (G \cap U \cap G_0) \cup (G \cap U \cap T_0) = G \cap U \cap T_0 \in \perp$, a contradiction), hence $G \cap U \cap G_0 \setminus (S \cup S_0)$ is not from \perp and there is a point $e \in G \cap U \cap G_0 \setminus (S \cup S_0) \subset E$, so $f(e) \in V$. On the other hand, $e \in E_0$, so $f(e) \in Y \setminus \overline{V}$, a contradiction. In the second case, when $E_0 \in \mathcal{O}$, $E_0 = E_0 \cap G \cap U$ is a non-empty open subset of G , so E_0 is not from \perp and there is a point $e \in E_0 \setminus S$ for which $f(e) \in Y \setminus \overline{V}$. Since $E_0 \subset G$, we have $e \in E$ and $f(e) \in V$, a contradiction.

Secondly suppose, the set E is open subset of U . Then there is a point $a \in A \cap E$ such that $\mathcal{E}_f(a) \cap (Y \setminus \overline{V}) \neq \emptyset$. Pick up $z \in \mathcal{E}_f(a) \cap (Y \setminus \overline{V})$. Then there is a set $E_0 \in \mathcal{E}$, $E_0 \subset E$ such that $f(E_0) \subset Y \setminus \overline{V}$ but $f(E) \subset V$, which is a contradiction. \square

Corollary 1. *Let Y be regular and $\mathcal{E} \subset \mathcal{O} \cup \mathcal{E}_\perp$. If a function f is \mathcal{E} -continuous on an open set H , then f is quasi continuous on H .*

Proof. Since f is \mathcal{E} -continuous, $f(h) \in \mathcal{E}_f(h)$ for any $h \in H$ and by the above theorem, \mathcal{E}_f is \mathcal{O} -minimal on H . Then any of its selections is quasi continuous, hence f is also quasi continuous. \square

Corollary 2. *Let Y be T_1 -regular and $\mathcal{E} \subset \mathcal{O} \cup \mathcal{E}_1$. If a closed valued multifunction F defined on X is c - u - \mathcal{E} -continuous and partially \mathcal{E} -bounded, then F has a selection which is quasi continuous except for a nowhere dense set.*

Theorem 4. *Let Y be a σ -compact regular space and let F be a closed valued multifunction with $\text{Dom}(F) = X$. Each of the following conditions ensures the existence of a selection of F which is both locally bounded and quasi continuous except for a nowhere dense set.*

- (1) F is c -upper Baire continuous.
- (2) X is Baire and F is c -upper quasi continuous.

Moreover, if X is Baire and F is c -usc, then F has a non-empty valued submultifunction which is both locally bounded and \mathcal{O} -minimal usco except for a nowhere dense set.

Proof. (1) Since F is c -upper Baire continuous, X is Baire. We will show $F^+(V)$ has the Baire property for any open set V with compact complement. If not, there is an open set U such that both the sets $X_0 := F^+(V)$ and $X \setminus X_0$ are of second category at any point from U . Let $x \in X_0 \cap U$. By the c -upper Baire continuity there is $E \in \mathcal{B}r$, $E \subset U$ such that $F(E) \subset V$. Since E is of second category with the Baire property, $E = (G \setminus I) \cup J$ for some G open and I, J of first category and $G \cap U \neq \emptyset$ (otherwise $E = ((G \setminus I) \cup J) \cap U = ((G \setminus I) \cap U) \cup (J \cap U) = J \cap U$ is of first category). The set $X \setminus X_0$ is of second category at any point from U , so $((G \cap U \cap (X \setminus X_0)) \setminus I$ is of second category, that means there is a point $e \in ((G \cap U \cap (X \setminus X_0)) \setminus I \subset E$. So $F(e) \not\subset V$, a contradiction with $F(E) \subset V$.

Let $Y = \bigcup_{k \in \mathbb{N}} C_k$, let C_k be compact and G non-empty open. Since $G \subset \bigcup_{k \in \mathbb{N}} F^-(C_k)$, there is m such that $F^-(C_m) = X \setminus F^+(Y \setminus C_m)$ has the Baire property and is of second category, so F is partially $\mathcal{B}r$ -bounded. By Theorem 2 and Corollary 1, F has a desirable selection.

(2) Since X is Baire, F is also c -upper Baire continuous and the proof follows from item (1).

Moreover, suppose X is Baire and F is c -usc. Hence F is c -upper Baire continuous and by item (2), F has a selection f which is quasi continuous and locally bounded on an open dense set H . Put $F_0 = \mathcal{B}r_f$. That means $f(h) \in F(h) \cap F_0(h)$ for all $h \in H$. It is clear that F_0 is both locally bounded and with closed graph, so it is usco on H and F_0 is \mathcal{O} -minimal on H . Hence for any $x \in H$ there is an open U_0 containing x such that $F_0(U_0) \subset C$, where C is compact. We will show that $F_0(x) \subset F(x)$. If not, there is a point $y \in F_0(x) \setminus F(x)$ and there are two disjoint open sets $V \supset F(x)$ and $W \ni y$ such that $\overline{W} \cap V = \emptyset$ (we use regularity of Y and the closed values of F).

A set $C \cap \overline{W}$ is non-empty compact and disjoint with $F(x)$ and since F is c -usc, there is an open set U containing x , $U \subset U_0$ such that $F(U) \subset Y \setminus (C \cap \overline{W})$. Since F_0 is minimal, there is a non-empty open set $H_0 \subset U$ such that $F_0(H_0) \subset W$. Hence $F_0(H_0) \subset C \cap \overline{W}$. So F and F_0 have disjoint values on H_0 , a contradiction with the fact that $f(h) \in F(h) \cap F_0(h)$ for all $h \in H$. Defining G as $G(x) = F_0(x)$ if $x \in H$ and $G(x)$ an arbitrary non-empty subset of $F(x)$ if $x \in X \setminus H$ we obtain a desirable submultifunction of F . \square

As we have mentioned, by [5] there is a multifunction F which is c -usc but not usc/lsc at any point. The question is, if there is some reasonable “small” or “big” submultifunction of F . A “small” variant is given in Theorem 4 by proving the existence of a submultifunction which is both \mathcal{O} -minimal usco and locally bounded except for a nowhere dense set. The open problem is a “big” variant, namely, to describe a “maximal” usco (usc, lsc) submultifunction of F . More generally, for a c -upper Baire continuous closed (compact) valued multifunction to describe its maximal submultifunction which is lower/upper quasi continuous or usco except for a nowhere dense set.

References

- [1] *J. Cao and W.B. Moors*: Quasicontinuous selections of upper continuous set-valued mappings. *Real Anal. Exchange* 31 (2005–2006), 63–71.
- [2] *L. Drewnowski and I. Labuda*: On minimal upper semicontinuous compact valued maps. *Rocky Mount. J. Math.* 20 (1990), 737–752.
- [3] *D. K. Ganguly and Mitra Chandrani*: Some remarks on B^* -continuous functions. *Anal. St. Univ. Ali. Cuza, Isai* 46 (2000), 331–336.
- [4] *D. K. Ganguly and P. Mallick*: On generalized continuous multifunctions and their selections. *Real Anal. Exchange* 33 (2007/2008), 449–456.
- [5] *L. Holá, V. Baláž and T. Neubrunn*: Remarks on c -continuous multifunctions. *Acta Math. Univ. Comeniana L-LI* (1987), 51–59.
- [6] *L. Holá and D. Holý*: Minimal usco maps, densely continuous forms and upper semicontinuous functions. *Rocky Mount. J. Math.* 39 (2009), 545–562.
- [7] *D. Holý and L. Matejíčka*: C -upper semicontinuous and C^* -upper semicontinuous multifunctions. *Tatra Mount. Math. Publ.* 34 (2006), 159–165.
- [8] *D. Holý and L. Matejíčka*: Quasicontinuous functions, minimal USCO maps and topology of pointwise convergence. *Math. Slovaca* (accepted).
- [9] *M. Matejdes*: Sur les sélectors des multifonctions. *Math. Slovaca* 37 (1987), 111–124.
- [10] *M. Matejdes*: Continuity of multifunctions. *Real Anal. Exchange* 19 (1993–94), 394–413.
- [11] *M. Matejdes*: Minimality of multifunctions. *Real Anal. Exchange* 32 (2007), 519–526.
- [12] *T. Neubrunn*: C -continuity and closed graphs. *Čas. pro přest. mat.* 110 (1985), 172–178.
- [13] *T. Neubrunn*: Quasi continuity. *Real Anal. Exchange* 14 (1988–89), 259–306.

Author’s address: Milan Matejdes, Department of Mathematics, Faculty of Applied Informatics, Tomas Bata University in Zlín, Nad Stráněmi 4511, 760 05 Zlín, Czech Republic, e-mail: matejdes@fai.utb.cz.