

Fidel Casarrubias-Segura; Fernando Hernández-Hernández; Angel  
Tamariz-Mascarúa

Martin's Axiom and  $\omega$ -resolvability of Baire spaces

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 51 (2010), No. 3, 519--540

Persistent URL: <http://dml.cz/dmlcz/140728>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2010

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## Martin's Axiom and $\omega$ -resolvability of Baire spaces

FIDEL CASARRUBIAS-SEGURA, FERNANDO  
HERNÁNDEZ-HERNÁNDEZ, ÁNGEL TAMARIZ-MASCARÚA

*Abstract.* We prove that, assuming MA, every crowded  $T_0$  space  $X$  is  $\omega$ -resolvable if it satisfies one of the following properties: (1) it contains a  $\pi$ -network of cardinality  $< \mathfrak{c}$  constituted by infinite sets, (2)  $\chi(X) < \mathfrak{c}$ , (3)  $X$  is a  $T_2$  Baire space and  $c(X) \leq \aleph_0$  and (4)  $X$  is a  $T_1$  Baire space and has a network  $\mathcal{N}$  with cardinality  $< \mathfrak{c}$  and such that the collection of the finite elements in it constitutes a  $\sigma$ -locally finite family.

Furthermore, we prove that the existence of a  $T_1$  Baire irresolvable space is equivalent to the existence of a  $T_1$  Baire  $\omega$ -irresolvable space, and each of these statements is equivalent to the existence of a  $T_1$  almost- $\omega$ -irresolvable space.

Finally, we prove that the minimum cardinality of a  $\pi$ -network with infinite elements of a space  $\text{Seq}(u_t)$  is strictly greater than  $\aleph_0$ .

*Keywords:* Martin's Axiom, Baire spaces, resolvable spaces,  $\omega$ -resolvable spaces, almost resolvable spaces, almost- $\omega$ -resolvable spaces, infinite  $\pi$ -network

*Classification:* Primary 54E52, 54A35; Secondary 54D10, 54A10

### 1. Introduction

Every space in this article is  $T_0$  and crowded (that is, without isolated points) and so it is infinite. A space  $X$  is *resolvable* if it contains two dense disjoint subsets. A space which is not resolvable is called *irresolvable*. Resolvable and irresolvable spaces were studied extensively first by Hewitt [14]. Later, El'kin and Malykhin published a number of papers on these subjects and their connections with various topological problems. One of the problems considered by Malykhin in [22] refers to the existence of irresolvable spaces satisfying the Baire Category Theorem. Kunen, Symański and Tall in [19] afterwards proved that there is such a space if and only if there is a space  $X$  on which every real-valued function is continuous at some point. (The question about the existence of a  $\pi$ -Hausdorff-space on which every real-valued function is continuous at some point was posed by M. Katětov in [16].) They also proved (see [18] as well):

1. if we assume  $V = L$ , there is no Baire irresolvable space,
2. the conditions “there is a measurable cardinal” and “there is a Baire irresolvable space” are equiconsistent.

Bolstein introduced in [5] the spaces  $X$  in which it is possible to define a real-valued function  $f$  with countable range and such that  $f$  is discontinuous at every

point of  $X$  (he called these spaces *almost resolvable*), and proved that every resolvable space satisfies this condition. It was proved in [12] that  $X$  is almost resolvable iff there is a function  $f : X \rightarrow \mathbb{R}$  such that  $f$  is discontinuous at every point of  $X$ . *Almost- $\omega$ -resolvable* spaces were introduced in [26]; these are spaces in which it is possible to define a real-valued function  $f$  with countable range, and such that  $r \circ f$  is discontinuous in every point of  $X$ , for every real-valued finite-to-one function  $r$ . It was proved in that article that for a Tychonoff space  $X$ , the space of real continuous functions with the box topology,  $C_{\square}(X)$ , is discrete if and only if  $X$  is almost- $\omega$ -resolvable. It was also proved that the existence of a measurable cardinal is equiconsistent with the existence of a Tychonoff space without isolated points which is not almost- $\omega$ -resolvable, and that, on the contrary, if  $V = L$  then every crowded space is almost- $\omega$ -resolvable. Later, it was pointed out in [2, Corollary 5.4] that a Baire space is resolvable if and only if it is almost resolvable; so,

**1.1 Theorem.** *A Baire almost- $\omega$ -resolvable space is resolvable.*

It is unknown if every Baire almost- $\omega$ -resolvable space is 3-resolvable. With respect to this problem we have the following theorems.

**1.2 Theorem** ([24]). *Gödel's axiom of constructibility,  $V = L$ , implies that every Baire space is  $\omega$ -resolvable.*

**1.3 Theorem** ([2]). *Every  $T_1$  Baire space such that each of its dense subsets is almost- $\omega$ -resolvable is  $\omega$ -resolvable.*

These last two results transform our problem to that of finding subclasses of Baire spaces such that each of its crowded dense subsets is almost- $\omega$ -resolvable, assuming axioms consistent with ZFC which contrast with  $V = L$ . Of course, a classic axiom with these characteristics is  $\text{MA} + \neg\text{CH}$ . This bet is strengthened by the following result due to V.I. Malykhin ([23, Theorem 1.2]):

**1.4 Theorem** [ $\text{MA}_{\text{countable}}$ ]. *Let a topology on a countable set  $X$  have a  $\pi$ -network of cardinality less than  $\mathfrak{c}$  consisting of infinite subsets. Then this topology is  $\omega$ -resolvable.*

It was proved in [2] that every space with countable tightness, every space with  $\pi$ -weight  $\leq \aleph_1$  and every  $\sigma$ -space are hereditarily almost- $\omega$ -resolvable. So, by Theorem 1.3, every  $T_1$  Baire space with either countable tightness or  $\pi$ -weight  $\leq \aleph_1$  or  $\sigma$  is  $\omega$ -resolvable.

In this article we are going to continue the study of almost- $\omega$ -resolvable and Baire resolvable spaces, and we will solve some problems related to these posed in [2]. Section 2 is devoted to establishing basic definitions and results. In Section 3 we prove that under  $\text{MA}$  every space with either  $\pi$ -weight  $< \mathfrak{c}$  or  $\chi(X) < \mathfrak{c}$  is  $\omega$ -resolvable. Furthermore, we are going to see in Section 4 that under  $\text{SH}$  every  $T_2$  Baire space with countable cellularity is  $\omega$ -resolvable. Section 5 is devoted to

analyse almost- $\omega$ -irresolvable spaces. We prove in this section that there is a  $T_1$  Baire irresolvable space iff there is a  $T_1$  Baire  $\omega$ -irresolvable space, iff there is a  $T_1$  almost- $\omega$ -irresolvable space. Finally in Section 6, we prove that the minimum cardinality of a  $\pi$ -network with infinite elements of a space  $\text{Seq}(u_t)$  is strictly greater than  $\aleph_0$ . Moreover, we propose several problems related to our matter through the article.

## 2. Basic definitions and preliminaries

A space  $X$  is *resolvable* if it is the union of two disjoint dense subsets. We say that  $X$  is *irresolvable* if it is not resolvable. For a cardinal number  $\kappa > 1$ , we say that  $X$  is  $\kappa$ -*resolvable* if  $X$  is the union of  $\kappa$  pairwise disjoint dense subsets.

The *dispersion character*  $\Delta(X)$  of a space  $X$  is the minimum of the cardinalities of non-empty open subsets of  $X$ . If  $X$  is  $\Delta(X)$ -resolvable, then we say that  $X$  is *maximally resolvable*. A space  $X$  is *hereditarily irresolvable* if every subspace of  $X$  is irresolvable. And  $X$  is *open-hereditarily irresolvable* if every open subspace of  $X$  is irresolvable.

We call a space  $(X, t)$  *maximal* if  $(X, t')$  contains at least one isolated point when  $t'$  strictly contains the topology  $t$ . And a space  $X$  is *submaximal* if every dense subset of  $X$  is open. Moreover, maximal spaces are submaximal, and these are hereditarily irresolvable spaces, which in turn are open-hereditarily irresolvable.

It is possible to prove that a space  $X$  is *almost resolvable* if and only if  $X$  is the union of a countable collection of subsets each of them with an empty interior [5].

It was proved in [26] that the following formulation can be given as a definition of almost- $\omega$ -resolvable space: A space  $X$  is called *almost- $\omega$ -resolvable* if  $X$  is the union of a countable collection  $\{X_n : n < \omega\}$  of subsets in such a way that for each  $m < \omega$ ,  $\text{int}(\bigcup_{i \leq m} X_i) = \emptyset$ . In particular, every almost- $\omega$ -resolvable space is almost resolvable, every  $\omega$ -resolvable space is almost- $\omega$ -resolvable, every almost resolvable space is infinite, and every  $T_1$  separable space is almost- $\omega$ -resolvable.

We are going to say that a space  $X$  is *hereditarily almost- $\omega$ -resolvable* if each crowded subspace of  $X$  is almost- $\omega$ -resolvable, and  $X$  is *dense-hereditarily almost- $\omega$ -resolvable* if each crowded dense subspace of  $X$  is almost- $\omega$ -resolvable.

Let  $X$  be a  $\kappa$ -resolvable (resp., almost-resolvable, almost- $\omega$ -resolvable) space. A  $\kappa$ -*resolution* (resp., an *almost resolution*, an *almost- $\omega$ -resolution*) for  $X$  is a partition  $\{V_\alpha : \alpha < \kappa\}$  (resp., a partition  $\{V_n : n < \omega\}$ ) of  $X$  such that each  $V_\alpha$  is a dense subset of  $X$  (resp.,  $\text{int}(V_n) = \emptyset$  for every  $n < \omega$ ,  $\text{int}(\bigcup_{i=0}^n V_i) = \emptyset$  for every  $n < \omega$ ).

Finally, a space  $X$  is *almost- $\omega$ -irresolvable* (resp.,  $\kappa$ -irresolvable) if  $X$  is not almost- $\omega$ -resolvable (resp.,  $X$  is not  $\kappa$ -resolvable). The hereditary version of almost- $\omega$ -irresolvability or  $\kappa$ -irresolvability is that which states that every crowded subspace of  $X$  is not almost- $\omega$ -resolvable, and, respectively, is not  $\kappa$ -resolvable.

**2.1 Example.** There are non- $T_0$  topological spaces which are almost resolvable but not almost- $\omega$ -resolvable. In fact, let  $X$  be an infinite set and  $x, y \in X$  with  $x \neq y$ . We define a collection  $\mathcal{T}$  of subsets of  $X$  as follows:  $A \in \mathcal{T}$  if either  $A$  is the empty set or  $x, y \in A$ . The family  $\mathcal{T}$  is a topology in  $X$  and  $(X, \mathcal{T})$  satisfies the required conditions.

**2.2 Example.** It was proved in Theorem 4.4 of [19] that the existence of an  $\omega_1$ -complete ideal  $\mathcal{I}$  over  $\omega_1$  which has a dense set of size  $\omega_1$  implies the existence of a  $T_2$  Baire strongly irresolvable topology  $\mathcal{T}$  on  $\omega_1$ . On the other hand, it was observed in [26, Corollary 4.9] that every Baire irresolvable space is not almost resolvable. Therefore,  $(\omega_1, \mathcal{T})$  is not almost resolvable.

**2.3 Example.** If there is a measurable cardinal  $\kappa$ , then there is a resolvable Baire space  $X$  which is not almost- $\omega$ -resolvable and  $\Delta(X) = \kappa$ . Indeed, let  $\kappa$  be a non-countable Ulam-measurable cardinal, and let  $p$  be a free ultrafilter on  $\kappa$   $\omega_1$ -complete. Let  $X = \kappa \cup \{p\}$ . We define a topology  $t$  for  $X$  as follows:  $A \in t \setminus \{\emptyset\}$  if and only if  $p \in A$  and  $A \cap \kappa \in p$ . This space is a Baire resolvable non-almost- $\omega$ -resolvable space with  $\Delta(X) = \alpha$ . Now, let  $\mathcal{T}$  be equal to  $\{A \subseteq X : A \cap \kappa \in p\}$ ;  $\mathcal{T}$  is a topology in  $X$  too, and  $(X, \mathcal{T})$  is  $T_1$  submaximal, Baire with  $\Delta(X) = \alpha$ , but it is not almost resolvable.

Related to the last examples we have:

**2.4 Question.** *Is there a  $T_2$  resolvable Baire space which is not almost- $\omega$ -resolvable?*

**2.5 Examples.** In ZFC, there are almost- $\omega$ -resolvable spaces which are not resolvable. Indeed, the union of Tychonoff crowded topologies in  $\mathbb{Q}$  generates a Tychonoff crowded topology. By Zorn's Lemma, we can consider a maximal Tychonoff topology  $\mathcal{T}$  in  $\mathbb{Q}$ . The space  $(\mathbb{Q}, \mathcal{T})$  is countable (so, almost- $\omega$ -resolvable) hereditarily irresolvable ([14, Theorems 15 and 26], [8, Example 3.3]).  $(\mathbb{Q}, \mathcal{T})$  is Tychonoff.

In [1], the authors construct by transfinite recursion a "concrete" (in the sense that we can say how its open sets look) example of a countable dense subset  $X$  of the space  $2^{\mathfrak{c}}$  which is irresolvable. Since  $X$  is countable, it is almost- $\omega$ -resolvable.

**2.6 Example.** For every cardinal number  $\kappa$ , there exists a Tychonoff space  $X$  which is almost- $\omega$ -resolvable, hereditarily irresolvable and  $\Delta(X) \geq \kappa$ . In fact, let  $\lambda$  be a cardinal number such that  $\kappa \leq \lambda$  and  $\text{cof}(\lambda) = \aleph_0$ . Let  $H, G$  and  $\tau$  be the topological groups and the topology in  $G$ , respectively, defined in [11, pp. 33 and 34], with  $|H| = \lambda$ . L. Feng proved there that  $(H, \tau|_H)$  is an irresolvable card-homogeneous (every open subset of  $H$  has the same cardinality as  $H$ ) Tychonoff space, and each subset  $S \subseteq H$  with cardinality strictly less than  $\lambda$  is a nowhere dense subset of  $H$ . Let  $(\lambda_n)_{n < \omega}$  be a sequence of cardinal numbers such that  $\lambda_n < \lambda_{n+1}$  for every  $n < \omega$  and  $\sup\{\lambda_n : n < \omega\} = \lambda$ . We take subsets  $H_n$  of  $H$  with the properties  $H_n \subseteq H_{n+1}$  and  $|H_n| = \lambda_n$  for each  $n < \omega$ , and

$H = \bigcup_{n < \omega} H_n$ . We have that each  $H_n$  is nowhere dense in  $H$ ; so  $\{H_n : n < \omega\}$  is an almost- $\omega$ -resolvable sequence on  $H$ . That is,  $H$  is almost- $\omega$ -resolvable. By the Hewitt Decomposition Theorem (see [14, Theorem 28]), there exists a non-empty open subset  $U$  of  $H$  which is hereditarily irresolvable. Besides,  $\Delta(U) = \Delta(H) \geq \kappa$  and  $U$  is almost- $\omega$ -resolvable.

**2.7 Examples.** The first example of a Hausdorff maximal group was constructed by Malykhin in [21] under Martin's Axiom. Malykhin also constructed in [23], in the BK model  $M_{\omega_1}$  (see [3]) a topological group topology  $\mathcal{T}'$  in the infinite countable Boolean group  $\Omega$  of all finite subsets of  $\omega$  with symmetric difference as the group operation, such that  $(\Omega, \mathcal{T}')$  is  $T_2$ , irresolvable and its weight is  $\omega_1$  (compare with Corollary 3.6 below). Moreover, in  $M_{\omega_1}$ ,  $\omega_1 < \mathfrak{c}$ . Moreover, he constructed in  $M_{\omega_1}$  a countable irresolvable dense subset in  $2^{\omega_1}$ . This space has of course weight  $\omega_1$ .

On the other hand, the class of resolvable spaces includes spaces with well known properties:

- 2.8 Theorem.** (1) *If  $X$  has a  $\pi$ -network  $\mathcal{N}$  such that  $|\mathcal{N}| \leq \Delta(X)$  and each  $N \in \mathcal{N}$  satisfies  $|N| \geq \Delta(X)$ , then  $X$  is maximally resolvable [9].*
- (2) *Hausdorff  $k$ -spaces are maximally resolvable [25].*
  - (3) *Countably compact regular  $T_1$  spaces are  $\omega$ -resolvable [7].*
  - (4) *Arc connected spaces are  $\omega$ -resolvable.*
  - (5) *Every biradial space is maximally resolvable [29].*
  - (6) *Every homogeneous space containing a non-trivial convergent sequence is  $\omega$ -resolvable [28].*
  - (7) *If  $G$  is an uncountable  $\aleph_0$ -bounded topological group, then  $G$  is  $\aleph_1$ -resolvable [29].*
  - (8)  *$T_1$  Baire spaces with either countable tightness or  $\pi$ -weight  $\leq \aleph_1$  are  $\omega$ -resolvable [2].*

The following basic results will be very helpful (see, for example, [6]).

- 2.9 Propositions.** (1) *If  $X$  is the union of  $\kappa$ -resolvable (resp., almost-resolvable, almost- $\omega$ -resolvable) subspaces, then  $X$  has the same property.*
- (2) *Every open and every regular closed subset of a  $\kappa$ -resolvable (resp., almost resolvable, almost- $\omega$ -resolvable) space shares this property.*
  - (3) *Let  $X$  be a space which contains a dense subset which is  $\kappa$ -resolvable (resp., almost resolvable, almost- $\omega$ -resolvable). Then,  $X$  satisfies this property too.*

The following results are easy to prove and are well known.

**2.10 Proposition.** *Let  $Y$  be a  $\kappa$ -resolvable (resp., almost-resolvable, almost- $\omega$ -resolvable) space. If  $f : X \rightarrow Y$  is a continuous and onto function, and for each*

open subset  $A$  of  $X$  the interior of  $f[A]$  is not empty, then  $X$  is  $\kappa$ -resolvable (resp., almost-resolvable, almost- $\omega$ -resolvable).

**2.11 Proposition.** *Let  $f : X \rightarrow Y$  be continuous and bijective. If  $X$  is  $\kappa$ -resolvable (resp., almost-resolvable, almost- $\omega$ -resolvable), so is  $Y$ .*

**2.12 Proposition.** (1) *If  $X$  is  $\kappa$ -resolvable (resp., almost resolvable, almost- $\omega$ -resolvable) and  $Y$  is an arbitrary topological space, then  $X \times Y$  is  $\kappa$ -resolvable (resp., almost resolvable, almost- $\omega$ -resolvable).*

(2) [2] *If  $X$  and  $Y$  are almost resolvable, then  $X \times Y$  is resolvable.*

(3) (O. Masaveu) *If  $X$  is the product space  $\prod_{\alpha < \kappa} X_\alpha$  where  $\kappa \geq \omega$  and each  $X_\alpha$  has more than one point, then  $X$  is  $2^\kappa$ -resolvable.*

The following lemmas will be useful later.

**2.13 Proposition.** *If  $X$  is a crowded space such that  $\text{cof}(|X|) = \aleph_0$  and every open subset of  $X$  has cardinality  $|X|$ , then  $X$  is almost- $\omega$ -resolvable.*

**2.14 Proposition.** *If  $X$  has tightness equal to  $\kappa$ , then each point  $x \in X$  is contained in a crowded subset of  $X$  of cardinality  $\leq \kappa$ .*

PROOF: Let  $x_0 \in X$  be an arbitrary fixed point. Since  $X$  is crowded,  $x_0 \in \text{cl}_X [X \setminus \{x_0\}]$ ; so there is a subset  $F_1 \subseteq X \setminus \{x_0\}$  of cardinality  $\leq \kappa$  such that  $x_0 \in \text{cl}_X F_1$ . If  $F_0 \cup F_1$  is crowded, where  $F_0 = \{x_0\}$ , then we have finished. Otherwise, for each isolated point  $x$  of  $F_0 \cup F_1$ , there is a subset  $F_x^2 \subseteq X \setminus (\{x_0\} \cup F_1)$  of cardinality  $\leq \kappa$  such that  $x \in \text{cl}_X F_x^2$ . Let  $F_2 = \bigcup_{x \in G_1} F_x^2$  where  $G_1$  is the set of isolated points of  $F_0 \cup F_1$ . Again, there are two possible situations: either  $F_0 \cup F_1 \cup F_2$  is a crowded subspace of cardinality  $\leq \kappa$  containing  $x_0$ , or  $G_2 = \{x \in F_2 : x \text{ is an isolated point of } F_0 \cup F_1 \cup F_2\}$  is not empty. In this last case, for each  $x \in G_2$  we take a subset  $F_x^3 \subseteq X \setminus (F_0 \cup F_1 \cup F_2)$  of cardinality  $\leq \kappa$  for which  $x \in \text{cl}_X F_x^3$ . We write  $F_3 = \bigcup_{x \in G_2} F_x^3$ . Continuing this process if necessary, we obtain either a finite sequence  $F_0, \dots, F_n$  of subsets of  $X$  such that  $x_0 \in F = \bigcup_{0 \leq i \leq n} F_i$  and  $F$  has cardinality  $\leq \kappa$  and is crowded, or we have to go further:  $x_0 \in \bar{F} = \bigcup_{n < \omega} F_n$ . In this last case too,  $F$  has cardinality  $\leq \kappa$  and is crowded.  $\square$

### 3. Martin’s Axiom, $\pi$ -netweight and $\omega$ -resolvable spaces

First, in this section we are going to present, by using Martin’s Axiom, a generalization of Theorem 1.4. As usual, if  $I$  and  $J$  are two sets,  $\text{Fn}(I, J)$  stands for the collection of the finite functions with domain contained in  $I$  and range contained in  $J$ . We define a partial order  $\leq$  in  $\text{Fn}(I, J)$  by letting  $p \leq q$  iff  $p \supseteq q$ . The partial order set  $(\text{Fn}(I, J), \leq)$  is ccc if and only if  $|J| \leq \aleph_0$  (Lemma 5.4, p. 205 in [17]).

Let  $(X, \tau)$  be a topological space. A collection  $\mathcal{N} \subseteq \mathcal{P}(X)$  is a  $\pi$ -network of  $X$  if each element  $U \in \tau \setminus \{\emptyset\}$  contains an element of  $\mathcal{N}$ .

**3.1 Definitions.** Let  $\kappa$  be an infinite cardinal.

- (1) A space  $X$  is almost- $\kappa$ -resolvable if  $X$  can be partitioned as  $X = \bigcup_{\alpha < \kappa'} X_\alpha$  where  $\omega \leq \kappa' \leq \kappa$ ,  $X_\alpha \neq \emptyset$ , and  $X_\alpha \cap X_\xi = \emptyset$  if  $\alpha \neq \xi$ , such that every non-empty open subset of  $X$  has a non-empty intersection with an infinite collection of elements in  $\{X_\alpha : \alpha < \kappa\}$ .
- (2) Let  $\mathcal{X} = \{X_\alpha : \alpha < \kappa\}$  be a partition of  $X$ . A collection  $\mathcal{N} = \{N_\xi : \xi < \tau\}$  of infinite subsets of  $\kappa$  is a  $\pi$ -network of  $\mathcal{X}$  if for each open set  $U$  of  $X$ ,  $\{\alpha < \kappa : X_\alpha \cap U \neq \emptyset\} \supseteq N_\xi$  for a  $\xi < \tau$ .
- (3) A space  $X$  is called precisely almost- $\kappa$ -resolvable if  $X$  contains a resolution with a  $\pi$ -network  $\mathcal{N}$  such that  $|\mathcal{N}| \leq \kappa$ .

The following well known result is due to K. Kuratowski.

**3.2 Lemma** (The disjoint refinement lemma). *Let  $\{A_\xi : \xi < \kappa\}$  be a collection of sets such that, for each  $\xi < \kappa$ ,  $|A_\xi| \geq \kappa$ . Then, there is a collection  $\{B_\xi : \xi < \kappa\}$  of sets satisfying:*

- (1)  $B_\xi \subseteq A_\xi$  for all  $\xi < \kappa$ ,
- (2)  $|B_\xi| = \kappa$  for all  $\xi < \kappa$ ,
- (3)  $B_\xi \cap B_\zeta = \emptyset$  for  $\xi, \zeta < \kappa$  with  $\xi \neq \zeta$ .

**3.3 Proposition.** *A space  $X$  is precisely almost- $\omega$ -resolvable if and only if  $X$  is  $\omega$ -resolvable.*

PROOF: Let  $X$  be a precisely almost- $\omega$ -resolvable space. Let  $\mathcal{X} = \{X_\xi : \xi < \tau\}$  be a precise partition of  $X$ , and  $\mathcal{M} = \{M_n : n < \omega\}$  be a  $\pi$ -network of  $\mathcal{X}$ . Because of Lemma 3.2, there are infinite and pairwise disjoint sets  $T_0, T_1, \dots, T_n, \dots$  such that  $T_i \subseteq M_i$  for all  $i < \omega$ .

For each  $n < \omega$ , we faithfully enumerate  $T_n$ :  $\{k_i^n : i < \omega\}$ . Now we define for each  $i < \omega$ ,  $D_i = \bigcup_{j < \omega} X_{k_i^j}$ . Each  $D_n$  is dense in  $X$  and  $D_i \cap D_j = \emptyset$  if  $i \neq j$ .

Moreover, if  $X$  is  $\omega$ -resolvable and  $\mathcal{D} = \{D_n : n < \omega\}$  is a collection of pairwise disjoint dense subsets of  $X$ , then  $\mathcal{D}$  is a precise partition of  $X$  and  $\mathcal{M} = \{\omega\}$  is a  $\pi$ -network of  $\mathcal{D}$ . □

When we assume Martin's Axiom, we can generalize Proposition 3.3:

**3.4 Theorem.** *Let  $\mathcal{X} = \{X_\alpha : \alpha < \tau\}$  be an almost- $\tau$ -resolvable partition of  $X$ . Let  $\mathcal{N} = \{N_\xi : \xi < \kappa\}$  be a  $\pi$ -network of  $\mathcal{X}$  such that  $\kappa < \mathfrak{c}$ . If we assume Martin's Axiom, then  $X$  is  $\omega$ -resolvable. In particular, MA implies that  $\omega$ -resolvability and almost- $\kappa$ -resolvability precise coincide when  $\kappa < \mathfrak{c}$ .*

PROOF: In this case, we put  $\mathbb{P} = (\text{Fn}(\kappa, \omega), \leq)$  where  $\leq$  is defined at the beginning of this section. For each  $k \in \omega$  and  $N \in \mathcal{N}$ , we take the set

$$D_N^k = \{p \in \mathbb{P} : \exists \xi \in N \text{ such that } p(\xi) = k\}.$$



It happens that each  $D_N^k$  is dense in  $\mathbb{P}$ . In fact, let  $q$  be an arbitrary element in  $\mathbb{P}$ . We can take  $\xi \in N \setminus \text{dom}(q)$  because  $N$  is infinite. The function  $p = q \cup \{(\xi, k)\}$  belongs to  $D_N^k$  and is less than  $q$ .

The partially ordered set  $\mathbb{P}$  is ccc and  $\mathcal{D} = \{D_N^k : k < \omega, N \in \mathcal{N}\}$  has cardinality strictly less than  $\mathfrak{c}$ . So, there exists a  $\mathcal{D}$ -generic filter  $G$  in  $\mathbb{P}$ . Take  $f = \bigcup G$ . Then  $f : \kappa \rightarrow \omega$  is onto and  $\kappa = \bigcup_{n < \omega} Y_n$  where  $Y_n = f^{-1}[\{n\}]$ .

Now, for each  $n < \omega$ , we consider the set  $X_n = \bigcup_{\alpha \in Y_n} X_\alpha$ . It is easy to prove that  $\{X_n : n < \omega\}$  is a partition of  $\bigcup_{n < \omega} X_n$ . Moreover, each  $X_n$  is a dense subset of  $X$ . Indeed, let  $n_0$  be a natural number. We are going to prove that  $X_{n_0}$  is dense. Let  $U$  be an open set of  $X$ . Because of the properties of  $\mathcal{N}$ , there is  $N_0 \in \mathcal{N}$  such that  $\{\alpha < \tau : X_\alpha \cap U \neq \emptyset\} \supseteq N_0$ . We take  $p \in D_{N_0}^{n_0} \cap G$ . It happens that there is a  $\xi \in N_0$  such that  $p(\xi) = n_0$ . Hence,  $f(\xi) = n_0$ . This means that  $\xi \in f^{-1}[\{n_0\}] = Y_{n_0}$ . By definition,  $X_\xi$  must have a non-empty intersection with  $U$ , and therefore  $U \cap X_{n_0} = U \cap \bigcup_{\alpha \in Y_{n_0}} X_\alpha \neq \emptyset$ . □

Assume that  $\{x_\xi : \xi < \tau\}$  is a faithful enumeration of a space  $X$ . If  $X$  possesses a  $\pi$ -network  $\mathcal{N}$  with infinite elements, the collection  $\{M_N : N \in \mathcal{N}\}$  where  $M_N = \{\xi < \tau : x_\xi \in N\}$ , is a  $\pi$ -network of the partition  $\{\{x_\xi\} : \xi < \tau\}$ . So the following result is a corollary of Theorem 3.4.

**3.5 Theorem.** *Let  $X$  be a crowded topological space with a  $\pi$ -network  $\mathcal{N}$  with cardinality  $\kappa < \mathfrak{c}$  and such that each element in  $\mathcal{N}$  is infinite. If we assume Martin’s Axiom, then  $X$  is an  $\omega$ -resolvable space.*

Recall that every biradial space is maximally resolvable. Moreover, every space with  $\pi w(X) \leq \Delta(X)$  is maximally resolvable (see [4]). With respect to these ideas we have:

**3.6 Corollary [MA].** *Every crowded space  $X$  with  $\pi$ -weight  $< \mathfrak{c}$  is  $\omega$ -resolvable. In particular, every space with weight  $< \mathfrak{c}$  is hereditarily  $\omega$ -resolvable.*

PROOF: Let  $\mathcal{N}$  be a  $\pi$ -base of  $X$  of cardinality  $< \mathfrak{c}$ . Since  $X$  is crowded and each element of  $\mathcal{N}$  is open in  $X$ , then  $|N| \geq \aleph_0$  for each  $N \in \mathcal{N}$ . On the other hand,  $\mathcal{N}$  is a  $\pi$ -network in  $X$ , so the conclusion follows. □

It is easy to see that if  $X$  has  $\pi$ -character and density  $\leq \kappa$ , then  $X$  has a  $\pi$ -base of cardinality  $\leq \kappa$ .

**3.7 Proposition [MA].** *If  $X$  is a space with density and  $\pi$ -character  $< \mathfrak{c}$ , then every dense subset of  $X$  is  $\omega$ -resolvable.*

PROOF: The space  $X$  has a  $\pi$ -base  $\mathcal{B}$  of cardinality  $< \mathfrak{c}$ . Let  $H$  be an arbitrary dense subset of  $X$ . It happens now that  $\mathcal{M} = \{N \cap H : N \in \mathcal{N}\}$  is a  $\pi$ -base of  $H$  and has cardinality  $< \mathfrak{c}$ . So, by Corollary 3.6,  $H$  is  $\omega$ -resolvable. □

For every space  $X$ ,  $\max\{t(X), \pi\chi(X)\} \leq \chi(X)$ , so, as a consequence of the last result, and related to Theorems 2.8(2) and 2.8(8), we have:

**3.8 Theorem** [MA]. *If  $X$  is a space such that  $\chi(x, X) < \mathfrak{c}$  for each  $x \in X$ , then  $X$  is hereditarily  $\omega$ -resolvable.*

PROOF: Let  $Y$  be a crowded subspace of  $X$ . The character of  $Y$  is strictly less than  $\mathfrak{c}$ ; thus, the tightness of  $Y$  is  $< \mathfrak{c}$ . Hence, each point  $y$  in  $Y$  is contained in a crowded subspace  $Y_y$  of  $Y$  of cardinality  $< \mathfrak{c}$  (Proposition 2.14). The density and character of each  $Y_y$  is strictly less than  $\mathfrak{c}$ . By Proposition 3.7,  $Y_y$  is  $\omega$ -resolvable. Then  $Y$  is  $\omega$ -resolvable (see Proposition 2.9(1)).  $\square$

The following result is a generalization of Theorems 3.5 and 3.8, which answers, affirmatively, a question posed by the referee. A collection  $\mathcal{N} \subseteq \mathcal{P}(X)$  is a  $\pi$ -network of  $X$  at the point  $x \in X$  if every open set of  $X$  containing  $x$  contains an element of  $\mathcal{N}$ . For each point  $x \in X$ , we define  $\pi nw^*(x, X) = \min\{|\mathcal{N}| : \mathcal{N} \text{ is a } \pi\text{-network of } X \text{ at } x \text{ and each element in } \mathcal{N} \text{ is infinite}\}$ . Of course, for each  $x \in X$ ,  $\pi nw^*(x, X) \leq \chi(x, X)$ . Since MA implies that  $\mathfrak{c}$  is a regular cardinal, we have that, by Theorem 3.5, MA implies that every space  $X$  containing a dense subset  $Y$  of cardinality  $\leq \kappa < \mathfrak{c}$  and such that for every  $y \in Y$ ,  $\pi nw^*(y, X) < \mathfrak{c}$ , is  $\omega$ -resolvable. This result can be ameliorated. Indeed, by using a similar proof to that of Proposition 2.14, if  $X$  is a space with  $\pi nw^*(x, X) < \mathfrak{c}$  for each  $x \in X$ , then each point  $x \in X$  is contained in a crowded subspace  $X_x$  of  $X$  of cardinality  $< \mathfrak{c}$  and having, for each  $y \in X_x$ ,  $\pi nw^*(y, X_x) < \mathfrak{c}$ . So:

**3.9 Corollary** [MA]. *Let  $X$  be a space such that for every  $x \in X$ ,  $\pi nw^*(x, X) < \mathfrak{c}$ . Then  $X$  is  $\omega$ -resolvable.*

We obtain another result with a slightly different mood of that of the previous corollary by defining for each point  $x \in X$  the number  $R(x, X) = \min\{|\Lambda| : \Lambda \text{ is a directed partially ordered set and there is a net } (x_\alpha)_{\alpha \in \Lambda} \text{ in } X \setminus \{x\} \text{ such that } (x_\alpha)_{\alpha \in \Lambda} \text{ converges to } x \text{ in } X\}$ . Indeed, following a similar argumentation to that given in the previous paragraph of Corollary 3.9, we obtain:

**3.10 Corollary** [MA]. *Let  $X$  be a space such that for every  $x \in X$ ,  $R(x, X) < \mathfrak{c}$ . Then  $X$  is  $\omega$ -resolvable.*

In Proposition 4.5 of [2] it was proved that every  $T_2$   $\sigma$ -space is almost- $\omega$ -resolvable. When  $X$  has a countable network, we can repeat that proof assuming only the weaker condition  $T_0$ . So every space with countable network is almost- $\omega$ -resolvable. With respect to  $\sigma$ -spaces, Proposition 4.5 in [2] and Martin's Axiom, Proposition 3.11 allows us to say something else which is, in some sense, stronger than Theorem 3.5:

**3.11 Proposition** [MA]. *Let  $\kappa$  be an infinite cardinal  $< \mathfrak{c}$ . Let  $X$  be a space with a network  $\mathcal{N}$  such that for each finite subcollection  $\mathcal{N}'$  of  $\mathcal{N}$ ,  $\bigcap \mathcal{N}'$  is infinite or empty, and for each  $x \in X$ ,  $|\{N \in \mathcal{N} : x \in N\}| \leq \kappa$ . Then,  $X$  is hereditarily  $\omega$ -resolvable.*

PROOF: The space  $X$  is the condensation of a crowded space  $Y$  ( $Y$  is  $X$  with the topology generated by  $\mathcal{N}$  as a base) which has character strictly less than  $\mathfrak{c}$  (see Proposition 2.11).  $\square$

Next, we obtain a result that we can locate between Theorem 3.5 which deals with  $\pi$ -networks and Corollary 3.6 which speaks of bases. First a definition and some remarks. A space  $X$  is called  $\sigma$ -locally finite if  $X$  can be written as  $\bigcup_{n < \omega} X_n$  where, for each  $n < \omega$ , the collection  $\{\{x\} : x \in X_n\}$  is locally finite in  $X$ . It can be proved that a  $\sigma$ -locally finite crowded space is hereditarily almost- $\omega$ -resolvable.

**3.12 Theorem** [MA]. *Let  $X$  be a crowded topological space with a network  $\mathcal{N}$  with cardinality  $\kappa < \mathfrak{c}$  and such that  $\mathcal{N}_0 = \{N \in \mathcal{N} : |N| < \aleph_0\}$  is  $\sigma$ -locally finite in  $\bigcup \mathcal{N}_0$ . Then  $X$  can be written as  $Y_0 \cup Y_1$  where  $Y_0$  is a (possibly empty) regular closed  $\omega$ -resolvable subspace and  $Y_1$  is an open (possibly empty) almost- $\omega$ -resolvable, hereditarily  $\omega$ -irresolvable space. Besides, if  $Y_1$  is not void, it contains a non-empty open subset which is hereditarily almost- $\omega$ -resolvable. Moreover, if  $X$  is a  $T_1$  Baire space, then  $X$  must be  $\omega$ -resolvable.*

PROOF: Let  $\mathcal{M}$  be the collection of all subspaces of  $X$  which are  $\omega$ -resolvable. Take  $Y_0 = \text{cl}_X \bigcup \mathcal{M}$  and  $Y_1 = X \setminus Y_0$ . Of course  $Y_0$  is closed and  $\omega$ -resolvable. Now, if  $Y_1$  is empty, we have already finished; if the contrary happens,  $Y_1$  is hereditarily  $\omega$ -irresolvable and the collection  $\mathcal{N}' = \{N \in \mathcal{N} : N \subseteq Y_1\}$  is a network in  $Y_1$  with cardinality  $< \mathfrak{c}$  and such that  $\mathcal{N}'_0 = \{N \in \mathcal{N}' : |N| < \aleph_0\}$  is  $\sigma$ -locally finite in  $\bigcup \mathcal{N}'_0$ . Of course  $\mathcal{N}'_0$  is not empty, because otherwise, by Theorem 3.5,  $Y_1$  would be  $\omega$ -resolvable, but this is not possible. Let  $Z$  be the subspace  $\bigcup_{N \in \mathcal{N}'_0} N$  of  $X$ . The space  $Z$  is  $\sigma$ -locally finite. Since  $Y_1$  is hereditarily  $\omega$ -irresolvable,  $Z$  is a dense subset of  $Y_1$ . Then,  $Y_1$  is almost- $\omega$ -resolvable. Furthermore, there must exist a non-empty open subset  $U$  of  $Y_1$  such that each element of  $\mathcal{N}'$  contained in  $U$  is finite because otherwise  $Y_1$  would be  $\omega$ -resolvable (again by Theorem 3.5). So,  $\text{int } Z$  is a non-empty open subset which is hereditarily almost- $\omega$ -resolvable.

Assume now that  $X$  is  $T_1$  and satisfies all the conditions of our proposition including the Baire property. In this case  $Y_1$  must be empty because if this is not the case, the subspace  $\text{int } Z$  of  $Y_1$  would be a  $T_1$  Baire hereditarily almost- $\omega$ -resolvable space. But this means, by Theorem 1.3, that  $\text{int } Z$  is  $\omega$ -resolvable, which is not possible.  $\square$

If we consider in the previous theorem  $\pi$ -networks instead of networks, we still get something interesting.

**3.13 Proposition** [MA]. *Let  $X$  be a crowded topological space with a  $\pi$ -network  $\mathcal{N}$  with cardinality  $\kappa < \mathfrak{c}$  and such that  $\mathcal{N}_0 = \{N \in \mathcal{N} : |N| < \aleph_0\}$  is  $\sigma$ -locally finite. Then  $X$  is equal to  $X_0 \cup X_1$  where  $X_0 \cap X_1 = \emptyset$ ,  $X_0$  is a regular closed (possibly empty) almost- $\omega$ -resolvable space and  $X_1$  is an open (possibly empty)  $\omega$ -resolvable subspace. In particular,  $X$  is, in this case, almost- $\omega$ -resolvable.*

PROOF: Let  $Y$  be the subspace  $\bigcup_{N \in \mathcal{N}_0} N$ . The space  $Y$  is  $\sigma$ -locally finite. If  $Y$  is empty, we obtain our result by Theorem 3.5. If  $Y$  is crowded, then it is almost- $\omega$ -resolvable (see Theorem 3.5 in [26]). If  $Y$  is not empty and is not crowded, we can find an ordinal number  $\alpha > 0$  and, for each  $\beta < \alpha$ , an  $\omega$ -resolvable subspace  $M_\beta$  of  $X$  such that  $X_0 = \text{cl}_X(Y \cup \text{cl}_X(\bigcup_{\beta < \alpha} M_\beta))$  is almost- $\omega$ -resolvable. In fact, let  $D_0$  be the set of isolated points in  $Y_0 = Y$ . For each  $x \in D_0$ , there is an open set  $A_x$  in  $X$  such that  $A_x \cap Y_0 = \{x\}$ . Observe that  $A_x \setminus \{x\}$  is a dense subset of  $A_x$  and it satisfies the conditions in Theorem 3.5, so it is  $\omega$ -resolvable. Thus,  $M_0 = \text{cl}_X(\bigcup_{x \in D_0} A_x)$  is an  $\omega$ -resolvable space. Assume that we have already constructed  $\omega$ -resolvable subspaces  $M_\beta$  of  $X$  with  $\beta < \gamma$ . Put  $Y_\gamma = Y \setminus \text{cl}_X(\bigcup_{\beta < \gamma} M_\beta)$ . If  $Y_\gamma$  is empty or crowded, we take  $\alpha = \gamma$ , and in this case  $\text{cl}_X(Y \cup \text{cl}_X(\bigcup_{\beta < \gamma} M_\beta))$  is almost- $\omega$ -resolvable because  $Y_\gamma$  is empty or crowded and  $\sigma$ -locally finite. If  $Y_\gamma$  is not empty and is not crowded, let  $D_\gamma$  be the set of isolated points in  $Y_\gamma$ . For each  $x \in D_\gamma$  there is an open set  $A_x$  in  $X$  such that  $A_x \cap Y_\gamma = \{x\}$  and  $A_x \cap \text{cl}_X(\bigcup_{\beta < \gamma} M_\beta) = \emptyset$ . Again  $A_x \setminus \{x\}$  is a dense subset of  $A_x$  and it is  $\omega$ -resolvable because of Theorem 3.5. Thus,  $M_\gamma = \text{cl}_X(\bigcup_{x \in D_\gamma} A_x)$  is an  $\omega$ -resolvable space. Continuing with this process we have to find an ordinal number  $\alpha$  for which  $X_0 = \text{cl}_X(Y \setminus \text{cl}_X(\bigcup_{\beta < \alpha} M_\beta))$  is almost- $\omega$ -resolvable.

Now, if  $X_1 = X \setminus X_0$  is not empty, then it is a crowded space and  $\mathcal{N}_1 = \{N \in \mathcal{N} : N \subseteq X_1\}$  is a  $\pi$ -network in  $X_1$  with infinite elements and  $|\mathcal{N}_1| < \mathfrak{c}$ . Then, again by Theorem 3.5,  $X_1$  is  $\omega$ -resolvable. Therefore,  $X = X_0 \cup X_1$ , and  $X_0, X_1$  satisfy the conditions of our proposition.  $\square$

**3.14 Questions.** (1) *Let  $X$  be a crowded space with cardinality  $< \mathfrak{c}$ . Does  $\text{MA} + \neg \text{CH}$  imply that  $X$  is almost- $\omega$ -resolvable?*

(2) *Is there a combinatorial axiom on  $\omega_1$  ensuring that every card-homogeneous topology in  $\omega_1$  is almost- $\omega$ -resolvable?*

(3) *Does  $\diamond$  imply that every card-homogeneous topology in  $\omega_1$  is almost- $\omega$ -resolvable?*

#### 4. Martin's Axiom, cellularity and $\omega$ -resolvable Baire spaces

It is well known that  $\text{MA}(\omega_1)$  implies that a Souslin line does not exist. That is,  $\text{MA}(\omega_1) \Rightarrow \text{SH}$ . We show that it is enough to assume SH in order to prove that every  $T_2$  space with countable cellularity is almost- $\omega$ -resolvable.

**4.1 Theorem [SH].** *Every crowded  $T_2$  space with countable cellularity is almost- $\omega$ -resolvable.*

PROOF: Let  $a_0 \in X$  and  $F_0 = \{a_0\}$ . Let  $\mathcal{C}_0$  be a maximal cellular family of open sets in  $X \setminus F_0$  containing at least two elements. Let  $X_0$  be equal to  $\bigcup \mathcal{C}_0$ . Assume that we have already constructed, by recursion, families  $\{\mathcal{C}_\alpha : \alpha < \gamma\}$ ,  $\{X_\alpha : \alpha < \gamma\}$  and  $\{F_\alpha : \alpha < \gamma\}$ , such that

(1) for all  $\alpha < \gamma$ ,  $\mathcal{C}_\alpha$  is a maximal cellular collection of open sets in  $X$ ;

- (2) if  $\alpha < \xi < \gamma$ , then  $\mathcal{C}_\xi$  properly refines  $\mathcal{C}_\alpha$ ;
- (3) if  $\alpha < \xi < \gamma$  and  $C \in \mathcal{C}_\alpha$ , then  $\mathcal{C}_\xi$  contains a maximal cellular family of proper open sets of  $C$  having more than one element;
- (4)  $X_\alpha = \bigcup \mathcal{C}_\alpha$  for each  $\alpha < \gamma$ ;
- (5) the family  $\{X_\alpha : \alpha < \gamma\}$  is a strictly decreasing  $\gamma$ -sequence of open sets in  $X$ ;
- (6)  $F_\alpha \neq \emptyset$  for every  $\alpha < \gamma$ ;
- (7)  $F_\alpha \subseteq (\bigcap_{\xi < \alpha} X_\xi) \setminus X_\alpha$  for all  $\alpha < \gamma$ ;
- (8)  $\text{int}(F_\alpha) = \emptyset$  for all  $\alpha < \gamma$ .

If  $\gamma$  is a successor ordinal, say  $\gamma = \xi + 1$ , take for each  $C \in \mathcal{C}_\xi$  a point  $a_C^\gamma \in C$ . Now, take a maximal cellular family of open proper subsets in  $C \setminus \{a_C^\gamma\}$  with more than one element,  $\mathcal{C}_C^\xi$  (this is possible because  $C$  is  $T_2$  and infinite). Put  $\mathcal{C}_\gamma = \bigcup_{C \in \mathcal{C}_\xi} \mathcal{C}_C^\xi$ ,  $X_\gamma = \bigcup \mathcal{C}_\gamma$  and  $F_\gamma = \{a_C^\gamma : C \in \mathcal{C}_\xi\}$ .

If  $\gamma$  is a limit ordinal, analyse the set  $\bigcap_{\xi < \gamma} X_\xi$ : if  $\text{int}(\bigcap_{\xi < \gamma} X_\xi) = \emptyset$ , declare our process finished; and if  $\text{int}(\bigcap_{\xi < \gamma} X_\xi)$  is not empty, take a point  $a_\gamma \in \text{int}(\bigcap_{\xi < \gamma} X_\xi)$  and take a maximal cellular family  $\mathcal{C}_\gamma$  with cardinality bigger than one of open proper subsets in  $\text{int}(\bigcap_{\xi < \gamma} X_\xi) \setminus F_\gamma$  where  $F_\gamma = \{a_\gamma\}$ . Put  $X_\gamma = \bigcup \mathcal{C}_\gamma$ .

In this way we can find an ordinal number  $\alpha_0$  and families  $\mathfrak{C} = \{\mathcal{C}_\alpha : \alpha < \alpha_0\}$ ,  $\mathcal{X} = \{X_\alpha : \alpha < \alpha_0\}$  and  $\mathcal{F} = \{F_\alpha : \alpha < \alpha_0\}$  satisfying properties from (1) to (8) above where  $\alpha_0$  is an ordinal number such that  $\text{int}(\bigcap_{\xi < \alpha_0} X_\xi) = \emptyset$  and for each  $\alpha < \alpha_0$ ,  $\text{int}(\bigcap_{\xi < \alpha} X_\xi) \neq \emptyset$ .

First, observe that  $\alpha_0$  must be a limit ordinal and every  $X_\alpha$  is an open set of  $X$ . Now, consider the collection  $\mathcal{Y} = \{Y_\alpha : \alpha < \alpha_0\}$  of subspaces of  $X$  where  $Y_0 = X \setminus X_0$ , and  $Y_\alpha = (\bigcap_{\xi < \alpha} X_\xi) \setminus X_\alpha$  if  $\alpha > 0$ . We have that  $F_\alpha \subseteq Y_\alpha$  and  $\text{int}(Y_\alpha) = \emptyset$  for every  $\alpha < \alpha_0$ .

The set  $\bigcup_{\alpha < \alpha_0} \mathcal{C}_\alpha$  with the order relation  $\subseteq$  is a tree  $T$  and each element in it has at least two immediate successors.

**Claim 1.** The height of  $T$ ,  $\alpha_0$ , is at most  $c(X)^+ = \omega_1$ .

In fact, if  $\alpha_0 > \omega_1$ , then  $\mathcal{C}_{\omega_1} \neq \emptyset$ . Take  $C_{\omega_1} \in \mathcal{C}_{\omega_1}$ . Let  $\mathcal{C} = \{C \in T : C \supseteq C_{\omega_1} \text{ and } C \neq C_{\omega_1}\}$ . Since  $T$  is a tree,  $\mathcal{C}$  is a well ordered set with order type  $\omega_1$ . We can rename  $\mathcal{C}$  as  $\{C_\alpha : \alpha < \omega_1\}$  where  $C_\alpha$  is the only element in  $\mathcal{C}_\alpha$  which belongs to  $\mathcal{C}$ . For each  $\alpha < \omega_1$ , there is  $A_{\alpha+1} \in \mathcal{C}_{\alpha+1}$  such that  $A_{\alpha+1} \subseteq C_\alpha$  and  $A_{\alpha+1} \cap C_{\alpha+1} = \emptyset$ . The set  $\mathcal{A} = \{A_{\alpha+1} : \alpha < \omega_1\}$  is an antichain in  $T$ . Indeed, let  $A_{\alpha+1}$  and  $A_{\xi+1}$  be two different elements of  $\mathcal{A}$ . Assume that  $\alpha < \xi$ . Hence,  $A_{\xi+1} \subseteq C_\xi$  and  $C_\xi \subseteq C_{\alpha+1}$ . But  $C_{\alpha+1} \cap A_{\alpha+1} = \emptyset$ . Therefore,  $A_{\alpha+1} \cap A_{\xi+1} = \emptyset$ . This means that  $c(X) > \aleph_0$ , which is a contradiction. We get that every chain and every antichain of  $T$  has cardinality  $\leq \aleph_0$ . Since we are assuming the Souslin's Hypothesis, there are no Souslin trees. Therefore  $\alpha_0 < \omega_1$ .

It is not difficult to prove that the set  $Z = X \setminus X_{\alpha_0}$  is equal to  $\bigcup_{\alpha < \alpha_0} Y_\alpha$  and that the collection  $\{Y_\alpha : \alpha < \alpha_0\}$  is a partition of  $Z$ .

**Claim 2.** The collection  $\{Y_\alpha : \alpha < \alpha_0\} \cup \{X_{\alpha_0}\}$  is an almost- $\omega$ -resolution for  $X$ ; that is,  $X$  is almost- $\omega$ -resolvable.

The collection  $\mathcal{Y} = \{Y_\alpha : \alpha < \alpha_0\} \cup \{X_{\alpha_0}\}$  is a countable partition of  $X$ . Assume that  $A$  is a non-empty open set of  $X$  and  $|\{\alpha < \alpha_0 : A \cap Y_\alpha \neq \emptyset\}| < \aleph_0$ . Assume that  $H = \{\alpha < \alpha_0 : A \cap Y_\alpha \neq \emptyset\}$  is equal to  $\{\xi_1, \dots, \xi_n\}$  with  $\xi_1 < \xi_2 < \dots < \xi_n$ .

If  $B = A \cap X_{\alpha_0} \neq \emptyset$ , then  $A \cap X_{\xi_n} = B$ . But  $A$  and  $X_{\xi_n}$  are open sets in  $X$ , so  $B$  is a non-empty open set in  $X$ , contradicting the fact that  $\text{int}(X_{\alpha_0}) = \emptyset$ . This means that  $A \cap X_{\alpha_0}$  must be empty.

Now, let  $B = A \cap Y_{\xi_n}$ .  $B$  is not empty and  $A \cap X_{\xi_{n-1}} = B$ . Thus,  $B$  is a non-empty open set in  $X$  which does not intersect any member of  $\mathcal{C}_{\xi_n}$ . If  $\xi_n = \alpha + 1$ ,  $\mathcal{C}_{\xi_n}$  is a maximal cellular collection of open sets contained in  $(\bigcup \mathcal{C}_\alpha) \setminus \{a_C^\alpha : C \in \mathcal{C}_\alpha\} = X_\alpha \setminus \{a_C^\alpha : C \in \mathcal{C}_\alpha\}$ . Hence,  $B \cap \{a_C^\alpha : C \in \mathcal{C}_\alpha\} \neq \emptyset$ . Let  $a_C^\gamma \in B$ . We have that  $M = (C \cap B) \setminus \{a_C^\gamma\}$  is an open set contained in  $X_\alpha \setminus \{a_C^\alpha : C \in \mathcal{C}_\alpha\}$  and no element in  $\mathcal{C}_\xi$  intersects  $M$ . By maximality of  $\mathcal{C}_\xi$ , we must have that  $M$  is empty; that is,  $C \cap B = \{a_C^\gamma\}$ , and this is not possible because  $X$  does not have isolated points.

Now assume that  $\xi_n$  is a limit ordinal. Since  $B$  is open and  $B \subseteq \bigcap_{\xi < \xi_n} X_\xi$ ,  $B$  must be contained in  $\text{int}(\bigcap_{\xi < \xi_n} X_\xi)$ . Since  $\{a_{\xi_n}\}$  is closed and  $B$  does not intersect any element of  $\mathcal{C}_{\xi_n}$  which is a maximal cellular family of open sets contained in the set  $\text{int}(\bigcap_{\xi < \xi_n} X_\xi) \setminus \{a_{\xi_n}\}$ ,  $B$  must be equal to  $\{a_{\xi_n}\}$ , which is again a contradiction.

Therefore,  $|\{\xi < \alpha_0 : A \cap Y_\xi \neq \emptyset\}|$  must be equal to  $\aleph_0$ . □

Since the cellularity of a space is a monotone function when it is applied on dense subspaces, and using Theorem 1.3, we conclude:

**4.2 Corollary** [SH]. *Every  $T_2$  Baire space with  $c(X) \leq \aleph_0$  is  $\omega$ -resolvable.*

Example 4.3 in [26] (see Example 2.3 above) gives us a space which is Baire,  $T_1$  with countable cellularity but it is not almost- $\omega$ -resolvable. This example is constructed assuming the existence of measurable cardinals. Moreover, there is a model  $M$  in which SH holds and there are measurable cardinals. So we cannot get anything stronger than our results of this section by assuming only  $T_1$ . Furthermore, we cannot erase the Baire condition in Corollary 4.2 because there is in ZFC a Tychonoff, countable irresolvable space (see Examples 2.5). Finally, in 2.2 we list an example of a space with cellularity  $\leq \aleph_1$  which is Baire and is not almost- $\omega$ -resolvable. This last example is given by assuming the existence of an  $\omega_1$ -complete ideal over  $\omega_1$  which has a dense set of cardinality  $\omega_1$ . Hence, it is natural to ask:

**4.3 Question.** *Does MA imply that every crowded  $T_2$  space of cellularity  $< \mathfrak{c}$  is almost- $\omega$ -resolvable?*

In this question, we cannot change “almost- $\omega$ -resolvable” for “resolvable” since there is in ZFC an irresolvable countable space.

## 5. Almost- $\omega$ -irresolvable spaces

A space is *almost- $\omega$ -irresolvable* if it is not almost- $\omega$ -resolvable. In a similar way we define almost irresolvable spaces.

**5.1 Proposition.** *If  $X$  is almost- $\omega$ -irresolvable, then there is a non-empty open subset  $U$  of  $X$  which is hereditarily almost- $\omega$ -irresolvable.*

PROOF: Let  $\mathcal{U}$  be the collection of all almost- $\omega$ -resolvable subspaces  $Y$  of  $X$ . The set  $Z = \text{cl}_X(\bigcup \mathcal{U})$  is almost- $\omega$ -resolvable and  $U = X \setminus Z$  is not empty and satisfies the requirements.  $\square$

**5.2 Proposition.** *If  $X$  is open hereditarily almost- $\omega$ -irresolvable, then  $X$  is a Baire space.*

PROOF: Let  $\{U_n : n < \omega\}$  be a sequence of open and dense subsets of  $X$ . We can choose this sequence to be  $\subseteq$ -decreasing. Denote by  $F$  the set  $\bigcap_{n < \omega} U_n$ . We claim that  $F$  is dense in  $X$ . In fact, if for a  $k < \omega$ ,  $\text{cl}_X F \supseteq U_k$ , then  $\text{cl}_X F \supseteq \text{cl}_X U_k = X$  and  $F$  is dense. Now, assume that for each  $n < \omega$ ,  $U_n \setminus \text{cl}_X F$  is not empty. In this case, the collection  $T = \{i < \omega : (U_i \setminus U_{i+1}) \cap (X \setminus \text{cl}_X F) \neq \emptyset\}$  is infinite. For each  $i \in T$ , we put  $T_i = (U_i \setminus U_{i+1}) \cap (X \setminus \text{cl}_X F)$ . The collection  $\{T_i : i < \omega\}$  forms an almost- $\omega$ -resolution of  $X \setminus \text{cl}_X F$ . But this is not possible.  $\square$

**5.3 Corollary.** *If there is an almost resolvable space  $X$  which is almost- $\omega$ -irresolvable, then there is a resolvable Baire open subspace  $U$  of  $X$  which is hereditarily almost- $\omega$ -irresolvable.*

PROOF: Let  $X$  be an almost-resolvable almost- $\omega$ -irresolvable space. The space  $X$  contains a non-empty open subspace  $U$  which is hereditarily almost- $\omega$ -irresolvable. By Proposition 5.2,  $U$  is a Baire space; so, it is resolvable being almost resolvable.  $\square$

**5.4 Corollary.** *There is an almost resolvable space  $X$  which is almost- $\omega$ -irresolvable if and only if there is an almost resolvable Baire space which is hereditarily almost- $\omega$ -irresolvable.*

As a consequence of the previous result, we have that almost resolvability and almost- $\omega$ -resolvability coincide in the class of spaces  $X$  in which every open subset is not a Baire space. Even more was obtained in [2, Corollary 5.21]: every space which does not contain a Baire open subspace is almost- $\omega$ -resolvable.

**5.5 Proposition.** *Let  $X$  be a  $T_1$  space. Then  $X$  is hereditarily resolvable if and only if  $X$  is hereditarily  $\omega$ -resolvable.*

PROOF: Let  $Y$  be a crowded subspace of  $X$  and assume that  $Y$  is not  $\omega$ -resolvable. Then, there is  $k \in \omega$  with  $k > 1$  such that  $X$  is  $k$ -resolvable but  $X$  is not  $(k + 1)$ -resolvable [15]. So there are  $D_0, \dots, D_{k-1}$  dense and pairwise disjoint subspaces of  $Y$ . But, then, each  $D_i$  is crowded and irresolvable, a contradiction.  $\square$

**5.6 Proposition.** *Let  $X$  have the property that every of its crowded subspaces is Baire. Then  $X$  is hereditarily  $\omega$ -resolvable iff  $X$  is hereditarily resolvable iff  $X$  is hereditarily almost- $\omega$ -resolvable iff  $X$  is hereditarily almost resolvable.*

Several results established in [2, Section 5] and [26, Section 4] relate Baire irresolvable spaces with the property of almost- $\omega$ -resolvability (see also [1, Section 3]). In the following theorem we obtain the most general possible result in the mood of these propositions.

**5.7 Theorem.** *For crowded  $T_1$  spaces and for a crowded-hereditarily topological property  $P$ , the following assertions are equivalent:*

- (1) every Baire space with  $P$  is  $\omega$ -resolvable,
- (2) every Baire space with  $P$  is resolvable,
- (3) every space with  $P$  is almost- $\omega$ -resolvable,
- (4) every space with  $P$  is almost resolvable.

PROOF: The implications (1)  $\Rightarrow$  (2) and (3)  $\Rightarrow$  (4) are evident.

(2)  $\Rightarrow$  (3): Assume that  $X$  is not almost- $\omega$ -resolvable and satisfies  $P$ . The space  $X$  contains an open and non-empty subset  $U$  which is hereditarily almost- $\omega$ -irresolvable. By Proposition 5.5,  $U$  is not hereditarily resolvable, so there is a crowded subspace  $Y$  which is not resolvable. Observe that  $Y$  is hereditarily almost- $\omega$ -irresolvable, then  $Y$  is an irresolvable Baire space because of Proposition 5.2. Since  $P$  is a crowded-hereditarily topological property,  $Y$  satisfies  $P$  too.

(4)  $\Rightarrow$  (2): Assume that  $X$  is a Baire space with  $P$ . By hypothesis,  $X$  is almost resolvable and every Baire almost resolvable space is resolvable (see [2, Corollary 5.4]).

(3)  $\Rightarrow$  (1): Assume that  $X$  is a Baire space with  $P$ . By hypothesis, every crowded subspace  $Y$  of  $X$  has  $P$  and so it is almost- $\omega$ -resolvable; hence  $X$  is  $\omega$ -resolvable because of Theorem 1.3.  $\square$

Taking  $P$  equal to “ $X$  is a crowded topological space”, we have:

**5.8 Corollary.** *For crowded  $T_1$  spaces, the following assertions are equivalent:*

- (1) every Baire space is  $\omega$ -resolvable,
- (2) every Baire space is resolvable,
- (3) every space is almost- $\omega$ -resolvable,
- (4) every space is almost resolvable.



A space is *locally homogeneous* if each of its points has a homogeneous neighborhood. For a cardinal number  $\kappa \geq 1$ , we will say that  $X$  is *exactly  $\kappa$ -resolvable*, in symbols  $E_\kappa R$ , if  $X$  is  $\kappa$ -resolvable but is not  $\kappa^+$ -resolvable. The space  $X$  is said to be  $OE_\kappa R$  if every non-empty open set in  $X$  is  $E_\kappa R$ . The concept and examples of  $E_n R$  spaces for  $n \in \omega$  have existed in the literature for some time (see, for example, [10] and [8]). It is clear that the  $OE_\kappa R$  spaces are  $E_\kappa R$ . The above definitions can be viewed as natural generalizations of the concepts of irresolvable and open-hereditarily irresolvable spaces since  $E_1 R$  and irresolvability are the same concept and  $OE_1 R$  and open-hereditarily irresolvability coincide.

It was proved in [1, Theorem 3.13] that every locally homogeneous irresolvable space such that its cardinality is not a measurable cardinal is of the first category. Also, Li Feng and O. Masaveu [13] proved that every crowded topological space  $X$  can be written as

$$X = \Omega \cup \text{cl}_X \left( \bigcup_{n=1}^{\infty} O_n \right),$$

where

- (1) for each  $n$ ,  $O_n$  is an open, possibly empty, subset of  $X$ ;
- (2) for each  $n$ , if  $O_n \neq \emptyset$ , then it is  $OE_n R$ ;
- (3) for  $n \neq m$ ,  $O_n \cap O_m = \emptyset$ ; and
- (4)  $\Omega$  is an open, possibly empty,  $\omega$ -resolvable subset of  $X$ .

Thus we obtain the following:

**5.9 Proposition.** *Every locally homogeneous Baire space of cardinality strictly less than the first measurable cardinal is resolvable.*

PROOF: Let  $X$  be a locally homogeneous Baire space. Write  $X$  as Feng and Masaveu say:  $X = \Omega \cup \text{cl}_X (\bigcup_{n=1}^{\infty} O_n)$ . Assume that  $O_1$  is not empty and take  $x \in O_1$ . There is a homogeneous neighborhood  $W$  of  $x$ . (Observe that  $W$  has to be contained in  $X \setminus \text{cl}_X (\Omega \cup \bigcup_{n>1} O_n) \subseteq \text{int}_X \text{cl}_X O_1$ ). On the other hand,  $O_1$  is open hereditarily irresolvable, so  $\text{int}_X W \cap O_1$  is irresolvable. Since  $\text{int}_X W \cap O_1$  is a non-empty open subset of  $W$ ,  $W$  is irresolvable. By Theorem 3.13 in [1],  $W$  is of first category. In particular the open and non-empty subset  $O_1 \cap \text{int}_X W$  of  $X$  is of first category in itself, but this is not possible because  $X$  is a Baire space. Hence,  $O_1 = \emptyset$  and  $X$  is resolvable.  $\square$

- 5.10 Questions.**
- (1) *Is every pseudocompact (resp., Čech-complete) Tychonoff space almost- $\omega$ -resolvable in ZFC?*
  - (2) *Is every Baire locally homogeneous space (resp., homogeneous space, topological group)  $\omega$ -resolvable?*
  - (3) *For each  $n > 1$ , is there a Baire  $OE_n R$  space?*

## 6. The infinite $\pi$ -network and $\text{Seq}(u_t)$ spaces

We define the *infinite  $\pi$ -networkweight* of a crowded space  $X$ ,  $\pi nw^*(X)$ , as the minimum infinite cardinal of a  $\pi$ -network with infinite elements. And  $\pi nw(X)$  is the minimum infinite cardinal of a  $\pi$ -network in  $X$ . It is easy to prove that  $\pi nw(X) = d(X)$  for every topological space  $X$ . Moreover, for a crowded space  $X$ , we have  $d(X) \leq \pi nw^*(X) \leq \min\{d(X) \cdot \sup\{\pi nw^*(x, X) : x \in X\}, d(X) \cdot R(X), \pi w(X)\}$ , where  $nw^*(x, X)$  and  $R(X)$  were defined before Corollaries 3.9 and 3.10. Besides, for every metrizable space  $X$  we have  $d(X) = w(X)$ . So, for a crowded metrizable space  $X$ , the equality  $\pi nw^*(X) = \pi nw(X)$  always holds. We have the same phenomenon for spaces of the form  $C_p(X)$ , the space of real continuous function defined on  $X$  with the pointwise convergent topology (here,  $X$  is not necessarily crowded). Indeed, for  $f \in C_p(X)$ , the sequence  $(f_n)_{n < \omega}$  where  $f_n = f + 1/n$ , converges to  $f$ . So, if  $D$  is a dense subset of  $C_p(X)$  with cardinality equal to  $d(C_p(X))$ , the collection  $\{\{f\} \cup \{f_i : i \geq n\} : f \in D, n < \omega\}$  is a  $\pi$ -network of cardinality  $d(C_p(X))$  constituted by infinite elements. So,  $\pi nw^*(C_p(X)) = \pi nw(C_p(X))$ . In particular, for every cardinal number  $\kappa$ ,  $\pi nw^*(\mathbb{R}^\kappa) = d(\mathbb{R}^\kappa)$ . The same can be said for spaces of the form  $C_p(X, 2)$  where  $X$  is an infinite zero-dimensional  $T_2$  space. In fact, we can take an infinite discrete subspace  $Y = \{x_n : n < \omega\}$  of  $X$ , and clopen subsets  $\{V_n : n < \omega\}$  such that, for each  $n < \omega$ ,  $Y \cap V_n = \{x_n\}$ . The characteristic functions  $\chi_{V_n}$  constitute a sequence which converge to the constant function 0. So, in this case too,  $\pi nw^*(C_p(X, 2)) = d(C_p(X, 2))$ .

We have already mentioned that in [1] a dense countable subset  $Y$  of  $2^c$  which is irresolvable was constructed in ZFC. This space has  $\pi nw(Y) = \aleph_0$ , but every of its countable  $\pi$ -networks has to have finite elements, because otherwise  $Y$  would be maximally resolvable (see Theorem 2.8(1)). The  $\text{Seq}(u_t)$  spaces considered below are also examples of spaces of this kind.

We recall that for a  $p \in \omega^*$ ,  $\chi(p) = \min\{|b| : b \text{ is a base for } p\}$ . Of course we can also define:  $\pi\chi(p) = \min\{|b| : b \text{ is a } \pi\text{-base for } p\}$  where a family of infinite sets  $\mathcal{G}$  in  $\omega$  is a  $\pi$ -base for  $p$  if every member of  $p$  contains an element of  $\mathcal{G}$ . It is not difficult to prove that for every  $p \in \omega^*$ ,  $\pi\chi(p) \leq \chi(p)$  and  $\pi\chi(p) > \aleph_0$ . In fact, assume that  $N_0, \dots, N_k, \dots$  are infinite subsets of  $\omega$ . By recursion, we can construct two sequences  $A = \{a_0, \dots, a_n, \dots\}$  and  $B = \{b_0, \dots, b_n, \dots\}$  such that the elements in  $A \cup B$  are pairwise different, and for each  $n < \omega$ ,  $a_n, b_n \in N_n$ . If  $A \in p$  then  $A$  is an element of  $p$  which does not contain any  $N_k$ . If  $A \notin p$ , then  $\omega \setminus A$  belongs to  $p$  and does not contain any  $N_k$ .

By  $\text{Seq}$  we mean the set of all finite sequences of natural numbers. More precisely, for each natural number  $n \in \omega$ , let  ${}^n\omega = \{t : t \text{ is a function and } t : n \rightarrow \omega\}$ . Then  $\text{Seq} = \bigcup_{n \in \omega} {}^n\omega$ . If  $t \in \text{Seq}$ , with domain  $k = \{0, 1, \dots, (k-1)\}$ , and  $n \in \omega$ , let  $t \frown n$  denote the function  $t \cup \{(k, n)\}$ . For every  $t \in \text{Seq}$  let  $u_t$  be a non-principal ultrafilter on  $\omega$ . By  $\text{Seq}(\{u_t : t \in \text{Seq}\})$  we denote the space with underlying set  $\text{Seq}$  and topology defined by declaring a set  $U \subseteq \text{Seq}$  to be open if

and only if

$$(\forall t \in U) \{n \in \omega : t \frown n \in U\} \in u_t.$$

For short, we write  $\text{Seq}(u_t)$  instead of  $\text{Seq}(\{u_t : t \in \text{Seq}\})$ . We also consider the case where there is a single non-principal ultrafilter  $p$  in  $\omega$  such that  $u_t = p$  for all  $t \in \text{Seq}$ , and in this case we write  $\text{Seq}(p)$  instead of  $\text{Seq}(u_t)$ .

We use the following notation of W. Lindgren and A. Szymanski [20]; put  $L_n = \{s \in \text{Seq} : \text{dom}(s) = n\}$ , and for any  $s \in \text{Seq}$  the *cone over  $s$*  is defined by  $C(s) = \{t \in \text{Seq} : s \subseteq t\}$ . In particular,  $L_0 = \{\emptyset\}$ . We add some other notations: For each  $s \in L_n$ ,  $T(s) = \{t \in L_{n+1} : s \subseteq t\}$ . Observe that for every  $s \in \text{Seq}$ ,  $C(s)$  is a clopen subset of  $\text{Seq}(u_t)$ .

It is well-known that for any choice of  $\{u_t : t \in \text{Seq}\} \subseteq \omega^*$ , the space  $\text{Seq}(u_t)$  is a zero-dimensional, extremally disconnected, Hausdorff space with no isolated points. By the way,  $\text{Seq}(p)$  is homogeneous and if  $p$  is Ramsey, there is a binary group operation  $+$  such that  $(\text{Seq}(p), +)$  is a topological group (see [27]).

**6.1 Proposition.** *Every  $\text{Seq}(u_t)$  space is  $\omega$ -resolvable.*

PROOF: In fact, let  $\{E_n : n < \omega\}$  be a partition of  $\omega$  where each  $E_n$  is infinite. Set  $D_n = \bigcup_{i \in E_n} L_i$ . Each  $D_n$  is dense in  $\text{Seq}(u_t)$  and  $D_n \cap D_m = \emptyset$  if  $n \neq m$ . □

**6.2 Proposition.** *Let  $\{u_t : t \in \text{Seq}\} \subseteq \omega^*$ . Then, the infinite  $\pi$ -netweight of  $\text{Seq}(u_t)$  is not countable.*

PROOF: For each  $n < \omega$ , each  $s \in L_n$ , and each sequence  $S$  of subcollections of the form

$$\begin{aligned} &\{B(s)\}, \{B(s, i_{n+1}) : i_{n+1} \in B(s)\}, \{B(s, i_{n+1}, i_{n+2}) : i_{n+1} \in B(s), \\ &\quad i_{n+2} \in B(s, i_{n+1})\}, \dots, \{B(s, i_{n+1}, \dots, i_{n+k+1}) : i_{n+1} \in B(s), \\ &\quad i_{n+1} \in B(s, i_{n+1}), \dots, i_{n+k+1} \in B(s, i_{n+1}, \dots, i_{n+k})\}, \dots \end{aligned}$$

where  $B(s) \in u_s$  and, if  $i_{n+1} \in B(s), i_{n+2} \in B(s, i_{n+1}), \dots, i_{n+k} \in B(s, i_{n+1}, \dots, i_{n+k-1}), B(s, i_{n+1}, \dots, i_{n+k}) \in u_t$  with  $t = s \frown \widehat{i_{n+1}} \dots \frown i_{n+k}$ , we define a set  $V(s, S)$  as follows:

$$\begin{aligned} V(s, S) = \{ &s\} \cup \{t \in \text{Seq}(p) : m \in \omega, t \in L_{n+m+1}, s \subseteq t, t(n+1) \in B(s), \\ &t(n+2) \in B(s, t(n+1)), \dots, \\ &t(n+m+1) \in B(s, t(n+1), t(n+2), \dots, t(n+m))\}. \end{aligned}$$

We call this set  $V(s, S)$  *cascade of  $\text{Seq}(p)$  defined by  $(s, S)$* . Moreover, we will called each sequence  $S$ , described as above, *fan on  $(s, (u_t))$* .

Of course, the collection of cascades forms a base of clopen sets for  $\text{Seq}(u_t)$ .

**Claim 1.** If  $\mathcal{N} = \{N_0, \dots, N_k, \dots\}$  is a countable set of infinite subsets of  $\text{Seq}(u_t)$ , then  $\mathcal{N}$  is not a  $\pi$ -network of  $\text{Seq}(u_t)$ .

We are going to prove Claim 1 in several lemmas.

**Claim 1.1.** If  $\mathcal{M}$  is a finite collection of subsets of  $\text{Seq}$ , then there is a non-empty open set  $A$  of  $\text{Seq}(u_t)$  such that  $M \setminus A \neq \emptyset$  for all  $M \in \mathcal{M}$ .

PROOF: Take  $s_0, \dots, s_n$  elements in  $\text{Seq}$  such that each  $M$  in  $\mathcal{M}$  contains one of this points. There is  $k < \omega$  such that  $s_i \in L_m$  implies  $m < k$  for all  $i \in \{0, \dots, n\}$ . Take  $s \in L_k$ . The cone  $C(s)$  is open and contains no element in  $\mathcal{M}$ .  $\square$

**Claim 1.2.** Assume that  $F \subseteq \text{Seq}(u_t)$  is such that  $|F \cap T(s)| \leq 1$  for every  $s \in \text{Seq}$ . Then,  $F$  is a proper closed subset of  $\text{Seq}(u_t)$ .

PROOF: Let  $P$  be the set  $\{s < \text{Seq} : F \cap T(s) \neq \emptyset\}$ . Let  $z_s$  be the only point belonging to  $F \cap T(s)$  for each  $s \in P$ . Let  $x \in \text{Seq}(u_t) \setminus F$ . Assume that  $x = (n_0, \dots, n_k)$  (the argument is similar if  $x = \emptyset$ ). Let

$$S = \{\{B(x)\}, \{B(x, i_0) : i_0 \in B(x)\}, \{B(x, i_0, i_1) : i_0 \in B(x), i_1 \in B(x, i_0)\}, \dots, \\ \{B(x, i_0, \dots, i_{k+1}) : i_0 \in B(x), i_1 \in B(x, i_1), \dots, i_{k+1} \in B(x, i_0, \dots, i_k)\}, \dots\}$$

be a fan on  $(x, (u_t))$ . We claim that the set  $V(x, S) \setminus F$  is an open set. Indeed, if  $y \in V(x, S) \setminus F$ ,  $y$  is of the form  $(n_0, \dots, n_k, i_0, \dots, i_{m+1})$  where  $m < \omega$ ,  $i_0 \in B(x), i_1 \in B(x, i_0), \dots, i_{m+1} \in B(x, i_0, i_1, \dots, i_m)$ .

The set  $\{l < \omega : (n_0, \dots, n_k, i_0, \dots, i_{m+1}, l) \in V(x, S) \setminus F\}$  is equal to

$$B(x, i_0, i_1, \dots, i_{m+1}) \setminus F.$$

Moreover, the set  $B(x, i_0, i_1, \dots, i_{m+1}) \cap F = G$  is either empty if  $F \cap T(x, i_0, i_1, \dots, i_{m+1}) = \emptyset$ , or  $G = \{z_{(x, i_0, i_1, \dots, i_{m+1})}\}$  if  $F \cap T(x, i_0, i_1, \dots, i_{m+1}) \neq \emptyset$ . Of course, in both cases,  $B(x, i_0, i_1, \dots, i_{m+1}) \setminus F$  belongs to  $u_t$  where  $t = x \hat{\ } i_0 \hat{\ } \dots \hat{\ } i_{m+1}$ . This means that  $V(x, S) \setminus F$  is open.  $\square$

**Claim 1.3.** Let  $\mathcal{M} = \{N \in \mathcal{N} : \forall s \in \text{Seq}(|N \cap T(s)| < \aleph_0)\}$ . Then, there is a non-empty open set  $A$  of  $\text{Seq}(u_t)$  such that  $N \setminus A \neq \emptyset$  for all  $N \in \mathcal{M}$ .

PROOF: First, we define in  $\text{Seq}$  a well order  $\sqsubseteq$  as follows:  $\emptyset$  is the  $\sqsubseteq$ -first element, and for two elements  $s$  and  $t$  different to  $\emptyset$ , we define  $s \sqsubset t$  if either  $s \in L_{n+1}$ ,  $t \in L_{m+1}$  and  $n < m$ , or  $n = m$  and  $s(n) < t(n)$ .

Because of Claim 1.1, we can assume that  $\mathcal{M}$  is infinite. We faithfully enumerate  $\mathcal{M}$  as  $\{M_0, M_1, \dots, M_k, \dots\}$ . Consider the set  $J = \{s \in \text{Seq} : \exists M \in \mathcal{M}$  such that  $T(s) \cap M \neq \emptyset\}$ . Because of the definition of  $\mathcal{M}$ , we must have  $|J| = \aleph_0$ . Hence, we can enumerate  $J$  as  $\{s_m : m < \omega\}$  in such a way that  $s_0 \sqsubset s_1 \sqsubset \dots \sqsubset s_n \sqsubset s_{n+1} \sqsubset \dots$ .

Let  $k_0$  be the first natural number  $m$  such that  $M_m \cap T(s_0) \neq \emptyset$ . We take  $z_0 \in M_{k_0} \cap T(s_0)$ . Assume that we have already defined two finite sequences  $k_0, \dots, k_l$  and  $z_0, \dots, z_l$  such that

- (1) for each  $i \in \{0, \dots, l - 1\}$ ,  $k_{i+1}$  is the first natural number  $m \in \omega \setminus \{k_0, \dots, k_i\}$  such that  $M_m \cap T(s_{i+1}) \neq \emptyset$ , and
- (2)  $z_{i+1} \in M_{k_{i+1}} \cap T(s_{i+1})$  for each  $i \in \{0, \dots, l - 1\}$ .

We define now  $k_{l+1}$  as the first natural number  $m \in \omega \setminus \{k_0, \dots, k_l\}$  such that  $M_m \cap T(s_{l+1}) \neq \emptyset$ . Take  $z_{l+1} \in M_{k_{l+1}} \cap T(s_{l+1})$ .

Observe that  $\{k_i : i < \omega\} = \omega$ . Indeed, assume that  $\{0, \dots, m\} \subseteq \{k_i : i < \omega\}$  and  $\{k_{i_0}, \dots, k_{i_m}\} = \{0, \dots, m\}$ . Let  $j$  be a natural number greater than  $k_{i_l}$  for all  $l \in \{0, \dots, m\}$  and such that  $M_{m+1} \cap T(s_j) \neq \emptyset$ . Then we must have  $m + 1 \in \{k_0, \dots, k_j\}$ .

We put  $F = \{z_i : i < \omega\}$ . The set  $F$  satisfies the conditions required in Claim 1.2; so,  $F$  is a proper closed subset of  $\text{Seq}(u_t)$ . Therefore,  $A = \text{Seq}(u_t) \setminus F$  is a non-empty open set which does not contain any of the sets  $M \in \mathcal{M}$ . □

**Claim 1.4.** Let  $\mathcal{O} = \mathcal{N} \setminus \mathcal{M} = \{N \in \mathcal{N} : \exists s \in \text{Seq}(|N \cap T(s)| \geq \aleph_0)\}$ . Then, there is an open set  $B$  of  $\text{Seq}(u_t)$  such that  $N \setminus B \neq \emptyset$  for all  $N \in \mathcal{O}$ .

PROOF: Let  $T = \{n < \omega : N_n \in \mathcal{O}\}$ . The open set  $B$  will be an open cascade  $V(s, S)$  defined by  $(s, S)$  where  $s = \emptyset$  and the fan

$$S = \{\{B(s)\}, \{B(s, i_1) : i_1 \in B(s)\}, \{B(s, i_1, i_2) : i_1 \in B(s), i_2 \in B(s, i_1)\}, \dots, \{B(s, i_1, \dots, i_{k+1}) : i_1 \in B(s), i_1 \in B(s, i_1), \dots, i_{k+1} \in B(s, i_1, \dots, i_k)\}, \dots\}$$

will be constructed by recursion.

Assume that we have already selected

$$\{\{B(s)\}, \{B(s, i_1) : i_1 \in B(s)\}, \{B(s, i_1, i_2) : i_1 \in B(s), i_2 \in B(s, i_1)\}, \dots, \{B(s, i_1, \dots, i_k) : i_1 \in B(s), i_2 \in B(s, i_1), \dots, i_k \in B(s, i_1, \dots, i_{k-1})\}\}$$

For each sequence  $i_1 \in B(s), i_2 \in B(s, i_1), \dots, i_{k+1} \in B(s, i_1, i_2, \dots, i_k)$ , consider the ultrafilter  $u_t$  where  $t = s \widehat{\cap} i_1 \widehat{\cap} \dots \widehat{\cap} i_k$ , and consider the set  $P(s, i_1, \dots, i_{k+1}) = \{n \in T : |N_n \cap T(s, i_1, \dots, i_k)| \geq \aleph_0\}$ . If  $P(s, i_1, \dots, i_{k+1})$  is empty, we choose  $B(s, i_1, \dots, i_{k+1})$  to be an arbitrary element of  $u_t$ . If  $P(s, i_1, \dots, i_{k+1})$  is not empty, there is  $B(s, i_1, \dots, i_{k+1}) \in u_t$  such that  $N_n \setminus B(s, i_1, \dots, i_{k+1}) \neq \emptyset$  for every  $n \in P(s, i_1, \dots, i_{k+1})$  because  $\pi\chi(u_t) > \aleph_0$ .

We have already finished the description of the recursive process that define the fan  $S$ . The set  $B = V(s, S)$  is the required open set.

We finished the proof of Claim 1 by saying that the open set  $A \cap B$ , where  $A$  was defined in the proof of Claim 1.3 and  $B$  in that of Claim 1.4, is not empty and does not contain any of the elements in  $\mathcal{N}$ .  $\square$

## REFERENCES

- [1] Alas O., Sanchis M., Tkačhenko M.G., Tkachuk V.V., Wilson R.G., *Irresolvable and submaximal spaces: Homogeneity versus  $\sigma$  discreteness and new ZFC examples*, Topology Appl. **107** (2000), 259–273.
- [2] Angoa J., Ibarra M., Tamariz-Mascarúa Á., *On  $\omega$ -resolvable and almost- $\omega$ -resolvable spaces*, Comment. Math. Univ. Carolin. **49** (2008), 485–508.
- [3] Bell M., Kunen K., *On the  $\pi$ -character of ultrafilters*, C.T. Math. Rep. Acad. Sci. Canada **3** (1981), 351–356.
- [4] Biernias J., Terepeta M., *A sufficient condition for maximal resolvability of topological spaces*, Comment. Math. Univ. Carolin. **41** (2004), 139–144.
- [5] Bolstein R., *Sets of points of discontinuity*, Proc. Amer. Math. Soc. **38** (1973), 193–197.
- [6] Comfort W.W., Feng L., *The union of resolvable spaces is resolvable*, Math. Japonica **38** (1993), 413–414.
- [7] Comfort W.W., García-Ferreira S., *Resolvability: a selective survey and some new results*, Topology Appl. **74** (1996), 149–167.
- [8] van Douwen E.K., *Applications of maximal topologies*, Topology Appl. **51** (1993), 125–139.
- [9] El’kin A.G., *On the maximal resolvability of products of topological spaces*, Soviet Math. Dokl. **10** (1969), 659–662.
- [10] El’kin A.G., *Resolvable spaces which are not maximally resolvable*, Moscow Univ. Math. Bull. **24** (1969), 116–118.
- [11] Feng L., *Strongly exactly  $n$ -resolvable spaces of arbitrary large dispersion character*, Topology Appl. **105** (2000), 31–36.
- [12] Foran J., Liebnitz P., *A characterization of almost resolvable spaces*, Rend. Circ. Mat. Palermo (2) **40** (1991), 136–141.
- [13] Feng L., Masaveu O., *Exactly  $n$ -resolvable spaces and  $\omega$ -resolvability*, Math. Japonica **50** (1999), 333–339.
- [14] Hewitt E., *A problem of set-theoretic topology*, Duke Math. J. **10** (1943), 306–333.
- [15] Illanes A., *Finite and  $\omega$ -resolvability*, Proc. Amer. Math. Soc. **124** (1996), 1243–1246.
- [16] Katětov M., *On topological spaces containing no disjoint dense sets*, Mat. Sb. **21** (1947), 3–12.
- [17] Kunen K., *Set Theory. An Introduction to Independence Proofs*, Studies in Logic and the Foundations of Mathematics, 102, North Holland, sixth impression, Amsterdam, London, New York, Tokyo, 1995.
- [18] Kunen K., Tall F., *On the consistency of the non-existence of Baire irresolvable spaces*, Topology Atlas, <http://at.yorku.ca/v/a/a/a/27.htm> (1998).
- [19] Kunen K., Szymansky A., Tall F., *Baire irresolvable spaces and ideal theory*, Ann. Math. Silesiana **2** (14) (1986), 98–107.
- [20] Lindgren W.F., Szymanski A.A., *A non-pseudocompact product of countably compact spaces via Seq*, Proc. Amer. Math. Soc. **125** (1997), 3741–3746.
- [21] Mal’khin V.I., *On extremally disconnected topological groups*, Soviet Math. Dokl. **16** (1975), 21–25.
- [22] Mal’khin V.I., *On the resolvability of the product of two spaces and a problem of Katětov*, Dokl. Akad. Nauk SSSR **222** (1975), 765–729.
- [23] Mal’khin V.I., *Irresolvable countable spaces of weight less than  $\mathfrak{c}$* , Comment. Math. Univ. Carolin. **40** (1999), no. 1, 181–185.

- [24] Pavlov O., *On resolvability of topological spaces*, Topology Appl. **126** (2002), 37-47.
- [25] Pytke'ev E.G., *On maximally resolvable spaces*, Proc. Steklov. Inst. Math. **154** (1984), 225-230.
- [26] Tamariz-Mascarúa Á., Villegas-Rodríguez H., *Spaces of continuous functions, box products and almost- $\omega$ -resolvable spaces*, Comment. Math. Univ. Carolin. **43** (2002), no. 4, 687-705.
- [27] Vaughan J.E., *Two spaces homeomorphic to  $Seq(p)$* , manuscript.
- [28] Villegas L.M., *On resolvable spaces and groups*, Comment. Math. Univ. Carolin. **36** (1995), 579-584.
- [29] Villegas L.M., *Maximal resolvability of some topological spaces*, Bol. Soc. Mat. Mexicana **5** (1999), 123-136.

F. Casarrubias-Segura, Á. Tamariz-Mascarúa:

DEPARTAMENTO DE MATEMÁTICAS, FACULTAD DE CIENCIAS, UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO, CIUDAD UNIVERSITARIA, MEXICO D.F., 04510, MÉXICO

*E-mail*: guli@servidor.unam.mx  
atamariz@servidor.unam.mx

F. Hernández-Hernández:

FACULTAD DE CIENCIAS FÍSICO-MATEMÁTICAS, UNIVERSIDAD MICHOACANA DE SAN NICOLÁS DE HIDALGO, MORELIA, MICHOACÁN, MÉXICO

*E-mail*: fhernandez@fismat.umich.mx

(Received August 12, 2009, revised March 17, 2010)