

Mihail G. Tkachenko

Cellularity and the index of narrowness in topological groups

Commentationes Mathematicae Universitatis Carolinae, Vol. 52 (2011), No. 2, 309--315

Persistent URL: <http://dml.cz/dmlcz/141492>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2011

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Cellularity and the index of narrowness in topological groups

M. TKACHENKO

Abstract. We study relations between the cellularity and index of narrowness in topological groups and their G_δ -modifications. We show, in particular, that the inequalities $\text{in}((H)_\tau) \leq 2^{\tau \cdot \text{in}(H)}$ and $c((H)_\tau) \leq 2^{2^{\tau \cdot \text{in}(H)}}$ hold for every topological group H and every cardinal $\tau \geq \omega$, where $(H)_\tau$ denotes the underlying group H endowed with the G_τ -modification of the original topology of H and $\text{in}(H)$ is the index of narrowness of the group H .

Also, we find some bounds for the *complexity* of continuous real-valued functions f on an arbitrary ω -narrow group G understood as the minimum cardinal $\tau \geq \omega$ such that there exists a continuous homomorphism $\pi: G \rightarrow H$ onto a topological group H with $w(H) \leq \tau$ such that $\pi \prec f$. It is shown that this complexity is not greater than 2^{2^ω} and, if G is weakly Lindelöf (or 2^ω -steady), then it does not exceed 2^ω .

Keywords: cellularity, G_δ -modification, index of narrowness, ω -narrow, weakly Lindelöf, \mathbb{R} -factorizable, complexity of functions

Classification: 54H11, 54A25, 54C30

1. Introduction

Passing to a subspace of a (compact) space can increase the cellularity of a space. Indeed, for every uncountable cardinal τ , the Tychonoff cube I^τ of weight τ contains a discrete subspace of cardinality τ , while the cellularity of the cube itself is countable. The same happens in (compact) topological groups — it suffices to replace the Tychonoff cube I^τ with \mathbb{T}^τ , where \mathbb{T} is the circle group with the usual multiplication and topology inherited from the complex plane \mathbb{C} .

However, the gap between the cellularity $c(G)$ of a topological group G and the cellularity of subgroups of G becomes considerably smaller. According to [1, Theorem 5.4.11], the inequality $c(H) \leq 2^{c(G)}$ holds for every subgroup H of G . In addition, if G is precompact, then every subgroup of G has countable cellularity.

Another important fact for our study was proved by I. Juhász in [3]: If X is a compact space and γ is a disjoint family of G_δ -sets in X , then the cardinality of γ is at most $2^{c(X)}$. This result shows that the cellularity of the G_δ -modification of X , say $(X)_\omega$, does not exceed $2^{c(X)}$. As usual, by G_δ -modification of X we mean the underlying set X which carries the topology whose base consists of G_δ -sets in X . Similarly, one defines the G_τ -modification of X , for any cardinal $\tau \geq \omega$, which will be denoted by $(X)_\tau$.

Our main concern here is to find a bound for the cellularity of the G_τ -modification of a topological group H in terms of the cellularity of H . This is done in Theorem 3.1 where we show that the inequalities $\text{in}((H)_\tau) \leq 2^{\tau \cdot \text{in}(H)}$ and $c((H)_\tau) \leq 2^{2^{\tau \cdot \text{in}(H)}}$ hold for every topological group H and every cardinal $\tau \geq \omega$, where $\text{in}(H)$ is the index of narrowness of H (see Section 2 below). This means, in particular, that every τ -narrow topological group H satisfies $c((H)_\tau) \leq 2^{2^\tau}$. It turns out that this bound is exact — in Example 3.4 we present an ω -narrow Abelian group H such that $c((H)_\omega) = 2^{2^\omega}$.

A topological group G is called \mathbb{R} -factorizable if every continuous real-valued function f on G can be represented as a composition of a continuous homomorphism of G to a *second countable* group H and a continuous real-valued function on H (see [7, Section 5] or [1, Chapter 8]). In other words, G is \mathbb{R} -factorizable if every continuous real-valued function on G has ‘countable complexity’. By [7, Proposition 5.3], every \mathbb{R} -factorizable group is ω -narrow, but ω -narrow groups need not be \mathbb{R} -factorizable according to [7, Example 5.14]. These facts give rise to the problem of finding bounds for the complexity of continuous real-valued functions on ω -narrow groups (see [6, Problem 3.3] or Problem 4.1 below).

We show in Theorem 4.2 that 2^{2^ω} is such a bound. However, we do not know whether this bound is exact. However, it is shown in Proposition 4.3 that 2^ω is a bound for the complexity of continuous real-valued functions on *weakly Lindelöf* topological groups, while Proposition 4.4 extends this fact to 2^ω -*steady* groups (the terms are explained in the next section).

2. Notation and terminology

Given a topological group G , we define the *index of narrowness* of G , $\text{in}(G)$, as the minimum infinite cardinal τ such that G can be covered by at most τ translates of every neighborhood of the identity. It is easy to verify that $\text{in}(G) \leq c(G)$ for every topological group G , where $c(G)$ is the cellularity of G (see [1, Proposition 5.2.1]). We say that G is τ -*narrow* if it satisfies $\text{in}(G) \leq \tau$.

Suppose that $p: G \rightarrow H$ is a continuous homomorphism. Given a continuous mapping $f: G \rightarrow X$ of the group G to a space X , we write $p \prec f$ if there exists a continuous mapping $h: H \rightarrow X$ such that $f = h \circ p$.

A space X is *weakly Lindelöf* if every open covering of X contains a countable subfamily whose union is dense in X . All Lindelöf spaces as well as all spaces of countable cellularity are weakly Lindelöf. By virtue of [1, Proposition 5.2.8], every weakly Lindelöf topological group is ω -narrow.

A topological group G is called τ -*steady* (see [1, Section 5.6]) if every continuous homomorphic image H of G with $\psi(H) \leq \tau$ satisfies $nw(H) \leq \tau$. By [1, Corollary 5.6.11], every τ -steady topological group is τ -narrow.

The *Nagami number* of a Tychonoff space X is $\text{Nag}(X)$ (see [1, Section 5.3]). Every topological group G with $\text{Nag}(G) \leq \tau$ is τ -steady and the class of τ -steady groups is productive according to [1, Theorem 5.6.4]. It is also clear that a continuous homomorphic image of a τ -steady group is τ -steady.

3. Cellularity and index of narrowness

Let us consider the behavior of the cellularity in topological groups when passing from a group H to $(H)_\omega$ or $(H)_\tau$, for an infinite cardinal τ . It is known that if H is σ -compact or, more generally, a Lindelöf Σ -group, then every family γ of G_δ -sets in H contains a countable subfamily λ such that $\bigcup \lambda$ is dense in $\bigcup \gamma$ (see [9, Theorem 2] or [7, Theorem 4.14]). Further, the cellularity of an ω -bounded group G cannot be greater than 2^ω [7, Theorem 4.29], and this bound is attained even if G is Lindelöf [2, Example 8]. An interesting complement to the former fact was found in [4]: If H is a Lindelöf topological group, then every family γ of G_δ -sets in H contains a subfamily λ with $|\lambda| \leq 2^\omega$ such that $\bigcup \lambda$ is dense in $\bigcup \gamma$. It is an open problem whether this result remains valid for the class of ω -narrow groups [4]. We also recall that if X is a compact space of countable cellularity, then the cellularity of the space $(X)_\omega$ does not exceed 2^ω [3]. It turns out that if H is a τ -narrow topological group, then the cellularity of $(G)_\tau$ does not exceed the second exponent of τ :

Theorem 3.1. *The inequalities $\text{in}((G)_\tau) \leq 2^{\tau \cdot \text{in}(G)}$ and $c((G)_\tau) \leq 2^{2^{\tau \cdot \text{in}(G)}}$ hold for every topological group G and every cardinal $\tau \geq \omega$. In particular, if G is τ -narrow, then $c((G)_\tau) \leq 2^{2^\tau}$.*

PROOF: First we show that $\text{in}((G)_\tau) \leq 2^\lambda$, where $\lambda = \tau \cdot \text{in}(G)$. Let O be a neighbourhood of the identity e in $(G)_\tau$. Then there exists a family $\gamma = \{U_\alpha : \alpha < \tau\}$ of open neighbourhoods of e in G such that $\bigcap \gamma \subseteq O$. By [7, Lemma 3.7], for every $\alpha < \tau$, one can find a continuous homomorphism $p_\alpha : G \rightarrow H_\alpha$ onto a topological group H_α with $w(H_\alpha) \leq \lambda$ and an open neighbourhood V_α of the identity in H_α such that $p_\alpha^{-1}(V_\alpha) \subseteq U_\alpha$. Denote by p the diagonal product of the homomorphisms p_α , $\alpha < \tau$. Then the homomorphism $p : G \rightarrow \prod_{\alpha < \tau} H_\alpha$ is continuous and the group $H = p(G) \subseteq \prod_{\alpha < \tau} H_\alpha$ satisfies $w(H) \leq \lambda$. Therefore, $|H| \leq 2^\lambda$. For every $\alpha < \tau$, there exists a continuous homomorphism $\pi_\alpha : H \rightarrow H_\alpha$ such that $p_\alpha = \pi_\alpha \circ p$. Then $W_\alpha = \pi_\alpha^{-1}(V_\alpha)$ is an open neighbourhood of the identity in H and $p^{-1}(W_\alpha) = p^{-1}\pi_\alpha^{-1}(V_\alpha) = p_\alpha^{-1}(V_\alpha) \subseteq U_\alpha$ for each $\alpha < \tau$. Hence the set $W = \bigcap_{\alpha < \tau} W_\alpha$ contains the identity of H and satisfies $p^{-1}(W) \subseteq O$. In particular, $\ker p \subseteq O$. Since $|H| \leq 2^\lambda$, we can find a subset A of G such that $p(A) = H$ and $|A| \leq 2^\lambda$. Then

$$G = A \cdot \ker p \subseteq A \cdot O \subseteq G,$$

that is, $A \cdot O = G$. This proves the inequality $\text{in}((G)_\tau) \leq 2^\lambda$.

By [7, Theorem 4.29], every topological group K satisfies $c(K) \leq 2^{\text{in}(K)}$. We apply this inequality with $(G)_\tau$ in place of K to conclude that $c((G)_\tau) \leq 2^{2^\lambda}$. \square

Corollary 3.2. *Every topological group G satisfies $c((G)_\tau) \leq 2^{2^{\tau \cdot c(G)}}$. In particular, $c((G)_\omega) \leq 2^{2^{c(G)}}$.*

PROOF: Since $\text{in}(G) \leq c(G)$ by [7, Proposition 3.3(b)], the conclusion follows from Theorem 3.1. \square

Let us show that the upper bounds for the cellularity given in Theorem 3.1 and Corollary 3.2 are exact. First, we need a lemma.

Lemma 3.3. *The free Abelian group $A_{\mathfrak{c}}$ with \mathfrak{c} generators admits a second countable Hausdorff precompact group topology, where $\mathfrak{c} = 2^\omega$.*

PROOF: Denote by \mathcal{T} the maximal precompact group topology on $A_{\mathfrak{c}}$ (i.e., the Bohr topology of $A_{\mathfrak{c}}$, see [1, Section 9.9]). Since $|A_{\mathfrak{c}}| = \mathfrak{c}$, \mathcal{T} contains a weaker metrizable group topology \mathcal{T}_ω by [1, Proposition 9.9.37]. Since every precompact group has countable cellularity, we conclude that the group $K = (A_{\mathfrak{c}}, \mathcal{T}_\omega)$ is Hausdorff, second countable, and precompact. \square

Example 3.4. *There exists a precompact Abelian topological group H such that $c((H)_\omega) = 2^{\mathfrak{c}}$.*

PROOF: We apply Uspenskij's result in [8]: For every infinite cardinal τ , there exists a subgroup G_τ of $(A_{\tau,d})^{2^\tau}$ such that $c(G_\tau) = 2^\tau$, where $A_{\tau,d}$ is the free Abelian group A_τ with τ generators endowed with the discrete topology (the construction in [9] makes the use of the free group F_τ with τ generators instead of A_τ , but a similar argument works as well for A_τ , see [1, Example 5.4.13]).

By Lemma 3.3, the free Abelian group $A = A_{\mathfrak{c}}$ admits a second countable, Hausdorff, precompact group topology \mathcal{T}_ω . Put $K = (A, \mathcal{T}_\omega)$ and $\lambda = 2^{\mathfrak{c}}$. It is clear that $(K)_\omega$ coincides with the discrete group A , say, A_d . Consider the identity isomorphism $\varphi: K^\lambda \rightarrow A_d^\lambda$ and let $H = \varphi^{-1}(G)$, where G is a subgroup of A_d^λ satisfying $c(G) = \lambda$. Then H is precompact being a subgroup of the precompact group K^λ and, therefore, $c(H) \leq \omega$. In addition, $\varphi: (K^\lambda)_\omega \rightarrow (A_d^\lambda)_\omega$ is a topological isomorphism and the topology of $(A_d^\lambda)_\omega$ is finer than that of A_d^λ . Therefore, the restriction of $\varphi: (K^\lambda)_\omega \rightarrow A_d^\lambda$ to the subgroup $(H)_\omega$ of $(K^\lambda)_\omega$ is a continuous isomorphism of $(H)_\omega$ onto G and, hence, $2^{\mathfrak{c}} = c(G) \leq c((H)_\omega)$. On the other hand, $c((H)_\omega) \leq 2^{\mathfrak{c}}$ by Theorem 3.1, so $c((H)_\omega) = 2^{\mathfrak{c}}$. \square

4. Complexity of continuous real-valued functions on ω -narrow groups

Since \mathbb{R} -factorizable groups form a proper subclass of ω -narrow groups, it is natural to consider the following problem (see also [6, Problem 3.3]):

Problem 4.1. *Let G be an ω -narrow topological group and f be a continuous real-valued function on G . Does there exist a continuous homomorphism $\pi: G \rightarrow K$ onto a topological group K with $w(K) \leq 2^\omega$ such that $\pi \prec f$?*

It turns out that the complexity of continuous real-valued functions on ω -narrow topological groups does not exceed $2^{\mathfrak{c}}$, where $\mathfrak{c} = 2^\omega$. We do not know, however, if this bound is exact.

Theorem 4.2. *Let f be a continuous real-valued function on an ω -narrow topological group G . Then there exists a continuous homomorphism $\pi: G \rightarrow H$ onto a topological group H satisfying $w(H) \leq 2^{\mathfrak{c}}$ such that $\pi \prec f$.*

PROOF: By [7, Theorem 4.29], the cellularity of G is not greater than \mathfrak{c} . Hence, according to [1, Theorem 8.1.18], one can find a continuous homomorphism $\varphi: G \rightarrow K$ onto a topological group K with $\psi(K) \leq \mathfrak{c}$ such that $\varphi \prec f$. Take a continuous real-valued function g on K satisfying $f = g \circ \varphi$. Clearly, the group K is ω -narrow as a continuous homomorphic image of the ω -narrow group G . We can now apply [7, Theorem 4.6] according to which $|K| \leq 2^{\text{in}(K) \cdot \psi(K)} \leq 2^{\mathfrak{c}}$. In particular, $nw(K) \leq |K| \leq 2^{\mathfrak{c}}$. Now we use the following weak form of Shakhmatov's theorem in [5] (with $\tau = 2^{\mathfrak{c}}$): If K is a topological group with $nw(K) \leq \tau$ and $g: K \rightarrow \mathbb{R}$ is a continuous function, then there exist a continuous isomorphism $i: K \rightarrow H$ onto a topological group H with $w(H) \leq \tau$ and a continuous function $h: H \rightarrow \mathbb{R}$ such that $g = h \circ i$.

$$\begin{array}{ccc}
 G & \xrightarrow{f} & \mathbb{R} \\
 \varphi \downarrow & \searrow \pi & \uparrow h \\
 K & \xrightarrow{i} & H
 \end{array}$$

Then the continuous homomorphism $\pi = i \circ \varphi$ of G onto H and the function h satisfy the equality $f = h \circ \pi$, i.e., $\pi \prec f$. Since $w(H) \leq 2^{\mathfrak{c}}$, this finishes the proof. □

The following result provides a partial solution to Problem 4.1 in the special case when H is weakly Lindelöf. As usual we denote by \mathfrak{c} the power of the continuum.

Proposition 4.3. *Let $f: G \rightarrow X$ be a continuous mapping, where G is a weakly Lindelöf topological group and X is a Tychonoff space with $w(X) \leq \mathfrak{c}$. Then there exists a continuous homomorphism $\pi: G \rightarrow L$ onto a topological group L with $w(L) \leq \mathfrak{c}$ such that $\pi \prec f$.*

PROOF: Clearly X is homeomorphic to a subspace of $\mathbb{R}^{\mathfrak{c}}$. Taking compositions of f with projections of $\mathbb{R}^{\mathfrak{c}}$ to the factors, we can assume that $X = \mathbb{R}$. Then by [1, Theorem 8.1.18], one can find a continuous homomorphism $\varphi: G \rightarrow K$ onto a topological group K of countable pseudocharacter and a continuous real-valued function $g: K \rightarrow \mathbb{R}$ such that $f = g \circ \varphi$. The group G is ω -narrow since it is weakly Lindelöf [7, Proposition 4.4], so K is also ω -narrow as a continuous homomorphic image of G . Therefore, $|K| \leq 2^{\text{in}(K) \cdot \psi(K)} = \mathfrak{c}$ by [7, Theorem 4.6]. In particular, $nw(K) \leq \mathfrak{c}$. By a theorem in [5], there exist a continuous isomorphism $i: K \rightarrow L$ onto a topological group L with $w(L) \leq \mathfrak{c}$ and a continuous function $h: L \rightarrow \mathbb{R}$ such that $g = h \circ i$. Hence the homomorphism $\pi = i \circ \varphi: G \rightarrow L$ is as required. □

We are now in the position to present another subclass of ω -narrow groups where Problem 4.1 is solved in the affirmative.

Proposition 4.4. *Let G be an ω -narrow topological group. If G is \mathfrak{c} -steady, then for every continuous real-valued function f on G there exists a continuous homomorphism $\pi: G \rightarrow H$ onto a topological group H with $w(H) \leq \mathfrak{c}$ such that $\pi \prec f$.*

PROOF: Given a continuous real-valued function f on G , we can find, as in the proof of Theorem 4.2, a continuous homomorphism $\varphi: G \rightarrow K$ onto a topological group K with $\psi(K) \leq \mathfrak{c}$ and a continuous real-valued function g on K such that $f = g \circ \varphi$. Since G is \mathfrak{c} -steady, the group K satisfies $nw(K) \leq \mathfrak{c}$. Applying Shakhmatov's theorem in [5] once again, we find a continuous isomorphism $i: K \rightarrow H$ of K onto a topological group H with $w(H) \leq \mathfrak{c}$ and a continuous real-valued function h on H such that $g = h \circ i$. Therefore, the continuous homomorphism $\pi = i \circ \varphi$ of G onto H satisfies $\pi \prec f$. \square

5. Open problems

There exist ω -narrow groups H satisfying $c(H) = \mathfrak{c}$ [9]. In fact, there are even Lindelöf groups with the same property [2, Example 8]. We do not know, however, whether large pairwise disjoint families of open sets in ω -narrow groups can be discrete:

Problem 5.1. *Does there exist an ω -narrow topological group which contains a discrete family γ of open sets with $|\gamma| = \mathfrak{c}$?*

Another related problem concerns regular closed subsets of Lindelöf groups:

Problem 5.2. *Is every regular closed subset of a Lindelöf topological group the intersection of at most 2^ω open sets?*

Example 3.4 leaves the following open problem.

Problem 5.3. *Let γ be a family of G_δ -sets in a precompact topological group K . Does there exist a subfamily γ_0 of γ such that $|\gamma_0| \leq \mathfrak{c}$ and $\bigcup \gamma_0$ is dense in $\bigcup \gamma$? What if the group K is ω -narrow?*

REFERENCES

- [1] Arhangel'skii A.V., Tkachenko M.G., *Topological Groups and Related Structures*, Atlantis Studies in Mathematics, Vol. I, Atlantis Press/World Scientific, Paris-Amsterdam, 2008.
- [2] Gartside P., Reznichenko E., Sipacheva O., *Mal'tsev and retral spaces*, Topology Appl. **80** (1997), 115–129.
- [3] Juhász I., *Cardinal Functions in Topology*, Math. Centre Tracts 34, Amsterdam, 1971.
- [4] Pasynkov B.A., *On the relative cellularity of Lindelöf subspaces of topological groups*, Topology Appl. **57** (1994), 249–258.
- [5] Shakhmatov D.B., *On condensations of universal topological algebras preserving continuity of operations and decreasing the weight*, Vestnik Moskov. Univ. Ser. Mat. Mekh. 1984, no. 2, 42–45 (in Russian).

- [6] Tkachenko M.G., *Subgroups, quotient groups and products of \mathbb{R} -factorizable groups*, Topology Proc. **16** (1991), 201–231.
- [7] Tkachenko M.G., *Introduction to topological groups*, Topology Appl. **86** (1998), 179–231.
- [8] V. V. Uspenskij, A topological group generated by a Lindelöf Σ -space has the Souslin property, *Soviet Math. Dokl.* **26** (1982), 166–169.
- [9] Uspenskij V.V., *On continuous images of Lindelöf topological groups*, Soviet Math. Dokl. **32** (1985), 802–806.

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD AUTÓNOMA METROPOLITANA, AV. SAN RAFAEL ATLIXCO 186, COL. VICENTINA, DEL. IZTAPALAPA, C.P. 09340, MEXICO D.F., MEXICO

E-mail: mich@xanum.uam.mx

(Received November 3, 2010)