

J. M. Soriano

Weakly coercive mappings sharing a value

Czechoslovak Mathematical Journal, Vol. 61 (2011), No. 1, 65–72

Persistent URL: <http://dml.cz/dmlcz/141518>

Terms of use:

© Institute of Mathematics AS CR, 2011

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

WEAKLY COERCIVE MAPPINGS SHARING A VALUE

J. M. SORIANO, Sevilla

(Received July 7, 2009)

Abstract. Some sufficient conditions are provided that guarantee that the difference of a compact mapping and a proper mapping defined between any two Banach spaces over \mathbb{K} has at least one zero. When conditions are strengthened, this difference has at most a finite number of zeros throughout the entire space. The proof of the result is constructive and is based upon a continuation method.

Keywords: zero point, continuation method, C^1 -homotopy, surjective implicit function theorem, proper mapping, compact mapping, coercive mapping, Fredholm mapping

MSC 2010: 58C30, 65H10

1. PRELIMINAIRES

Let X and Y be two Banach spaces. If $u: D(F) \subseteq X \rightarrow Y$ is a continuous mapping, then one way of solving the equation

$$(1) \quad u(x) = 0$$

is to embed (1) in a continuum of problems

$$(2) \quad H(x, t) = 0 \quad (0 \leq t \leq 1),$$

which can easily be resolved when $t = 0$. When $t = 1$, the problem (2) becomes (1). In the case when it is possible to continue the solution for all t in $[0, 1]$ then (1) is solved. This method is called continuation with respect to a parameter [1]–[9].

In this paper some sufficient conditions are provided in order to guarantee that the difference of a compact and a proper weakly coercive C^1 -mapping has at least

This work is partially supported by D.G.E.S. Pb 96-1338-CO 2-01 and the Junta de Andalucia.

one zero. If these conditions are conveniently strengthened this difference has at most a finite number of zeros on X . Other conditions, sufficient to guarantee the existence of fixed points, have been given by the author in a finite-dimensional setting, see for example [7], and in an infinite-dimensional setting, see for example [8]. A continuation method was used in the proofs of these papers. The proofs supply the existence of implicitly defined continuous mappings whose ranges reach zero points [1]–[9]. A continuation method is also used here. The key is the use of the surjective implicit function theorem [10], and the properties of proper and Fredholm C^1 -mappings (see [9]).

We briefly recall some theorems and concepts to be used.

Definitions [26], [27]. Henceforth we will assume that X and Y are Banach spaces over \mathbb{K} , where $\mathbb{K} = \mathbb{R}$ or $\mathbb{K} = \mathbb{C}$.

A mapping $F: D(F) \subseteq X \rightarrow Y$ is called *weakly coercive* if and only if $\|F(x)\| \rightarrow \infty$ as $\|x\| \rightarrow \infty$.

A mapping $F: D(F) \subseteq X \rightarrow Y$ is said to be *compact* whenever it is continuous and the image $F(B)$ is relatively compact (i.e. its closure $\overline{F(B)}$ is compact in Y) for every bounded subset $B \subset D(F)$.

A mapping F is said to be *proper* whenever the pre-image $F^{-1}(K)$ of every compact subset $K \subset Y$ is also a compact subset of $D(F)$.

The symbol *dim* means dimension, *codim* means codimension, *ker* means kernel, $R(L)$ stands for the range of the mapping L .

That $L: X \rightarrow Y$ is a *linear Fredholm* mapping means that L is linear and continuous and both the numbers $\dim(\ker(L))$ and $\text{codim}(R(L))$ are finite, and therefore $\ker(L) = X_1$ is a Banach space and has the topological complement X_2 , since $\dim(X_1)$ is finite. The integer number $\text{Ind}(L) = \dim(\ker(L)) - \text{codim}(R(L))$ is called the *index* of L .

Let $F: D(F) \subseteq X \rightarrow Y$. If $D(F)$ is open, then the mapping F is said to be a *Fredholm* mapping if and only if F is a C^1 -mapping and $F'(x): X \rightarrow Y$ is a Fredholm linear mapping for all $x \in D(F)$. If $\text{Ind}(F'(x))$ is constant with respect to $x \in D(F)$, then we call this number the index of F and write it as $\text{Ind}(F)$.

X, Y are called *isomorphic* if and only if there is a linear homeomorphism (*isomorphism*) $L: X \rightarrow Y$.

Let $\mathcal{F}(X, Y)$ denote the set of all linear Fredholm mappings $A: X \rightarrow Y$.

Let $\mathcal{L}(X, Y)$ denote the set of all linear continuous mappings $L: X \rightarrow Y$.

Let $\text{Isom}(X, Y)$ denote the set of all isomorphisms $L: X \rightarrow Y$.

Let $F: D(F) \subseteq X \rightarrow Y$ with $D(F)$ open be a C^1 -mapping. The point $u \in X$ is called a *regular point* of F if and only if $F'(u) \in \mathcal{L}(X, Y)$ maps onto Y . A point

$v \in Y$ is called a *regular value* of F if and only if the pre-image $F^{-1}(v)$ is empty or consists solely of regular points.

Theorem 1. The Surjective Implicit Function Theorem. [10, Section 4–13, Theorem 4–H]. *Let X, Y, Z be Banach spaces over $\mathbb{K} = \mathbb{R}$ or $\mathbb{K} = \mathbb{C}$, and let*

$$F: U(u_0, v_0) \subseteq X \times Y \rightarrow Z$$

be a C^1 -mapping on an open neighbourhood of the point (u_0, v_0) . Suppose that

- (i) $F(u_0, v_0) = 0$, and
- (ii) $F_v(u_0, v_0): Y \rightarrow Z$ is surjective.

Then the following assertion is true:

Let $r > 0$. There is a number $\varrho > 0$ such that, for each given $u \in X$ with $\|u - u_0\| < \varrho$, the equation

$$F(u, v) = 0$$

has a solution v such that $\|v - v_0\| < r$.

Theorem 2 [9, Section 7–9, Theorem 7–33]. *Let $g: D(g) \subseteq X \rightarrow Y$ be a compact mapping, where $a \in D(g)$ and $D(g)$ is open. If the derivative $g'(a)$ exists, then $g'(a) \in \mathcal{L}(X, Y)$ is also a compact mapping.*

Theorem 3 [9, Section 8–4, Example 8–16]. *Let $S \in \mathcal{F}(X, Y)$. The perturbed mapping $S + C$ verifies $S + C \in \mathcal{F}(X, Y)$ and $\text{Ind}(S + C) = \text{Ind}(S)$ provided $C \in \mathcal{L}(X, Y)$ and C is a compact mapping.*

2. WEAKLY COERCIVE MAPPINGS SHARING A VALUE

Clearly, if we define $u := f - g$, then u has a zero if and only if f and g share a value, that is, there is $x \in X$ with $f(x) = g(x)$. We thereby establish our result in terms of f, g .

Theorem 4. *Let $f, g: D \subseteq X \rightarrow Y$ be two C^1 -mappings, where X and Y are two Banach spaces over $\mathbb{K} = \mathbb{R}$ or $\mathbb{K} = \mathbb{C}$, and D is open.*

We assume

- (i) f is a compact mapping, g is a proper mapping and $tf(x) - g(x)$ is weakly coercive, jointly in both coordinates.
- (ii) The mapping g has a zero, x_0 .
- (iii) For any fixed t , belonging to $[0, 1]$, the zero of Y is a regular value of the mapping $tf(x) - g(x)$.

Then the following assertion holds

- (a) f and g share at least one value, i.e., there is at least one x^* such that $f(x^*) = g(x^*)$.

If in addition the condition

- (iv) g is a Fredholm mapping of index zero is satisfied, then in addition we have
- (b) f and g share at most a finite number of values on D , and at least one value.

Proof.

(a) Conclusion (a) will be proved in this section.

(a1) Let us define a mapping

$$H: D \times [0, 1] \subseteq X \times [0, 1] \rightarrow Y, \quad \text{where } H(x, t) := tf(x) - g(x).$$

We will prove here that H is a proper mapping, which will imply that $H^{-1}(0)$ is a compact set, since $\{0\} \subset Y$ is a compact set and H is proper.

Let \mathcal{C} be any fixed compact subset of Y , and let a sequence be fixed such that $(H(x_n, t_n))_{n \geq 1}$ belongs to \mathcal{C} . It suffices to show that the sequence $((x_n, t_n))_{n \geq 1}$ contains a convergent subsequence $((x_{n''}, t_{n''}))_{n'' \geq 1}$, which will imply that $H^{-1}(\mathcal{C})$ is relatively compact, and since

$$(x_{n''}, t_{n''}) \rightarrow (u, t) \quad \text{as } n'' \rightarrow \infty,$$

H is continuous and \mathcal{C} compact, therefore $H(u, t) \in \mathcal{C}$, that is, $(u, t) \in H^{-1}(\mathcal{C})$, and hence $H^{-1}(\mathcal{C})$ is compact.

Since the set \mathcal{C} is bounded and the mapping H is weakly coercive, $((x_n, t_n))_{n \geq 1}$ is a bounded sequence. Consequently $(x_n)_{n \geq 1}$ and $(t_n)_{n \geq 1}$ are bounded sequences. Since f is a compact mapping and $(x_n)_{n \geq 1}$ a bounded sequence, there exists a subsequence $(x_{n'})_{n' \geq 1}$ such that

$$f(x_{n'}) \rightarrow w' \quad \text{as } n' \rightarrow \infty$$

for some $w' \in Y$, and furthermore, since $(t_n)_{n \geq 1}$ is a bounded sequence of real numbers, there is $t \in \mathbb{R}$ with

$$t_{n'} \rightarrow t \quad \text{as } n' \rightarrow \infty.$$

Therefore

$$t_{n'} f(x_{n'}) \rightarrow tw' := w \quad \text{as } n' \rightarrow \infty$$

for some $w \in Y$.

All $H(x_{n'}, t_{n'})$, $n' \geq 1$ lie in the compact set \mathcal{C} and therefore, there is a subsequence $(H(x_{n''}, t_{n''}))_{n'' \geq 1}$ such that

$$H(x_{n''}, t_{n''}) \rightarrow v \quad \text{as} \quad n'' \rightarrow \infty$$

for some $v \in \mathcal{C}$. Therefore

$$g(x_{n''}) \rightarrow w - v \quad \text{as} \quad n'' \rightarrow \infty.$$

Since g is proper, $(x_{n''})$ has a convergent subsequence

$$x_{n'''} \rightarrow u \quad \text{as} \quad n''' \rightarrow \infty.$$

On the other hand,

$$t_{n'''} \rightarrow t \quad \text{as} \quad n''' \rightarrow \infty,$$

hence

$$(x_{n'''}, t_{n'''}) \rightarrow (u, t) \quad \text{as} \quad n''' \rightarrow \infty$$

as required.

(a2) Let us suppose that

$$(3) \quad H(x_a, t_a) = 0$$

for a fixed (x_a, t_a) . We will prove that there is $\varrho > 0$ such that for any $t \in (t_a - \varrho, t_a + \varrho)$, there exists $x = x(t)$ such that $H(x(t), t) = 0$.

Since zero is a regular value for the mappings $\{tf(x) - g(x)\}$ by hypothesis (iii),

$$H_x(x, t) = tf'(x) - g'(x) \in \mathcal{L}(X, Y)$$

is surjective for every pair (x, t) such that $H(x, t) = 0$. In particular, the mapping

$$H_x(x_a, t_a) = t_a f'(x_a) - g'(x_a) \in \mathcal{L}(X, Y)$$

is surjective, which together with identity (3) and Theorem 1 implies the existence of $\varrho > 0, r > 0$ such that for any $t \in (t_a - \varrho, t_a + \varrho)$ there is $x(t)$ with $H(x(t), t) = 0$ and $\|x(t) - x_a\| < r$.

(a3) We will prove that for every t in $[0, 1]$ there exists $x(t)$ such that $H(x(t), t) = 0$.

Let M denote the set of all t such that there is a solution $x(t)$. By assumption (ii) this set is not empty. By (a2), the set is relatively open. Finally, since the set is the projection of $H^{-1}(0)$ into the second component (i.e. the t component), and since the set is the image of a compact set by the continuous function projection,

it is therefore compact, and hence also closed. However, a relatively open, closed, non-empty subset of $[0, 1]$ is the whole interval, since $[0, 1]$ is connected.

Thus

$$H(x(1), 1) = f(x(1)) - g(x(1)) = 0 \Rightarrow f(x(1)) = g(x(1)),$$

which is conclusion (a) of the theorem, where $x^* := x(1)$.

(b) We include the hypothesis (iv) in Section (b).

(b1) We will see that the C^1 -mapping $u: D \subseteq X \rightarrow Y$, $u := f - g$ is a proper Fredholm mapping of index zero.

Since f is a compact mapping, Theorem 2 implies that for each $x \in X$, the mapping $f'(x) \in \mathcal{L}(X, Y)$ is also a compact mapping.

Since $f'(x)$ is a linear compact mapping and given that $g'(x)$ is a linear Fredholm mapping of index zero, Theorem 3 implies that

$$u'(x) = f'(x) - g'(x) \in \mathcal{L}(X, Y), \quad \forall x \in D$$

is a Fredholm linear mapping and $\text{Ind}(u'(x)) = \text{Ind}(g'(x)) = 0, \forall x \in D$. Therefore the non-linear C^1 -mapping u is a Fredholm mapping of index zero.

Furthermore, Section (a1) implies that u is a proper mapping. In fact, this is the particular case in which $t = 1$, i.e. $u(x) = H(x, 1)$.

(b2) We will see that, if any $x \in D$ exists which verifies $u(x) = 0$, then u is a local C^1 -diffeomorphism at x .

Let $x \in D$ exist such that $u(x) = 0$. Since zero is a regular value of u , the linear Fredholm mapping $u'(x)$ maps onto Y , and since $\text{Ind}(u'(x)) = 0$, therefore $\dim(\ker(u'(x))) = 0$. Thus $u'(x) \in \text{Isom}(X, Y)$. By the Local Inverse Theorem [10], u is a local diffeomorphism at x .

(b3) We will prove here that f and g share at most a finite number of values on D .

Since u is proper, $u^{-1}(0)$ is a compact set. If there were an infinite sequence

$$(x_n)_{n \geq 1} \subset D \quad \text{with } x_n \neq x_m \text{ when } n \neq m$$

verifying $u(x_n) = 0, \forall n \in \mathbb{N}$, there would be a subsequence $(x_{n'})_{n' \geq 1} \subset u^{-1}(0)$, which would converge at a point $x \in u^{-1}(0)$, and x would be a non-isolated zero of u . However, since u is a local diffeomorphism at x , given in Section (b2), x is an isolated zero of u . This is a contradiction. Hence there is not an infinite number of zeros of u on D . Thus f and g share at most a finite number of values on D . \square

Example. Let us consider an integral mapping

$$(Au)(x) = \int_a^b F(x, y, u(y)) dy, \quad \forall x \in [a, b],$$

where $-\infty < a < b < +\infty$. Define

$$Q := \{(x, y, u) \in \mathbb{R}^3: x, y \in [a, b] \text{ and } |u| < r \text{ for fixed } r > 0\}.$$

Suppose that the function

$$F: \{(x, y, u) \in \mathbb{R}^3: x, y \in [a, b] \text{ and } |u| \leq r\} \rightarrow \mathbb{R}$$

is twice continuously differentiable. Define $X := C[a, b]$ and $M := \{u \in X: \|u\| < r\}$, where $\|u\| = \max_{a \leq y \leq b} |u(y)|$. It can be easily proved that the mapping $A: M \rightarrow X$ is compact, and it is twice continuously differentiable.

Let B be the mapping $B: M \times [0, 1] \rightarrow X$, $B(u, t) = u$ which is $C^\infty(M \times [0, 1], X)$, is proper, has a zero, and since it is defined only on a bounded set, it is trivially weakly coercive.

The mapping $H(u, t) = t(Au) - (Bu): M \times [0, 1] \rightarrow X$ is weakly coercive, since it is defined only on a bounded set. Theorem 4 implies that if zero is a regular value, then there is $u \in M$ such that

$$(Au)(x) = (Bu)(x), \quad \forall x \in [a, b].$$

If we do not know that zero is a regular value, it is possible to prove the existence of $u \in M$ with $(Au)(x)$ as near to $u(x)$, $\forall x \in [a, b]$ as wanted in the following way:

Since $B'(u, t) \in \mathcal{L}(X \times \mathbb{R}, X)$ is surjective and $\dim(\ker(B'(u, t))) = 1$ for $\forall(u, t) \in M \times [0, 1]$, B is a Fredholm mapping of index one.

Define a mapping $A^*: M \times [0, 1] \rightarrow X$, $A^*(u, t) := t(Au)$ that is differentiable and compact. Theorem 2 implies that $(A^*)'(u, t)$ is compact. Theorem 3 implies that $H'(u, t)$ is a linear Fredholm mapping of index one for $\forall(u, t) \in M \times [0, 1]$, and hence $H(u, t)$ is a Fredholm mapping of index one. Thus we obtain from the Sard-Smale theorem [9, Theorem 4.K] that the set of regular values of the proper mapping H is open and dense in X .

Acknowledgment. I am deeply indebted to the referee for helpful suggestions.

References

- [1] *E. L. Allgower*: A survey of homotopy methods for smooth mappings. Allgower, Glashoff, and Peitgen (eds.) Springer-Verlag, Berlin, 1981, pp. 2–29.
- [2] *E. L. Allgower, K. Glashoff and H. Peitgen (eds.)*: Proceedings of the Conference on Numerical Solution of Nonlinear Equations. Bremen, July 1980, Lecture Notes in Math. 878. Springer-Verlag, Berlin, 1981.
- [3] *E. L. Allgower and K. Georg*: Numerical Continuation Methods. Springer Series in Computational Mathematics 13, Springer-Verlag, New York, 1990.
- [4] *J. C. Alexander and J. A. York*: Homotopy Continuation Method: numerically implementable topological procedures. Trans. Amer. Math. Soc. *242* (1978), 271–284.
- [5] *C. B. Garcia and T. I. Li*: On the number of solutions to polynomial systems of non-linear equations. SIAM J. Numer. Anal. *17* (1980), 540–546.
- [6] *C. B. Garcia and W. I. Zangwill*: Determining all solutions to certain systems of non-linear equations. Math. Operations Research *4* (1979), 1–14.
- [7] *J. M. Soriano*: Global minimum point of a convex function. Appl. Math. Comput. *55* (1993), 213–218.
- [8] *J. M. Soriano*: Continuous embeddings and continuation methods. Nonlinear Anal. Theory Methods Appl. *70* (2009), 4118–4121.
- [9] *E. Zeidler*: Nonlinear Functional Analysis and its applications I. Springer-Verlag, New York, 1992.
- [10] *E. Zeidler*: Applied Functional Analysis. Springer-Verlag, Applied Mathematical Sciences 109, New York, 1995.

Author's address: J. M. Soriano, Departamento de Análisis Matemático, Facultad de Matemáticas, Universidad de Sevilla, Apto. 1160, Sevilla 41080, Spain, e-mail: soriano@us.es.