

Yingbo Han; Shuxiang Feng

The first eigenvalue of spacelike submanifolds in indefinite space form R_p^{n+p}

Archivum Mathematicum, Vol. 47 (2011), No. 2, 77--82

Persistent URL: <http://dml.cz/dmlcz/141555>

Terms of use:

© Masaryk University, 2011

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

THE FIRST EIGENVALUE OF SPACELIKE SUBMANIFOLDS IN INDEFINITE SPACE FORM R_p^{n+p}

YINGBO HAN AND SHUXIANG FENG

ABSTRACT. In this paper, we prove that the first eigenvalue of a complete spacelike submanifold in R_p^{n+p} with the bounded Gauss map must be zero.

1. INTRODUCTION

Let M^n be a complete noncompact Riemannian manifold and $\Omega \subset M^n$ be a domain with compact closure and nonempty boundary $\partial\Omega$. The Dirichlet eigenvalue $\lambda_1(\Omega)$ of Ω is defined by

$$\lambda_1(\Omega) = \inf \left(\frac{\int_{\Omega} |\nabla f|^2 dM}{\int_M f^2 dM} : f \in L_{1,0}^2(\Omega) \setminus \{0\} \right),$$

where dM is the volume element on M^n and $L_{1,0}^2(\Omega)$ the completion of C_0^∞ with respect to the norm

$$\|\varphi\|_\Omega^2 = \int_M \varphi^2 dM + \int_M |\nabla \varphi|^2 dM.$$

If $\Omega_1 \subset \Omega_2$ are bounded domains, then $\lambda_1(\Omega_1) \geq \lambda_1(\Omega_2) \geq 0$. Thus one may define the first Dirichlet eigenvalue of M^n as the following limit

$$\lambda_1(M) = \lim_{r \rightarrow \infty} \lambda_1(B(p, r)) \geq 0,$$

where $B(p, r)$ is the geodesic ball of M^n with radius r centered at p . It is clear that the definition of $\lambda_1(M)$ does not depend on the center point p . It is interesting to ask that for what geometries a noncompact manifold M^n has zero first eigenvalue. Cheng and Yau [1] showed that $\lambda_1(M) = 0$ if M^n has polynomial volume growth.

In [5], B. Wu proved the following result.

Theorem A. *Let M^n be a complete spacelike hypersurface in R_1^{n+1} whose Gauss map is bounded, then $\lambda_1(M) = 0$.*

In this note, we discover that Wu's result still holds for higher codimensional complete spacelike submanifolds in R_p^{n+p} . In fact, we prove

2010 *Mathematics Subject Classification*: primary 53C42; secondary 53B30.

Key words and phrases: spacelike submanifolds, the first eigenvalue.

The first author is supported by NSFC grant No. 10971029 and NSFC-TianYuan Fund. No. 11026062.

Received September 17, 2010. Editor O. Kowalski.

Theorem 3.1. *Let M^n be a complete spacelike submanifold in R_p^{n+p} whose Gauss map is bounded, then $\lambda_1(M) = 0$.*

2. THE GEOMETRY OF PSEUDO-GRASSMANNIAN

In this section we review some basic properties about the geometry of pseudo-Grassmannian. For details one referred to see [6, 3].

Let R_p^{n+p} be the $(n+p)$ -dimensional pseudo-Euclidean space with index p , where, for simplicity, we assume that $n \geq p$. The case $n < p$ can be treated similarly. We choose a pseudo-Euclidean frame field $\{e_1, \dots, e_{n+p}\}$ such that the pseudo-Euclidean metric of R_p^{n+p} is given by $ds^2 = \sum_i (\omega_i)^2 - \sum_\alpha \omega_\alpha = \sum_A \varepsilon_A (\omega_A)^2$, where $\{\omega_1, \dots, \omega_{n+p}\}$ is the dual frame field of $\{e_1, \dots, e_{n+p}\}$, $\varepsilon_i = 1$ and $\varepsilon_\alpha = -1$. Here and in the following we shall use the following convention on the ranges of indices:

$$1 \leq i, j, \dots \leq n; \quad n+1 \leq \alpha, \beta, \dots \leq n+p; \quad 1 \leq A, B, \dots \leq n+p.$$

The structure equations of R_p^{n+p} are given by

$$\begin{aligned} de_A &= - \sum_B \varepsilon_A \omega_{AB} e_B, \\ d\omega_A &= - \sum_B \omega_{AB} \wedge \omega_B, \quad \omega_{AB} + \omega_{BA} = 0, \\ d\omega_{AB} &= - \sum_C \varepsilon_C \omega_{AC} \wedge \omega_{CB}. \end{aligned}$$

Let $G_{n,p}^p$ be the pseudo-Grassmannian of all spacelike n -subspace in R_p^{n+p} , and $\widetilde{G}_{n,p}^p$ be the pseudo-Grassmannian of all timelike p -subspace in R_p^{n+p} . They are specific Cartan-Hadamard manifolds, and the canonical Riemannian metric on $G_{n,p}^p$ and $\widetilde{G}_{n,p}^p$ is

$$ds_G = ds_{\widetilde{G}} = \sum_{i\alpha} (\omega_{\alpha i})^2.$$

Let 0 be the origin of R_p^{n+p} . Let $SO^0(n+p, p)$ denote the identity component of the Lorentzian group $O(n+p, p)$. $SO^0(n+p, p)$ can be viewed as the manifold consisting of all pseudo-Euclidean frames $(0; e_i, e_\alpha)$, and $SO^0(n+p, p)/SO(n) \times SO(p)$ can be viewed as $G_{n,p}^p$ or $\widetilde{G}_{n,p}^p$. Any element in $G_{n,p}^p$ can be represented by a unit simple n -vector $e_1 \wedge \dots \wedge e_n$, while any element in $\widetilde{G}_{n,p}^p$ can be represented by a unit simple p -vector $e_{n+1} \wedge \dots \wedge e_{n+p}$. They are unique up to an action of $SO(n) \times SO(p)$. The Hodge star $*$ provides an one to one correspondence between $G_{n,p}^p$ and $\widetilde{G}_{n,p}^p$. The product \langle, \rangle on $G_{n,p}^p$ for $e_1 \wedge \dots \wedge e_n, v_1 \wedge \dots \wedge v_n$ is defined by

$$\langle e_1 \wedge \dots \wedge e_n, v_1 \wedge \dots \wedge v_n \rangle = \det(\langle e_i, v_j \rangle).$$

The product on $\widetilde{G}_{n,p}^p$ can be defined similarly.

Now we fix a standard pseudo-Euclidean frame e_i, e_α for R_p^{n+p} , and take $g_0 = e_1 \wedge \dots \wedge e_n \in G_{n,p}^p$, $\widetilde{g}_0 = *g_0 = e_{n+1} \wedge \dots \wedge e_{n+p} \in \widetilde{G}_{n,p}^p$. Then we can span the

spacelike n -subspace g in a neighborhood of g_0 by n spacelike vectors f_i :

$$f_i = e_i + \sum_{\alpha} z_{i\alpha} e_{\alpha},$$

where $(z_{i\alpha})$ are the local coordinates of g . By an action of $SO(n) \times SO(p)$ we can assume that

$$(z_{i\alpha}) = \begin{pmatrix} \mu_1 & & & \\ & \ddots & & \\ & & \mu_p & \\ & & 0 & \end{pmatrix}.$$

From [3] we know that the normal geodesic $g(t)$ between g_0 and g has local coordinates

$$(z_{i\alpha}) = \begin{pmatrix} \tanh(\lambda_1 t) & & & \\ & \ddots & & \\ & & \tanh(\lambda_p t) & \\ & & 0 & \end{pmatrix},$$

for real numbers $\lambda_1 \dots \lambda_p$ such that $\sum_{i=1}^p \lambda_i^2 = 1$. This means that $g(t)$ is spanned by $f_1(t) = e_1 + \tanh(\lambda_1 t)e_{n+1}, \dots, f_p(t) = e_p + \tanh(\lambda_p t)e_{n+p}, f_{p+1} = e_{p+1}, \dots, f_n = e_n$. Consequently, $g(t)$ can also be represented by a unit simple n -vector as following:

$$g(t) = (\cosh(\lambda_1 t)e_1 + \sinh(\lambda_1 t)e_{n+1}) \wedge \dots \wedge (\cosh(\lambda_p t)e_1 + \sinh(\lambda_p t)e_{n+p}) \wedge e_{p+1} \wedge \dots \wedge e_n.$$

Set $\lambda_{\alpha} = \lambda_{\alpha-n}$, then it is clear that

$$\begin{aligned} & \cosh(\lambda_1 t)e_1 + \sinh(\lambda_1 t)e_{n+1}, \dots, \cosh(\lambda_p t)e_1 + \sinh(\lambda_p t)e_{n+p}, e_{p+1}, \dots, e_n, \\ & \sinh(\lambda_{n+1} t)e_1 + \cosh(\lambda_{n+1} t)e_{n+1}, \dots, \sinh(\lambda_{n+p} t)e_p + \cosh(\lambda_{n+p} t)e_{n+p} \end{aligned}$$

is again a pseudo-Euclidean frame for R_p^{n+p} , so we have

$$\begin{aligned} \widetilde{g(t)} = *g(t) &= (\sinh(\lambda_{n+1} t)e_1 + \cosh(\lambda_{n+1} t)e_{n+1}) \wedge \dots \wedge (\sinh(\lambda_{n+p} t)e_p \\ &+ \cosh(\lambda_{n+p} t)e_{n+p}) \in \widetilde{G_{n,p}^p}. \end{aligned}$$

Thus we have

$$\langle g_0, g \rangle = (-1)^p \langle *g_0, *g \rangle = (-1)^p \langle \widetilde{g_0}, \widetilde{g} \rangle = \prod_{\alpha} \cosh(\lambda_{\alpha} t).$$

In this note, we also need the following lemma,

Lemma 2.1 ([4]). *Let $\mu_1 \geq 1, \dots, \mu_p \geq 1$ and $\prod_{\alpha} \mu_{\alpha} = C$. Then $\sum_{\alpha} \cosh^2(\lambda_{\alpha}) \leq C^2 + p - 1$, and the equality holds if and only if $\mu_{i_0} = C$ for some $1 \leq i_0 \leq p$ and $\mu_i = 1$ for any $i \neq i_0$.*

3. MAIN RESULTS FOR SPACE-LIKE SUBMANIFOLDS

In this note, we get the following result:

Theorem 3.1. *Let M^n be a complete space-like submanifold in R_p^{n+p} whose Gauss map is bounded, then we have $\lambda_1(M) = 0$.*

Proof. We choose a local frames e_1, \dots, e_{n+p} in R_p^{n+p} such that restricted to M^n , e_1, \dots, e_n are tangent to M^n , e_{n+1}, \dots, e_{n+p} are normal to M^n , the Gauss map is defined by $e_{n+1} \wedge \dots \wedge e_{n+p}: M^n \rightarrow \widetilde{G_{n,p}^p}$. Let us fix p -vector and n -vector $a_{n+1} \wedge \dots \wedge a_{n+p} \in \widetilde{G_{n,p}^p}$, $a_1 \wedge \dots \wedge a_n \in G_{n,p}^p$, where $\langle a_\alpha, a_\beta \rangle = -\delta_{\alpha\beta}$ and $\langle a_i, a_j \rangle = \delta_{ij}$. We defined the projection $\Pi: M^n \rightarrow R_a^n$ by

$$(1) \quad \Pi(x) = x + \sum_{\alpha=n+1}^{n+p} \langle x, a_\alpha \rangle a_\alpha,$$

where \langle, \rangle is the standard indefinite inner product on R_p^{n+p} and R_a^n the totally geodesic Euclidean n -space determined by $a = a_{n+1} \wedge \dots \wedge a_{n+p}$ which is defined by

$$(2) \quad R_a^n = \{x \in R_p^{n+p} : \langle x, a_{n+1} \rangle = \dots = \langle x, a_{n+p} \rangle = 0\}.$$

It is clear from (1) that

$$(3) \quad d\Pi(X) = X + \sum_{\alpha=n+1}^{n+p} \langle X, a_\alpha \rangle a_\alpha$$

for any tangent vector field on M^n and consequently,

$$(4) \quad |d\Pi(X)|^2 = |X|^2 + \sum_{\alpha=n+1}^{n+p} \langle X, a_\alpha \rangle^2.$$

From the equation (4), we know that the map $\Pi: M^n \rightarrow R_a^n$ increases the distance. If a map, from a complete Riemannian manifold M_1 into another Riemannian manifold M_2 of same dimension, increases the distance, then it is a covering map and M_2 is complete (in [2, VIII, Lemma 8.1]). Hence Π is a covering map, but R_a^n being simply connected this means that Π is in fact a diffeomorphism between M^n and R_a^n , and thus M^n is noncompact. Now assume that the Gauss map $e_{n+1} \wedge \dots \wedge e_{n+p}: M^n \rightarrow \widetilde{G_{n,p}^p}$ is bounded, then there exists $\rho > 0$ such that

$$(5) \quad 1 \leq (-1)^p \langle e_{n+1} \wedge \dots \wedge e_{n+p}, a_{n+1} \wedge \dots \wedge a_{n+p} \rangle \leq \rho.$$

From Section 2 we know that by an action of $SO(n) \times SO(p)$ we can assume that

$$\begin{aligned} e_{n+1} &= \sinh(\lambda_{n+1}t)a_1 + \cosh(\lambda_{n+1}t)a_{n+1}, \dots, e_{n+p} \\ &= \sinh(\lambda_{n+p}t)a_1 + \cosh(\lambda_{n+p}t)a_{n+p}, \end{aligned}$$

where $\sum_\alpha \lambda_\alpha^2 = 1$ and $t \in R$.

Write

$$(6) \quad a_\alpha = a^\top - \sum_{\beta=n+1}^{n+p} \langle a_\alpha, e_\beta \rangle e_\beta,$$

where a_α^\top denote the component of a_α which is tangent to M^n , and $\alpha = n+1, \dots, n+p$. Since $\langle a_\alpha, a_\beta \rangle = -\delta_{\alpha\beta}$, we have

$$(7) \quad -1 = |a_\alpha^\top|^2 - \sum_{\beta=n+1}^{n+p} \langle a_\alpha, e_\beta \rangle^2 = |a_\alpha^\top|^2 - \cosh^2(\lambda_\alpha t),$$

where $\alpha = n+1, \dots, n+p$. It follows from Lemma 2.1 and Eq. (5), (7), we have

$$(8) \quad 1 + \sum_{\alpha=n+1}^{n+p} |a_\alpha^\top|^2 = \sum_{\alpha=n+1}^{n+p} \cosh^2(\lambda_\alpha t) - p + 1 \leq \prod \cosh^2(\lambda_\alpha t) \leq \rho^2.$$

From Eq.(4) and (8), we have

$$(9) \quad |d\Pi(X)|^2 = |X|^2 + \sum_{\alpha=n+1}^{n+p} \langle X, a_\alpha^\top \rangle^2 \leq |X|^2 \left(1 + \sum_{\alpha=n+1}^{n+p} |a_\alpha^\top|^2\right) \leq \rho^2 |X|^2.$$

for any tangent vector field on M^n . Let $B(p, r)$ is the geodesic ball of M^n with radius r centered at $p \in M^n$. We claim that $\Pi(B(p, r)) \subset \tilde{B}(\tilde{p}, \rho r)$, where $\tilde{B}(\tilde{p}, \rho r)$ denotes the geodesic ball of R_a^n with radius ρr centered at $\tilde{p} = \Pi(p)$. In fact, for any $\tilde{q} \in \Pi(B(p, r))$ let $q \in B(p, r)$ be the unique point such that $\Pi(q) = \tilde{q}$, and $\gamma: [a, b] \rightarrow M^n$ is the minimal geodesic joining p and q , then from (9) we have

$$\tilde{d}(\tilde{p}, \tilde{q}) \leq L(\Pi \circ r) = \int_a^b |d\Pi(\gamma'(t))| dt \leq \rho \int_a^b |\gamma'(t)| dt = \rho L(\gamma) = \rho d(p, q) \leq \rho r,$$

where \tilde{d} and d denote the distance in R_a^n and M^n , respectively. This prove our claim.

Let dV denotes the n -dimensional volume element on R_a^n . Using (3) and (6) it follows that

$$\begin{aligned} \Pi^*(dV)(X_1, \dots, X_n) &= \det(d\Pi(X_1), \dots, d\Pi(X_n), a_{n+1}, \dots, a_{n+p}) \\ &= \det(X_1, \dots, X_n, a_{n+1}, \dots, a_{n+p}) \\ &= (-1)^p \langle e_{n+1} \wedge \dots \wedge e_{n+p}, a_{n+1} \wedge \dots \wedge a_{n+p} \rangle \\ &\quad \det(X_1, \dots, X_n, e_{n+1}, \dots, e_{n+p}) \\ &= (-1)^p \langle e_{n+1} \wedge \dots \wedge e_{n+p}, a_{n+1} \wedge \dots \wedge a_{n+p} \rangle \\ &\quad dM(X_1, \dots, X_n) \end{aligned}$$

for any tangent vector fields X_1, \dots, X_n of M^n . In other words,

$$(10) \quad \Pi^*(dV) = (-1)^p \langle e_{n+1} \wedge \dots \wedge e_{n+p}, a_{n+1} \wedge \dots \wedge a_{n+p} \rangle dM \geq dM.$$

Since $\Pi(B(p, r)) \subset \tilde{B}(\tilde{p}, \rho r)$ and $\Pi: M^n \rightarrow R_a^n$ is diffeomorphism, it follows from Eq. (10) that

$$\begin{aligned} \rho^n r^n \omega_n &= \text{Vol}(\tilde{B}(\tilde{p}, \rho r)) \geq \text{Vol}(\Pi(B(p, r))) = \int_{\Pi(B(p, r))} dV \\ (11) \quad &= \int_{B(p, r)} \Pi^* dV \geq \int_{B(p, r)} dM = \text{Vol}(B(p, r)), \end{aligned}$$

where ω_n denotes the volume of unit ball in Euclidean n -space. (11) means that the order of the volume growth of M^n is not larger than n , thus by [1] we see that $\lambda_1(M) = 0$. \square

REFERENCES

- [1] Cheng, S. Y., Yau, S. T., *Differential equations on Riemannian manifolds and geometric applications*, Comm. Pure Appl. Math. **28** (1975), 333–354.
- [2] Kobayashi, S., Nomizu, K., *Foundations of differential geometry*, John Wiley & Sons, Inc., 1969.
- [3] Wong, Y. C., *Euclidean n -space in pseudo-Euclidean spaces and differential geometry of Cartan domain*, Bull. Amer. Math. Soc. (N.S.) **75** (1969), 409–414.
- [4] Wu, B. Y., *On the volume and Gauss map image of spacelike submanifolds in de Sitter space form*, J. Geom. Phys. **53** (2005), 336–344.
- [5] Wu, B. Y., *On the first eigenvalue of spacelike hypersurfaces in Lorentzian space*, Arch. Math. (Brno) **42** (2006), 233–238.
- [6] Xin, Y. L., *A rigidity theorem for spacelike graph of higher codimension*, Manuscripta Math. **103** (2000), 191–202.

COLLEGE OF MATHEMATICS AND INFORMATION SCIENCE,
XINYANG NORMAL UNIVERSITY,
XINYANG 464000, HENAN, P. R. CHINA
E-mail: yingbhan@yahoo.com.cn fsxhyb@yahoo.com.cn