

Qingliu Yao

Solvability of a class of elastic beam equations with strong Carathéodory nonlinearity

Applications of Mathematics, Vol. 56 (2011), No. 6, 543--555

Persistent URL: <http://dml.cz/dmlcz/141766>

Terms of use:

© Institute of Mathematics AS CR, 2011

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

SOLVABILITY OF A CLASS OF ELASTIC BEAM EQUATIONS
WITH STRONG CARATHÉODORY NONLINEARITY

QINGLIU YAO, Nanjing

(Received January 26, 2009)

Abstract. We study the existence of a solution to the nonlinear fourth-order elastic beam equation with nonhomogeneous boundary conditions

$$\begin{cases} u^{(4)}(t) = f(t, u(t), u'(t), u''(t), u'''(t)), & \text{a.e. } t \in [0, 1], \\ u(0) = a, u'(0) = b, u(1) = c, u''(1) = d, \end{cases}$$

where the nonlinear term $f(t, u_0, u_1, u_2, u_3)$ is a strong Carathéodory function. By constructing suitable height functions of the nonlinear term $f(t, u_0, u_1, u_2, u_3)$ on bounded sets and applying the Leray-Schauder fixed point theorem, we prove that the equation has a solution provided that the integration of some height function has an appropriate value.

Keywords: nonlinear ordinary differential equation, boundary value problem, existence, fixed point theorem

MSC 2010: 34B15, 34B16

1. INTRODUCTION

In this paper we consider the nonlinear elastic beam equation with nonhomogeneous Dirichlet boundary conditions on the left and nonhomogeneous Navier boundary conditions on the right

$$(P) \quad \begin{cases} u^{(4)}(t) = f(t, u(t), u'(t), u''(t), u'''(t)), & \text{a.e. } t \in [0, 1], \\ u(0) = a, u'(0) = b, u(1) = c, u''(1) = d. \end{cases}$$

In the homogeneous case, (P) corresponds to the elastic beam equation rigidly fixed at the left and simply supported at the right.

Throughout this paper, $\|u\| = \max_{0 \leq t \leq 1} |u(t)|$ for $u \in C[0, 1]$ and

$$p(t) = \frac{1}{4}(2a + 2b - 2c + d)t^3 + \frac{1}{4}(6c - 6a - 6b - d)t^2 + bt + a.$$

Thus, $p(0) = a$, $p'(0) = b$, $p(1) = c$, $p''(1) = d$, $p'''(t) \equiv 3a + 3b - 3c + \frac{3}{2}d$.

Denote $\gamma_i = \max_{0 \leq t \leq 1} |p^{(i)}(t)|$, $i = 0, 1, 2, 3$, and

$$\begin{aligned} \mu_0 &= \max_{0 \leq t \leq 1} p(t), & \nu_0 &= \min_{0 \leq t \leq 1} p(t), & \mu_1 &= \max_{0 \leq t \leq 1} p'(t), & \nu_1 &= \min_{0 \leq t \leq 1} p'(t), \\ \mu_2 &= \max_{0 \leq t \leq 1} p''(t), & \nu_2 &= \min_{0 \leq t \leq 1} p''(t), & \mu_3 &= \nu_3 = 3a + 3b - 3c + \frac{3}{2}d. \end{aligned}$$

Let us introduce constants

$$\begin{aligned} K &= \frac{65536}{39 + 55\sqrt{33}} \approx 184.6340, & M &= \frac{3}{17 - 12\sqrt{2}} \approx 101.9117, \\ k_0 &= 1, & k_1 &= \frac{65536}{45(39 + 55\sqrt{33})} \approx 4.1030, & k_2 &= \frac{8192}{39 + 55\sqrt{33}} \approx 23.0793, \\ & & k_3 &= \frac{40960}{39 + 55\sqrt{33}} \approx 115.3963, \\ m_0 &= 1, & m_1 &= \frac{1}{9[17 - 12\sqrt{2}]} \approx 3.7745, & m_2 &= \frac{\sqrt{3}}{3[17 - 12\sqrt{2}]} \approx 19.6129, \\ & & m_3 &= \frac{3}{17 - 12\sqrt{2}} \approx 101.9117. \end{aligned}$$

In addition, let $G(t, s)$ be the Green function of the linear homogeneous boundary value problem $u^{(4)}(t) = 0$, $0 \leq t \leq 1$, $u(0) = u'(0) = u(1) = u''(1) = 0$, that is,

$$G(t, s) = \begin{cases} \frac{1}{12}(1-t)s^2[3(1-s) - (1-t)^2(3-s)], & 0 \leq s \leq t \leq 1, \\ \frac{1}{12}t^2(1-s)[3(1-t) - (1-s)^2(3-t)], & 0 \leq t \leq s \leq 1. \end{cases}$$

Clearly, $G(t, s) > 0$, $0 < t, s < 1$.

The nonlinear beam equation (P) has been studied by some authors when $f: [0, 1] \times \mathbb{R}^4 \rightarrow \mathbb{R}$ is continuous, for example, see [1], [4], [5], [7], [9], [10], [11], [14]. Particularly, Agarwal [1] introduced the maximum height

$$\Phi(r_{0-3}) = \max\{|f(t, u_0, u_1, u_2, u_3)|: 0 \leq t \leq 1, |u_i| \leq 2r_i, i = 0, 1, 2, 3\}$$

of the nonlinear term $f(u_0, u_1, u_2, u_3)$ on the bounded set $[0, 1] \times [-2r_0, 2r_0] \times [-2r_1, 2r_1] \times [-2r_2, 2r_2] \times [-2r_3, 2r_3]$ and proved the following local existence theorem.

Theorem 1.1. *Suppose that $f: [0, 1] \times \mathbb{R}^4 \rightarrow \mathbb{R}$ is continuous and there exist four positive numbers $r_i > 0$, $i = 0, 1, 2, 3$, such that $r_i \geq \gamma_i$ and*

$$\Phi(r_{0-3}) \leq K \min\{k_i^{-1}r_i, i = 0, 1, 2, 3\}.$$

Then equation (P) has at least one solution $u^ \in C^3[0, 1]$ and $\|(u^*)^{(i)}\| \leq 2r_i$, $i = 0, 1, 2, 3$.*

Theorem 1.1 shows that the equation (P) has a solution if there exist four positive numbers $r_i > 0$, $i = 0, 1, 2, 3$, such that the maximum height $\Phi(r_{0-3})$ has an appropriate value.

The aim of this paper is to study the existence of a solution to the equation (P) when $f: [0, 1] \times \mathbb{R}^4 \rightarrow \mathbb{R}$ is a strong Carathéodory function. Here, $f: [0, 1] \times \mathbb{R}^4 \rightarrow \mathbb{R}$ is referred to as a strong Carathéodory function if

- (C1) for a.e. $t \in [0, 1]$, $f(t, \cdot, \cdot, \cdot, \cdot): \mathbb{R}^4 \rightarrow \mathbb{R}$ is continuous;
- (C2) for all $(u_0, u_1, u_2, u_3) \in \mathbb{R}^4$, $f(\cdot, u_0, u_1, u_2, u_3): [0, 1] \rightarrow \mathbb{R}$ is measurable;
- (C3) for every $r > 0$ there exists a nonnegative function $j_r \in L^1[0, 1]$, such that

$$|f(t, u_0, u_1, u_2, u_3)| \leq j_r(t), \quad (t, u_0, u_1, u_2, u_3) \in [0, 1] \times [-r, r]^4.$$

In the ordinary way, $f(t, u_0, u_1, u_2, u_3)$ is called a Carathéodory function if $f: [0, 1] \times \mathbb{R}^4 \rightarrow \mathbb{R}$ only satisfies the conditions (C1) and (C2).

If $f: (0, 1) \times \mathbb{R}^4 \rightarrow \mathbb{R}$ is continuous and satisfies the assumption (C3), then it is a strong Carathéodory function. In this case, the function $f(t, u_0, u_1, u_2, u_3)$ may be singular at $t = 0$, $t = 1$.

Therefore, we allow that there exists a zero measure set $E \subset [0, 1]$ such that the nonlinear term $f(t, u_0, u_1, u_2, u_3)$ is discontinuous or singular at $t \in E$.

It is impossible to describe the growth change of a singular function on a bounded set by the maximum height $\Phi(r_{0-3})$. Therefore, we must introduce new tool in order to deal with the singular equation (P).

In this paper, we introduce a height function $\varphi(t, r_{0-3})$ to estimate the growth change of the nonlinear term $f(t, u_0, u_1, u_2, u_3)$ on bounded set. Here, the height function is

$$\varphi(t, r_{0-3}) = \max\{|f(t, u_0, u_1, u_2, u_3)|: -r_i + \nu_i \leq u_i \leq r_i + \mu_i, i = 0, 1, 2, 3\}.$$

When $f: [0, 1] \times \mathbb{R}^4 \rightarrow \mathbb{R}$ is a strong Carathéodory function, the height function $\varphi(\cdot, r_{0-3})$ is integrable on $[0, 1]$.

In geometry, the height function $\varphi(t, r_{0-3})$ is the maximum height of the nonlinear term $f(t, u_0, u_1, u_2, u_3)$ on the set $\{t\} \times [-r_0 + \nu_0, r_0 + \mu_0] \times [-r_1 + \nu_1, r_1 + \mu_1] \times [-r_2 + \nu_2, r_2 + \mu_2] \times [-r_3 + \nu_3, r_3 + \mu_3]$.

By applying the height function, we obtain a new result. More precisely, we prove the following local existence theorem.

Theorem 1.2. *Suppose that $f: [0, 1] \times \mathbb{R}^4 \rightarrow \mathbb{R}$ is a strong Carathéodory function and there exist four positive numbers $r_i > 0$, $i = 0, 1, 2, 3$, such that one of the following conditions is satisfied:*

- (1) $\max_{0 \leq t \leq 1} \int_0^1 |(\partial^i / \partial t^i) G(t, s)| \varphi(s, r_{0-3}) ds \leq r_i$, $i = 0, 1, 2, 3$,
- (2) $\int_0^1 \varphi(t, r_{0-3}) dt \leq M \min\{m_i^{-1} r_i, i = 0, 1, 2, 3\}$.

Then the equation (P) has at least one solution $u^ \in C^3[0, 1]$ and $\|(u^*)^{(i)} - p^{(i)}\| \leq r_i$, $i = 0, 1, 2, 3$. In addition, the solution u^* is nontrivial if $a^2 + b^2 + c^2 + d^2 > 0$ or $\int_0^1 |f(t, 0, 0, 0, 0)| dt > 0$.*

Theorem 1.2 shows that the equation (P) has a solution if there exist four positive numbers $r_i > 0$, $i = 0, 1, 2, 3$, such that the integration of the height function $\varphi(t, r_{0-3})$ has an appropriate value.

Obviously, Theorem 1.2 can deal with more complex cases than Theorem 1.1. We will verify that Theorem 1.2 extends Theorem 1.1 and illustrate that the extension is true by an example. The localization idea of this work comes from the papers [2], [8], [12], [13], [14], [15].

2. PRELIMINARIES

After direct computations, we get

$$\max_{0 \leq t, s \leq 1} G(t, s) = G(2 - \sqrt{2}, 2 - \sqrt{2}) = \frac{1}{3}[17 - 12\sqrt{2}].$$

Computing the partial derivatives of $G(t, s)$ in t , we have

$$\begin{aligned} \frac{\partial}{\partial t} G(t, s) &= \begin{cases} \frac{1}{4}t(1-s)[2-3t-(2-t)(1-s)^2], & 0 \leq t \leq s \leq 1, \\ \frac{1}{4}s^2[(1-t)^2(3-s)-(1-s)], & 0 \leq s \leq t \leq 1; \end{cases} \\ \frac{\partial^2}{\partial t^2} G(t, s) &= \begin{cases} \frac{1}{2}(1-s)[1-3t-(1-t)(1-s)^2], & 0 \leq t \leq s \leq 1, \\ \frac{1}{2}s^2(1-t)(s-3), & 0 \leq s \leq t \leq 1; \end{cases} \\ \frac{\partial^3}{\partial t^3} G(t, s) &= \begin{cases} \frac{1}{2}(1-s)(s^2-2s-2), & 0 \leq t < s \leq 1, \\ \frac{1}{2}s^2(3-s), & 0 \leq s < t \leq 1. \end{cases} \end{aligned}$$

This yields

$$\begin{aligned} \max_{0 \leq t, s \leq 1} \left| \frac{\partial}{\partial t} G(t, s) \right| &= \left| \frac{\partial}{\partial t} G\left(1, \frac{2}{3}\right) \right| = \frac{1}{27}, \\ \max_{0 \leq t, s \leq 1} \left| \frac{\partial^2}{\partial t^2} G(t, s) \right| &= \left| \frac{\partial^2}{\partial t^2} G\left(0, \frac{3 - \sqrt{3}}{3}\right) \right| = \frac{\sqrt{3}}{9}, \\ \sup_{0 \leq t, s \leq 1} \left| \frac{\partial^3}{\partial t^3} G(t, s) \right| &= \lim_{s \rightarrow 1} \left| \frac{\partial^3}{\partial t^3} G(1, s) \right| = 1. \end{aligned}$$

Therefore,

$$\begin{aligned} M &= \frac{3}{17 - 12\sqrt{2}} = \left[\max_{0 \leq t, s \leq 1} G(t, s) \right]^{-1}, \\ m_1 &= \frac{1}{9[17 - 12\sqrt{2}]} = M \max_{0 \leq t, s \leq 1} \left| \frac{\partial}{\partial t} G(t, s) \right|, \\ m_2 &= \frac{\sqrt{3}}{3[17 - 12\sqrt{2}]} = M \max_{0 \leq t, s \leq 1} \left| \frac{\partial^2}{\partial t^2} G(t, s) \right|, \\ m_3 &= \frac{3}{17 - 12\sqrt{2}} = M \max_{0 \leq t, s \leq 1} \left| \frac{\partial^3}{\partial t^3} G(t, s) \right|. \end{aligned}$$

In this paper, $C^3[0, 1]$ is the Banach space with the norm

$$\|u\| = \max\{\|u\|, \|u'\|, \|u''\|, \|u'''\|\}.$$

If $u \in C^3[0, 1]$ and the assumption (C3) holds, then there exists a nonnegative function $j_{\|u\|+1} \in L^1[0, 1]$ such that

$$|f(t, u(t), u'(t), u''(t), u'''(t))| \leq j_{\|u\|+1}(t), \quad 0 \leq t \leq 1.$$

Define an operator T as follows, for $u \in C^3[0, 1]$ and $0 \leq t \leq 1$:

$$(Tu)(t) = \int_0^1 G(t, s) f(s, u(s) + p(s), u'(s) + p'(s), u''(s) + p''(s), u'''(s) + p'''(s)) ds.$$

For convenience, we use the abbreviation

$$F(t, u(t) + p(t)) = f(t, u(t) + p(t), u'(t) + p'(t), u''(t) + p''(t), u'''(t) + p'''(t)).$$

Thus

$$(Tu)(t) = \int_0^1 G(t, s) F(s, u(s) + p(s)) ds, \quad 0 \leq t \leq 1.$$

For $u \in C^3[0, 1]$ and $0 \leq t \leq 1$, let

$$(Au)(t) = f(t, u(t) + p(t), u'(t) + p'(t), u''(t) + p''(t), u'''(t) + p'''(t)),$$

$$(Bu)(t) = \int_0^1 G(t, s)u(s) \, ds,$$

$$(Cu)(t) = \int_0^1 \frac{\partial}{\partial t} G(t, s)u(s) \, ds,$$

$$(Du)(t) = \int_0^1 \frac{\partial^2}{\partial t^2} G(t, s)u(s) \, ds,$$

$$(Eu)(t) = \int_0^1 \frac{\partial^3}{\partial t^3} G(t, s)u(s) \, ds.$$

Proposition 2.1. *The operator $A: C^3[0, 1] \rightarrow L^1[0, 1]$ is continuous.*

Proof. Let $u \in C^3[0, 1]$. Since $u + p \in C^3[0, 1]$, there exist four sequences $\{\xi_n^i\}_{n=1}^\infty$, $i = 0, 1, 2, 3$, of simple functions such that

$$\lim_{n \rightarrow \infty} \xi_n^i(t) = u^{(i)}(t) + p^{(i)}(t), \quad \text{a.e. } t \in [0, 1], \quad i = 0, 1, 2, 3$$

(see (11.35), [6]). By the assumption (C2), $f(t, \xi_n^0(t), \xi_n^1(t), \xi_n^2(t), \xi_n^3(t))$ is a measurable function on $[0, 1]$ for each $n = 1, 2, \dots$. By the assumption (C1),

$$\lim_{n \rightarrow \infty} f(t, \xi_n^0(t), \xi_n^1(t), \xi_n^2(t), \xi_n^3(t)) = (Au)(t), \quad \text{a.e. } t \in [0, 1].$$

Since the limit of measurable functions is measurable (see (11.18), [6]), we assert that $(Au)(t)$ is a measurable function on $[0, 1]$.

By making use of the above abbreviation, we have

$$(Au)(t) = F(t, u(t) + p(t)), \quad 0 \leq t \leq 1.$$

Let $r_u = \|u\| + \|p\| + 1$. Since

$$\max\{\|u + p\|, \|u' + p'\|, \|u'' + p''\|, \|u''' + p'''\|\} \leq \|u\| + \|p\| + 1$$

and (C3), there exists $j_{r_u} \in L^1[0, 1]$ such that

$$|F(t, u(t) + p(t))| \leq j_{r_u}(t), \quad 0 \leq t \leq 1.$$

Hence $Au \in L^1[0, 1]$. It follows that $A: C^3[0, 1] \rightarrow L^1[0, 1]$.

Let $u_n \in C^3[0, 1]$ and $\|u_n - u\| \rightarrow 0$. Then $\lim_{n \rightarrow \infty} u_n^{(i)}(t) = u^{(i)}(t)$, $0 \leq t \leq 1$, $i = 0, 1, 2, 3$. By (C1),

$$\lim_{n \rightarrow \infty} F(t, u_n(t) + p(t)) = F(t, u(t) + p(t)), \quad \text{a.e. } t \in [0, 1].$$

Since $\|u_n - u\| \rightarrow 0$, there exists a positive integer $N(u)$ such that, for any $n \geq N(u)$, $\|u_n + p\| \leq \|u\| + \|p\| + 1$. So, for any $n \geq N(u)$,

$$|F(t, u_n(t) + p(t))| \leq j_{r_u}(t), \quad 0 \leq t \leq 1.$$

By the Lebesgue dominated convergence theorem (see (12.24), [6]), we have

$$\begin{aligned} & \lim_{n \rightarrow \infty} \int_0^1 |(Au_n)(t) - (Au)(t)| dt \\ &= \lim_{n \rightarrow \infty} \int_0^1 |F(t, u_n(t) + p(t)) - F(t, u(t) + p(t))| ds \\ &= \int_0^1 \lim_{n \rightarrow \infty} |F(t, u_n(t) + p(t)) - F(t, u(t) + p(t))| ds = 0. \end{aligned}$$

It follows that $A: C^3[0, 1] \rightarrow L^1[0, 1]$ is continuous. \square

Proposition 2.2. $B, C, D, E: L^1[0, 1] \rightarrow C[0, 1]$ are completely continuous operators.

Proof. Obviously, $B, C, D, E: L^1[0, 1] \rightarrow C[0, 1]$ are bounded linear operators. Applying the Arzela-Ascoli theorem, we prove easily that B, C, D, E are completely continuous operators. \square

Proposition 2.3. For any $u \in L^1[0, 1]$ and a.e. $t \in [0, 1]$,

$$(Bu)'(t) = (Cu)(t), \quad (Cu)'(t) = (Du)(t), \quad (Du)'(t) = (Eu)(t).$$

Proof. By the generalized mean value theorem in the nonsmooth analysis (see Theorem 2.3.7, [3]), we have, for every $i = 0, 1, 2$,

$$\left| \frac{\partial^i}{\partial t^i} G(t + \Delta t, s) - \frac{\partial^i}{\partial t^i} G(t, s) \right| \leq \max_{0 \leq t, s \leq 1} \left| \frac{\partial^{i+1}}{\partial t^{i+1}} G(t, s) \right| |\Delta t| \leq |\Delta t|.$$

From this inequality we obtain the estimate, for $i = 0, 1, 2$,

$$\left| \frac{1}{\Delta t} \left[\frac{\partial^i}{\partial t^i} G(t + \Delta t, s) - \frac{\partial^i}{\partial t^i} G(t, s) \right] u(s) \right| \leq |u(s)|, \quad \text{a.e. } t \in [0, 1].$$

Since $u \in L^1[0, 1]$, the Lebesgue dominated convergence theorem yields

$$\begin{aligned}
 (Bu)'(t) &= \lim_{\Delta t \rightarrow 0} \frac{1}{\Delta t} [(Bu)(t + \Delta t) - (Bu)(t)] \\
 &= \lim_{\Delta t \rightarrow 0} \frac{1}{\Delta t} \int_0^1 [G(t + \Delta t, s) - G(t, s)] u(s) \, ds \\
 &= \int_0^1 \lim_{\Delta t \rightarrow 0} \frac{1}{\Delta t} [G(t + \Delta t, s) - G(t, s)] u(s) \, ds \\
 &= \int_0^1 \frac{\partial}{\partial t} G(t, s) u(s) \, ds = (Cu)(t).
 \end{aligned}$$

Similarly, $(Cu)'(t) = (Du)(t)$, $(Du)'(t) = (Eu)(t)$. □

Lemma 2.4. $T: C^3[0, 1] \rightarrow C^3[0, 1]$ is a completely continuous operator.

Proof. By Proposition 2.3, we have the compositions

$$T = B \circ A, \quad (T(\cdot))' = C \circ A, \quad (T(\cdot))'' = D \circ A, \quad (T(\cdot))''' = E \circ A.$$

By Proposition 2.1 and 2.2, we assert that $T, (T(\cdot))', (T(\cdot))'', (T(\cdot))''' : C^3[0, 1] \rightarrow C[0, 1]$ are completely continuous operators. Therefore, $T: C^3[0, 1] \rightarrow C^3[0, 1]$ is a completely continuous operator. □

3. THE PROOF OF THEOREM 1.2

Let $V_{r_{0-3}} = \{u \in C^3[0, 1]: \|u^{(i)}\| \leq r_i, i = 0, 1, 2, 3\}$. We need to prove $T: V_{r_{0-3}} \rightarrow V_{r_{0-3}}$.

If $u \in V_{r_{0-3}}$ then $\|u^{(i)}\| \leq r_i, i = 0, 1, 2, 3$. This implies that, for $0 \leq t \leq 1$,

$$\begin{aligned}
 -r_0 + \nu_0 &\leq u(t) + p(t) \leq r_0 + \mu_0, \\
 -r_1 + \nu_1 &\leq u'(t) + p'(t) \leq r_1 + \mu_1, \\
 -r_2 + \nu_2 &\leq u''(t) + p''(t) \leq r_2 + \mu_2, \\
 -r_3 + \nu_3 &\leq u'''(t) + p'''(t) \leq r_3 + \mu_3.
 \end{aligned}$$

By the definition of $\varphi(t, r_{0-3})$, we have

$$|F(t, u(t) + p(t))| \leq \varphi(t, r_{0-3}), \quad t \in [0, 1].$$

Applying this fact and the assumption (1), we get

$$\begin{aligned}
\|Tu\| &\leq \max_{0 \leq t \leq 1} \int_0^1 G(t, s) |F(s, u(s) + p(s))| ds \\
&\leq \max_{0 \leq t \leq 1} \int_0^1 G(t, s) \varphi(s, r_{0-3}) ds \leq r_0, \\
\|(Tu)'\| &\leq \max_{0 \leq t \leq 1} \int_0^1 \left| \frac{\partial}{\partial t} G(t, s) \right| |F(s, u(s) + p(s))| ds \\
&\leq \max_{0 \leq t \leq 1} \int_0^1 \left| \frac{\partial}{\partial t} G(t, s) \right| \varphi(s, r_{0-3}) ds \leq r_1, \\
\|(Tu)''\| &\leq \max_{0 \leq t \leq 1} \int_0^1 \left| \frac{\partial^2}{\partial t^2} G(t, s) \right| |F(s, u(s) + p(s))| ds \\
&\leq \max_{0 \leq t \leq 1} \int_0^1 \left| \frac{\partial^2}{\partial t^2} G(t, s) \right| \varphi(s, r_{0-3}) ds \leq r_2, \\
\|(Tu)'''\| &\leq \max_{0 \leq t \leq 1} \int_0^1 \left| \frac{\partial^3}{\partial t^3} G(t, s) \right| |F(s, u(s) + p(s))| ds \\
&\leq \max_{0 \leq t \leq 1} \int_0^1 \left| \frac{\partial^3}{\partial t^3} G(t, s) \right| \varphi(s, r_{0-3}) ds \leq r_3.
\end{aligned}$$

Consequently, $T: V_{r_{0-3}} \rightarrow V_{r_{0-3}}$.

If the assumption (2) holds, then

$$\begin{aligned}
\max_{0 \leq t \leq 1} \int_0^1 G(t, s) \varphi(s, r_{0-3}) ds &\leq \max_{0 \leq t, s \leq 1} G(t, s) \int_0^1 \varphi(s, r_{0-3}) ds \\
&\leq m_0 M^{-1} \cdot M m_0^{-1} r_0 = r_0, \\
\max_{0 \leq t \leq 1} \int_0^1 \left| \frac{\partial}{\partial t} G(t, s) \right| \varphi(s, r_{0-3}) ds &\leq \max_{0 \leq t, s \leq 1} \left| \frac{\partial}{\partial t} G(t, s) \right| \int_0^1 \varphi(s, r_{0-3}) ds \\
&\leq m_1 M^{-1} \cdot M m_1^{-1} r_1 = r_1, \\
\max_{0 \leq t \leq 1} \int_0^1 \left| \frac{\partial^2}{\partial t^2} G(t, s) \right| \varphi(s, r_{0-3}) ds &\leq \max_{0 \leq t, s \leq 1} \left| \frac{\partial^2}{\partial t^2} G(t, s) \right| \int_0^1 \varphi(s, r_{0-3}) ds \\
&\leq m_2 M^{-1} \cdot M m_2^{-1} r_2 = r_2, \\
\max_{0 \leq t \leq 1} \int_0^1 \left| \frac{\partial^3}{\partial t^3} G(t, s) \right| \varphi(s, r_{0-3}) ds &\leq \max_{0 \leq t, s \leq 1} \left| \frac{\partial^3}{\partial t^3} G(t, s) \right| \int_0^1 \varphi(s, r_{0-3}) ds \\
&\leq m_3 M^{-1} \cdot M m_3^{-1} r_3 = r_3.
\end{aligned}$$

Therefore, the assumption (1) holds and $T: V_{r_{0-3}} \rightarrow V_{r_{0-3}}$.

Now, we prove that the equation (P) has a solution.

Since $V_{r_{0-3}}$ is a bounded, closed and convex set, by Lemma 2.4 and the Leray-Schauder fixed point theorem there exists $\bar{u} \in V_{r_{0-3}}$ such that $T\bar{u} = \bar{u}$. Let $u^* = \bar{u} + p$. Then $u^* \in C^3[0, 1]$, $\|(u^*)^{(i)} - p^{(i)}\| \leq r_i$, $i = 0, 1, 2, 3$, and

$$\begin{aligned} u^*(t) - p(t) &= \bar{u}(t) = (T\bar{u})(t) \\ &= \int_0^1 G(t, s) f(s, \bar{u}(s) + p(s), \bar{u}'(s) + p'(s), \bar{u}''(s) + p''(s), \bar{u}'''(s) + p'''(s)) \, ds \\ &= \int_0^1 G(t, s) f(s, u^*(s), (u^*)'(s), (u^*)''(s), (u^*)'''(s)) \, ds. \end{aligned}$$

Let $e = p'''(t) \equiv 3a + 3b - 3c + \frac{3}{2}d$. Then, for $0 \leq t \leq 1$,

$$\begin{aligned} (u^*)''' &= e + \int_0^1 \frac{\partial^3}{\partial t^3} G(t, s) f(s, u^*(s), (u^*)'(s), (u^*)''(s), (u^*)'''(s)) \, ds \\ &= e + \frac{1}{2} \int_0^t s^2(3-s)F(s, u^*(s)) \, ds \\ &\quad + \frac{1}{2} \int_t^1 (1-s)(s^2 - 2s - 2)F(s, u^*(s)) \, ds. \end{aligned}$$

Applying the properties of the indefinite integral (see (18.3), [6]), we get

$$(u^*)^{(4)}(t) = f(t, u^*(t), (u^*)'(t), (u^*)''(t), (u^*)'''(t)), \quad \text{a.e. } t \in [0, 1].$$

Noticing that $G(0, s) = G(1, s) = (\partial/\partial t)G(0, s) = (\partial^2/\partial t^2)G(1, s) = 0$, $0 \leq s \leq 1$, we have $u^*(0) = p(0) = a$, $(u^*)'(0) = p'(0) = b$, $u^*(1) = p(1) = c$, $(u^*)''(1) = p''(1) = d$. Therefore, u^* is a solution of the equation (P).

If $a^2 + b^2 + c^2 + d^2 > 0$ or $\int_0^1 |f(t, 0, 0, 0, 0)| \, dt > 0$, then $u^*(t) \not\equiv 0$. In other words, the solution u^* is nontrivial.

The proof is completed. □

4. REMARK AND EXAMPLE

R e m a r k 4.1. Theorem 1.1 is a special case of Theorem 1.2 (1).

Assume that the conditions of Theorem 1.1 are satisfied. Then $f: [0, 1] \times \mathbb{R}^4 \rightarrow \mathbb{R}$ is continuous and there exist four positive numbers $r_i \geq \gamma_i$, $i = 0, 1, 2, 3$, such that $\Phi(r_{0-3}) \leq K \min\{k_i^{-1}r_i, i = 0, 1, 2, 3\}$.

By the definitions, $|\mu_i| \leq \gamma_i$ and $|\nu_i| \leq \gamma_i$, $i = 0, 1, 2, 3$. So,

$$\begin{aligned} &[-r_0 + \nu_0, r_0 + \mu_0] \times [-r_1 + \nu_1, r_1 + \mu_1] \times [-r_2 + \nu_2, r_2 + \mu_2] \times [-r_3 + \nu_3, r_3 + \mu_3] \\ &\quad \subset [-2r_0, 2r_0] \times [-2r_1, 2r_1] \times [-2r_2, 2r_2] \times [-2r_3, 2r_3]. \end{aligned}$$

Using this fact, we get

$$\varphi(t, r_{0-3}) \leq \Phi(r_{0-3}) \leq K \min\{k_i^{-1}r_i, i = 0, 1, 2, 3\}, \quad 0 \leq t \leq 1.$$

From [1] we have $\max_{0 \leq t \leq 1} \int_0^1 |(\partial^i / \partial t^i)G(t, s)| ds \leq K^{-1}k_i, i = 0, 1, 2, 3$. It follows that, for $i = 0, 1, 2, 3$,

$$\begin{aligned} \max_{0 \leq t \leq 1} \int_0^1 |(\partial^i / \partial t^i)G(t, s)| \varphi(s, r_{0-3}) ds &\leq K k_i^{-1} r_i \max_{0 \leq t \leq 1} \int_0^1 |(\partial^i / \partial t^i)G(t, s)| ds \\ &\leq K k_i^{-1} r_i \cdot K^{-1} k_i = r_i. \end{aligned}$$

By Theorem 1.2 (1), the equation (P) has one solution $u^* \in C^3[0, 1]$ and $\|(u^*)^{(i)} - p^{(i)}\| \leq r_i, i = 0, 1, 2, 3$. So,

$$\|(u^*)^{(i)}\| \leq \|(u^*)^{(i)} - p^{(i)}\| + \|p^{(i)}\| \leq r_i + \gamma_i \leq 2r_i.$$

Theorem 1.1 is proved.

Example 4.2. Consider the nonlinear beam equation with homogeneous boundary condition

$$\begin{cases} u^{(4)}(t) = \frac{(u^2(t) + 2)e^{\frac{3}{2}((u''(t) + u'''(t)) \cos(u'(t) + \pi t))}}{180\sqrt{|t - \frac{1}{2}|}}, & t \in [0, \frac{1}{2}) \cup (\frac{1}{2}, 1], \\ u(0) = u'(0) = u(1) = u''(1) = 0. \end{cases}$$

In this equation, $p = 0$ and the nonlinear term is

$$f(t, u_0, u_1, u_2, u_3) = \frac{(u_0^2 + 2)e^{\frac{3}{2}(u_2 + u_3) \cos(u_1 + \pi t)}}{180\sqrt{|t - \frac{1}{2}|}}.$$

Thus, $f(t, u_0, u_1, u_2, u_3)$ is a strong Carathéodory function and it is discontinuous on the set $\{\frac{1}{2}\} \times \mathbb{R}$.

Let $r_0 = r_1 = r_2 = r_3 = 1$. Since $\mu_i = \nu_i = 0, i = 0, 1, 2, 3$, we have

$$\begin{aligned} \varphi(t, r_{0-3}) &\leq \sup \left\{ \left| \frac{(u_0^2 + 2)e^{\frac{3}{2}(u_2 + u_3) \cos(u_1 + \pi t)}}{180\sqrt{|t - \frac{1}{2}|}} \right| : \begin{array}{l} -1 \leq u_0 \leq 1, \quad -1 \leq u_1 \leq 1, \\ -1 \leq u_2 \leq 1, \quad -1 \leq u_3 \leq 1 \end{array} \right\} \\ &\leq \frac{e^3}{60\sqrt{|t - \frac{1}{2}|}}. \end{aligned}$$

Further,

$$\int_0^1 \varphi(t, r_{0-3}) dt \leq \frac{e^3}{60} \int_0^1 \frac{dt}{\sqrt{|t - \frac{1}{2}|}} = \frac{e^3}{30} \int_0^{1/2} \frac{dt}{\sqrt{t}} = \frac{\sqrt{2}e^3}{30}$$

$$\approx 0.9468 < 1 = Mm_3^{-1} = M \min\{m_i^{-1}r_i, i = 0, 1, 2, 3\}.$$

By Theorem 1.2(2), the equation has one solution $u^* \in C^3[0, 1]$ such that $\|(u^*)^{(i)}\| \leq 1$, $i = 0, 1, 2, 3$. Since $f(t, 0, 0, 0, 0) = \frac{2}{180}|t - \frac{1}{2}|^{-1/2} \cos \pi t \neq 0$, $t \in [0, \frac{1}{2}) \cup (\frac{1}{2}, 1]$, the solution u^* is nontrivial.

The conclusion cannot be derived from Theorem 1.1 and the existing literature because of the singularity of the nonlinear term.

The example illustrates that Theorem 1.2 is an effective and practical tool for the discontinuous equation (P).

References

- [1] *R. P. Agarwal*: On fourth order boundary value problems arising in beam analysis. *Differ. Integral Equ.* *2* (1989), 91–110.
- [2] *R. P. Agarwal, D. O'Regan, V. Lakshmikantham*: Singular $(p, n - p)$ focal (n, p) higher order boundary value problems. *Nonlinear Anal., Theory Methods Appl.* *42* (2000), 215–228.
- [3] *F. H. Clarke*: Optimization and Nonsmooth Analysis. John Wiley & Sons, New York, 1983.
- [4] *M. B. M. Elgindi, Z. Guan*: On the global solvability of a class of fourth-order nonlinear boundary value problems. *Int. J. Math. Math. Sci.* *20* (1997), 257–262.
- [5] *C. P. Gupta*: Existence and uniqueness theorems for the bending of an elastic beam equation. *Appl. Anal.* *26* (1988), 289–304.
- [6] *E. Hewitt, K. Stromberg*: Real and Abstract Analysis. Springer, Berlin-Heidelberg-New York, 1975.
- [7] *R. Ma*: Existence and uniqueness theorems for some fourth-order nonlinear boundary value problems. *Int. J. Math. Math. Sci.* *23* (2000), 783–788.
- [8] *D. O'Regan*: Singular Dirichlet boundary value problems. I: Superlinear and nonresonant case. *Nonlinear Anal., Theory Methods Appl.* *29* (1997), 221–245.
- [9] *Q. Yao*: A local existence theorem for nonlinear elastic beam equations fixed at left and simply supported at right. *J. Nat. Sci. Nanjing Norm. Univ.* *8* (2006), 1–4.
- [10] *Q. Yao*: Existence of n positive solutions to a singular beam equation rigidly fixed at left and simply supported at right. *J. Zhengzhou Univ., Nat. Sci. Ed.* *40* (2008), 1–5.
- [11] *Q. Yao*: Positive solutions of nonlinear elastic beam equation rigidly fastened on the left and simply supported on the right. *Nonlinear Anal., Theory Methods Appl.* *69* (2008), 1570–1580.
- [12] *Q. Yao*: Solution and positive solution to a class of semilinear third-order two-point boundary value problem. *Appl. Math. Lett.* *17* (2004), 1171–1175.
- [13] *Q. Yao*: Solvability of a fourth-order beam equation with all-order derivatives. *Southeast Asian Bull. Math.* *32* (2008), 563–571.
- [14] *Q. Yao*: Solvability of singular beam equations fixed at left and simply supported at right. *J. Lanzhou Univ., Nat. Sci.* *44* (2008), 115–118. (In Chinese.)

- [15] *Q. Yao*: Successive iteration and positive solution for a discontinuous third-order boundary value problem. *Comput. Math. Appl.* 53 (2007), 741–749.
- [16] *J. Wang*: Solvability of singular nonlinear two-point boundary value problems. *Nonlinear Anal., Theory Methods Appl.* 24 (1995), 555–561.

Author's address: *Qingliu Yao*, Department of Applied Mathematics, Nanjing University of Finance and Economics, Nanjing 210003, P. R. China, e-mail: yaoqingliu2002@hotmail.com.