

Applications of Mathematics

Hans-Görg Roos; Martin Schopf

Analysis of finite element methods on Bakhvalov-type meshes for linear convection-diffusion problems in 2D

Applications of Mathematics, Vol. 57 (2012), No. 2, 97--108

Persistent URL: <http://dml.cz/dmlcz/142030>

Terms of use:

© Institute of Mathematics AS CR, 2012

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

ANALYSIS OF FINITE ELEMENT METHODS ON
BAKHVALOV-TYPE MESHES FOR LINEAR
CONVECTION-DIFFUSION PROBLEMS IN 2D*

HANS-GÖRG ROOS, Dresden, MARTIN SCHOPF, Dresden

(Received January 22, 2010)

Abstract. So far optimal error estimates on Bakhvalov-type meshes are only known for finite difference and finite element methods solving linear convection-diffusion problems in the one-dimensional case. We prove (almost) optimal error estimates for problems with exponential boundary layers in two dimensions.

Keywords: finite element method, singular perturbation, convection-diffusion problem, Bakhvalov-type meshes, layer-adapted meshes

MSC 2010: 65N30

1. INTRODUCTION

We shall examine the finite element method for the numerical solution of the singularly perturbed linear elliptic boundary value problem

$$(1.1a) \quad Lu \equiv -\varepsilon \Delta u + b \cdot \nabla u + cu = f \quad \text{in } \Omega = (0, 1) \times (0, 1),$$

$$(1.1b) \quad u = 0 \quad \text{on } \partial\Omega,$$

where $\varepsilon \ll 1$ is a small positive parameter, b , c and f are smooth. Assuming

$$(1.2) \quad -b = (-b_1, -b_2) > (\beta_1, \beta_2) > 0 \quad \text{on } \bar{\Omega}$$

with constants β_1, β_2 , the solution of (1.1) typically has exponential boundary layers at $x = 0$ and $y = 0$ and a corner layer at $(0, 0)$. Additionally weak corner singularities

*This paper was written during a stay of the first author at the Charles University in Prague supported by the Nečas Center for Mathematical Modeling.

could exist but we assume still some compatibility such that the problem has a classical solution with $u \in C^{3,\alpha}(\bar{\Omega})$. Without loss of generality we can as well assume

$$c - \frac{1}{2} \operatorname{div} b \geq c_0 > 0,$$

where c_0 is a constant.

We want to solve (1.1) with linear or bilinear finite elements on a layer-adapted mesh. Let us introduce the ε -weighted $H^1(\Omega)$ norm by

$$\|v\|_\varepsilon^2 := \varepsilon |v|_1^2 + \|v\|_0^2.$$

First, for Shishkin meshes (see Section 2 for a detailed discussion of various meshes) it was proved that

$$(1.3) \quad \|u - u^N\|_\varepsilon \leq CN^{-1} \ln N,$$

for bilinear elements in [11], for linear elements in [1]. Here $N + 1$ is the number of mesh points used in every coordinate direction to define a tensor-product mesh, thus the degrees of freedom are of order $O(N^2)$. We remark that throughout the paper C will denote a generic positive constant that is independent of ε and of the mesh.

Later in [8] it was proved for Shishkin-type meshes (generalizing a result from [3]) that

$$(1.4) \quad \|u - u^N\|_\varepsilon \leq CN^{-1} \max |\psi'|,$$

here ψ denotes the mesh characterizing function (see Section 2). For a Bakhvalov-Shishkin mesh with $\psi(t) = 1 - 2(1 - N^{-1})t$ and a modified Bakhvalov-Shishkin mesh due to Vulanovic with $\psi(t) = \exp(-t/(q - t))$ and $q = 1/2 + 1/(2 \ln N)$ the factor $|\psi'|$ is uniformly bounded for $t \in [0, 1/2]$. Consequently, these meshes are examples of optimal meshes with

$$(1.5) \quad \|u - u^N\|_\varepsilon \leq CN^{-1}.$$

For Bakhvalov-type meshes, however, it is an open question whether or not (1.5) holds in 2D. For the one-dimensional case, see [9]. The ingredients used in the proof in [9] cannot be used in 2D, therefore in this paper we present a new approach for verifying (1.5).

Throughout the paper we shall assume $|\varepsilon \ln \varepsilon| \leq CN^{-1}$ as in general is satisfied for discretizations of convection-dominated problems.

2. LAYER-ADAPTED MESHES AND SOLUTION DECOMPOSITION

Let N , our discretization parameter, be an even positive integer. We introduce the mesh points

$$0 = x_0 < x_1 < \dots < x_{N-1} < x_N = 1, \quad 0 = y_0 < y_1 < \dots < y_{N-1} < y_N = 1$$

and consider a tensor-product mesh with mesh points (x_i, y_j) . Because both meshes have the same structure we only describe the meshes in the x -direction (for the mesh in the y -direction take $\beta_1 := \beta_2$).

The mesh is graded in $[0, x_{N/2}]$ but equidistant in $[x_{N/2}, 1]$. The graded part of the mesh is based on a mesh generating function φ with $\varphi(0) = 0$, $\varphi(1/2) = \ln(1/\varepsilon)$, moreover we assume φ to be continuous, monotonically increasing and differentiable. Set

$$(2.1) \quad x_i = \begin{cases} \frac{\sigma\varepsilon}{\beta_1} \varphi(t_i) & \text{with } t_i = i/N & \text{for } i = 0, 1, \dots, N/2, \\ 1 - (1 - x_{N/2})2(N - i)/N & \text{for } i = N/2 + 1, \dots, N. \end{cases}$$

Here σ is some positive constant which characterizes the order of the smallness of the layer term in $x_{N/2}$. A Bakhvalov-type mesh (B-mesh) is given by

$$(2.2) \quad \varphi(t) := -\ln[1 - 2(1 - \varepsilon)t].$$

Remark 1. For Shishkin-type meshes (S-type meshes), introduced and analyzed in [8], we require $\varphi(1/2) = \ln N$. Especially, a Bakhvalov-Shishkin mesh (B-S-mesh) is given by $\varphi(t) := -\ln[1 - 2(1 - N^{-1})t]$; Shishkin's original mesh, however, by the definition $\varphi(t) := 2(\ln N)t$. For a survey concerning layer-adapted meshes see [6].

Following [8] the mesh characterizing function ψ is defined by

$$(2.3) \quad \psi := \exp(-\varphi).$$

Consequently, we have for $t \in [0, 1/2]$

$$\psi(t) = \begin{cases} 1 - 2(1 - N^{-1})t & \text{for a B-S-mesh,} \\ 1 - 2(1 - \varepsilon)t & \text{for a B-mesh.} \end{cases}$$

Now in [8] the following result is proved:

If $\sigma = 2$ and ψ satisfies

$$(2.4) \quad \frac{\max |\psi'|}{\psi} \leq CN,$$

then for linear or bilinear elements the finite element error can be estimated by

$$\|u - u^N\|_\varepsilon \leq CN^{-1} \max |\psi'|.$$

On a B-S-mesh the condition (2.4) is satisfied but on a Bakhvalov-type mesh it is not. Therefore we shall present in Section 3 a modification of the analysis of [8] which allows to handle Bakhvalov-type meshes.

We assume that the exact solution of (1.1) can be decomposed as follows:

$$(2.5) \quad u = S + E_1 + E_2 + E_{12} = S + E.$$

The smooth part S is characterized by bounds uniform with respect to ε for certain derivatives, while E_1 , E_2 , E_{12} describe the layers at $x = 0$, $y = 0$ and the corner layer at $(0, 0)$, respectively. Precisely we suppose

$$(2.6a) \quad \left| \frac{\partial^{i+j} S}{\partial x^i \partial y^j} (x, y) \right| \leq C,$$

$$(2.6b) \quad \left| \frac{\partial^{i+j} E_1}{\partial x^i \partial y^j} (x, y) \right| \leq C \varepsilon^{-i} e^{-\beta_1 x / \varepsilon},$$

$$(2.6c) \quad \left| \frac{\partial^{i+j} E_2}{\partial x^i \partial y^j} (x, y) \right| \leq C \varepsilon^{-j} e^{-\beta_2 y / \varepsilon},$$

$$(2.6d) \quad \left| \frac{\partial^{i+j} E_{12}}{\partial x^i \partial y^j} (x, y) \right| \leq C \varepsilon^{-(i+j)} e^{-\beta_1 x / \varepsilon} e^{-\beta_2 y / \varepsilon},$$

for all $(x, y) \in \bar{\Omega}$ and $0 \leq i + j \leq 2$. See [10, Theorem III.1.26] for conditions that guarantee the existence of such a decomposition.

These pointwise estimates bounds are stronger than needed, see Remark 3.108 in [10].

3. ERROR ESTIMATION ON A BAKHVALOV-TYPE MESH

Let us study the discretization of the problem (1.1) with bilinear or linear (draw additionally diagonals to decompose rectangles into triangles) finite elements on the Bakhvalov-type mesh characterized by (2.1) with (2.2).

First let us notice

$$x_{N/2-1} = -\frac{\sigma \varepsilon}{\beta_1} \ln \left(\varepsilon + \frac{2(1-\varepsilon)}{N} \right)$$

and draw the important conclusion

$$(3.1) \quad |E_1(x_{N/2-1}, \cdot)| \leq c \left(\varepsilon + \frac{2(1-\varepsilon)}{N} \right)^\sigma \leq cN^{-\sigma}.$$

Moreover, the condition (2.4) is satisfied on the interval $[0, x_{N/2-1}]$ because ψ' is uniformly bounded and $1.5N^{-1}$ is a lower bound for the minimal value of ψ .

Next we observe that the largest mesh size of the graded part of the Bakhvalov-type mesh is $h_{N/2} = x_{N/2} - x_{N/2-1}$ which satisfies

$$(3.2) \quad h_{N/2} = \frac{\sigma\varepsilon}{\beta_1} \ln\left(1 + \frac{2(1-\varepsilon)}{\varepsilon N}\right) \geq \kappa\varepsilon$$

(the positive constant κ depends on σ , β_1 and the constant C in $\varepsilon \leq CN^{-1}$).

Remark 2. Instead of the Bakhvalov-type mesh given by (2.1) with (2.2) we could also study generalized Bakhvalov-type meshes given by (2.1) with (2.3). The mesh characterizing function ψ should be monotonically decreasing and differentiable with $\psi(0) = 1$ and $\psi(1/2) = \varepsilon$. In the analysis which follows we need the following three ingredients: the smallness of E_1 in the sense of (3.1) for $x \geq x_{N/2-1}$, condition (2.4) and $h_{N/2} \geq \kappa\varepsilon$. These three properties are guaranteed if there exist positive constants μ_1, μ_2 such that

$$\mu_1 N^{-1} \leq \psi(1/2 - N^{-1}) \leq \mu_2 N^{-1}$$

for $\varepsilon \leq CN^{-1}$.

Next we decompose Ω into 4 subdomains as illustrated in Fig. 1: $\bar{\Omega} = \Omega_{11} \cup \Omega_{12} \cup \Omega_{21} \cup \Omega_{22}$.




Figure 1. Subregions of Ω .

Additionally we introduce

$$\Omega_{22}^* := [x_{N/2}, 1] \times [y_{N/2}, 1],$$

in Ω_{22}^* our mesh is uniform with mesh size of order $O(N^{-1})$.

Let us denote the piecewise linear or bilinear interpolant of a given continuous function w by w^I . Then our convergence analysis is based on the following results for the interpolation error:

Lemma 1. *Let us assume that u allows the decomposition (2.6) and a Bakhvalov-type mesh is given with $\sigma \geq 2$. Then*

$$(3.3a) \quad \|E - E^I\|_{\infty, \Omega_{22}} \leq CN^{-2}, \quad \|E - E^I\|_{0, \Omega \setminus \Omega_{22}} \leq C\varepsilon^{1/2}N^{-1},$$

$$(3.3b) \quad \varepsilon^{1/2}|E - E^I|_1 \leq CN^{-1},$$

$$(3.3c) \quad N^{-1}|S - S^I|_1 + \|S - S^I\|_0 \leq CN^{-2}.$$

Proof. For the smooth part S the estimates are standard applications of the known anisotropic interpolation estimates.

Concerning the layer components let us, for instance, consider E_1 and start to estimate the interpolation error. While in Ω_{11} and Ω_{12} the result is well known (see [8] or [2]), in Ω_{22} and Ω_{21} the pointwise smallness of E_1 is used in combination with $\|E^I\|_{\infty} \leq C\|E\|_{\infty}$ and $|\text{meas}(\Omega_{21})| \leq C\varepsilon \ln N$.

Next consider $\|(E_1 - E_1^I)_x\|_0$, for instance. We have

$$\int_{\Omega} (E_1 - E_1^I)_x^2 \leq \int_{x \leq x_{N/2-1}} (E_1 - E_1^I)_x^2 + 2T$$

with

$$T = \int_{x_{N/2-1} \leq x \leq x_{N/2}} (E_1^I)_x^2 + \int_{x \geq x_{N/2}} (E_1^I)_x^2 + \int_{x \geq x_{N/2-1}} (E_1)_x^2.$$

Again, the estimate for $x \leq x_{N/2-1}$ is well known (see [8] or [2]). To estimate the integral with E_1 we just use the estimate in the solution decomposition, in $x \geq x_{N/2}$ we apply an inverse inequality combined with the smallness of E_1 to estimate the second term of T .

The first term of T is bounded by

$$\left| \int_{x_{N/2-1} \leq x \leq x_{N/2}} (E_1^I)_x^2 \right| \leq C \frac{1}{h_{N/2}^2} h_{N/2} \|E_1^I\|_{\infty, x \geq x_{N/2-1}}^2 \leq C \frac{N^{-4}}{\varepsilon},$$

because $h_{N/2} \geq \kappa\varepsilon$. Summarizing, the result follows. \square

With linear or bilinear finite elements on our given Bakhvalov-type mesh and the corresponding finite element space $V^N \subset H_0^1(\Omega)$, the finite element method reads:

Find $u^N \in V^N$ with

$$(3.4) \quad a(u^N, v) = (f, v) \quad \forall v \in V^N.$$

The bilinear form $a(\cdot, \cdot)$ is given by

$$a(w, v) := \varepsilon(\nabla w, \nabla v) + (b \cdot \nabla w + cw, v);$$

due to our assumptions the bilinear form is uniformly V -elliptic with respect to the ε -weighted H^1 norm: one has

$$a(w, w) \geq \alpha \|w\|_\varepsilon^2 \quad \text{for all } w \in H_0^1(\Omega)$$

with some positive α independent of ε .

With the Lagrange interpolant $u^I \in V^N$ of u studied already in the previous lemma we introduce the splitting of the error into the components

$$(3.5) \quad \eta := u^I - u, \quad v^N := u^I - u^N$$

and start the error estimate from

$$(3.6) \quad \alpha \|u^I - u^N\|_\varepsilon^2 \leq a(u^I - u^N, u^I - u^N) = a(u^I - u, u^I - u^N) = a(\eta, v^N).$$

Write

$$a(\eta, v^N) = \varepsilon(\nabla \eta, \nabla v^N) + (b \cdot \nabla \eta, v^N) + (c\eta, v^N).$$

Based on Lemma 1 it is easy to estimate the first and the third term of that representation just using the Cauchy-Schwarz inequality.

It is important that for the crucial convection term we can split the interpolation error into $(S - S^I) + (E - E^I)$ and, using integration by parts based on $v^N \in H_0^1(\Omega)$, estimate

$$(b \cdot \nabla(S - S^I), v^N) \quad \text{and} \quad (E - E^I, b \cdot \nabla v^N).$$

Again Cauchy-Schwarz and Lemma 1 yield immediately

$$|(b \cdot \nabla(S - S^I), v^N)| \leq CN^{-1} \|v^N\|_0 \leq CN^{-1} \|v^N\|_\varepsilon.$$

Finally we have to estimate the convective term for a layer part E . Set

$$T = \int_{\Omega} (E - E^I) b \cdot \nabla v^N.$$

Next we split T in several parts:

$$\begin{aligned} T &= \int_{\Omega_{22}^*} (E - E^I) b \cdot \nabla v^N + \int_{\Omega \setminus \Omega_{22}} (E - E^I) b \cdot \nabla v^N \\ &\quad + \int_{\Omega_{22} \setminus \Omega_{22}^*} (E - E^I) b \cdot \nabla v^N. \end{aligned}$$

Then we get applying an inverse inequality on Ω_{22}^*

$$|T_1| = \left| \int_{\Omega_{22}^*} (E - E^I) b \cdot \nabla v^N \right| \leq CN^{-1} \|v^N\|_0.$$

For the integral on $\Omega \setminus \Omega_{22}$ we use (3.3a):

$$|T_2| = \left| \int_{\Omega \setminus \Omega_{22}} (E - E^I) b \cdot \nabla v^N \right| \leq C\varepsilon^{1/2} N^{-1} |v^N|_1.$$

Finally we get

$$\begin{aligned} |T_3| &= \left| \int_{\Omega_{22} \setminus \Omega_{22}^*} (E - E^I) b \cdot \nabla v^N \right| \\ &\leq C \|E - E^I\|_{\infty, \Omega_{22}} (\text{meas}(\Omega_{22} \setminus \Omega_{22}^*))^{1/2} |v^N|_1. \end{aligned}$$

Introducing

$$Q(N, \varepsilon) := \max \left\{ 1, N^{-1} \left(\ln \frac{1}{\varepsilon} \right)^{1/2} \right\},$$

we proved finally with

$$|T_3| \leq CQ(N, \varepsilon) N^{-1} \varepsilon^{1/2} |v^N|_1$$

the following result:

Theorem 1. *Let us assume that u allows the decomposition (2.6) and a Bakhvalov-type mesh is given with $\sigma \geq 2$. Suppose, additionally, $|\varepsilon \ln \varepsilon| \leq CN^{-1}$. Then the finite element error on the given Bakhvalov-type mesh satisfies the almost optimal estimate*

$$(3.7) \quad \|u - u^N\|_\varepsilon \leq CQ(N, \varepsilon) N^{-1}.$$

Remark that practically $Q(N, \varepsilon)$ is bounded: If we assume $N \geq 10$ and $\varepsilon \geq 10^{-100}$, then

$$Q(N, \varepsilon) \leq \sqrt{\ln 10}.$$

4. REMARKS TO SUPERCLOSENESS AND SDFEM

For bilinear elements on Shishkin-type meshes one can prove for $\sigma \geq 2.5$ and a decomposition of the solution with bounded derivatives up to order three the supercloseness result

$$(4.1) \quad \|u^I - u^N\|_\varepsilon \leq CN^{-2}(\max |\psi'| + \ln^{1/4} N)^2,$$

see [4] and [2]. For our Bakhvalov-type mesh we can derive a similar result:

$$(4.2) \quad \|u^I - u^N\|_\varepsilon \leq CQ(N, \varepsilon)N^{-2} \ln^{1/2} N.$$

The following ingredients are used to prove that estimate:

- The use of the Lin identities for improving the estimates for

$$(b \cdot \nabla(S - S^I), v^N) \quad \text{and} \quad (\nabla(S - S^I), \nabla v^N)$$

analogously as in [4] and [2].

- The use of the Lin identities to estimate, e.g., the expression $\varepsilon(\nabla(E_1 - E_1^I), \nabla v^N)$ in $\Omega_{11} \cup \Omega_{12}$ analogously as in [4] and [2].
- The use of the smallness of E_1 for $x \geq x_{N/2-1}$ to improve the order with respect to N^{-1} in the estimate for $\varepsilon^{1/2}|E_1 - E_1^I|_1$ in that part of the domain.
- For the convective term and the layer components we use:
 - (i) in Ω_{22}^* simple the smallness with respect to N^{-1} and an inverse inequality as before,
 - (ii) in $\Omega \setminus \Omega_{22}$ an improved version with respect to the order of N^{-1} of (3.3a),
 - (iii) in $\Omega_{22} \setminus \Omega_{22}^*$ the same technique as before (but E_1 is smaller due to the choice of σ).

Based on the supercloseness result for the Galerkin method it is possible to analyze streamline diffusion stabilization (see [12] for Shishkin meshes and [2] for general S-type meshes). Let us add to the Galerkin bilinear form the stabilization term

$$a_{\text{stab}}(w, v) := N^{-1} \sum_{T \subset \Omega_{22}^*} (-\varepsilon \Delta w + b \cdot \nabla w + cw, b \cdot \nabla v)_T$$

(T denotes some element in Ω_{22}^*). Then the error analysis requires additionally to the Galerkin terms to estimate

$$a_{\text{stab}}(u - u^I, v^N) = N^{-1} \sum_{T \subset \Omega_{22}^*} (-\varepsilon \Delta u + b \cdot \nabla(u - u^I) + c(u - u^I), b \cdot \nabla v^N)_T.$$

For the smooth part S again Lin identities can be used, moreover for the layer part its smallness in Ω_{22}^* .

We renounce to present details, because the analysis is very similar to the analysis on Shishkin-type meshes.

5. A NUMERICAL COMPARISON

For our numerical experiments we consider the following test problem.

$$(5.1) \quad \begin{aligned} -\varepsilon\Delta u - 2u_x - 3u_y + u &= f & \text{in } \Omega = (0, 1)^2, \\ u &= 0 & \text{on } \partial\Omega, \end{aligned}$$

where we choose the right-hand side f in such a way that

$$u(x, y) = 2 \sin(1 - x)(1 - e^{-2x/\varepsilon})(1 - y)^2(1 - e^{-3y/\varepsilon})$$

is the exact solution of (5.1), which exhibits typical boundary and corner layer behavior. We take $\varepsilon = 10^{-4}, 10^{-8}, 10^{-12}$, which is sufficiently small to arise the phenomena of singular perturbation.

We observe for this test problem convergence of order one in the ε -weighted H^1 -norm on sequences of B-meshes (cf. Tab. 1) as on B-S-meshes (cf. Tab. 2), as well. As predicted by (3.7) the last two columns of each table suggest that these results are uniform in ε . Moreover, even the constant in $\|u^I - u^N\|_\varepsilon \leq CN^{-1}$ for B-meshes and for B-S-meshes has almost the same value.

With respect to supercloseness we observe practically $\|u^I - u^N\|_\varepsilon \leq CN^{-2}$ on both mesh types, here (cf. Tabs. 3 and 4).

Our numerical experiments indicate: B-meshes as well as B-S-meshes are particularly suitable for solving singularly perturbed problems, the differences between these two kinds of meshes being extremely small.

N	$\varepsilon = 10^{-8}$			$\varepsilon = 10^{-4}$	$\varepsilon = 10^{-12}$
	error	rate	$\ u - u^N\ _\varepsilon N$	error	error
8	3.0985e-1	1.0242	2.4788	3.0828e-1	3.1035e-1
16	1.5235e-1	1.0062	2.4376	1.5212e-1	1.5242e-1
32	7.5847e-2	1.0016	2.4271	7.5807e-2	7.5856e-2
64	3.7882e-2	1.0004	2.4244	3.7872e-2	3.7883e-2
128	1.8935e-2	1.0001	2.4237	1.8932e-2	1.8936e-2
256	9.4670e-3		2.4236	9.4657e-3	9.4671e-3

Table 1. Error on B-meshes in ε -weighted H^1 -norm, $\sigma = 2$.

N	$\varepsilon = 10^{-8}$			$\varepsilon = 10^{-4}$	$\varepsilon = 10^{-12}$
	error	rate	$\ u - u^N\ _\varepsilon N$	error	error
8	2.6679e-1		2.1343	2.6678e-1	2.6679e-1
16	1.4220e-1	0.9078	2.2752	1.4219e-1	1.4220e-1
32	7.3393e-2	0.9542	2.3486	7.3390e-2	7.3393e-2
64	3.7279e-2	0.9773	2.3859	3.7278e-2	3.7279e-2
128	1.8788e-2	0.9886	2.4048	1.8787e-2	1.8788e-2
256	9.4367e-3	0.9934	2.4158	9.4298e-3	9.4367e-3

Table 2. Error on B-S-meshes in ε -weighted H^1 -norm, $\sigma = 2$.

N	$\varepsilon = 10^{-8}$				$\varepsilon = 10^{-4}$	$\varepsilon = 10^{-12}$
	$\ \cdot\ _0$		$\ \cdot\ _\varepsilon$		$ \cdot _\varepsilon$	
	$\ v^N\ _0$	rate	$\ v^N\ _\varepsilon$	rate	$\ v^N\ _\varepsilon$	$\ v^N\ _\varepsilon$
8	5.2819e-3		2.7778e-2		2.8330e-2	2.7654e-2
16	1.8080e-3	1.5466	7.5677e-3	1.8760	7.7037e-3	7.5316e-3
32	5.3038e-4	1.7693	2.0100e-3	1.9127	2.0356e-3	2.0022e-3
64	1.4569e-4	1.8641	5.2262e-4	1.9433	5.2532e-4	5.2107e-4
128	3.8679e-5	1.9133	1.3403e-4	1.9632	1.3262e-4	1.3372e-4
256	1.0062e-5	1.9426	3.4084e-5	1.9754	3.2857e-5	3.4025e-5
512	2.5831e-6	1.9617	8.6204e-6	1.9833	8.2202e-6	8.6091e-6

Table 3. Supercloseness on B-meshes, $v^N = u^I - u^N$, $\sigma = 2.5$.

N	$\varepsilon = 10^{-8}$				$\varepsilon = 10^{-4}$	$\varepsilon = 10^{-12}$
	$\ \cdot\ _0$		$\ \cdot\ _\varepsilon$		$ \cdot _\varepsilon$	
	$\ v^N\ _0$	rate	$\ v^N\ _\varepsilon$	rate	$\ v^N\ _\varepsilon$	$\ v^N\ _\varepsilon$
8	7.2833e-3		2.3936e-2		2.3934e-2	2.3936e-2
16	1.9857e-3	1.8750	6.9827e-3	1.7774	6.9811e-3	6.9827e-3
32	5.4546e-4	1.8641	1.9216e-3	1.8615	1.9192e-3	1.9216e-3
64	1.4683e-4	1.8933	5.0990e-4	1.9140	5.0728e-4	5.0990e-4
128	3.8730e-5	1.9226	1.3225e-4	1.9469	1.2992e-4	1.3226e-4
256	1.0055e-5	1.9456	3.3839e-5	1.9666	3.2488e-5	3.3839e-5
512	2.5798e-6	1.9625	8.5861e-6	1.9786	8.1918e-6	8.5864e-6

Table 4. Supercloseness on B-S-meshes, $v^N = u^I - u^N$, $\sigma = 2.5$.

References

- [1] *M. Dobrowolski, H.-G. Roos*: A priori estimates for the solution of convection-diffusion problems and interpolation on Shishkin meshes. *Z. Anal. Anwend.* *16* (1997), 1001–1012.
- [2] *S. Franz*: Singularly perturbed problems with characteristic layers. PHD. TU Dresden, 2008.
- [3] *T. Linß*: Analysis of a Galerkin finite element method on a Bakhvalov-Shishkin mesh for a linear convection-diffusion problem. *IMA J. Numer. Anal.* *20* (2000), 621–632.
- [4] *T. Linß*: Uniform superconvergence of a Galerkin finite element method on Shishkin-type meshes. *Numer. Methods Partial Differential Equations* *16* (2000), 426–440.
- [5] *T. Linß, M. Stynes*: Asymptotic analysis and Shishkin-type decomposition for an elliptic convection-diffusion problem. *J. Math. Anal. Appl.* *261* (2001), 604–632.
- [6] *T. Linß*: Layer-adapted Meshes for Reaction-convection-diffusion Problems. Lecture Notes in Mathematics, Vol. 1985. Springer, Berlin, 2010.
- [7] *E. O’Riordan, G. Shishkin*: A technique to prove parameter-uniform convergence for a singularly perturbed convection-diffusion equation. *J. Comput. Appl. Math.* *206* (2007), 136–145.
- [8] *H.-G. Roos, T. Linß*: Sufficient conditions for uniform convergence on layer-adapted grids. *Computing* *63* (1999), 27–45.
- [9] *H.-G. Roos*: Error estimates for linear finite elements on Bakhvalov-type meshes. *Appl. Math.* *51* (2006), 63–72.
- [10] *H.-G. Roos, M. Stynes, L. Tobiska*: Robust Numerical Methods for Singularly Perturbed Differential Equations. Springer, Berlin, 2008.
- [11] *M. Stynes, E. O’Riordan*: A uniformly convergent Galerkin method on a Shishkin mesh for a convection-diffusion problem. *J. Math. Anal. Appl.* *214* (1997), 36–54.
- [12] *M. Stynes, L. Tobiska*: The SDFEM for a convection-diffusion problem with a boundary layer: optimal error analysis and enhancement of accuracy. *SIAM J. Numer. Anal.* *41* (2003), 1620–1642.

Authors’ address: *H.-G. Roos, M. Schopf*, Technical University of Dresden, Dresden, Germany, e-mail: Hans-Goerg.Roos@tu-dresden.de, Martin.Schopf@tu-dresden.de.