

Nikolay Moshchevitin

Exponents for three-dimensional simultaneous Diophantine approximations

Czechoslovak Mathematical Journal, Vol. 62 (2012), No. 1, 127–137

Persistent URL: <http://dml.cz/dmlcz/142046>

Terms of use:

© Institute of Mathematics AS CR, 2012

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

EXPONENTS FOR THREE-DIMENSIONAL SIMULTANEOUS
DIOPHANTINE APPROXIMATIONS

NIKOLAY MOSHCHEVITIN, Moskva

(Received November 9, 2010)

Abstract. Let $\Theta = (\theta_1, \theta_2, \theta_3) \in \mathbb{R}^3$. Suppose that $1, \theta_1, \theta_2, \theta_3$ are linearly independent over \mathbb{Z} . For Diophantine exponents

$$\alpha(\Theta) = \sup\{\gamma > 0: \limsup_{t \rightarrow +\infty} t^\gamma \psi_\Theta(t) < +\infty\},$$
$$\beta(\Theta) = \sup\{\gamma > 0: \liminf_{t \rightarrow +\infty} t^\gamma \psi_\Theta(t) < +\infty\}$$

we prove

$$\beta(\Theta) \geq \frac{1}{2} \left(\frac{\alpha(\Theta)}{1 - \alpha(\Theta)} + \sqrt{\left(\frac{\alpha(\Theta)}{1 - \alpha(\Theta)} \right)^2 + \frac{4\alpha(\Theta)}{1 - \alpha(\Theta)}} \right) \alpha(\Theta).$$

Keywords: Diophantine approximations, Diophantine exponents, Jarník's transference principle

MSC 2010: 11J13

1. DIOPHANTINE EXPONENTS

Let $\Theta = (\theta_1, \dots, \theta_n)$ be a real vector. We deal with the function

$$\psi_\Theta(t) = \min_{x \leq t} \max_{1 \leq i \leq n} \|\theta_i x\|.$$

Here the minimum is taken over positive integers x and $\|\cdot\|$ stands for the distance to the nearest integer.

Research is supported by RFBR grant No. 09-01-00371a.

Suppose that at least one of the numbers $\theta_1, \dots, \theta_n$ is irrational. Then $\psi_\Theta(t) > 0$ for all $t \geq 1$. The *uniform* Diophantine exponent $\alpha(\Theta)$ is defined as the supremum of the set

$$\{\gamma > 0: \limsup_{t \rightarrow +\infty} t^\gamma \psi_\Theta(t) < +\infty\}.$$

It is a well-known fact that for all Θ one has

$$\frac{1}{n} \leq \alpha(\Theta) \leq 1.$$

The *ordinary* Diophantine exponent $\beta(\Theta)$ is defined as the supremum of the set

$$\{\gamma > 0: \liminf_{t \rightarrow +\infty} t^\gamma \psi_\Theta(t) < +\infty\}.$$

Obviously

$$(1) \quad \beta(\Theta) \geq \alpha(\Theta).$$

2. FUNCTIONS

For each $\alpha \in [\frac{1}{3}, 1)$, define

$$g_1(\alpha) = \frac{\alpha}{1 - \alpha}$$

and

$$g_2(\alpha) = \frac{\alpha(1 - \alpha) + \sqrt{\alpha(\alpha^3 + 6\alpha^2 - 7\alpha + 4)}}{2(2\alpha^2 - 2\alpha + 1)}.$$

The value $g_2(\alpha)$ is the largest root of the equation

$$(2\alpha^2 - 2\alpha + 1)x^2 + \alpha(\alpha - 1)x - \alpha = 0.$$

Note that

$$g_2(1/3) = g_2(1) = 1,$$

and for $1/3 < \alpha < 1$ one has $g_2(\alpha) > 1$. Let α_0 be the unique real root of the equation

$$x^3 - x^2 + 2x - 1 = 0.$$

In the interval $1/3 < \alpha < \alpha_0$ one has

$$(2) \quad g_2(\alpha) > \max(1, g_1(\alpha)).$$

In the interval $\alpha_0 \leq \alpha < 1$ we see that

$$g_2(\alpha) \leq g_1(\alpha).$$

We define one more function. Put

$$(3) \quad g_3(\alpha) = \frac{1}{2} \left(\frac{\alpha}{1-\alpha} + \sqrt{\left(\frac{\alpha}{1-\alpha} \right)^2 + \frac{4\alpha}{1-\alpha}} \right).$$

Simple calculation shows that

$$(4) \quad g_3(\alpha) > \max(g_1(\alpha), g_2(\alpha)) \quad \forall \alpha \in \left(\frac{1}{3}, 1 \right).$$

3. JARNÍK'S RESULT

In a fundamental paper [1] V. Jarník proved the following theorem.

Theorem 1. *Let $\psi(t)$ be a continuous function in t , decreasing to zero as $t \rightarrow +\infty$. Suppose that the function $t\psi(t)$ increases to infinity as $t \rightarrow +\infty$. Let $\varrho(t)$ be the inverse function to the function $t\psi(t)$. Put*

$$\varphi^{[\psi]}(t) = \psi \left(\varrho \left(\frac{1}{6\psi(t)} \right) \right).$$

Suppose that $n \geq 2$ and among numbers $\theta_1, \dots, \theta_n$ there exist at least two numbers which, together with 1, are linearly independent over \mathbb{Z} . Suppose that

$$\psi_{\Theta}(t) \leq \psi(t)$$

for all t large enough. Then there exist infinitely many integers x such that

$$\max_{1 \leq j \leq n} \|x\theta_j\| \leq \varphi^{[\psi]}(x).$$

The next Jarník's result on Diophantine exponents is an obvious corollary of Theorem 1.

Theorem 2. *Suppose that $n \geq 2$ and among numbers $\theta_1, \dots, \theta_n$ there exist at least two numbers which, together with 1, are linearly independent over \mathbb{Z} . Then*

$$\beta(\Theta) \geq \alpha(\Theta)g_1(\alpha(\Theta)).$$

To obtain Theorem 2 from Theorem 1 one takes $\psi(t) = t^{-\alpha}$ with $\alpha < \alpha(\Theta)$.

On the other hand, V. Jarník [1] proved that there exists a collection of numbers $\Theta = (\theta_1, \dots, \theta_n)$ such that $1, \theta_1, \dots, \theta_n$ are linearly independent over \mathbb{Z} and

$$\beta(\Theta) < \frac{\alpha(\Theta)}{1 - \alpha(\Theta)}.$$

In the case $n = 2$ the lower bound in Jarník's Theorem 2 is optimal. The following result was proved by M. Laurent [2].

Theorem 3. *For any $\alpha, \beta > 0$ satisfying*

$$\frac{1}{2} \leq \alpha \leq 1, \quad \beta \geq \alpha g_1(\alpha)$$

there exists a vector $\Theta = (\theta_1, \theta_2) \in \mathbb{R}^2$ such that

$$\alpha(\Theta) = \alpha, \quad \beta(\Theta) = \beta.$$

This result is a corollary of a general theorem concerning four two-dimensional Diophantine exponents.

Note that in the case $n \geq 3$ the bound in Theorem 2 in the range $1/n \leq \alpha < \frac{1}{2}$ is weaker than the trivial bound (1).

N. Moshchevitin [3] (see also [4], Section 5.2) improved Jarník's result in the case $n = 3$ and for $\alpha \in (\frac{1}{3}, \alpha_0)$. He obtained

Theorem 4. *Suppose that $m = 1$, $n = 3$ and the collection $\Theta = (\theta_1, \theta_2, \theta_3)$ consists of numbers which, together with 1, are linearly independent over \mathbb{Z} . Then*

$$\beta(\Theta) \geq \alpha(\Theta)g_2(\alpha(\Theta)).$$

In the case $n = 3$, Theorems 2 and 4 together give an estimate which is better than the trivial estimate (1) for all admissible values of $\alpha(\Theta)$.

4. NEW RESULT

In this paper we give a new lower bound for $\beta(\Theta)$ in terms of $\alpha(\Theta)$. From (4) it follows that this bound is better than all the previous bounds (Theorems 2 and 4) for all admissible values of $\alpha(\Theta)$.

Theorem 5. Suppose that $m = 1$, $n = 3$ and the vector $\Theta = (\theta_1, \theta_2, \theta_3)$ consists of numbers linearly independent, together with 1, over \mathbb{Z} . Then

$$\beta(\Theta) \geq \alpha(\Theta)g_3(\alpha(\Theta)).$$

Sections 5, 6, 7 below contain auxiliary results. Theorem 5 is proved in Section 8.

5. BEST APPROXIMATIONS

For each integer x , put

$$\zeta(x) = \max_{1 \leq j \leq n} \|\theta_j x\|.$$

A positive integer x is said to be a *best approximation* if

$$\zeta(x) = \min_{x'} \zeta(x'),$$

where the minimum is taken over all $x' \in \mathbb{Z}$ such that

$$0 < x' \leq x.$$

Consider the case when all numbers 1 and θ_j , $1 \leq j \leq n$ are linearly independent over \mathbb{Z} . Then all best approximations lead to sequences

$$\begin{aligned} x_1 < x_2 < \dots < x_\nu < x_{\nu+1} < \dots, \\ \zeta(x_1) > \zeta(x_2) > \dots > \zeta(x_\nu) > \zeta(x_{\nu+1}) > \dots \end{aligned}$$

We use the notation

$$\zeta_\nu = \zeta(x_\nu).$$

Choose $y_{1,\nu}, \dots, y_{n,\nu} \in \mathbb{Z}$ such that

$$\|\theta_j x_\nu\| = |\theta_j \mathbf{x}_\nu - y_{j,\nu}|.$$

We define

$$\mathbf{z}_\nu = (x_\nu, y_{1,\nu}, \dots, y_{n,\nu}) \in \mathbb{Z}^{n+1}.$$

If $\psi(t)$ is a continuous function decreasing to 0 as $t \rightarrow \infty$, with

$$\psi_\Theta(t) \leq \psi(t),$$

then one easily sees that

$$(5) \quad \zeta_\nu \leq \psi(x_{\nu+1}).$$

Some useful fact about best approximations can be found in [4].

6. TWO-DIMENSIONAL SUBSPACES

Lemma 1. *Suppose that all vectors of the best approximations \mathbf{z}_l , $\nu \leq l \leq k$ lie in a certain two-dimensional linear subspace $\pi \subset \mathbb{R}^4$. Consider the two-dimensional lattice $\Lambda = \pi \cap \mathbb{Z}^4$ with the two-dimensional fundamental volume $\det \Lambda$. Then for all l from the interval $\nu \leq l \leq k - 1$ one has*

$$(6) \quad C_1 \det \Lambda \leq \zeta_l x_{l+1} \leq 2 \det \Lambda$$

where $C_1 = \left(2\sqrt{3(1 + (|\theta_1| + \frac{1}{2})^2 + (|\theta_2| + \frac{1}{2})^2 + (|\theta_3| + \frac{1}{2})^2)}\right)^{-1}$. In particular,

$$(7) \quad \det \Lambda \geq \frac{\min(\zeta_\nu x_{\nu+1}, \zeta_{k-1} x_k)}{2}.$$

P r o o f. The parallelepiped

$$\Omega_l = \left\{ \mathbf{z} = (x, y_1, y_2, y_3) : |x| < x_{l+1}, \max_{1 \leq j \leq 3} |\theta_j x - y_j| < \zeta_l \right\}$$

has no non-zero integer points inside for every l . Consider the two-dimensional $\mathbf{0}$ -symmetric convex body

$$\Xi_l = \Omega_l \cap \pi.$$

One can see that the two-dimensional Lebesgue measure $\mu(\Xi_l)$ of Ξ_l admits the following lower and upper bounds:

$$(8) \quad 2\zeta_l x_{l+1} \leq \mu(\Xi_l) \leq 4\sqrt{3\left(1 + \left(|\theta_1| + \frac{1}{2}\right)^2 + \left(|\theta_2| + \frac{1}{2}\right)^2 + \left(|\theta_3| + \frac{1}{2}\right)^2\right)} \zeta_l x_{l+1}.$$

We see that there is no non-zero point of Λ inside Ξ_l and that there are two linearly independent points $\mathbf{z}_l, \mathbf{z}_{l+1} \in \Lambda$ on the boundary of Ξ_l . So obviously

$$(9) \quad 2 \det \Lambda \leq \mu(\Xi_l).$$

From the Minkowski convex body theorem it follows that

$$(10) \quad \mu(\Xi_l) \leq 4 \det \Lambda.$$

Now (6) follows from (8, 9, 10). Lemma is proved. □

7. THREE-DIMENSIONAL SUBSPACES

Consider three consecutive best approximation vectors \mathbf{z}_{l-1} , \mathbf{z}_l , \mathbf{z}_{l+1} . Suppose that these vectors are linearly independent. Consider the three-dimensional linear subspace

$$\Pi_l = \text{span}(\mathbf{z}_{l-1}, \mathbf{z}_l, \mathbf{z}_{l+1}).$$

Consider the lattice

$$\Gamma_l = \Pi_l \cap \mathbb{Z}^4$$

with the fundamental volume $\det \Gamma_l$. Let Δ be the three-dimensional volume of the three-dimensional simplex \mathcal{S} with vertices $\mathbf{0}$, \mathbf{z}_{l-1} , \mathbf{z}_l , \mathbf{z}_{l+1} . We see that

$$(11) \quad \Delta \geq \frac{\det \Gamma_l}{6}.$$

Consider determinants

$$(12) \quad \Delta_1 = - \begin{vmatrix} x_{l-1} & y_{2,l-1} & y_{3,l-1} \\ x_l & y_{2,l} & y_{3,l} \\ x_{l+1} & y_{2,l+1} & y_{3,l+1} \end{vmatrix}, \quad \Delta_2 = \begin{vmatrix} x_{l-1} & y_{1,l-1} & y_{3,l-1} \\ x_l & y_{1,l} & y_{3,l} \\ x_{l+1} & y_{1,l+1} & y_{3,l+1} \end{vmatrix},$$

$$\Delta_3 = - \begin{vmatrix} x_{l-1} & y_{1,l-1} & y_{2,l-1} \\ x_l & y_{1,l} & y_{2,l} \\ x_{l+1} & y_{1,l+1} & y_{2,l+1} \end{vmatrix}.$$

The absolute values of these determinants are equal to the three-dimensional volumes of the projections of the simplex \mathcal{S} onto the three-dimensional coordinate subspaces ($\{y_1 = 0\}$, $\{y_2 = 0\}$ and $\{y_3 = 0\}$ respectively) multiplied by 6.

Note that for $j = 1, 2, 3$ one has

$$(13) \quad |\Delta_j| \leq 6\zeta_{l-1}\zeta_l x_{l+1}.$$

Lemma 2. *Among determinants (12) there exists a determinant with absolute value $\geq C_2 \Delta$, where $C_2 = 2/(2 + \max_{1 \leq i \leq 3} |\theta_i|)$.*

Proof. Consider the determinant

$$\Delta_0 = \begin{vmatrix} y_{1,l-1} & y_{2,l-1} & y_{3,l-1} \\ y_{1,l} & y_{2,l} & y_{3,l} \\ y_{1,l+1} & y_{2,l+1} & y_{3,l+1} \end{vmatrix}$$

and the vector

$$\mathbf{w} = (\Delta_0, \Delta_1, \Delta_2, \Delta_3) \in \mathbb{Z}^4.$$

We see that \mathbf{w} is orthogonal to the subspace Π_l , that is

$$\Delta_0 x_j + \Delta_1 y_{1,j} + \Delta_2 y_{2,j} + \Delta_3 y_{3,j} = 0, \quad j = l-1, l, l+1.$$

So

$$\Delta_0 = - \sum_{i=1}^3 \Delta_i \frac{y_{i,l}}{x_l} = - \sum_{i=1}^3 \Delta_i \left(\frac{y_{i,l}}{x_l} - \theta_i \right) - \sum_{i=1}^3 \Delta_i \theta_i.$$

As $|y_{i,l}/x_l - \theta_i| \leq 1$ we see that

$$(14) \quad |\Delta_0| \leq \left(1 + \max_{1 \leq i \leq 3} |\theta_i|\right) (|\Delta_1| + |\Delta_2| + |\Delta_3|).$$

However,

$$(15) \quad 36\Delta^2 = \Delta_0^2 + \Delta_1^2 + \Delta_2^2 + \Delta_3^2.$$

From (14), (15) we deduce the inequality

$$\Delta \leq \frac{1}{6} \left(2 + \max_{1 \leq i \leq 3} |\theta_i|\right) (|\Delta_1| + |\Delta_2| + |\Delta_3|),$$

and the lemma follows. □

8. PROOF OF THEOREM 5

Take $\alpha < \alpha(\Theta)$. Then

$$(16) \quad \zeta_l \leq x_{l+1}^{-\alpha}$$

for all l large enough.

Consider best approximation vectors $\mathbf{z}_\nu = (x_\nu, y_{1,\nu}, y_{2,\nu}, y_{3,\nu})$. From the condition that the numbers $1, \theta_1, \theta_2, \theta_3$ are linearly independent over \mathbb{Z} we see that there exist infinitely many pairs of indices $\nu < k, \nu \rightarrow +\infty$ such that

- both the triples

$$\mathbf{z}_{\nu-1}, \mathbf{z}_\nu, \mathbf{z}_{\nu+1}; \quad \mathbf{z}_{k-1}, \mathbf{z}_k, \mathbf{z}_{k+1}$$

consist of linearly independent vectors;

- there exists a two-dimensional linear subspace π such that

$$\mathbf{z}_l \in \pi, \quad \nu \leq l \leq k; \quad \mathbf{z}_{\nu-1} \notin \pi, \quad \mathbf{z}_{k+1} \notin \pi;$$

- the vectors

$$\mathbf{z}_{\nu-1}, \mathbf{z}_{\nu}, \mathbf{z}_k, \mathbf{z}_{k+1}$$

are linearly independent.

Consider the two-dimensional lattice

$$\Lambda = \pi \cap \mathbb{Z}^4.$$

By Lemma 1, its two-dimensional fundamental volume $\det \Lambda$ satisfies

$$(17) \quad \det \Lambda \asymp_{\Theta} \zeta_{\nu} x_{\nu+1} \asymp_{\Theta} \zeta_{k-1} x_k.$$

Consider the two dimensional orthogonal complement π^{\perp} to π and the lattice

$$\Lambda^{\perp} = \pi^{\perp} \cap \mathbb{Z}^4.$$

It is well-known that

$$(18) \quad \det \Lambda^{\perp} = \det \Lambda.$$

Consider the lattices

$$\Gamma_{\nu} = (\text{span}(\mathbf{z}_{\nu-1}, \mathbf{z}_{\nu}, \mathbf{z}_{\nu+1})) \cap \mathbb{Z}^4, \quad \Gamma_k = (\text{span}(\mathbf{z}_{k-1}, \mathbf{z}_k, \mathbf{z}_{k+1})) \cap \mathbb{Z}^4$$

and primitive integer vectors $\mathbf{w}_{\nu}, \mathbf{w}_k \in \mathbb{Z}^4$ which are orthogonal to $\Pi_{\nu} = \text{span}(\mathbf{z}_{\nu-1}, \mathbf{z}_{\nu}, \mathbf{z}_{\nu+1})$, $\Pi_k = \text{span}(\mathbf{z}_{k-1}, \mathbf{z}_k, \mathbf{z}_{k+1})$ respectively. Obviously

$$\mathbf{w}_{\nu}, \mathbf{w}_k \in \Lambda^{\perp}.$$

Put

$$b = \frac{1}{2} \left(-\frac{\alpha}{1-\alpha} + \sqrt{\left(\frac{\alpha}{1-\alpha}\right)^2 + \frac{4\alpha}{1-\alpha}} \right) \in (0, 1), \quad a = 1 - b,$$

so

$$\frac{\alpha}{1-\alpha} + b = g_3(\alpha).$$

Then

$$\det \Lambda^{\perp} \leq |w_{\nu}| \cdot |w_k|,$$

where $|\cdot|$ stands for the Euclidean norm, and so we obtain that either

$$(19) \quad \det \Gamma_{\nu} = |\mathbf{w}_{\nu}| \geq (\det \Lambda^{\perp})^a = (\det \Lambda)^a$$

or

$$(20) \quad \det \Gamma_k = |\mathbf{w}_k| \geq (\det \Lambda^\perp)^b = (\det \Lambda)^b$$

(using (18)).

If (19) holds then by Lemma 2, (13), (11) and (17) we see that

$$\zeta_{\nu-1} \zeta_\nu x_{\nu+1} \gg |\Delta_j| \gg_\Theta \det \Gamma_\nu \gg_\Theta (\det \Lambda)^a \gg (\zeta_\nu x_{\nu+1})^a$$

(here Δ_j is the determinant from Lemma 2 applied to the lattice $\Gamma = \Gamma_\nu$). From the definition of a and (16) we see that

$$x_{\nu+1} \gg_\Theta x_\nu^{g_3(\alpha)}.$$

We apply (16) again to obtain

$$\zeta_\nu \ll_\Theta x_\nu^{-\alpha g_3(\alpha)}.$$

If (20) holds then by Lemma 2, (13), (11) and (17) we see that

$$\zeta_{k-1} \zeta_k x_{k+1} \gg |\Delta_{j'}| \gg_\Theta \det \Gamma_k \gg_\Theta (\det \Lambda)^b \gg (\zeta_{k-1} x_k)^b$$

(here $\Delta_{j'}$ is the determinant from Lemma 2 applied to the lattice $\Gamma = \Gamma_k$). From the definition of b and (16) we see that

$$x_{k+1} \gg_\Theta x_k^{g_3(\alpha)}.$$

We apply (16) again to obtain

$$\zeta_k \ll_\Theta x_k^{-\alpha g_3(\alpha)}.$$

Theorem 5 is proved. □

Acknowledgement and a remark. The author thanks the anonymous referee for useful and important suggestions. Here we would like to note that the referee pointed out that it is possible to get a simpler proof of Theorem 5 by means of W. M. Schmidt's inequality on heights of rational subspaces (see [5]). For a rational subspace $U \subset \mathbb{R}^n$ its height $H(U)$ is defined as the co-volume of the lattice $U \cap \mathbb{Z}^n$. Schmidt showed that for any two rational subspaces $U, V \in \mathbb{R}^n$ one has

$$H(U \cap V)H(U + V) \ll_n H(U)H(V).$$

To prove our Theorem 5 one can use this inequality for

$$U = \text{span}(\mathbf{z}_{\nu-1}, \mathbf{z}_\nu, \mathbf{z}_{\nu+1}), \quad V = \text{span}(\mathbf{z}_{k-1}, \mathbf{z}_k, \mathbf{z}_{k+1}).$$

References

- [1] *V. Jarník*: Contribution à la théorie des approximations diophantiennes linéaires et homogènes. Czech. Math. J. 4 (1954), 330–353. (In Russian, French summary.)
- [2] *M. Laurent*: Exponents of Diophantine approximations in dimension two. Can. J. Math. 61 (2009), 165–189.
- [3] *N. G. Moshchevitin*: Contribution to Vojtěch Jarník. Preprint available at arXiv: 0912.2442v3.
- [4] *N. G. Moshchevitin*: Khintchine’s singular Diophantine systems and their applications. Russ. Math. Surv. 65 (2010), 433–511; Translation from Uspekhi Mat. Nauk. 65 (2010), 43–126.
- [5] *W. M. Schmidt*: On heights of algebraic subspaces and Diophantine approximations. Ann. Math. (2) 85 (1967), 430–472.

Author’s address: Nikolay G. Moshchevitin, Dept. of Number Theory, Fac. Mathematics and Mechanics, Moscow State University, Leninskie Gory 1, 119992 Moscow, Russia, e-mail: moshchevitin@gmail.com, moshchevitin@rambler.ru.