

Thomas Mejstrik
Some remarks on Nagumo's theorem

Czechoslovak Mathematical Journal, Vol. 62 (2012), No. 1, 235–242

Persistent URL: <http://dml.cz/dmlcz/142053>

Terms of use:

© Institute of Mathematics AS CR, 2012

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

SOME REMARKS ON NAGUMO'S THEOREM

THOMAS MEJSTRIK, Wien

(Received December 16, 2010)

Cordially dedicated to Prof. Adrian Constantin

Abstract. We provide a simpler proof for a recent generalization of Nagumo's uniqueness theorem by *A. Constantin*: On Nagumo's theorem. Proc. Japan Acad., Ser. A 86 (2010), 41–44, for the differential equation $x' = f(t, x)$, $x(0) = 0$ and we show that not only is the solution unique but the Picard successive approximations converge to the unique solution. The proof is based on an approach that was developed in *Z. S. Athanassov*: Uniqueness and convergence of successive approximations for ordinary differential equations. Math. Jap. 35 (1990), 351–367. Some classical existence and uniqueness results for initial-value problems for ordinary differential equations are particular cases of our result.

Keywords: ordinary differential equation, uniqueness

MSC 2010: 34A12, 34A34

1. INTRODUCTION

Nagumo's remarkable theorem [8] for the Cauchy problem

$$(1.1) \quad x'(t) = f(t, x(t))$$

with the initial data

$$(1.2) \quad x(0) = 0,$$

where $a > 0$ and $f: [0, a] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is continuous states that (1.1)–(1.2) has a unique solution if

$$(1.3) \quad |f(t, x) - f(t, y)| \leq \frac{|x - y|}{t}$$

for $t \in (0, a]$ and $x, y \in \mathbb{R}^n$ with $|x|, |y| \leq M$ for some $M > 0$. This result improves considerably the classical Lipschitz condition. It also motivates recent investigations in a variety of directions [1], [9], [10]. Among the various generalizations that appeared in the research literature, the most far-reaching one was recently obtained in [5]. It states that uniqueness holds if $f: [0, a] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is continuous, with

$$(1.4) \quad \frac{f(t, x)}{u'(t)} \rightarrow 0$$

as $t \downarrow 0$, uniformly in $|x| \leq M$ for some $M > 0$, and satisfies

$$(1.5) \quad |f(t, x) - f(t, y)| \leq \frac{u'(t)}{u(t)} \omega(|x - y|)$$

for $t \in (0, a]$ and $x, y \in \mathbb{R}^n$ with $|x|, |y| \leq M$, where u is an absolutely continuous function on $[0, a]$ with $u(0) = 0$ and $u'(t) > 0$ a.e. on $[0, a]$, and where ω belongs to the class \mathcal{F} of strictly increasing functions $\omega: [0, \infty) \rightarrow [0, \infty)$ with $\omega(0) = 0$ and such that

$$(1.6) \quad \int_0^r \frac{\omega(s)}{s} ds \leq r, \quad r > 0.$$

Notice that any strictly increasing continuous function $\omega: [0, \infty) \rightarrow [0, \infty)$ with $\omega(s) \leq s$ for $s \geq 0$ belongs to the class \mathcal{F} . There are also functions $\omega \in \mathcal{F}$ for which $\omega(r_n) > r_n$ for all $n \geq 1$, along an appropriate sequence $r_n \downarrow 0$, cf. [5].

The object of this note is to give a simpler proof of this uniqueness result and to show that the hypotheses ensure not only uniqueness but also the convergence of the successive approximations. To this end we adapt to the present context an approach that was developed in [2] to deal with the classical Nagumo theorem.

2. ALTERNATIVE PROOF OF UNIQUENESS

The aim of this section is to provide a simpler proof of the uniqueness result in [5]. We first derive a useful property of functions in the class \mathcal{F} .

Lemma 2.1. *If $\omega \in \mathcal{F}$ then $\omega(s) \leq es$ for $s \geq 0$.*

Proof. For $s > 0$ we have

$$(2.1) \quad s \geq \int_0^s \frac{\omega(r)}{r} dr \geq \int_{s/e}^s \frac{\omega(r)}{r} dr \geq \omega\left(\frac{s}{e}\right) \int_{s/e}^s \frac{1}{r} dr = \omega\left(\frac{s}{e}\right)$$

which yields the statement. □

Remark 2.2. The previous result might seem to indicate that we should simply set $\omega(s) = es$ in (1.5) and dispense altogether with the class \mathcal{F} . However, in Nagumo's classical theorem (with $u(t) = t$ and $\omega(s) = s$) the growth of the coefficient $1/t$ as $t \downarrow 0$ is optimal: for any $\alpha > 1$ there exist continuous functions f satisfying (1.3) with the right-hand side multiplied by α but for which (1.1)–(1.2) has nontrivial solutions [2]. Thus replacing $\omega(s)$ by $s \mapsto es$ is not an option.

A key role in our approach is the following Gronwall-type integral inequality (see [3], [6] for the classical Gronwall inequality and [4], [7] for generalizations in directions different to ours).

Lemma 2.3. *Let $u: [0, a] \rightarrow \mathbb{R}$ be absolutely continuous, nondecreasing and such that $u(t) > 0$ for $t > 0$. If $v: [0, a] \rightarrow \mathbb{R}$ is continuous, nonnegative, such that $v(t) = o(u(t))$ as $t \rightarrow 0^+$, and*

$$v(t) \leq \int_0^t \frac{\omega(v(s))}{u(s)} u'(s) \, ds, \quad 0 < t \leq a,$$

for some $\omega \in \mathcal{F}$, then v must be identically zero.

Proof. From Lemma 2.1 it follows that the integral is well-defined. Assume v is not the zero function. From $v(t)/u(t) \rightarrow 0$ as $t \rightarrow 0^+$ it follows that there exists $0 < \delta \leq a$ such that $v(t) \leq u(t)$ for $0 < t \leq \delta$. Let

$$\varepsilon = \frac{v(t_0)}{u(t_0)} = \sup_{0 < t \leq \delta} \left\{ \frac{v(t)}{u(t)} \right\} > 0$$

with $t_0 \in (0, \delta]$. We deduce that

$$\begin{aligned} \varepsilon u(t_0) = v(t_0) &\leq \int_0^{t_0} \omega(v(s)) \frac{u'(s)}{u(s)} \, ds < \int_0^{t_0} \omega(\varepsilon u(s)) \frac{u'(s)}{u(s)} \, ds \\ &= \int_{\varepsilon u(0)}^{\varepsilon u(t_0)} \frac{\omega(r)}{r} \, dr \leq \int_0^{\varepsilon u(t_0)} \frac{\omega(r)}{r} \, dr \leq \varepsilon u(t_0), \end{aligned}$$

which is a contradiction. Thus v is identically zero. □

This enables us to give a simple proof of the main result of [5]:

Theorem 2.4. *If f is continuous and satisfies (1.4) and (1.5), then (1.1)–(1.2) has a unique solution.*

P r o o f. The local existence of a solution is guaranteed by Peano's theorem. Let $x(t)$, $y(t)$ be two solutions of (1.1)–(1.2) for $0 < t \leq a$. In view of (1.4), given $\varepsilon > 0$, there exists $\delta = \delta(\varepsilon) > 0$ such that $|f(s, x)| \leq \varepsilon u'(s)$ for $0 < s \leq \delta$ and $|x| \leq M$. For $0 < t \leq \delta$ we have

$$|x(t) - y(t)| \leq \int_0^t |f(s, x(s)) - f(s, y(s))| \, ds \leq 2\varepsilon \int_0^t u'(s) \, ds \leq 2\varepsilon u(t)$$

so that $|x(t) - y(t)| = o(u(t))$ as $t \rightarrow 0^+$. Since

$$\begin{aligned} |x(t) - y(t)| &\leq \int_0^t |f(s, x(s)) - f(s, y(s))| \, ds \leq \int_0^t \frac{u'(s)}{u(s)} \omega(|x(s) - y(s)|) \, ds \\ &= \int_0^t \frac{u'(s)}{u(s)} \omega(|x(s) - y(s)|) \, ds, \end{aligned}$$

Lemma 2.3 yields $|x(t) - y(t)| \equiv 0$. □

3. CONVERGENCE OF THE SUCCESSIVE APPROXIMATIONS

The successive approximations for the problem (1.1)–(1.2) are defined by the sequence of functions

$$(3.1) \quad x_i(t) = \int_0^t f(s, x_{i-1}(s)) \, ds, \quad i \geq 1,$$

$x_0(t)$ being a continuous function on $[0, a]$ such that $x_0(0) = 0$ and $|x_0(t)| \leq M$ for $0 \leq t \leq a$. It turns out that the hypotheses (1.4) and (1.5) guarantee not only uniqueness but also the convergence of the successive approximations.

Theorem 3.1. *If the hypotheses of Theorem 2.4 are satisfied, then there exists a sufficiently small interval $0 \leq t \leq c$, $c > 0$, on which the successive approximations exist and converge uniformly to the unique solution of (1.1)–(1.2).*

P r o o f. We first prove that the successive approximations $\{x_i(t)\}_{i \geq 0}$ are well defined. From (1.4) it follows that, given $\varepsilon > 0$, there exists $\delta = \delta(\varepsilon) \in (0, a]$ such that

$$|f(s, x)| \leq \frac{\varepsilon}{2} u'(s), \quad 0 < s \leq \delta, \quad |x| \leq M.$$

Then it follows by (1.4) that for $t \in [0, \delta]$

$$|x_1(t)| \leq \int_0^t |f(s, x_0(s))| \, ds \leq \frac{\varepsilon}{2} \int_0^t u'(s) \, ds \leq \frac{\varepsilon}{2} u(a).$$

Taking

$$\varepsilon = \frac{2M}{u(a)}$$

and

$$c = \min \left\{ \delta, \frac{2M}{u(a)} \right\},$$

we obtain

$$|x_1(t)| \leq M \quad \text{for } 0 \leq t \leq c.$$

Suppose now that for $j \geq 1$ the continuous function $x_{j-1}(t)$ is well defined on $[0, c]$ and satisfies $x_{j-1}(0) = 0$. We then see that $f(t, x_{j-1}(t))$ is well defined, continuous and the integral in (3.1) exists, and its norm does not exceed $\frac{\varepsilon}{2}u(a)$ by our choice of c . This implies that $x_j(t)$ is also continuous and satisfies

$$x_j(0) = 0, \quad |x_j(t)| \leq M \quad \text{for } 0 \leq t \leq c.$$

It follows that that the successive approximations are well defined and uniformly bounded on $[0, c]$.

Now we prove that the family $\{x_j(t)\}$ is equicontinuous. Let $0 \leq t_1 < t_2 \leq c$ and $j \geq 1$ be given. Then

$$|x_j(t_2) - x_j(t_1)| = \left| \int_{t_1}^{t_2} f(s, x_{j-1}(s)) \, ds \right| \leq \int_{t_1}^{t_2} \varepsilon u'(s) \, ds.$$

From this and the previous calculations it follows that $\{x_j(t)\}$ is equicontinuous and uniformly bounded on $[0, c]$. Then, by the Arzela-Ascoli theorem, there exists a subsequence $\{x_{j_k}(t)\}$ which converges uniformly on $[0, c]$ to a continuous function $g(t)$ as $j_k \rightarrow \infty$. Since

$$x_{j_k+1}(t) = \int_0^t f(s, x_{j_k}(s)) \, ds,$$

by continuity of f , the sequence $\{x_{j_k+1}(t)\}$ converges uniformly to

$$\tilde{g}(t) = \int_0^t f(s, g(s)) \, ds.$$

We shall prove that on $[0, c]$ we have

$$(3.2) \quad \lim_{j \rightarrow \infty} |x_{j+1}(t) - x_j(t)| = 0.$$

By (3.1) this yields $g(t) = \tilde{g}(t)$ on $[0, c]$. This means that $g(t)$ is a solution of the equation. Since this solution is unique by Theorem 2.4, every subsequence of

$\{x_j(t)\}$ which is convergent will tend to the same solution $g(t)$, and this shows that $\{x_j(t)\}$ converges to $g(t)$ on $[0, c]$. Because of the uniform boundedness and the equicontinuity of the sequence this convergence is uniform.

To prove (3.2) we define on $[0, c]$ functions

$$\begin{aligned} y_j(t) &:= |x_{j+1}(t) - x_j(t)|, \quad j \geq 1, \\ m(t) &:= \sup_{0 \leq s \leq t} \frac{|x_2(s) - x_1(s)|}{u(s)}, \\ z_1(t) &:= m(t)u(t). \end{aligned}$$

Then for $t \in [0, c]$ we have

$$0 \leq m(t) \leq \varepsilon$$

so that

$$0 \leq z_1(t) \leq \varepsilon u(t).$$

Also

$$\begin{aligned} y_j(t) = |x_{j+1}(t) - x_j(t)| &\leq \int_0^t |f(s, x_j(s)) - f(s, x_{j-1}(s))| \, ds \leq \varepsilon \int_0^t u'(s) \, ds \\ &\leq \varepsilon u(t), \end{aligned}$$

while

$$y_1(t) \leq \sup_{0 \leq s \leq t} \left\{ \frac{|x_2(s) - x_1(s)|}{u(s)} \right\} u(t) = m(t)u(t) = z_1(t).$$

Define now on $[0, c]$ functions z_j with $j \geq 1$ as follows:

$$z_{j+1}(t) := \int_0^t \frac{u'(s)}{u(s)} \omega(z_j(s)) \, ds.$$

Since $0 \leq z_1(t) \leq \varepsilon u(t)$ and $u' \in L^1[0, a]$, the function z_2 is continuous on $[0, c]$ with

$$0 \leq z_2(t) \leq \int_0^t \frac{u'(s)}{u(s)} \omega(\varepsilon u(s)) \, ds = \int_{\varepsilon u(0)}^{\varepsilon u(t)} \frac{\omega(r)}{r} \, dr \leq \varepsilon u(t).$$

By induction we show that for $j \geq 1$

$$(3.3) \quad 0 \leq z_j(t) \leq \varepsilon u(t), \quad t \in [0, c].$$

On the other hand,

$$\begin{aligned} y_2(t) = |x_3(t) - x_2(t)| &\leq \int_0^t |f(s, x_2(s)) - f(s, x_1(s))| \, ds \leq \int_0^t \frac{u'(s)}{u(s)} \omega(y_1(s)) \\ &\leq \int_0^t \frac{u'(s)}{u(s)} \omega(z_1(s)) \, ds = z_2(t), \end{aligned}$$

and by induction one gets for $j \geq 1$ and $t \in [0, c]$ that

$$(3.4) \quad y_j(t) = |x_{j+1}(t) - x_j(t)| \leq z_j(t).$$

We now prove by induction that for $j \geq 1$ and $t \in [0, c]$ we have

$$(3.5) \quad 0 \leq z_{j+1}(t) \leq z_j(t).$$

Indeed,

$$\begin{aligned} z_1(t) - z_2(t) &= z_1(t) - \int_0^t \frac{u'(s)}{u(s)} \omega(z_1(s)) \, ds = z_1(t) - \int_0^t \frac{u'(s)}{u(s)} \omega(m(s)u(s)) \, ds \\ &\geq z_1(t) - \int_0^t \frac{u'(s)}{u(s)} \omega(m(t)u(s)) \, ds = z_1(t) - \int_{m(t)u(0)}^{m(t)u(t)} \frac{\omega(r)}{r} \, dr \\ &\geq z_1(t) - \int_0^{m(t)u(t)} \frac{\omega(r)}{r} \, dr \geq z_1(t) - z_1(t) = 0. \end{aligned}$$

Now assume

$$z_j(t) \leq z_{j-1}(t), \quad t \in [0, c].$$

Then

$$z_{j+1}(t) = \int_0^t \frac{u'(s)}{u(s)} \omega(z_j(s)) \, ds \leq \int_0^t \frac{u'(s)}{u(s)} \omega(z_{j-1}(s)) \, ds = z_j(t)$$

throughout $[0, c]$.

From (3.5) we infer that on $[0, c]$ the sequence $\{z_j(t)\}$ is decreasing and has a limit $z(t) \geq 0$ as $j \rightarrow \infty$. By Lebesgue's dominated convergence theorem we get

$$\begin{aligned} z(t) &= \lim_{j \rightarrow \infty} z_{j+1}(t) = \lim_{j \rightarrow \infty} \int_0^t \frac{u'(s)}{u(s)} \omega(z_j(s)) \, ds = \int_0^t \lim_{j \rightarrow \infty} \left\{ \frac{u'(s)}{u(s)} \omega(z_j(s)) \right\} \, ds \\ &= \int_0^t \frac{u'(s)}{u(s)} \omega\left(\lim_{j \rightarrow \infty} z_j(s)\right) \, ds = \int_0^t \frac{u'(s)}{u(s)} \omega(z(s)) \, ds. \end{aligned}$$

Since $z(t) = o(u(t))$ for $t \downarrow 0$, cf. (3.3), by Lemma 2.1 it follows that $z \equiv 0$. From this and (3.4) we deduce (3.2) and the proof is complete. \square

Acknowledgement. The author is grateful to the referee for useful suggestions.

References

- [1] *R. P. Agarwal, S. Heikkilä*: Uniqueness and well-posedness results for first order initial and boundary value problems.. *Acta Math. Hung.* *94* (2002), 67–92.
- [2] *Z. S. Athanassov*: Uniqueness and convergence of successive approximations for ordinary differential equations. *Math. Jap.* *35* (1990), 351–367.
- [3] *R. Bellman*: *Stability Theory of Differential Equations*. New York-London: McGraw-Hill Book Company, 166.
- [4] *A. Constantin*: Solutions globales d'équations différentielles perturbées. *C. R. Acad. Sci., Paris, Sér. I* *320* (1995), 1319–1322.
- [5] *A. Constantin*: On Nagumo's theorem. *Proc. Japan Acad., Ser. A* *86* (2010), 41–44.
- [6] *W. A. Coppel*: *Stability and Asymptotic Behavior of Differential Equations*. Boston: D. C. Heath and Company, 1965, pp. 166.
- [7] *O. Lipovan*: A retarded Gronwall-like inequality and its applications. *J. Math. Anal. Appl.* *252* (2000), 389–401.
- [8] *M. Nagumo*: Eine hinreichende Bedingung für die Unität der Lösung von Differentialgleichungen erster Ordnung. *Japanese Journ. of Math.* *3* (1926), 107–112.
- [9] *R. Negră*: On a class of backward stochastic differential equations and applications to the stochastic resonance. “Recent advances in stochastic modeling and data analysis”, pp. 26–33, World Sci. Publ., Hackensack, NJ, 2007.
- [10] *C. Sonoc*: On the pathwise uniqueness of solutions of stochastic differential equations. *Port. Math.* *55* (1998), 451–456.

Author's address: Thomas Meistrick, University of Vienna, Vienna, Austria, e-mail: tommsch@gmx.at.