

Paweł Krupski

Errata to hyperspaces of various locally connected subcontinua

*Acta Universitatis Carolinae. Mathematica et Physica*, Vol. 41 (2000), No. 1, 81

Persistent URL: <http://dml.cz/dmlcz/142705>

**Terms of use:**

© Univerzita Karlova v Praze, 2000

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

# ERRATA TO HYPERSPACES OF VARIOUS LOCALLY CONNECTED SUBCONTINUA

PAWEŁ KRUPSKI

University of Wrocław\*

1. The harmonic comb of  $n$ -disks  $C_n$  should be defined as follows:

$$C_n = [0, 1] \times [-1, 0]^{n-1} \cup \left( \sum_{i=1}^{\infty} \left[ s_i, \frac{1}{i} \right] \times [0, 1] \times [-1, 0]^{n-2} \right) \cup \left( \{0\} \times [0, 1] \times [-1, 0]^{n-2} \right) \subset [-1, 1]^n.$$

2. In the proof of Theorem 2 one should change coordinates of the end-points of segments  $A_i$  and  $B_i$  as well as the formula for  $f(\mathbf{x})$  in the following way:

Denote by  $A_i$  the segment from the point  $(\frac{1}{i}, 0)$  to  $(\frac{1}{i+1}, \frac{1}{2}x_i)$  in  $R^2$  and by  $B_i$  the segment from  $(\frac{1}{i+1}, \frac{1}{2}x_i)$  to  $(\frac{1}{i+1}, 0)$ .

Define an embedding  $f: Q \rightarrow C([-1, 1]^2) \subset C(X)$  by

$$f(\mathbf{x}) = (\{0\} \times [0, 1]) \cup \text{cl} \left( \bigcup_i (A_i \cup B_i) \right) \cup ([0, 1] \times \{1\}) \cup (\{1\} \times [0, 1]).$$

3. In the proof of Proposition 1 one should replace “the cube  $[-1, 0]^n$ ” by: cube  $[0, 1] \times [-1, 0]^{n-1}$ .

---

\* Mathematical Institute, University of Wrocław, Pl. Grunwaldzki 2/4, 50-384 Wrocław, Poland  
E-mail: krupski@math.uni.wroc.pl