

Xi Fu; Liulan Li; Xiantao Wang

A characterization of Fuchsian groups acting on complex hyperbolic spaces

*Czechoslovak Mathematical Journal*, Vol. 62 (2012), No. 2, 517–525

Persistent URL: <http://dml.cz/dmlcz/142843>

## Terms of use:

© Institute of Mathematics AS CR, 2012

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A CHARACTERIZATION OF FUCHSIAN GROUPS ACTING  
ON COMPLEX HYPERBOLIC SPACES

XI FU, Shaoxing, LIULAN LI, Hengyang,  
XIANTAO WANG, Changsha

(Received February 25, 2011)

*Abstract.* Let  $G \subset \mathbf{SU}(2, 1)$  be a non-elementary complex hyperbolic Kleinian group. If  $G$  preserves a complex line, then  $G$  is  $\mathbb{C}$ -Fuchsian; if  $G$  preserves a Lagrangian plane, then  $G$  is  $\mathbb{R}$ -Fuchsian;  $G$  is Fuchsian if  $G$  is either  $\mathbb{C}$ -Fuchsian or  $\mathbb{R}$ -Fuchsian. In this paper, we prove that if the traces of all elements in  $G$  are real, then  $G$  is Fuchsian. This is an analogous result of Theorem V.G. 18 of B. Maskit, Kleinian Groups, Springer-Verlag, Berlin, 1988, in the setting of complex hyperbolic isometric groups. As an application of our main result, we show that  $G$  is conjugate to a subgroup of  $\mathbf{S}(U(1) \times U(1, 1))$  or  $\mathbf{SO}(2, 1)$  if each loxodromic element in  $G$  is hyperbolic. Moreover, we show that the converse of our main result does not hold by giving a  $\mathbb{C}$ -Fuchsian group.

*Keywords:*  $\mathbb{R}$ -Fuchsian group,  $\mathbb{C}$ -Fuchsian group, complex line,  $\mathbb{R}$ -plane, trace

*MSC 2010:* 30F40, 20H10

1. INTRODUCTION

It is known that a Kleinian group  $G$  is Fuchsian if there exists a  $G$ -invariant disc  $\mathbb{D}$  in the Riemann sphere  $\widehat{\mathbb{C}} = \mathbb{C} \cup \infty$ . If we regard  $\mathbb{D}$  as  $\mathbb{H}^2$ , then  $G$  is a subgroup of  $\mathbf{SL}(2, \mathbb{R})$ . The following result due to Maskit is from Theorem V.G. 18 of [5].

**Theorem A.** *Let  $G \subset \mathbf{SL}(2, \mathbb{C})$  be a non-elementary Kleinian group in which  $\mathrm{tr}^2(f) \geq 0$  for all  $f \in G$ . Then  $G$  is Fuchsian.*

---

The research was partly supported by NSFs of China (No. 11071063) and Shaoxing College of Arts and Sciences (No. 20125009), NSF of Hunan (No. 10JJ4005), Hunan Provincial Education Department (No. 11B019) and the construct program of the key discipline in Hunan province.

This result shows that if the traces of all elements in  $G$  are real then  $G$  preserves a hyperbolic plane which is totally geodesic in  $\mathbb{H}^3$ . In this note, we will prove a similar result in the setting of complex hyperbolic Kleinian groups of  $\mathbf{SU}(2, 1)$ . Our result is as follows, whose proof will be given in Section 3.

**Theorem 1.1.** *Let  $G \subset \mathbf{SU}(2, 1)$  be a non-elementary complex hyperbolic Kleinian group in which  $\mathrm{tr}(f) \in \mathbb{R}$  for all  $f \in G$ . Then  $G$  is Fuchsian.*

Note that a loxodromic element in  $\mathbf{SU}(2, 1)$  is hyperbolic if and only if its trace is real. The proof of Theorem 1.1 easily yields

**Corollary 1.2.** *Let  $G \subset \mathbf{SU}(2, 1)$  be a non-elementary group. If each loxodromic element in  $G$  is hyperbolic, then  $G$  is conjugate to a subgroup of  $\mathbf{S}(U(1) \times U(1, 1))$  or  $\mathbf{SO}(2, 1)$ .*

As an application of Theorem 1.1, in Section 4, two Fuchsian groups are constructed: one is  $\mathbb{C}$ -Fuchsian and the other is  $\mathbb{R}$ -Fuchsian. We also give a  $\mathbb{C}$ -Fuchsian group which shows that the converse of Theorem 1.1 is not true.

## 2. COMPLEX HYPERBOLIC GEOMETRY

**2.1. Complex hyperbolic space.** Let  $\mathbb{C}^{2,1}$  be the complex vector space of dimension 3 equipped with a non-degenerate, indefinite Hermitian form  $\langle \cdot, \cdot \rangle$  of signature  $(2, 1)$  defined to be

$$\langle z, w \rangle = w^* J z = z_1 \bar{w}_3 + z_2 \bar{w}_2 + z_3 \bar{w}_1$$

with the matrix

$$J = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}.$$

We consider the subspaces

$$\begin{aligned} V_- &= \{\mathbf{z} \in \mathbb{C}^{2,1} : \langle \mathbf{z}, \mathbf{z} \rangle < 0\}, \\ V_0 &= \{\mathbf{z} \in \mathbb{C}^{2,1} - \{0\} : \langle \mathbf{z}, \mathbf{z} \rangle = 0\} \end{aligned}$$

and the canonical projection

$$\mathbb{P}: \mathbb{C}^{2,1} - \{0\} \rightarrow \mathbb{C}P^2$$

onto the complex projective space. The complex hyperbolic space  $\mathbf{H}_{\mathbb{C}}^2$  is defined to be  $\mathbb{P}(V_-)$  and its boundary  $\partial\mathbf{H}_{\mathbb{C}}^2$  is  $\mathbb{P}(V_0)$ . That is,

$$\mathbf{H}_{\mathbb{C}}^2 = \{(z_1, z_2) \in \mathbb{C}^2 : 2\Re(z_1) + |z_2|^2 < 0\}$$

and

$$\partial\mathbf{H}_{\mathbb{C}}^2 - \{\infty\} = \{(z_1, z_2) \in \mathbb{C}^2 : 2\Re(z_1) + |z_2|^2 = 0\}.$$

Given a point  $z \in \mathbb{C}^2 \subset \mathbb{C}P^2$ , we can lift  $z = (z_1, z_2)$  to a point  $\mathbf{z}$  in  $\mathbb{C}^{2,1}$ , called the standard lift of  $z$ , where

$$\mathbf{z} = \begin{pmatrix} z_1 \\ z_2 \\ 1 \end{pmatrix}.$$

There are two distinguished points in  $V_0$  which are denoted by  $\mathbf{0}$  and  $\infty$ , respectively. They are

$$\mathbf{0} = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \text{ and } \infty = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}.$$

**2.2. Isometries.** Denote by  $\mathbf{U}(2, 1)$  the group of unitary matrices for the Hermitian product  $\langle \cdot, \cdot \rangle$ . Each such matrix  $A$  satisfies the relation  $A^{-1} = JA^*J$ , where  $A^*$  is the Hermitian transpose of  $A$ . The full group of holomorphic isometries of  $\mathbf{H}_{\mathbb{C}}^2$  is the projective unitary group  $\mathbf{PU}(2, 1) = \mathbf{U}(2, 1)/\mathbf{U}(1)$ , where  $\mathbf{U}(1) = \{e^{i\theta}I : \theta \in [0, 2\pi)\}$  and  $I$  is the  $3 \times 3$  identity matrix. In this paper, we shall consider the group  $\mathbf{SU}(2, 1)$  of matrices which are unitary with respect to  $\langle \cdot, \cdot \rangle$  and have determinant 1. Following [3], holomorphic isometries of  $\mathbf{H}_{\mathbb{C}}^2$  are classified as follows.

- (1) An isometry is *elliptic* if it fixes at least one point of  $\mathbf{H}_{\mathbb{C}}^2$ ;
- (2) an isometry is *parabolic* if it fixes exactly one point of  $\partial\mathbf{H}_{\mathbb{C}}^2$ ;
- (3) an isometry is *loxodromic* if it fixes exactly two points of  $\partial\mathbf{H}_{\mathbb{C}}^2$ .

See [1], [3], [4], [7] for more details about complex hyperbolic geometry and complex hyperbolic Kleinian groups.

**2.3. Totally geodesic manifolds and Fuchsian groups.** Unlike the real hyperbolic space, there are two kinds of totally geodesic manifolds with codimension 2 in  $\mathbf{H}_{\mathbb{C}}^2$ . In the first place there are *complex lines* which have constant curvature  $-1$ . Every complex line  $L$  is the image of the complex line

$$L_0 = \{(z_1, z_2) \in \mathbf{H}_{\mathbb{C}}^2 : z_2 = 0\}$$

under some element of  $\mathbf{SU}(2, 1)$ . The subgroup of  $\mathbf{SU}(2, 1)$  stabilizing  $L$  is thus conjugate to the subgroup  $\mathbf{S}(U(1) \times U(1, 1)) \subset \mathbf{SU}(2, 1)$ . Secondly, we have totally

real *Lagrangian planes* which have constant curvature  $-\frac{1}{4}$ . Every Lagrangian plane is the image of the standard real Lagrangian plane

$$R_{\mathbb{R}} = \{(z_1, z_2) \in \mathbf{H}_{\mathbb{C}}^2 : z_i = x_i \in \mathbb{R}, 2x_1 + x_2^2 < 0\}$$

under some element of  $\mathbf{SU}(2, 1)$ . The group stabilizing  $R_{\mathbb{R}}$  is denoted by  $\mathbf{SO}(2, 1)$ , which is the subgroup of  $\mathbf{SU}(2, 1)$  comprising elements with real entries. We say a group  $G$  is *non-elementary* if there are two loxodromic elements in  $G$  with distinct fixed points. Following [2], for any non-elementary complex hyperbolic Kleinian group  $G \subset \mathbf{SU}(2, 1)$ ,

- (1)  $G$  is called  *$\mathbb{C}$ -Fuchsian* if it preserves a complex line;
- (2)  $G$  is called  *$\mathbb{R}$ -Fuchsian* if it preserves a Lagrangian plane;
- (3) otherwise,  $G$  is called *non-Fuchsian*.

We call a non-elementary Kleinian group  $G$  *Fuchsian* if  $G$  is either  $\mathbb{C}$ -Fuchsian or  $\mathbb{R}$ -Fuchsian.

**2.4. Cartan's angular invariant and the cross-ratio variety.** Let  $z_1, z_2, z_3$  be three distinct points in  $\partial\mathbf{H}_{\mathbb{C}}^2$  with lifts  $\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3$ , respectively. Cartan's angular invariant  $\mathbb{A}$  is defined to be

$$\mathbb{A}(z_1, z_2, z_3) = \arg(-\langle \mathbf{z}_1, \mathbf{z}_2 \rangle \langle \mathbf{z}_2, \mathbf{z}_3 \rangle \langle \mathbf{z}_3, \mathbf{z}_1 \rangle).$$

It is known that  $\mathbb{A}$  is invariant under the elements of  $\mathbf{SU}(2, 1)$ . The following is a useful property of  $\mathbb{A}$  which was proved by Goldman, see Section 7.1 of [3].

**Theorem B.** *Let  $z_1, z_2, z_3$  be three distinct points of  $\partial\mathbf{H}_{\mathbb{C}}^2$  and let  $\mathbb{A} = \mathbb{A}(z_1, z_2, z_3)$  denote their angular invariant. Then*

- (1)  $\mathbb{A} \in [-\frac{1}{2}\pi, \frac{1}{2}\pi]$ ;
- (2)  $\mathbb{A} = \pm\frac{1}{2}\pi$  if and only if  $z_1, z_2, z_3$  all lie on a chain;
- (3)  $\mathbb{A} = 0$  if and only if  $z_1, z_2, z_3$  all lie on an  $\mathbb{R}$ -circle.

Here we call the boundary of a complex line a *chain* and the boundary of a Lagrangian plane an  *$\mathbb{R}$ -circle*.

**Proposition 2.1.** *Let  $G \subset \mathbf{SU}(2, 1)$  be a non-elementary complex hyperbolic Kleinian group. Then  $G$  is  $\mathbb{C}$ -Fuchsian ( $\mathbb{R}$ -Fuchsian) if and only if the fixed points of all loxodromic elements in  $G$  are contained in a chain (an  $\mathbb{R}$ -circle).*

*Proof.* First, it is obvious that if  $G$  is  $\mathbb{C}$ -Fuchsian ( $\mathbb{R}$ -Fuchsian) then any loxodromic element  $U$  in  $G$  must preserve the invariant complex line (the Lagrangian

plane) and so its fixed points must be on the boundary chain (the  $\mathbb{R}$ -circle). Conversely, suppose  $G$  is non-elementary and contains loxodromic elements  $U$  and  $V$  with distinct fixed points. Suppose the fixed points of all loxodromic elements of  $G$  lie on a chain (an  $\mathbb{R}$ -circle). In particular, there is a unique complex line  $L$  (a unique Lagrangian plane  $R$ ) such that the fixed points of  $U$  and  $V$  lie in  $\partial L$  ( $\partial R$ ). Let  $A$  be any element of  $G$ . Then the fixed points of  $AUA^{-1}$  and  $AVA^{-1}$  lie on the boundary of the complex line  $A(L)$  (the Lagrangian plane  $A(R)$ ). By hypothesis, they also lie on the boundary of  $L$  ( $R$ ). Since four distinct points lie on at most one chain ( $\mathbb{R}$ -circle), we see that  $A$  sends  $L$  ( $R$ ) to itself (as a set). This is true for all elements of  $G$ , and so  $G$  is  $\mathbb{C}$ -Fuchsian ( $\mathbb{R}$ -Fuchsian).  $\square$

Let  $z_1, z_2, z_3, z_4$  be four distinct points of  $\partial\mathbf{H}_{\mathbb{C}}^2$  and  $\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3, \mathbf{z}_4$  their corresponding lifts in  $V_0 \subset \mathbb{C}^{2,1}$ , respectively. Then their *complex cross ratio* is defined to be

$$\mathbb{X} = [z_1, z_2, z_3, z_4] = \frac{\langle \mathbf{z}_3, \mathbf{z}_1 \rangle \langle \mathbf{z}_4, \mathbf{z}_2 \rangle}{\langle \mathbf{z}_4, \mathbf{z}_1 \rangle \langle \mathbf{z}_3, \mathbf{z}_2 \rangle}.$$

It is easy for us to know that  $\mathbb{X}$  is neither 0 nor  $\infty$ . By changing the order of the four points we can define the following three different cross-ratios:

$$\mathbb{X}_1 = [z_1, z_2, z_3, z_4], \quad \mathbb{X}_2 = [z_1, z_3, z_2, z_4] \quad \text{and} \quad \mathbb{X}_3 = [z_2, z_3, z_1, z_4].$$

The following lemma which is crucial for us follows from Propositions 5.12, 5.13 and 5.14 of [6].

**Lemma 2.2.** *Let  $z_1, z_2, z_3, z_4$  be four distinct points of  $\partial\mathbf{H}_{\mathbb{C}}^2$ . Then all  $z_i$  ( $i = 1, 2, 3, 4$ ) lie on a chain or an  $\mathbb{R}$ -circle if and only if all  $\mathbb{X}_j$  ( $j = 1, 2, 3$ ) are real.*

*Proof.* It follows from

$$\mathbb{X}_1 = [z_1, z_2, z_3, z_4] = \frac{\langle \mathbf{z}_3, \mathbf{z}_1 \rangle \langle \mathbf{z}_4, \mathbf{z}_2 \rangle}{\langle \mathbf{z}_4, \mathbf{z}_1 \rangle \langle \mathbf{z}_3, \mathbf{z}_2 \rangle} = \frac{\langle \mathbf{z}_1, \mathbf{z}_2 \rangle \langle \mathbf{z}_2, \mathbf{z}_3 \rangle \langle \mathbf{z}_3, \mathbf{z}_1 \rangle |\langle \mathbf{z}_2, \mathbf{z}_4 \rangle|^2}{\langle \mathbf{z}_1, \mathbf{z}_2 \rangle \langle \mathbf{z}_2, \mathbf{z}_4 \rangle \langle \mathbf{z}_4, \mathbf{z}_1 \rangle |\langle \mathbf{z}_2, \mathbf{z}_3 \rangle|^2}$$

that

$$\begin{aligned} \arg(\mathbb{X}_1) &= \arg(-\langle \mathbf{z}_1, \mathbf{z}_2 \rangle \langle \mathbf{z}_2, \mathbf{z}_3 \rangle \langle \mathbf{z}_3, \mathbf{z}_1 \rangle) - \arg(-\langle \mathbf{z}_1, \mathbf{z}_2 \rangle \langle \mathbf{z}_2, \mathbf{z}_4 \rangle \langle \mathbf{z}_4, \mathbf{z}_1 \rangle) \\ &= \mathbb{A}(z_1, z_2, z_3) - \mathbb{A}(z_1, z_2, z_4). \end{aligned}$$

Since all  $z_i$  ( $i = 1, 2, 3, 4$ ) lie on a chain or an  $\mathbb{R}$ -circle, by Theorem B we know that  $\mathbb{X}_1$  is real. Similar discussions yield that  $\mathbb{X}_2$  and  $\mathbb{X}_3$  are real.

Now we prove the sufficiency. It suffices to consider the case that all  $\mathbb{X}_j$  ( $j = 1, 2, 3$ ) are positive since if one of  $\mathbb{X}_j$  is negative, then by [6, Proposition 5.1] we know that all  $z_i$  lie on a chain. It follows that

$$\mathbb{A}(z_1, z_2, z_4) = \mathbb{A}(z_1, z_2, z_3), \quad \mathbb{A}(z_1, z_3, z_2) = \mathbb{A}(z_1, z_3, z_4)$$

and

$$\mathbb{A}(z_2, z_3, z_4) = \mathbb{A}(z_2, z_3, z_1).$$

According to the definition of Cartan's angular invariant, we have

$$\mathbb{A}(z_1, z_2, z_3) = -\mathbb{A}(z_1, z_3, z_2).$$

By [3, Lemma 7.1.10] and Theorem B, it is easy for us to prove that all  $z_i$  lie on an  $\mathbb{R}$ -circle.  $\square$

### 3. THE PROOF OF THEOREM 1.1

We prove this result by contradiction. Suppose that  $G$  is non-Fuchsian. Since  $G$  is non-elementary, by Proposition 2.1 we can find two loxodromic elements  $U, V \in G$  such that  $A_u, A_v, R_u$  and  $R_v$  lie neither on a chain nor an  $\mathbb{R}$ -circle and

$$\{A_u, R_u\} \cap \{A_v, R_v\} = \emptyset,$$

where  $A_w, R_w$  denote the attracting and repelling fixed points of the loxodromic element  $W \in G$ , respectively. Without loss of generality, we may assume that

$$U = \begin{pmatrix} r & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1/r \end{pmatrix}$$

and

$$V = \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & j \end{pmatrix} \begin{pmatrix} s & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1/s \end{pmatrix} \begin{pmatrix} \bar{j} & \bar{f} & \bar{c} \\ \bar{h} & \bar{e} & \bar{b} \\ \bar{g} & \bar{d} & \bar{a} \end{pmatrix},$$

where  $\begin{pmatrix} a & b & c \\ d & e & f \\ g & h & j \end{pmatrix} \in \mathbf{SU}(2, 1)$ ,  $ajgc \neq 0$ ,  $r, s > 1$  and  $r \neq s$  (if  $r = s$ , we can use  $V^2$  instead of  $V$ ). Applying Lemma 2.2, we know that at least one of  $\mathbb{X}_j$  ( $j = 1, 2, 3$ ) is not real, where

$$\mathbb{X}_1 = [A_v, A_u, R_u, R_v], \quad \mathbb{X}_2 = [A_v, R_u, A_u, R_v] \text{ and } \mathbb{X}_3 = [A_u, R_u, A_v, R_v].$$

By [6, Proposition 6.4], we have

$$\begin{aligned} \operatorname{tr}(UV) &= r + s + r^{-1} + s^{-1} + \mathbb{X}_1(r^{-1} - 1)(s^{-1} - 1) + \overline{\mathbb{X}}_1(r - 1)(s - 1) \\ &\quad + \mathbb{X}_2(r - 1)(s^{-1} - 1) + \overline{\mathbb{X}}_2(r^{-1} - 1)(s - 1) - 1 \end{aligned}$$

and

$$\begin{aligned} \operatorname{tr}[U, V] = & 3 - \Re[(\aleph_1 + \aleph_2)(r-1)(r^{-1}-1)(s-1)(s^{-1}-1)] \\ & + [1 - 2\Re(\aleph_1 + \aleph_2)][(r-1)^2(s-1)^2 + (r^{-1}-1)^2(s^{-1}-1)^2] \\ & + |\aleph_1(r-1)(s-1) + \overline{\aleph_1}(r^{-1}-1)(s^{-1}-1)| \\ & + |\aleph_2(r^{-1}-1)(s-1) + \overline{\aleph_2}(r-1)(s^{-1}-1)|^2 \\ & + (|\aleph_2|^2 - |\aleph_1|^2\aleph_3)(r^2 - 2r + 2r^{-1} - r^{-2})(s^2 - 2s + 2s^{-1} - s^{-2}). \end{aligned}$$

Now, we divide our proof into four cases.

*Case I.*  $\aleph_3$  is not real.

By computation, we have

$$\Im(\operatorname{tr}[U, V]) = |\aleph_1|^2(r - r^{-1})(r + r^{-1} - 2)(s - s^{-1})(s + s^{-1} - 2)\Im(\aleph_3),$$

which implies that  $\operatorname{tr}[U, V]$  is not real.

*Case II.*  $\aleph_1$  is real and  $\aleph_2$  is not real.

In this case,

$$\Im(\operatorname{tr}(UV)) = (r^{-1} - s^{-1})(r-1)(s-1)\Im(\aleph_2).$$

Since  $r, s > 1$  and  $r \neq s$ ,  $\Im(\operatorname{tr}(UV)) \neq 0$ . Therefore  $\operatorname{tr}(UV)$  is not real.

*Case III.*  $\aleph_2$  is real and  $\aleph_1$  is not real.

Then

$$\Im(\operatorname{tr}(UV)) = (r^{-1}s^{-1} - 1)(r-1)(s-1)\Im(\aleph_1).$$

It follows that  $\operatorname{tr}(UV)$  is not real.

*Case IV.* Neither  $\aleph_1$  nor  $\aleph_2$  are real.

If  $\Im[\overline{\aleph_1}(r-1) + \overline{\aleph_2}(r^{-1}-1)] = 0$ , then  $\Im(\aleph_2) = r\Im(\aleph_1)$ . So

$$\Im(\operatorname{tr}(UV)) = (r-1)(s-1)r^{-1}s^{-1}(1-r^2)\Im(\aleph_1) \neq 0.$$

Hence  $\operatorname{tr}(UV)$  is not real.

If  $\Im[\overline{\aleph_1}(r-1) + \overline{\aleph_2}(r^{-1}-1)] \neq 0$ , according to the definition of the cross-ratio variety, we know that  $\aleph_j$  ( $j = 1, 2, 3$ ) is independent of the value of  $s$  and  $r$ . Then there must exist a sufficiently large integer  $m$  such that

$$\begin{aligned} & \Im[\aleph_1(r^{-1}-1)(s^{-m}-1) + \aleph_2(r-1)(s^{-m}-1)] \\ & + \Im[\overline{\aleph_1}(r-1)(s^m-1) + \overline{\aleph_2}(r^{-1}-1)(s^m-1)] \neq 0. \end{aligned}$$

This implies that  $\operatorname{tr}(UV^m)$  is not real. □



#### 4. THREE EXAMPLES

**Example 4.1.** Let

$$G_1 = \left\langle A = \begin{pmatrix} 1 & 0 & i \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, B = \begin{pmatrix} 0 & 0 & i \\ 0 & 1 & 0 \\ i & 0 & 1 \end{pmatrix} \right\rangle.$$

Then  $G_1$  is  $\mathbb{C}$ -Fuchsian and each element in  $G_1$  has real trace.

*Proof.* It is obvious that  $G_1$  is a  $\mathbb{C}$ -Fuchsian group which keeps the complex line  $L_0 = \{(z_1, z_2) \in \mathbf{H}_{\mathbb{C}}^2 : z_2 = 0\}$  invariant. We only need to show that every element in  $G_1$  has real trace. Let  $M$  be an element having the following form

$$M = \begin{pmatrix} a & 0 & ib \\ 0 & 1 & 0 \\ ic & 0 & d \end{pmatrix},$$

where  $a, b, c, d \in \mathbb{Z}$ . Since the generators of  $G_1$  and their inverses have this form it is clear that this form is preserved under matrix multiplication. This implies that each element in  $G_1$  has real trace.  $\square$

**Example 4.2.** Let

$$G_2 = \mathbf{SO}(2, 1; \mathbb{Z}).$$

Then  $G_2$  is  $\mathbb{R}$ -Fuchsian and each element in  $G_2$  has real trace.

It is known that the converse to Maskit's theorem is clearly true (the trace of every element in a Fuchsian subgroup of  $\mathbf{SL}(2, \mathbb{C})$  is real), the converse to Theorem 1.1 is true for  $\mathbb{R}$ -Fuchsian groups, but false for  $\mathbb{C}$ -Fuchsian groups. The following is a  $\mathbb{C}$ -Fuchsian group but does not comprise only matrices with real trace.

**Example 4.3.**

$$G_3 = \left\langle A = \begin{pmatrix} -i & 0 & 1 \\ 0 & -1 & 0 \\ 0 & 0 & -i \end{pmatrix}, B = \begin{pmatrix} 0 & 0 & i \\ 0 & 1 & 0 \\ i & 0 & 1 \end{pmatrix} \right\rangle.$$

**Acknowledgement.** Most of this work was completed while the first author was a visiting research student at the University of Durham. He is grateful to Professor John R. Parker for his valuable help. All of the authors heartily thank the referee for careful reading of this paper as well as for many useful comments and suggestions. In particular, the proof of Proposition 2.1 was given by him or her.

### References

- [1] *A. F. Beardon*: The Geometry of Discrete Groups. Graduate Texts in Mathematics, Vol. 91, Springer, New York, 1983.
- [2] *S. S. Chen, L. Greenberg*: Hyperbolic spaces. *Contribut. to Analysis, Collect. of Papers dedicated to Lipman Bers (1974)*, 49–87.
- [3] *W. M. Goldman*: Complex Hyperbolic Geometry. Oxford: Clarendon Press, 1999.
- [4] *S. Kamiya*: Notes on elements of  $U(1, n; \mathbb{C})$ . *Hiroshima Math. J.* 21 (1991), 23–45.
- [5] *B. Maskit*: Kleinian Groups. Springer-Verlag, Berlin, 1988.
- [6] *J. R. Parker, I. D. Platis*: Complex hyperbolic Fenchel-Nielsen coordinates. *Topology* 47 (2008), 101–135.
- [7] *J. R. Parker*: Notes on Complex Hyperbolic Geometry. Cambridge University Press, Preprint, 2004.

*Authors' addresses*: Xi Fu, Department of Mathematics, Shaoxing College of Arts and Sciences, Shaoxing, Zhejiang 312000, People's Republic of China, e-mail: [foxi1000@yahoo.com.cn](mailto:foxi1000@yahoo.com.cn); Liulan Li, Department of Mathematics and computational science, Hengyang Normal University, Hengyang, Hunan 421008, People's Republic of China, e-mail: [lanlimail2008@yahoo.com.cn](mailto:lanlimail2008@yahoo.com.cn); Xiantao Wang, Department of Mathematics, Hunan Normal University, Changsha, Hunan 410081, People's Republic of China, e-mail: [xtwang@hunnu.edu.cn](mailto:xtwang@hunnu.edu.cn).