

Florian Luca

On a problem of Bednarek

Communications in Mathematics, Vol. 20 (2012), No. 2, 79--80

Persistent URL: <http://dml.cz/dmlcz/143140>

Terms of use:

© University of Ostrava, 2012

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>



VOLUME

20/2012

No. 2

ISSN 1804-1388

COMMUNICATIONS IN MATHEMATICS

Editor-in-Chief

Olga Rossi, The University of Ostrava & La Trobe University, Melbourne

Division Editors

Ilka Agricola, Philipps-Universität Marburg

Attila Bérczes, University of Debrecen

Anthony Bloch, University of Michigan

George Bluman, The University of British Columbia, Vancouver

Yong-Xin Guo, Eastern Liaoning University, Dandong

Haizhong Li, Tsinghua University, Beijing

Vilém Novák, The University of Ostrava

Štefan Porubský, Academy of Sciences of the Czech Republic, Prague

Geoff Prince, La Trobe University, Melbourne

Thomas Vetterlein, Johannes Kepler University Linz

Technical Editors

Jan Štěpnička, The University of Ostrava

Jan Šustek, The University of Ostrava

Available online at
<http://cm.osu.cz>



UNIVERSITAS
OSTRAVIENSIS

On a problem of Bednarek

Florian Luca

Abstract. We answer a question of Bednarek proposed at the 9th Polish, Slovak and Czech conference in Number Theory.

There are several problems in the literature concerning various arithmetic properties of the digit sum number function, see e.g. [1] and the references given there. In this paper, we deal with a particular problem. Namely, at the 9th Polish, Slovak and Czech Conference on Number Theory, June 11–14, 2012, W. Bednarek (via A. Schinzel) asked the following question.

Question *Is there a positive integer n divisible by $\underbrace{\overline{11\dots 1}}_{k \text{ times}}$ whose digit sum is less than k ?*

Here, we prove that the answer is no in a slightly more general setting. For integers $N \geq 1$ and $b \geq 2$, let $N = \overline{d_m d_{m-1} \dots d_0}_{(b)}$ be the base b representation of N , where $d_0, \dots, d_m \in \{0, 1, \dots, b-1\}$ with $d_m \neq 0$. We have the following result.

Theorem *If $n \geq 2$ is a multiple of $\underbrace{\overline{11\dots 1}}_{k \text{ times}}_{(b)}$, then the sum of its base b digits is greater than or equal to k .*

Proof. We may assume that $k \geq 2$, otherwise there is nothing to prove. Write

$$n = \sum_{i=0}^m d_i b^i,$$

where d_0, \dots, d_m are in $\{0, 1, \dots, b-1\}$ with $d_m \neq 0$. We may also assume that $d_0 \neq 0$. Put $N = (b^k - 1)/(b - 1)$. Then $b^k \equiv 1 \pmod{N}$. Thus,

$$n \equiv \sum_{j=0}^{k-1} c_j b^j \pmod{N},$$

where

$$c_j = \sum_{\substack{0 \leq i \leq m \\ i \equiv j \pmod{k}}} d_i.$$

It is clear that $c_0 + \cdots + c_{k-1}$ is the sum of the digits of n .
For each $\ell \in \{0, 1, \dots, k-1\}$, put

$$r_{j,\ell} = j + \ell - k \left\lfloor \frac{j + \ell}{k} \right\rfloor,$$

and consider the integer

$$m_\ell = \sum_{j=0}^{k-1} c_j b^{r_{j,\ell}}.$$

Note that since $b^k \equiv 1 \pmod{N}$, it follows that

$$m_\ell \equiv \sum_{j=0}^{k-1} b^{j+\ell} c_j \pmod{N} \equiv b^\ell n \pmod{N} \equiv 0 \pmod{N},$$

and since $c_j > 0$ for some j , we get that $m_\ell \geq N$. Summing this up for all $\ell \in \{0, 1, \dots, k-1\}$, we get

$$kN \leq \sum_{\ell=0}^{k-1} \sum_{j=0}^{k-1} b^{r_{j,\ell}} c_j = \sum_{j=0}^{k-1} c_j \sum_{\ell=0}^{k-1} b^{r_{j,\ell}} = N \sum_{j=0}^{k-1} c_j,$$

so $\sum_{j=0}^{k-1} c_j \geq k$, which is what we wanted to prove. \square

Note After this paper was submitted, we learned that Bednarek's question was also asked by Zhi-Wei Sun in [3], who solved the particular case when the modulus b is a prime. We also learned that the main result of this paper was obtained independently by Pan in [2].

Acknowledgements I thank the referee for comments which improved the quality of this note and Professor A. Schinzel for advice. This work was supported in part by Project PAPIIT 104512 and a Marcos Moshinsky fellowship.

References

- [1] F. Luca: Arithmetic properties of positive integers with a fixed digit sum. *Revista Matemática Iberoamericana* 22 (2006) 369–412.
- [2] H. Pan: On the m -ary expansion of a multiple of $(m^k - 1)/(m - 1)$. *Publ. Math. Debrecen*, to appear.
- [3] Z.-W. Sun: On divisibility concerning binomial coefficients. *J. Austral. Math. Soc.*, in press, arXiv:1005.1054.

Author's address:

FLORIAN LUCA: CENTRO DE CIENCIAS DE MATEMÁTICAS, UNIVERSIDAD NACIONAL AUTONOMA DE MÉXICO, C.P. 58089, MORELIA, MICHOACÁN, MÉXICO

E-mail: fluca@matmor.unam.mx

Received: 15 June, 2012

Accepted for publication: 24 August, 2012

Communicated by: Attila Bérczes