

M. Ferrer; S. Hernández; V. Uspenskij  
The dual space of precompact groups

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 54 (2013), No. 2, 239--244

Persistent URL: <http://dml.cz/dmlcz/143271>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2013

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## The dual space of precompact groups

M. FERRER, S. HERNÁNDEZ, V. USPENSKIJ

*Dedicated to the 120th birthday anniversary of Eduard Čech.*

*Abstract.* For any topological group  $G$  the dual object  $\widehat{G}$  is defined as the set of equivalence classes of irreducible unitary representations of  $G$  equipped with the Fell topology. If  $G$  is compact,  $\widehat{G}$  is discrete. In an earlier paper we proved that  $\widehat{G}$  is discrete for every metrizable precompact group, i.e. a dense subgroup of a compact metrizable group. We generalize this result to the case when  $G$  is an almost metrizable precompact group.

*Keywords:* compact group, precompact group, representation, Pontryagin–van Kampen duality, compact-open topology, Fell dual space, Fell topology, Kazhdan property (T)

*Classification:* Primary 43A40; Secondary 22A25, 22C05, 22D35, 43A35, 43A65, 54H11

### 1. Introduction

For a topological group  $G$  let  $\widehat{G}$  be the set of equivalence classes of irreducible unitary representations of  $G$ . The set  $\widehat{G}$  can be equipped with a natural topology, the so-called Fell topology (see Section 2 for a definition).

A topological group  $G$  is *precompact* if it is isomorphic (as a topological group) to a subgroup of a compact group  $H$  (we may assume that  $G$  is dense in  $H$ ). If  $H$  is compact, then  $\widehat{H}$  is discrete. If  $G$  is a dense subgroup of  $H$ , the natural mapping  $\widehat{H} \rightarrow \widehat{G}$  is a bijection but in general need not be a homeomorphism. Moreover, for every countable non-metrizable precompact group  $G$  the space  $\widehat{G}$  is not discrete [12, Theorem 5.1], and every non-metrizable compact group  $H$  has a dense subgroup  $G$  such that  $\widehat{G}$  is not discrete [12, Theorem 5.2]. (The Abelian case was considered in [5, 6, 14]). On the other hand, if  $G$  is a precompact metrizable group, then  $\widehat{G}$  is discrete [12, Theorem 4.1]. (The Abelian case was considered in [2], [4]). The aim of the present paper is to generalize this result to the almost metrizable case:  $\widehat{G}$  is discrete for every almost metrizable precompact topological group  $G$ . A topological group  $G$  is *almost metrizable* if it has a compact subgroup  $K$  such that the quotient space  $G/K$  is metrizable. According to Pasyнков's theorem [1, 4.3.20], a topological group is almost metrizable if and only if it is feathered in the sense of Arhangel'skii.

---

The first and second listed authors acknowledge partial support by the Spanish Ministry of Science, grant MTM2008-04599/MTM.

We reduce the almost metrizable case to the metrizable case considered in [12, Theorem 4.1].

## 2. Preliminaries: Fell topologies

All topological spaces and groups that we consider are assumed to be Hausdorff. For a (complex) Hilbert space  $\mathcal{H}$  the unitary group  $U(\mathcal{H})$  of all linear isometries of  $\mathcal{H}$  is equipped with the strong operator topology (this is the topology of pointwise convergence). With this topology,  $U(\mathcal{H})$  is a topological group.

A *unitary representation*  $\rho$  of the topological group  $G$  is a continuous homomorphism  $G \rightarrow U(\mathcal{H})$ , where  $\mathcal{H}$  is a complex Hilbert space. A closed linear subspace  $E \subseteq \mathcal{H}$  is an *invariant* subspace for  $\mathcal{S} \subseteq U(\mathcal{H})$  if  $ME \subseteq E$  for all  $M \in \mathcal{S}$ . If there is a closed subspace  $E$  with  $\{0\} \subsetneq E \subsetneq \mathcal{H}$  which is invariant for  $\mathcal{S}$ , then  $\mathcal{S}$  is called *reducible*; otherwise  $\mathcal{S}$  is *irreducible*. An *irreducible representation* of  $G$  is a unitary representation  $\rho$  such that  $\rho(G)$  is irreducible.

If  $\mathcal{H} = \mathbb{C}^n$ , we identify  $U(\mathcal{H})$  with the *unitary group of order  $n$* , that is, the compact Lie group of all complex  $n \times n$  matrices  $M$  for which  $M^{-1} = M^*$ . We denote this group by  $\mathbb{U}(n)$ .

Two unitary representations  $\rho : G \rightarrow U(\mathcal{H}_1)$  and  $\psi : G \rightarrow U(\mathcal{H}_2)$  are *equivalent* if there exists a Hilbert space isomorphism  $M : \mathcal{H}_1 \rightarrow \mathcal{H}_2$  such that  $\rho(x) = M^{-1}\psi(x)M$  for all  $x \in G$ . The *dual object* of a topological group  $G$  is the set  $\widehat{G}$  of equivalence classes of irreducible unitary representations of  $G$ .

If  $G$  is a precompact group, the Peter-Weyl Theorem (see [15]) implies that all irreducible unitary representation of  $G$  are finite-dimensional and determine an embedding of  $G$  into the product of unitary groups  $\mathbb{U}(n)$ .

If  $\rho : G \rightarrow U(\mathcal{H})$  is a unitary representation, a complex-valued function  $f$  on  $G$  is called a *function of positive type* (or *positive-definite function*) *associated with*  $\rho$  if there exists a vector  $v \in \mathcal{H}$  such that  $f(g) = (\rho(g)v, v)$  (here  $(\cdot, \cdot)$  denotes the inner product in  $\mathcal{H}$ ). We denote by  $P'_\rho$  the set of all functions of positive type associated with  $\rho$ . Let  $P_\rho$  be the convex cone generated by  $P'_\rho$ , that is, the set of sums of elements of  $P'_\rho$ .

Let  $G$  be a topological group,  $\mathcal{R}$  a set of equivalence classes of unitary representations of  $G$ . The *Fell topology* on  $\mathcal{R}$  is defined as follows: a typical neighborhood of  $[\rho] \in \mathcal{R}$  has the form

$$W(f_1, \dots, f_n, C, \epsilon) = \{[\sigma] \in \mathcal{R} : \exists g_1, \dots, g_n \in P_\sigma \forall x \in C |f_i(x) - g_i(x)| < \epsilon\},$$

where  $f_1, \dots, f_n \in P_\rho$  (or  $\in P'_\rho$ ),  $C$  is a compact subspace of  $G$ , and  $\epsilon > 0$ . In particular, the Fell topology is defined on the dual object  $\widehat{G}$ . If  $G$  is locally compact, the Fell topology on  $\widehat{G}$  can be derived from the Jacobson topology on the primitive ideal space of  $C^*(G)$ , the  $C^*$ -algebra of  $G$  [7, Section 18], [3, Remark F.4.5].

Every onto homomorphism  $f : G \rightarrow H$  of topological groups gives rise to a continuous injective dual map  $\hat{f} : \hat{H} \rightarrow \hat{G}$ . A mapping  $h : X \rightarrow Y$  between topological spaces is *compact-covering* if for every compact set  $L \subset Y$  there exists a compact set  $K \subset X$  such that  $h(K) = L$ .

**Lemma 2.1.** *If  $f : G \rightarrow H$  is a compact-covering onto homomorphism of topological groups, the dual map  $\hat{f} : \hat{H} \rightarrow \hat{G}$  is a homeomorphic embedding.*

PROOF: This easily follows from the definition of Fell topology. □

Let  $\pi$  be a unitary representation of a topological group  $G$  on a Hilbert space  $\mathcal{H}$ . Let  $F \subseteq G$  and  $\epsilon > 0$ . A unit vector  $v \in \mathcal{H}$  is called  $(F, \epsilon)$ -invariant if  $\|\pi(g)v - v\| < \epsilon$  for every  $g \in F$ .

A topological group  $G$  has *property (T)* if and only if there exists a pair  $(Q, \epsilon)$  (called a *Kazhdan pair*), where  $Q$  is a compact subset of  $G$  and  $\epsilon > 0$ , such that for every unitary representation  $\rho$  having a unit  $(Q, \epsilon)$ -invariant vector there exists a non-zero invariant vector. Equivalently,  $G$  has property (T) if and only if the trivial representation  $1_G$  is isolated in  $\mathcal{R} \cup \{1_G\}$  for every set  $\mathcal{R}$  of equivalence classes of unitary representations of  $G$  without non-zero invariant vectors [3, Proposition 1.2.3].

Compact groups have property (T) [3, Proposition 1.1.5], but countable Abelian precompact groups do not have property (T) [12, Theorem 6.1].

We refer to Fell’s papers [9], [10], the classical text by Dixmier [7] and the recent monographs by de la Harpe and Valette [13], and Bekka, de la Harpe and Valette [3] for basic definitions and results concerning Fell topologies and property (T).

### 3. Almost metrizable groups

If  $A$  is a subset of a topological space  $X$ , the *character*  $\chi(A, X)$  of  $A$  in  $X$  is the least cardinality of a base of neighborhoods of  $A$  in  $X$ . (If this definition leads to a finite value of  $\chi(A, X)$ , we replace it by  $\omega$ , the first infinite cardinal, and similarly for other cardinal invariants.) If  $A$  is a closed subset of a compact space  $X$ , the character  $\chi(A, X)$  equals the *pseudocharacter*  $\psi(A, X)$  — the least cardinality of a family  $\gamma$  of open subsets of  $X$  such that  $\bigcap \gamma = A$ . In particular, if  $A$  is a closed  $G_\delta$ -subset of a compact space  $X$ , then  $\chi(A, X) = \omega$ .

If  $K$  is a compact subgroup of a topological group, then  $G/K$  is metrizable if and only if  $\chi(K, G) = \omega$  [1, Lemma 4.3.19]. Let  $G$  be an almost metrizable topological group,  $\mathcal{K}$  the collection of all compact subgroups  $K \subset G$  such that  $\chi(K, G) = \omega$ . Then for every neighborhood  $O$  of the neutral element there is  $K \in \mathcal{K}$  such that  $K \subset O$  [1, Proposition 4.3.11]. We now show that if  $G$  is additionally  $\omega$ -narrow, then  $K$  can be chosen normal (in the algebraic sense). Recall that a topological group  $G$  is  $\omega$ -narrow [1] if for every neighborhood  $U$  of the neutral element there exists a countable set  $A \subset G$  such that  $AU = G$ .

**Lemma 3.1.** *Let  $G$  be an  $\omega$ -narrow almost metrizable group,  $\mathcal{N}$  the collection of all normal (= invariant under inner automorphisms) compact subgroups  $K$  of  $G$  such that the quotient group  $G/K$  is metrizable (equivalently,  $\chi(K, G) = \omega$ ).*

Then for every neighborhood  $O$  of the neutral element there exists  $K \in \mathcal{N}$  such that  $K \subset O$ .

PROOF: Let  $L \subset O$  be a compact subgroup of  $G$  such that the quotient space  $G/L = \{xL : x \in G\}$  is metrizable. It suffices to prove that  $K = \bigcap \{gLg^{-1} : g \in G\}$ , the largest normal subgroup of  $G$  contained in  $L$ , belongs to  $\mathcal{N}$ .

There exists a compatible metric on  $G/L$  which is invariant under the action of  $G$  by left translations. To construct such a metric, consider a countable base  $U_1, U_2, \dots$  of neighborhoods of  $L$  in  $G$ . We may assume that for each  $n$  we have  $U_n = U_n^{-1} = U_nL$  and  $U_{n+1}^2 \subset U_n$ . Let  $\gamma_n = \{gU_n : g \in G\}$ . The open cover  $\gamma_n$  of  $G$  is invariant under left  $G$ -translations and under right  $L$ -translations, and  $\gamma_{n+1}$  is a barycentric refinement of  $\gamma_n$ . The pseudometric on  $G$  that can be constructed in a canonical way from the sequence  $(\gamma_n)$  of open covers (see [8, Theorem 8.1.10]) gives rise to a compatible  $G$ -invariant metric on  $G/L$ . A similar construction was used in [1, Lemma 4.3.19].

If an  $\omega$ -narrow group transitively acts on a metric space  $X$  by isometries, then  $X$  is separable [1, 10.3.2]. Thus  $X = G/L$  is separable. Let  $Y$  be a dense countable subset of  $X$ . Then  $K = \{g \in G : gx = x \text{ for every } x \in X\} = \{g \in G : gx = x \text{ for every } x \in Y\}$  is a  $G_\delta$ -subset of  $L$ , hence  $\chi(K, L) = \omega$ . It follows that  $\chi(K, G) \leq \chi(K, L)\chi(L, G) = \omega$  ([8, Exercise 3.1.E]).  $\square$

#### 4. Main theorem

**Theorem 4.1.** *If  $G$  is a precompact almost metrizable group, then  $\widehat{G}$  is discrete.*

PROOF: Let  $\rho$  be an irreducible unitary representation of  $G$ . We must prove that  $[\rho]$  is isolated in  $\widehat{G}$ . It suffices to find a discrete open subset  $D \subset \widehat{G}$  such that  $[\rho] \in D$ .

Precompact groups are  $\omega$ -narrow, so Lemma 3.1 applies to  $G$ . Let  $\mathcal{N}$ , as above, be the collection of all normal compact subgroups  $K \subset G$  such that  $\chi(K, G) = \omega$ . Then  $\mathcal{N}$  is closed under countable intersections, and it follows from Lemma 3.1 that for every  $G_\delta$ -subset  $A$  of  $G$  containing the neutral element there exists  $K \in \mathcal{N}$  such that  $K \subset A$ . In particular, there exists  $K \in \mathcal{N}$  such that  $K$  lies in the kernel of  $\rho$ . Let  $D \subset \widehat{G}$  be the set of all classes  $[\sigma] \in \widehat{G}$  such that  $K$  is contained in the kernel of  $\sigma$ . Then  $[\rho] \in D$ . It suffices to verify that  $D$  is open and discrete.

Step 1. We verify that  $D$  is open. Let  $\mathcal{R}$  be the set of equivalence classes of all finite-dimensional unitary representations (which may be reducible) of  $K$  without non-zero invariant vectors. Let  $\tau_n$  be the trivial  $n$ -dimensional representation  $1_K \oplus \dots \oplus 1_K$  ( $n$  summands) of  $K$ ,  $n = 1, 2, \dots$ . In the notation of Section 2,  $P_{\tau_n}$  does not depend on  $n$  and is the set of non-negative constant functions on  $K$ . It follows that in the space  $\mathcal{S} = \mathcal{R} \cup \{[\tau_n] : n = 1, 2, \dots\}$ , equipped with the Fell topology, the points  $[\tau_n]$  are indistinguishable: any open set containing one of these points contains all the others. Since  $K$  has property (T),  $[\tau_1] = [1_K]$  is not in the closure of  $\mathcal{R}$ . Therefore  $\mathcal{R}$  is closed in  $\mathcal{S}$  and  $\mathcal{S} \setminus \mathcal{R}$  is open in  $\mathcal{S}$ .

We claim that for every irreducible unitary representation  $\sigma$  of  $G$  the class of the restriction  $\sigma|_K$  belongs to  $\mathcal{S}$ . In other words, the claim is that  $\sigma|_K$  is trivial if it admits a non-zero invariant vector. Let  $V$  be the (finite-dimensional) space of the representation  $\sigma$ . For  $g \in G$  and  $x \in V$  we write  $gx$  instead of  $\sigma(g)x$ . The space  $V' = \{x \in V : gx = x \text{ for all } g \in K\}$  of all  $K$ -invariant vectors is  $G$ -invariant. Indeed, if  $x \in V'$ ,  $g \in G$  and  $h \in K$ , then  $g^{-1}hgx = x$  because  $g^{-1}hg \in K$  and  $x$  is  $K$ -invariant. It follows that  $hgx = gx$  which proves that  $gx \in V'$ . Since  $\sigma$  is irreducible, either  $V' = \{0\}$  or  $V' = V$ . Accordingly, either  $\sigma|_K$  admits no non-zero invariant vectors or else is trivial.

We have just proved that the restriction map  $r : \widehat{G} \rightarrow \mathcal{S}$  is well-defined. Clearly  $r$  is continuous, and therefore  $D = r^{-1}(\mathcal{S} \setminus \mathcal{R})$  is open in  $\widehat{G}$ .

Step 2. We verify that  $D$  is discrete. Let  $p : G \rightarrow G/K$  be the quotient map. Then  $D$  is the image of the dual map  $\widehat{p} : \widehat{G/K} \rightarrow \widehat{G}$ . According to [12, Theorem 4.1], the dual space of a metrizable precompact group is discrete. Thus  $\widehat{G/K}$  is discrete. Since  $p$  is a perfect map, it is compact-covering, and Lemma 2.1 implies that  $\widehat{p} : \widehat{G/K} \rightarrow \widehat{G}$  is a homeomorphic embedding. Therefore,  $D = \widehat{p}(\widehat{G/K})$  is discrete.  $\square$

## REFERENCES

- [1] Arhangel'skii A., Tkachenko M., *Topological Groups and Related Structures*, Atlantis Press, Amsterdam-Paris, 2008.
- [2] Aussenhofer L., *Contributions to the duality theory of Abelian topological groups and to the theory of nuclear groups*, Dissertation, Tübingen 1998; *Dissertationes Math. (Rozprawy Mat.)* **384** (1999).
- [3] Bekka B., de la Harpe P., Valette A., *Kazhdan's Property (T)*, Cambridge University Press, Cambridge, 2008.
- [4] Chasco M.J., *Pontryagin duality for metrizable groups*, *Arch. Math.* **70** (1998), 22–28.
- [5] Comfort W.W., Raczkowski S.U., Trigos-Arrieta F.J., *The dual group of a dense subgroup*, *Czechoslovak Math. J.* **54** (129) (2004), 509–533.
- [6] Dikranjan D., Shakhmatov D., *Quasi-convex density and determining subgroups of compact Abelian groups*, *J. Math. Anal. Appl.* **363** (2010), no. 1, 42–48.
- [7] Dixmier J., *Les  $C^*$ -algèbres et leurs représentations*, Gauthier-Villars, Paris, 1969.
- [8] Engelking R., *General Topology*, revised and completed edition, Heldermann Verlag, Berlin, 1989.
- [9] Fell J.M.G., *The dual spaces of  $C^*$ -algebras*, *Trans. Amer. Math. Soc.* **94** (1960), 365–403.
- [10] Fell J.M.G., *Weak containment and induced representations of groups*, *Canad. J. Math.* **14** (1962), 237–268.
- [11] Ferrer M.V., Hernández S., *Dual topologies on groups*, *Topology Appl.*, to appear.
- [12] Ferrer M.V., Hernández S., Uspenskij V., *Precompact groups and property (T)*, arXiv:1112.1350
- [13] de la Harpe P., Valette A., *La propriété (T) de Kazhdan pour les groupes localement compacts*, *Astérisque* **175**, Soc. Math. France, 1989.
- [14] Hernández S., Macario S., Trigos-Arrieta F.J., *Uncountable products of determined groups need not be determined*, *J. Math. Anal. Appl.* **348** (2008), no. 2, 834–842.

- [15] Hofmann K.H., Morris S.A., *The Structure of Compact Groups: A Primer for Students - a Handbook for the Expert*, De Gruyter Studies in Mathematics, 25, Walter de Gruyter, Berlin-New York, 2006.

UNIVERSITAT JAUME I, INSTITUTO DE MATEMÁTICAS DE CASTELLÓN, CAMPUS DE RIU SEC, 12071 CASTELLÓN, SPAIN

*E-mail:* mferrer@mat.uji.es

UNIVERSITAT JAUME I, INIT AND DEPARTAMENTO DE MATEMÁTICAS, CAMPUS DE RIU SEC, 12071 CASTELLÓN, SPAIN

*E-mail:* hernande@mat.uji.es

DEPARTMENT OF MATHEMATICS, OHIO UNIVERSITY, 321 MORTON HALL, ATHENS, OHIO 45701, USA

*E-mail:* uspenski@ohio.edu

(Received December 12, 2012)