

Applications of Mathematics

Changjin Xu; Qianhong Zhang; Maoxin Liao

Existence and global attractivity of positive periodic solutions for a delayed competitive system with the effect of toxic substances and impulses

Applications of Mathematics, Vol. 58 (2013), No. 3, 309--328

Persistent URL: <http://dml.cz/dmlcz/143280>

Terms of use:

© Institute of Mathematics AS CR, 2013

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

EXISTENCE AND GLOBAL ATTRACTIVITY OF POSITIVE
PERIODIC SOLUTIONS FOR A DELAYED COMPETITIVE SYSTEM
WITH THE EFFECT OF TOXIC SUBSTANCES AND IMPULSES

CHANGJIN XU, QIANHONG ZHANG, Guiyang, MAOXIN LIAO, Hengyang

(Received April 28, 2011)

Abstract. In this paper, a class of non-autonomous delayed competitive systems with the effect of toxic substances and impulses is considered. By using the continuation theorem of coincidence degree theory, we derive a set of easily verifiable sufficient conditions that guarantees the existence of at least one positive periodic solution, and by constructing a suitable Lyapunov functional, the uniqueness and global attractivity of the positive periodic solution are established.

Keywords: competitive system, toxic substance, periodic solution, impulse, coincidence degree theory

MSC 2010: 34K13, 34K25

1. INTRODUCTION

In recent years, the dynamical behavior of a competitive system has been one of the dominant themes in both ecology and mathematical ecology due to its universal existence and importance. Most widely studied competitive systems are mainly continuous or discrete [2], [3], [5], [7], [10], [11], [13], [14], [15]. Recently there has been a new category of competitive systems, which are neither purely continuous-time nor purely discrete-time ones; these are called impulsive competitive system. This category of impulsive competitive systems displays a combination of characteristics of both the continuous-time and discrete-time systems [4], [8].

This work is supported by National Natural Science Foundation of China (No. 11261010 and No. 10961008), Soft Science and Technology Program of Guizhou Province (No. 2011LKC2030), Natural Science and Technology Foundation of Guizhou Province (J[2012]2100), Governor Foundation of Guizhou Province ([2012]53) and Doctoral Foundation of Guizhou University of Finance and Economics (2010).

In 2003, Song and Chen [14] proposed a delay two-species competitive system in which two species have toxic inhibitory effects on each other:

$$(1.1) \quad \begin{cases} \frac{dx}{dt} = x(t)[K_1(t) - \alpha_1(t)x(t) - \beta_1(t)y(t) - \gamma_1(t)x(t)y(t - \tau_1(t))], \\ \frac{dy}{dt} = y(t)[K_2(t) - \alpha_2(t)y(t) - \beta_2(t)x(t) - \gamma_2(t)x(t - \tau_2(t))y(t)], \end{cases}$$

where $x(t), y(t)$ stand for the population densities of two competing species, respectively. $K_i(t)$ ($i = 1, 2$) are the intrinsic growth rates of the two competing species; $\alpha_i(t)$ ($i = 1, 2$) denote the coefficients of interspecific competition; $K_i(t)/\alpha_i(t)$ ($i = 1, 2$) are the environmental carrying capacities of two competing species; γ_1 and γ_2 stand for, respectively, the rates of toxic inhibition of the species x by the species y and vice versa. For more details about the model, one can see [12]. By applying the coincidence degree theory, Song and Chen [12] established the existence of a positive periodic solution for system (1.1).

Considering the impulsive effects and periodic perturbations, Liu et al. [9] investigated the periodic impulsive delay competitive system with the effect of toxic substances

$$(1.2) \quad \begin{cases} \frac{dx}{dt} = x(t)[K_1(t) - \alpha_1(t)x(t) - \beta_1(t)y(t) - \gamma_1(t)x(t)y(t - \tau_1(t))], & t \neq t_k, \\ \frac{dy}{dt} = y(t)[K_2(t) - \alpha_2(t)y(t) - \beta_2(t)x(t) - \gamma_2(t)x(t - \tau_2(t))y(t)], & t \neq t_k, \\ x(t_k^+) = x(t_k) + p, & t = t_k, \\ y(t_k^+) = (1 + b_k)y(t_k), & t = t_k \end{cases}$$

with an initial condition $(x(s), y(s)) = \varphi(s) = (\varphi_1(s), \varphi_2(s))$ for $-\tau \leq s \leq 0, \varphi(0) > 0, \varphi \in PC([- \tau, 0], \mathbb{R}_+^2)$, where $\tau = \max_{1 \leq i \leq 2} \max_{t \in [0, \omega]} \{\tau_i(t)\}$; $K_i(t), \alpha_i(t), \beta_i(t), \gamma_i(t), \tau_i(t)$ ($i = 1, 2$) are continuous ω -periodic functions, and $\alpha_i(t), \beta(t), \gamma_i(t)$ ($i = 1, 2$) are positive and $\tau_i(t)$ ($i = 1, 2$) are nonnegative. The intrinsic growth rates $K_i(t)$ ($i = 1, 2$) are not necessarily positive and may be negative. Also $k \in \mathbb{N}$ and \mathbb{N} is the set of positive integers. The jump conditions reflect the possibility of impulsive effects on the species x and y . $p > 0$ is the impulsive stocking amount of the species x at time $t = t_k$, which implies that the populations are subject to impulsive stocking at a constant rate p . The term $b_k y(t_k) < 0$ ($k \in \mathbb{N}$) represents the impulsive harvesting amount of the species y at time $t = t_k$, while $b_k y(t_k) > 0$ which represents the perturbations may stand for the impulsive stocking amount of the species y at time $t = t_k$. By applying the theory of impulsive differential equations and some analysis techniques, Liu et al. [9] obtained a set of sufficient conditions for the permanence and partial extinction of system (1.2).

Considering that the harvest of many populations is not continuous, the harvest can be viewed as an annual harvest pulse. To describe a system more accurately, we should consider using impulsive differential equations. Then system (1.1) is revised into the following form:

$$(1.3) \quad \begin{cases} \frac{dx}{dt} = x(t)[K_1(t) - \alpha_1(t)x(t) - \beta_1(t)y(t) - \gamma_1(t)x(t)y(t - \tau_1(t))], & t \neq t_k, \\ \frac{dy}{dt} = y(t)[K_2(t) - \alpha_2(t)y(t) - \beta_2(t)x(t) - \gamma_2(t)x(t - \tau_2(t))y(t)], & t \neq t_k, \\ \Delta x(t_k) = x(t_k^+) - x(t_k^-) = \varrho_{1k}x(t_k), & t = t_k, \\ \Delta y(t_k) = y(t_k^+) - y(t_k^-) = \varrho_{2k}y(t_k), & t = t_k, \end{cases}$$

where $\Delta x(t_k) = x(t_k^+) - x(t_k^-)$ and $\Delta y(t_k) = y(t_k^+) - y(t_k^-)$ are the impulses at moments t_k and $t_1 < t_2 < \dots$ is a strictly increasing sequence such that $\lim_{k \rightarrow \infty} t_k = +\infty$.

The principal object of this article is by using Mawhin's continuation theorem of coincidence degree theory and by constructing Lyapunov functions to investigate the stability and existence of periodic solutions of (1.3). To the best of the authors' knowledge, it is the first time one deals with the existence and stability of periodic solutions of (1.3).

In order to obtain our main results, throughout the paper we always assume that the following conditions are fulfilled:

- (H1) $K_i(t)$, $\alpha_i(t)$, $\beta_i(t)$, $\gamma_i(t)$ ($i = 1, 2$), are all continuous ω periodic, i.e., $K_i(t + \omega) = K_i(t)$, $\alpha_i(t + \omega) = \alpha_i(t)$, $\beta_i(t + \omega) = \beta_i(t)$ ($i = 1, 2$), $\gamma_i(t + \omega) = \gamma_i(t)$ for any $t \in \mathbb{R}$.
- (H2) $K_i(t)$, $\alpha_i(t)$, $\beta_i(t)$, $\gamma_i(t)$ ($i = 1, 2$) are all positive.
- (H3) $\varrho_{ik} \geq 0$ for all $k \in \mathbb{N}$ and there exists a positive integer q such that $t_{k+q} = t_k + \omega$, $\varrho_{ik+q} = \varrho_{ik}$ ($i = 1, 2; k = 1, 2, 3, \dots$).
- (H4) At least one of the following four conditions $\bar{\alpha}_1\bar{\alpha}_2 \neq \bar{\beta}_1\bar{\beta}_2$, $\bar{\alpha}_1\bar{\gamma}_2 \neq \bar{\beta}_2\bar{\gamma}_1$, $\bar{\alpha}_2\bar{\gamma}_2 \neq \bar{\beta}_1\bar{\gamma}_2$, $\bar{\gamma}_2^2 \neq \bar{\gamma}_1\bar{\gamma}_2$ holds.

The organization of the paper is as follows. In Section 2, we introduce some notation and definitions, and state some preliminary results needed in later sections. We then establish, in Section 3, some simple criteria for the existence of positive periodic solutions of system (1.3) by using the continuation theorem of the coincidence degree theory proposed by Gaines and Mawhin [6]. In Section 4, the uniqueness and global attractivity of the positive periodic solution are presented. In Section 5, an illustrative example is given to demonstrate the correctness of the results obtained.

2. PRELIMINARIES

We shall introduce some notation and definitions, and state some preliminary results. Consider the impulsive system

$$(2.1) \quad \begin{cases} \dot{x}(t) = f(t, x), & t \neq t_k, \quad k = 1, 2, \dots, \\ \Delta x(t)|_{t=t_k} = I_k(x(t_k^-)), \end{cases}$$

where $x \in \mathbb{R}^n$, $f: \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is continuous and $f(t+\omega, x) = f(t, x)$; $I_k: \mathbb{R}^n \rightarrow \mathbb{R}^n$ are continuous, and there exists a positive integer q such that $t_{k+q} = t_k + \omega$, $I_{k+q}(x) = I_k(x)$ with $t_k \in \mathbb{R}$, $t_{k+1} > t_k$, $\lim_{k \rightarrow \infty} t_k = \infty$, $\Delta x(t)|_{t=t_k} = x(t_k^+) - x(t_k^-)$. For $t_k \neq 0$ ($k = 1, 2, \dots$), $[0, \omega] \cap \{t_k\} = \{t_1, t_2, \dots, t_q\}$. As we know, $\{t_k\}$ are called the points of jump.

Let us recall some definitions for the Cauchy problem

$$(2.2) \quad \begin{cases} \dot{x}(t) = f(t, x), & t \in [0, \omega], \quad t \neq t_k, \\ \Delta x(t)|_{t=t_k} = I_k(x(t_k^-)), x(0) = x_0. \end{cases}$$

Definition 1.1. A map $x: [0, \omega] \rightarrow \mathbb{R}^n$ is said to be a solution of (2.2), if it satisfies the following conditions:

- (i) $x(t)$ is a piecewise continuous map with first-class discontinuity points at $t_k \cap [0, \omega]$, and at each discontinuity point it is continuous from the left;
- (ii) $x(t)$ satisfies (2.2).

Definition 1.2. A map $x: [0, \omega] \rightarrow \mathbb{R}^n$ is said to be an ω periodic solution of (2.1), if

- (i) $x(t)$ satisfies (i) and (ii) of Definition 1 in the interval $[0, \omega]$;
- (ii) $x(t)$ satisfies $x(t + \omega - 0) = x(t - 0)$, $t \in \mathbb{R}$.

Obviously, if $x(t)$ is a solution of (2.2) defined on $[0, \omega]$ such that $x(0) = x(\omega)$, then by the periodicity of (2.2) in t , the function $x^*(t)$ defined by

$$x^*(t) = \begin{cases} x(t - j\omega), & t \in [j\omega, (j+1)\omega] \setminus \{t_k\}, \\ x^*(t) \text{ is left continuous at } t = t_k \end{cases}$$

is an ω periodic solution of (2.1).

For system (1.3), finding the periodic solutions is equivalent to finding solutions of the following boundary value problem:

$$(2.3) \quad \begin{cases} \frac{dx}{dt} = x(t)[K_1(t) - \alpha_1(t)x(t) - \beta_1(t)y(t) - \gamma_1(t)x(t)y(t - \tau_1(t))], \\ \quad t \neq t_k, t \in [0, \omega], k = 1, 2, \dots, q, \\ \frac{dy}{dt} = y(t)[K_2(t) - \alpha_2(t)y(t) - \beta_2(t)x(t) - \gamma_2(t)x(t - \tau_2(t))y(t)], \\ \quad t \neq t_k, t \in [0, \omega], k = 1, 2, \dots, q, \\ \Delta x(t_k) = x(t_k^+) - x(t_k^-) = \varrho_{1k}x(t_k), t = t_k, x(0) = x(\omega), k = 1, 2, \dots, q, \\ \Delta y(t_k) = y(t_k^+) - y(t_k^-) = \varrho_{2k}y(t_k), t = t_k, y(0) = y(\omega), k = 1, 2, \dots, q. \end{cases}$$

3. EXISTENCE OF POSITIVE PERIODIC SOLUTIONS

In this section, based on Mawhin's continuation theorem, we shall study the existence of at least one periodic solution of (1.3). To do so, we shall make some preparations.

Let X, Y be normed vector spaces, $L: \text{Dom } L \subset X \rightarrow Y$ a linear mapping, $N: X \rightarrow Y$ a continuous mapping. The mapping L will be called a Fredholm mapping of index zero if $\dim \text{Ker } L = \text{codim } \text{Im } L < +\infty$ and $\text{Im } L$ is closed in Y . If L is a Fredholm mapping of index zero and there exist continuous projections $P: X \rightarrow X$ and $Q: Y \rightarrow Y$ such that $\text{Im } P = \text{Ker } L$, $\text{Im } L = \text{Ker } Q = \text{Im}(I - Q)$, it follows that $L|_{\text{Dom } L \cap \text{Ker } P}: (I - P)X \rightarrow \text{Im } L$ is invertible. We denote the inverse of that map by K_P . If Ω is an open bounded subset of X , the mapping N will be called L -compact on $\overline{\Omega}$ if $QN(\overline{\Omega})$ is bounded and $K_P(I - Q)N: \overline{\Omega} \rightarrow X$ is compact. Since $\text{Im } Q$ is isomorphic to $\text{Ker } L$, there exist isomorphisms $J: \text{Im } Q \rightarrow \text{Ker } L$.

Now we introduce Mawhin's continuation theorem [6] as follows.

Lemma 3.1 ([6] Continuation Theorem). *Let L be a Fredholm mapping of index zero and let N be L -compact on $\overline{\Omega}$. Suppose*

- (a) *for each $\lambda \in (0, 1)$, every solution x of $Lx = \lambda Nx$ is such that $x \notin \partial\Omega$;*
- (b) *$QNx \neq 0$ for each $x \in \text{Ker } L \cap \partial\Omega$, and $\deg\{JQN, \Omega \cap \text{Ker } L, 0\} \neq 0$.*

Then the equation $Lx = Nx$ has at least one solution lying in $\text{Dom } L \cap \overline{\Omega}$.

For convenience and simplicity of the following discussion, we use the notation below throughout the paper:

$$\bar{f} = \frac{1}{\omega} \int_0^\omega f(t) dt, \quad f^L = \min_{t \in [0, \omega]} f(t), \quad f^M = \max_{t \in [0, \omega]} f(t), \quad |\bar{f}| = \frac{1}{\omega} \int_0^\omega |f(t)| dt,$$

where $f(t)$ is a ω continuous periodic function. For any non-negative integer p , let $C^{(p)}[0, \omega; t_1, t_2, \dots, t_q] = \{x: [0, \omega] \rightarrow \mathbb{R}^m | x^{(p)}(t) \text{ exist for } t \neq t_1, \dots, t_q; x^{(p)}(t^+) \text{ and } x^{(p)}(t^-) \text{ exist at } t_1, t_2, \dots, t_q; \text{ and } x^{(j)}(t_k) = x^{(j)}(t_k^-), k = 1, \dots, m, j = 0, 1, 2, \dots, p\}$ with the norm $\|x\|_p = \max\left\{\sup_{t \in [0, \omega]} \|x^{(j)}(t)\|\right\}_{j=1}^p$, where $\|\cdot\|$ is any norm in \mathbb{R}^m . It is easy to see that $C^{(p)}[0, \omega; t_1, t_2, \dots, t_q]$ is a Banach space.

Now we are in a position to state and prove the existence of periodic solutions of (2.3).

Theorem 3.1. *Let B_3 and B_9 be defined by (3.12) and (3.20), respectively. In addition to (H1)–(H4), assume further that*

$$(H5) \quad \bar{K}_2\omega + \sum_{k=1}^q \ln(1 + \varrho_{2k}) > \max\{\bar{\gamma}_2\omega \exp(B_3), \bar{\gamma}_2\omega \exp(B_9)\},$$

then the system (1.2) has at least one ω periodic solution.

Proof. According to the discussion above in Section 2, we only need to prove that the boundary value problem (2.3) has a solution. Since the solutions of (2.3) remain positive for all $t \geq 0$, we let $u_1(t) = \ln[x(t)]$, $u_2(t) = \ln[y(t)]$. Then system (2.3) can be transformed to

$$(3.1) \quad \begin{cases} \dot{u}_1(t) = K_1(t) - \alpha_1(t) \exp(u_1(t)) - \beta_1(t) \exp(u_2(t)) \\ \quad - \gamma_1(t) \exp(u_1(t)) \exp(u_2(t - \tau_1(t))), \\ \quad t \neq t_k, t \in [0, \omega], k = 1, 2, \dots, q, \\ \dot{u}_2(t) = K_2(t) - \alpha_2(t) \exp(u_2(t)) - \beta_2(t) \exp(u_1(t)) \\ \quad - \gamma_2(t) \exp(u_1(t - \tau_2(t))) \exp(u_2(t)), \\ \quad t \neq t_k, t \in [0, \omega], k = 1, 2, \dots, q, \\ \Delta u_i(t_k) = \ln(1 + \varrho_{ik}), t = t_k, k = 1, 2, \dots, q; \\ u_1(0) = u_1(\omega), u_2(0) = u_2(\omega). \end{cases}$$

It is easy to see that if system (3.1) has an ω periodic solution $(u_1^*(t), u_2^*(t))^T$, then $(x^*(t), y^*(t))^T = (e^{u_1^*(t)}, e^{u_2^*(t)})^T$ is a positive solution of system (1.3). Therefore, to complete the proof, it suffices to show that system (3.1) has at least one ω periodic solution.

In order to use the continuation theorem of coincidence degree theory to establish the existence of a solution of (3.1), we take

$$X = \{u \in C[0, \omega; t_1, t_2, \dots, t_q]\}, Y = X \times \mathbb{R}^{2 \times (q+1)}.$$

Then X is a Banach space with the norm $\|\cdot\|_0$, and Y is also a Banach space with the norm $\|z\| = \|x\|_0 + \|y\|$, $x \in X$, $y \in \mathbb{R}^{2 \times (q+1)}$.

Let

$$\text{Dom } L = \{u = (u_1, u_2)^T \in C[0, \omega]; t_1, t_2, \dots, t_q\},$$

$$L: \text{Dom } L \subset X \rightarrow Y, x \rightarrow (x', \Delta u(t_1), \Delta u(t_2), \dots, \Delta u(t_q), 0),$$

$$N: X \rightarrow Y,$$

$$Nu = \left(\begin{aligned} & \left(K_1(t) - \alpha_1(t) \exp(u_1(t)) - \beta_1(t) \exp(u_2(t)) - \gamma_1(t) \exp(u_1(t)) \exp(u_2(t - \tau_1(t))) \right) \\ & \left(K_2(t) - \alpha_2(t) \exp(u_2(t)) - \beta_2(t) \exp(u_1(t)) - \gamma_2(t) \exp(u_1(t - \tau_2(t))) \exp(u_2(t)) \right) \end{aligned} \right), \\ \left(\ln(1 + \varrho_{11}) \right), \left(\ln(1 + \varrho_{21}) \right), \dots, \left(\ln(1 + \varrho_{31}) \right), \left(\ln(1 + \varrho_{32}) \right), 0 \Big).$$

Obviously,

$$\text{Ker } L = \{u: u(t) = h \in \mathbb{R}^2, t \in [0, \omega]\},$$

$$\text{Im } L = \left\{ z = (f, a_1, a_2, \dots, a_q, d) \in Y: \int_0^\omega f(s) ds + \sum_{k=1}^q a_k + d = 0 \right\} \\ = X \times \mathbb{R}^{2 \times q} \times \{0\},$$

$$\dim \text{Ker } L = 2 = \text{codim Im } L.$$

So, $\text{Im } L$ is closed in Y , L is a Fredholm mapping of index zero. Define two projections

$$Px = \frac{1}{\omega} \int_0^\omega x(t) dt,$$

$$Qz = Q(f, a_1, a_2, \dots, a_q, d) = \left(\frac{1}{\omega} \left[\int_0^\omega f(s) ds + \sum_{k=1}^q a_k + d \right], 0, 0, \dots, 0 \right).$$

It is easy to show that P and Q are continuous and satisfy $\text{Im } P = \text{Ker } L, \text{Im } L = \text{Ker } Q = \text{Im}(I - Q)$.

Further, through an easy computation, we can find that the inverse K_P of L , $K_P: \text{Im } L \rightarrow \text{Ker } P \cap \text{Dom } L$ has the following form:

$$K_P(z) = \int_0^t f(s) ds + \sum_{t_k < t} a_k - \frac{1}{\omega} \int_0^\omega \int_0^t f(s) ds dt - \sum_{k=1}^q a_k.$$

Moreover, it is easy to check that

$$\begin{aligned}
QN u &= \left(\left(\frac{1}{\omega} \int_0^t F_1(s) \, ds + \frac{1}{\omega} \sum_{k=1}^q \ln(1 + \varrho_{1k}) \right), 0, 0, \dots, 0 \right), \\
K_P(I - Q)Nu &= \left(\int_0^t F_1(s) \, ds + \sum_{t > t_k} \ln(1 + \varrho_{1k}) \right) \\
&\quad - \left(\frac{1}{\omega} \int_0^\omega \int_0^t F_1(s) \, ds \, dt + \sum_{k=1}^q \ln(1 + \varrho_{1k}) \right) \\
&\quad - \left(\frac{t}{\omega} - \frac{1}{2} \right) \int_0^\omega F_1(s) \, ds + \sum_{k=1}^q \ln(1 + \varrho_{1k}) \\
&\quad - \left(\int_0^t F_2(s) \, ds + \sum_{t > t_k} \ln(1 + \varrho_{2k}) \right) \\
&\quad - \left(\frac{1}{\omega} \int_0^\omega \int_0^t F_2(s) \, ds \, dt + \sum_{k=1}^q \ln(1 + \varrho_{2k}) \right) \\
&\quad - \left(\frac{t}{\omega} - \frac{1}{2} \right) \int_0^\omega F_2(s) \, ds + \sum_{k=1}^q \ln(1 + \varrho_{2k}) \right),
\end{aligned}$$

where

$$\begin{aligned}
F_1(s) &= K_1(s) - \alpha_1(s) \exp(u_1(s)) - \beta_1(s) \exp(u_2(s)) \\
&\quad - \gamma_1(s) \exp(u_1(s)) \exp(u_2(s - \tau_1(s))), \\
F_2(s) &= K_2(s) - \alpha_2(s) \exp(u_2(s)) - \beta_2(s) \exp(u_1(s)) \\
&\quad - \gamma_2(s) \exp(u_1(s - \tau_2(s))) \exp(u_2(s)).
\end{aligned}$$

Obviously, QN and $K_P(I - Q)N$ are continuous. Since X is a finite-dimensional Banach space, using the Ascoli-Arzelà theorem, it is not difficult to show that $\overline{K_P(I - Q)N(\overline{\Omega})}$ is compact for any open bounded set $\Omega \subset X$. Moreover, $QN(\overline{\Omega})$ is bounded. Thus, N is L -compact on $\overline{\Omega}$ with any open bounded set $\Omega \subset X$.

Now we are at the point to search for an appropriate open, bounded subset Ω for the application of the continuation theorem. Corresponding to the operator equation

$Lu = \lambda Nu$, $\lambda \in (0, 1)$, we have

$$(3.2) \quad \left\{ \begin{array}{l} \dot{u}_1(t) = \lambda[K_1(t) - \alpha_1(t) \exp(u_1(t)) - \beta_1(t) \exp(u_2(t)) \\ \quad - \gamma_1(t) \exp(u_1(t)) \exp(u_2(t - \tau_1(t)))] \\ \quad t \neq t_k, t \in [0, \omega], k = 1, 2, \dots, q, \\ \dot{u}_2(t) = \lambda[K_2(t) - \alpha_2(t) \exp(u_2(t)) - \beta_2(t) \exp(u_1(t)) \\ \quad - \gamma_2(t) \exp(u_1(t - \tau_2(t))) \exp(u_2(t))] \\ \quad t \neq t_k, t \in [0, \omega], k = 1, 2, \dots, q, \\ \Delta u_i(t_k) = \lambda \ln(1 + \varrho_{ik}), i = 1, 2; \\ \quad k = 1, 2, \dots, q; u_1(0) = u_1(\omega), u_2(0) = u_2(\omega). \end{array} \right.$$

Suppose that $u(t) = (u_1(t), u_2(t))^T \in X$ is an arbitrary solution of system (3.2) for a certain $\lambda \in (0, 1)$. Integrating both sides of (3.2) over the interval $[0, \omega]$ with respect to t , we obtain

$$(3.3) \quad \left\{ \begin{array}{l} \int_0^\omega f_1(t) dt = \sum_{k=1}^q \ln(1 + \varrho_{1k}) + \int_0^\omega K_1(t) dt, \\ \int_0^\omega f_2(t) dt = \sum_{k=1}^q \ln(1 + \varrho_{2k}) + \int_0^\omega K_2(t) dt, \end{array} \right.$$

where

$$\begin{aligned} f_1(t) &= \alpha_1(t) \exp(u_1(t)) + \beta_1(t) \exp(u_2(t)) + \gamma_1(t) \exp(u_1(t)) \exp(u_2(t - \tau_1(t))), \\ f_2(t) &= \alpha_2(t) \exp(u_2(t)) + \beta_2(t) \exp(u_1(t)) + \gamma_2(t) \exp(u_1(t - \tau_2(t))) \exp(u_2(t)). \end{aligned}$$

From (3.2) and (3.3), we can obtain

$$(3.4) \quad \int_0^\omega |\dot{u}_1(t)| dt < 2\bar{K}_1\omega + \sum_{k=1}^q \ln(1 + \varrho_{1k}),$$

$$(3.5) \quad \int_0^\omega |\dot{u}_2(t)| dt < 2\bar{K}_2\omega + \sum_{k=1}^q \ln(1 + \varrho_{2k}).$$

Let

$$(3.6) \quad u_i(\xi_i) = \min_{t \in [0, \omega]} u_i(t), u_i(\eta_i) = \max_{t \in [0, \omega]} u_i(t), i = 1, 2.$$

Then, by (3.3), we get

$$\int_0^\omega f_1(t) dt < 2\bar{K}_1\omega + \sum_{k=1}^q \ln(1 + \varrho_{1k}),$$

which leads to

$$\int_0^\omega \alpha_1(t) \exp(u_1(\xi_1)) dt < 2\bar{K}_1\omega + \sum_{k=1}^q \ln(1 + \varrho_{1k}),$$

$$\int_0^\omega \beta_1(t) \exp(u_2(\xi_2)) dt < 2\bar{K}_1\omega + \sum_{k=1}^q \ln(1 + \varrho_{1k}).$$

Thus

$$(3.7) \quad u_1(\xi_1) < \ln \left[\frac{2\bar{K}_1\omega + \sum_{k=1}^q \ln(1 + \varrho_{1k})}{\bar{\alpha}_1\omega} \right],$$

$$(3.8) \quad u_2(\xi_2) < \ln \left[\frac{2\bar{K}_1\omega + \sum_{k=1}^q \ln(1 + \varrho_{1k})}{\bar{\beta}_1\omega} \right].$$

In the sequel, we consider two cases.

(a) If $u_1(\eta_1) \geq u_2(\eta_2)$, then it follows from (3.3) that

$$\overline{(\alpha_1 + \beta_1)}\omega \exp(u_1(\eta_1)) + \bar{\gamma}_1\omega \exp(2u_1(\eta_1)) \geq \bar{K}_1\omega + \sum_{k=1}^q \ln(1 + \varrho_{1k}),$$

which leads to

$$(3.9) \quad u_1(\eta_1) > \ln \left[\frac{-\overline{(\alpha_1 + \beta_1)}\omega + \sqrt{[\overline{(\alpha_1 + \beta_1)}\omega]^2 + 4\bar{\gamma}_1\omega(\bar{K}_1\omega + \sum_{k=1}^q \ln(1 + \varrho_{1k}))}}{2\bar{\gamma}_1\omega} \right].$$

It follows from (3.7) and (3.9) that

$$(3.10) \quad u_1(t) \leq u_1(\xi_1) + \int_0^\omega |\dot{u}_1(t)| dt$$

$$\leq \ln \left[\frac{2\bar{K}_1\omega + \sum_{k=1}^q \ln(1 + \varrho_{1k})}{\bar{\alpha}_1\omega} \right] + 2\bar{K}_1\omega + \sum_{k=1}^q \ln(1 + \varrho_{1k}) := B_1,$$

$$(3.11) \quad u_1(t) \geq u_1(\eta_1) - \int_0^\omega |\dot{u}_1(t)| dt$$

$$\geq \ln \left[\frac{-\overline{(\alpha_1 + \beta_1)}\omega + \sqrt{[\overline{(\alpha_1 + \beta_1)}\omega]^2 + 4\bar{\gamma}_1\omega(\bar{K}_1\omega + \sum_{k=1}^q \ln(1 + \varrho_{1k}))}}{2\bar{\gamma}_1\omega} \right]$$

$$- 2\bar{K}_1\omega - \sum_{k=1}^q \ln(1 + \varrho_{1k}) := B_2.$$

From (3.10) and (3.11) we have

$$(3.12) \quad \sup_{t \in [0, \omega]} |u_1(t)| < \max\{|B_1|, |B_2|\} := B_3.$$

From (3.3) we obtain

$$\bar{\alpha}_2 \omega \exp(u_2(\eta_2)) + \bar{\beta}_2 \omega \exp(B_3) + \bar{\gamma}_2 \omega \exp(B_3) \exp(u_2(\eta_2)) \geq \bar{K}_2 \omega + \sum_{k=1}^q \ln(1 + \varrho_{2k}).$$

Then

$$(3.13) \quad u_2(\eta_2) \geq \ln \left[\frac{\bar{K}_2 \omega + \sum_{k=1}^q \ln(1 + \varrho_{2k}) - \bar{\gamma}_2 \omega \exp(B_3)}{\bar{\alpha}_2 \omega + \bar{\gamma}_2 \omega \exp(B_3)} \right].$$

Thus

$$(3.14) \quad \begin{aligned} u_2(t) &\leq u_2(\xi_2) + \int_0^\omega |\dot{u}_2(t)| dt \\ &\leq \ln \left[\frac{2\bar{K}_1 \omega + \sum_{k=1}^q \ln(1 + \varrho_{1k})}{\bar{\beta}_1 \omega} \right] + 2\bar{K}_2 \omega \\ &\quad + \sum_{k=1}^q \ln(1 + \varrho_{1k}) := B_4, \end{aligned}$$

$$(3.15) \quad \begin{aligned} u_2(t) &\geq u_2(\eta_2) - \int_0^\omega |\dot{u}_2(t)| dt \\ &\geq \ln \left[\frac{\bar{K}_2 \omega + \sum_{k=1}^q \ln(1 + \varrho_{2k}) - \bar{\gamma}_2 \omega \exp(B_3)}{\bar{\alpha}_2 \omega + \bar{\gamma}_2 \omega \exp(B_3)} \right] \\ &\quad - 2\bar{K}_2 \omega - \sum_{k=1}^q \ln(1 + \varrho_{2k}) := B_5. \end{aligned}$$

It follows from (3.14) and (3.15) that

$$(3.16) \quad \sup_{t \in [0, \omega]} |u_2(t)| < \max\{|B_4|, |B_5|\} := B_6.$$

(b) If $u_1(\eta_1) < u_2(\eta_2)$, then it follows from (3.3) that

$$\overline{(\alpha_1 + \beta_1)} \omega \exp(u_2(\eta_2)) + \bar{\gamma}_1 \omega \exp(2u_2(\eta_2)) \geq \bar{K}_1 \omega + \sum_{k=1}^q \ln(1 + \varrho_{1k}),$$

which leads to

$$(3.17) \quad u_2(\eta_2) > \ln \left[\frac{-\overline{(\alpha_1 + \beta_1)} \omega + \sqrt{[\overline{(\alpha_1 + \beta_1)} \omega]^2 + 4\bar{\gamma}_1 \omega (\bar{K}_1 \omega + \sum_{k=1}^q \ln(1 + \varrho_{1k}))}}{2\bar{\gamma}_1 \omega} \right].$$

It follows from (3.7) and (3.9) that

(3.18)

$$\begin{aligned} u_2(t) &\leq u_2(\xi_1) + \int_0^\omega |\dot{u}_2(t)| dt \\ &\leq \ln \left[\frac{2\bar{K}_2\omega + \sum_{k=1}^q \ln(1 + \varrho_{2k})}{\bar{\alpha}_2\omega} \right] + 2\bar{K}_2\omega + \sum_{k=1}^q \ln(1 + \varrho_{2k}) := B_7, \end{aligned}$$

(3.19)

$$\begin{aligned} u_2(t) &\geq u_2(\eta_2) - \int_0^\omega |\dot{u}_2(t)| dt \\ &\geq \ln \left[\frac{-\overline{(\alpha_1 + \beta_1)}\omega + \sqrt{[\overline{(\alpha_1 + \beta_1)}\omega]^2 + 4\bar{\gamma}_1\omega(\bar{K}_1\omega + \sum_{k=1}^q \ln(1 + \varrho_{1k}))}}{2\bar{\gamma}_1\omega} \right] \\ &\quad - 2\bar{K}_2\omega - \sum_{k=1}^q \ln(1 + \varrho_{2k}) := B_8. \end{aligned}$$

From (3.10) and (3.11) we derive

$$(3.20) \quad \sup_{t \in [0, \omega]} |u_2(t)| < \max\{|B_7|, |B_8|\} := B_9.$$

From (3.3) we have

$$\bar{\alpha}_2\omega \exp(B_9) + \bar{\beta}_2\omega \exp(u_1(\eta_1)) + \bar{\gamma}_2\omega \exp(B_9) \exp(u_1(\eta_1)) \geq \bar{K}_2\omega + \sum_{k=1}^q \ln(1 + \varrho_{2k}).$$

Then

$$(3.21) \quad u_1(\eta_1) \geq \ln \left[\frac{\bar{K}_2\omega + \sum_{k=1}^q \ln(1 + \varrho_{2k}) - \bar{\gamma}_2\omega \exp(B_9)}{\bar{\alpha}_2\omega + \bar{\gamma}_2\omega \exp(B_9)} \right].$$

Thus

$$(3.22) \quad \begin{aligned} u_1(t) &\leq u_1(\xi_1) + \int_0^\omega |\dot{u}_1(t)| dt \\ &\leq \ln \left[\frac{2\bar{K}_1\omega + \sum_{k=1}^q \ln(1 + \varrho_{1k})}{\bar{\alpha}_1\omega} \right] + 2\bar{K}_1\omega + \sum_{k=1}^q \ln(1 + \varrho_{1k}) := B_{10}, \end{aligned}$$

$$(3.23) \quad \begin{aligned} u_1(t) &\geq u_1(\eta_1) - \int_0^\omega |\dot{u}_1(t)| dt \\ &\geq \ln \left[\frac{\bar{K}_2\omega + \sum_{k=1}^q \ln(1 + \varrho_{2k}) - \bar{\gamma}_2\omega \exp(B_9)}{\bar{\alpha}_2\omega + \bar{\gamma}_2\omega \exp(B_9)} \right] \\ &\quad - 2\bar{K}_1\omega - \sum_{k=1}^q \ln(1 + \varrho_{1k}) := B_{11}. \end{aligned}$$

It follows from (3.22) and (3.23) that

$$(3.24) \quad \sup_{t \in [0, \omega]} |u_1(t)| < \max\{|B_{10}|, |B_{11}|\} := B_{12}.$$

Obviously, B_i ($i = 3, 6, 9, 12$) are independent of $\lambda \in (0, 1)$. Similarly to the proof of Theorem 2.1 of [16], we can easily find a sufficiently large $M > 0$ so that if we denote

$$\Omega = \{u(t) = (u_1(t), u_2(t))^T \in x: \|u\| < M, u(t_k^+) \in \Omega, k = 1, 2, \dots, q\},$$

it is clear that Ω satisfies the requirement (a) in Lemma 3.1.

When $(u_1(t), u_2(t))^T \in \partial\Omega \cap \text{Ker } L = \partial\Omega \cap \mathbb{R}^2$, $u = \{(u_1, u_2)^T\}$ is a constant vector in \mathbb{R}^2 with $\|u\| = \|(u_1(t), u_2(t))^T\| = M$, then we have

$$QN u = \left(\begin{array}{c} \left(\begin{array}{c} \bar{K}_1 - \bar{\alpha}_1 \exp(u_1) - \bar{\beta}_1 \exp(u_2) - \bar{\gamma}_1 \exp(u_1) \exp(u_2) \\ + \frac{1}{\omega} \sum_{k=1}^q \ln(1 + \varrho_{1k}) \\ \bar{K}_2 - \bar{\alpha}_2 \exp(u_2) - \bar{\beta}_2 \exp(u_1) - \bar{\gamma}_2 \exp(u_1) \exp(u_2) \\ + \frac{1}{\omega} \sum_{k=1}^q \ln(1 + \varrho_{2k}) \end{array} \right), 0, \dots, 0 \end{array} \right) \neq 0.$$

Letting $J: \text{Im } Q \rightarrow \text{Ker } L, (r, 0, \dots, 0, 0) \rightarrow r$, by direct calculation we get

$$\begin{aligned} & \deg\{JQN(u_1, u_2)^T; \Omega \cap \text{ker } L; 0\} \\ &= \text{sign det} \begin{pmatrix} -(\bar{\alpha}_1 + \bar{\gamma}_2 e^{u_2})e^{u_1} & -(\bar{\beta}_1 + \bar{\gamma}_1 e^{u_1})e^{u_2} \\ -(\bar{\beta}_2 + \bar{\gamma}_2 e^{u_2})e^{u_1} & -(\bar{\alpha}_2 + \bar{\gamma}_2 e^{u_1})e^{u_2} \end{pmatrix} \\ &= \text{sign}\{(\bar{\alpha}_1 \bar{\alpha}_2 - \bar{\beta}_1 \bar{\beta}_2) + (\bar{\alpha}_1 \bar{\gamma}_2 - \bar{\beta}_2 \bar{\gamma}_1)e^{u_1} + (\bar{\alpha}_2 \bar{\gamma}_2 - \bar{\beta}_1 \bar{\gamma}_2)e^{u_2} \\ & \quad + (\bar{\gamma}_2^2 - \bar{\gamma}_1 \bar{\gamma}_2)e^{u_1 + u_2}\} \neq 0. \end{aligned}$$

This proves that condition (b) in Lemma 3.1 is satisfied. By now, we have proved that Ω verifies all requirements of Lemma 3.1, hence it follows that $Lu = Nu$ has at least one solution $(u_1(t), u_2(t))^T$ in $\text{Dom } L \cap \bar{\Omega}$, that is to say, (3.1) has at least one ω periodic solution in $\text{Dom } L \cap \bar{\Omega}$. Then we know that $(x(t), y(t))^T = (e^{u_1(t)}, e^{u_2(t)})^T$ is an ω periodic solution of system (2.3) with strictly positive components. This completes the proof. \square

4. UNIQUENESS AND GLOBAL STABILITY OF PERIODIC SOLUTIONS

Under the hypotheses (H1), (H2), (H3), we consider the following ordinary differential equation without impulses:

$$(4.1) \quad \begin{cases} \dot{z}_1(t) = z_1(t) \left[K_1(t) - \alpha_1(t) \prod_{0 < t_k < t} (1 + \varrho_{1k}) z_1(t) - \beta_1(t) \prod_{0 < t_k < t} (1 + \varrho_{2k}) z_2(t) \right. \\ \qquad \qquad \qquad \left. - \gamma_1(t) \prod_{0 < t_k < t} (1 + \varrho_{1k}) z_1(t) \prod_{0 < t_k < t} (1 + \varrho_{2k}) z_2(t - \tau_1(t)) \right], \\ \dot{z}_2(t) = z_2(t) \left[K_2(t) - \alpha_2(t) \prod_{0 < t_k < t} (1 + \varrho_{2k}) z_2(t) - \beta_2(t) \prod_{0 < t_k < t} (1 + \varrho_{1k}) z_1(t) \right. \\ \qquad \qquad \qquad \left. - \gamma_2(t) \prod_{0 < t_k < t} (1 + \varrho_{1k}) z_1(t - \tau_2(t)) \prod_{0 < t_k < t} (1 + \varrho_{2k}) z_2(t) \right] \end{cases}$$

with the initial conditions $z_i(0) > 0$, $i = 1, 2$.

Let $\tau = \max_{1 \leq i \leq 2} \{ \max_{t \in [0, \omega]} \tau_i(t) \}$. The following lemmas will be helpful in the proofs of our results. The proof of Lemma 4.1 is similar to that of Theorem 1 in [17], and will be omitted.

Lemma 4.1. *Assume that (H1), (H2), (H3) hold. Then*

- (i) *if $z(t) = (z_1(t), z_2(t))^T$ is a solution of (4.1) on $[0, +\infty)$, then $x_i(t) = \prod_{0 < t_k < t} (1 + \varrho_{ik}) z_i(t)$ ($i = 1, 2$) is a solution of (2.3) on $[-\tau, +\infty)$;*
- (ii) *if $x(t) = (x_1(t), x_2(t))^T$ is a solution of (2.3) on $[0, +\infty)$, then $z_i(t) = \prod_{0 < t_k < t} (1 + \varrho_{ik})^{-1} x_i(t)$ ($i = 1, 2$) is a solution of (4.1) on $[-\tau, +\infty)$.*

Lemma 4.2. *Let $z(t) = (z_1(t), z_2(t))^T$ denote any positive solution of system (4.1) with initial conditions $z_i(0) > 0$ ($i = 1, 2$). Then there exists a $T_3 > 0$ such that*

$$0 < z_i(t) \leq M_i \quad (i = 1, 2) \text{ for } t \geq T_3,$$

where

$$M_1 > M_1^* = \frac{K_1^M}{\alpha_1^L \prod_{0 < t_k < t} (1 + \varrho_{1k})},$$

$$M_2 > M_2^* = \frac{K_2^M}{\alpha_2^L \prod_{0 < t_k < t} (1 + \varrho_{2k})},$$

Proof. From the first equation of (4.1), we can obtain

$$(4.2) \quad \begin{aligned} \dot{z}_1(t) &\leq z_1(t) \left[K_1(t) - \alpha_1(t) \prod_{0 < t_k < t} (1 + \varrho_{1k}) z_1(t) \right], \\ &\leq z_1(t) \left[K_1^M - \alpha_1^L \prod_{0 < t_k < t} (1 + \varrho_{1k}) z_1(t) \right]. \end{aligned}$$

By (4.2), we can derive

(A1) If $z_1(0) \leq M_1$, then $z_1(t) \leq M_1$, $t \geq 0$.

(A2) If $z_1(0) > M_1$, let $-\theta_1 = M_1 [K_1^M - \alpha_1^L \prod_{0 < t_k < t} (1 + \varrho_{1k}) M_1]$ ($\theta_1 > 0$). Then there exists $\varepsilon_1 > 0$ such that $t \in [0, \varepsilon_1)$, then $z_1(t) > M_1$, and also we have

$$\dot{z}_1(t) < -\theta_1 < 0.$$

From what has been discussed above, we can easily conclude that if $z_1(0) > M_1$, then $z_1(t)$ is strictly monotone decreasing with speed at least θ_1 . Therefore there exists a $T_1 > 0$ such that if $t > T_1$, then $z_1(t) \leq M_1$.

From the second equation of (4.1), we can obtain

$$(4.3) \quad \begin{aligned} \dot{z}_2(t) &\leq z_2(t) \left[K_2(t) - \alpha_2(t) \prod_{0 < t_k < t} (1 + \varrho_{2k}) z_2(t) \right] \\ &\leq z_2(t) \left[K_2^M - \alpha_2^L \prod_{0 < t_k < t} (1 + \varrho_{2k}) z_2(t) \right]. \end{aligned}$$

By (4.3), we can derive

(B1) If $z_2(0) \leq M_2$, then $z_2(t) \leq M_2$, $t \geq 0$.

(B2) If $z_2(0) > M_2$, let $-\theta_2 = M_2 [K_2^M - \alpha_2^L \prod_{0 < t_k < t} (1 + \varrho_{2k}) M_2]$ ($\theta_2 > 0$). Then there exists $\varepsilon_2 > 0$ such that $t \in [0, \varepsilon_2)$, then $z_2(t) > M_2$, and also we have

$$\dot{z}_2(t) < -\theta_2 < 0.$$

From what has been discussed above, we can easily conclude that if $z_2(0) > M_2$, then $z_2(t)$ is strictly monotone decreasing with speed at least θ_2 . Therefore there exists a $T_2 > 0$ such that if $t > T_2$, then $z_2(t) \leq M_2$. The proof is complete. \square

Lemma 4.3. *Let (H1), (H2), (H3) hold. Assume that the following condition holds.*

$$(H6) \quad K_1^L > \beta_1^M \prod_{0 < t_k < t} (1 + \varrho_{2k}) M_2, \quad K_2^L - \beta_2^M \prod_{0 < t_k < t} (1 + \varrho_{2k}) M_1.$$

Then there exist positive constants $T > 0$ and m_i ($i = 1, 2$) such that for all $t > T$,

$$m_i < z_i(t) \quad (i = 1, 2) \quad \text{for } t \geq T,$$

where

$$m_1 < m_1^* = \frac{K_1^L - \beta_1^M \prod_{0 < t_k < t} (1 + \varrho_{2k}) M_2}{\alpha_1^M \prod_{0 < t_k < t} (1 + \varrho_{1k}) + \gamma_1^M \prod_{0 < t_k < t} (1 + \varrho_{1k}) \prod_{0 < t_k < t} (1 + \varrho_{2k}) M_2},$$

$$m_2 < m_2^* = \frac{K_2^L - \beta_2^M \prod_{0 < t_k < t} (1 + \varrho_{2k}) M_1}{\alpha_2^M \prod_{0 < t_k < t} (1 + \varrho_{2k}) + \gamma_2^M \prod_{0 < t_k < t} (1 + \varrho_{1k}) \prod_{0 < t_k < t} (1 + \varrho_{2k}) M_1}.$$

Proof. By the first equation of (4.1), It is easy to obtain that for $t \geq T_3$,

$$\begin{aligned} \dot{z}_1(t) \geq z_1(t) & \left[K_1^L - \alpha_1^M \prod_{0 < t_k < t} (1 + \varrho_{1k}) z_1(t) - \beta_1^M \prod_{0 < t_k < t} (1 + \varrho_{2k}) M_2 \right. \\ & \left. - \gamma_1^M \prod_{0 < t_k < t} (1 + \varrho_{1k}) z_1(t) \prod_{0 < t_k < t} (1 + \varrho_{2k}) M_2 \right], \end{aligned}$$

where T_3 is defined in Lemma 4.2.

(C1) If $z_1(T_3) \geq m_1$, then $z_1(t) \geq m_1$, $t \geq T_3$.

(C2) If $z_1(T_3) < m_1$, let us denote

$$\begin{aligned} \mu_1 = z_1(T_3) & \left[K_1^L - \alpha_1^M \prod_{0 < t_k < t} (1 + \varrho_{1k}) m_1 - \beta_1^M \prod_{0 < t_k < t} (1 + \varrho_{2k}) M_2 \right. \\ & \left. - \gamma_1^M \prod_{0 < t_k < t} (1 + \varrho_{1k}) m_1 \prod_{0 < t_k < t} (1 + \varrho_{2k}) M_2 \right]. \end{aligned}$$

Then there exists $\varepsilon_3 > 0$ such that if $t \in [T_3, T_3 + \varepsilon_3)$, then $z_1(t) > m_1$, and also we have

$$\dot{z}_2(t) > \mu_1 > 0.$$

Then we know that if $z_1(T_3) < m_1$, $z_1(t)$ will strictly monotonically increase with speed μ_1 . Thus there exists $T_4 > T_3$ such that if $t \geq T_4$, then $z_1(t) \geq m_1$.

By the second equation of (4.1), It is easy to obtain that for $t \geq T_3$,

$$\begin{aligned} \dot{z}_2(t) \geq z_2(t) & \left[K_2^L - \alpha_2^M \prod_{0 < t_k < t} (1 + \varrho_{2k}) z_2(t) - \beta_2^M \prod_{0 < t_k < t} (1 + \varrho_{2k}) M_1 \right. \\ & \left. - \gamma_2^M \prod_{0 < t_k < t} (1 + \varrho_{1k}) M_1 \prod_{0 < t_k < t} (1 + \varrho_{2k}) z_2(t) \right], \end{aligned}$$

where T_3 is defined in Lemma 4.2.

(D1) If $z_2(T_3) \geq m_2$, then $z_2(t) \geq m_2$, $t \geq T_3$.

(D2) If $z_2(T_3) < m_2$, let us denote

$$\begin{aligned} \mu_2 = z_2(T_3) & \left[K_2^L - \alpha_2^M \prod_{0 < t_k < t} (1 + \varrho_{2k}) m_2 - \beta_2^M \prod_{0 < t_k < t} (1 + \varrho_{2k}) M_1 \right. \\ & \left. - \gamma_2^M \prod_{0 < t_k < t} (1 + \varrho_{1k}) M_1 \prod_{0 < t_k < t} (1 + \varrho_{2k}) m_2 \right]. \end{aligned}$$

Then there exists $\varepsilon_4 > 0$ such that if $t \in [T_3, T_3 + \varepsilon_4)$, then $z_2(t) > m_1$, and also we have

$$\dot{z}_2(t) > \mu_2 > 0.$$

Then we know that if $z_2(T_3) < m_2$, $z_2(t)$ will strictly monotonically increase with speed μ_2 . Thus there exists $T_5 > T_3$ such that if $t \geq T_5$, then $z_2(t) \geq m_2$.

Set $T = \max\{T_4, T_5\}$, then we have

$$z_i(t) > m_i \quad (i = 1, 2) \quad \text{for } t \geq T.$$

□

In the sequel, we formulate the uniqueness and global stability of the ω periodic solution $x^*(t)$ in Theorem 4.1. It is immediate that if $x^*(t)$ is globally asymptotically stable, then $x^*(t)$ is in fact unique.

Theorem 4.1. *In addition to (H1)–(H6), assume further that*

$$(H7) \quad \liminf_{t \rightarrow \infty} A_i(t) > 0,$$

where

$$\begin{aligned} A_1 &= -\alpha_1(t) \prod_{0 < t_k < t} (1 + \varrho_{1k}) - 2m_2 \gamma_1(t) \prod_{0 < t_k < t} (1 + \varrho_{1k}) \prod_{0 < t_k < t} (1 + \varrho_{2k}) + \beta_2(t) \prod_{0 < t_k < t} (1 + \varrho_{1k}), \\ A_2 &= -\alpha_2(t) \prod_{0 < t_k < t} (1 + \varrho_{2k}) - 2m_1 \gamma_2(t) \prod_{0 < t_k < t} (1 + \varrho_{1k}) \prod_{0 < t_k < t} (1 + \varrho_{2k}) + \beta_1(t) \prod_{0 < t_k < t} (1 + \varrho_{2k}). \end{aligned}$$

Then system (2.3) has a unique positive ω periodic solution $x^*(t) = (x_1^*(t), x_2^*(t))^T$ which is globally asymptotically stable.

Proof. According to the conclusion of Theorem 3.1, we only need to show the global asymptotic stability of the positive periodic solution of (2.3). Let $x^*(t) = (x_1^*(t), x_2^*(t))^T$ be a positive ω periodic solution of system (2.3), let $x(t) = (x_1(t), x_2(t))^T$ be any positive solution of system (2.3). Then $z^*(t) = (z_1^*(t), z_2^*(t))^T$, where $(z_1^*(t) = \prod_{0 < t_k < t} (1 + \varrho_{1k})x_1^*(t), z_2^*(t) = \prod_{0 < t_k < t} (1 + \varrho_{2k})x_2^*(t)$, is the positive ω periodic solution of (4.1), and $z(t)$ is the positive solution of (4.1). It follows from Lemma 4.2 and Lemma 4.3 that there exist positive constants $T > 0$, M_i and m_i (defined by Lemma 4.2 and Lemma 4.2, respectively) such that for all $t > T$,

$$m_i < z_i^*(t) \leq M_i, \quad m_i < z_i(t) \leq M_i, \quad i = 1, 2.$$

Define

$$(4.4) \quad V(t) = |\ln z_1^*(t) - \ln z_1(t)| + |\ln z_2^*(t) - \ln z_2(t)|.$$

Calculating the upper-right derivative of $V(t)$ along the solution of (4.1), it follows for $t \geq T$ that

$$\begin{aligned} D^+V(t) &= \sum_{i=1}^2 \left(\frac{z_i^{*'}(t)}{z_i^*(t)} - \frac{z_i'(t)}{z_i(t)} \right) \operatorname{sgn}(z_i^*(t) - z_i(t)) \\ &= \operatorname{sgn}(z_1^*(t) - z_1(t)) \left[-\alpha_1(t) \prod_{0 < t_k < t} (1 + \varrho_{1k})(z_1^*(t) - z_1(t)) \right. \\ &\quad - \beta_1(t) \prod_{0 < t_k < t} (1 + \varrho_{2k})(z_2^*(t) - z_2(t)) - \gamma_1(t) \prod_{0 < t_k < t} (1 + \varrho_{1k})(z_1^*(t) - z_1(t)) \\ &\quad \left. \times \prod_{0 < t_k < t} (1 + \varrho_{2k})(z_2^*(t - \tau_1(t)) - z_2(t - \tau_1(t))) \right] \\ &\quad + \operatorname{sgn}(z_2^*(t) - z_2(t)) \left[-\alpha_2(t) \prod_{0 < t_k < t} (1 + \varrho_{2k})(z_2^*(t) - z_2(t)) \right. \\ &\quad - \beta_2(t) \prod_{0 < t_k < t} (1 + \varrho_{1k})(z_1^*(t) - z_1(t)) - \gamma_2(t) \prod_{0 < t_k < t} (1 + \varrho_{1k}) \\ &\quad \left. \times (z_1^*(t - \tau_1(t)) - z_1(t - \tau_2(t))) \prod_{0 < t_k < t} (1 + \varrho_{2k})(z_2^*(t) - z_2(t)) \right] \\ &\leq \sum_{i=1}^2 A_i |z_i^*(t) - z_i(t)| \quad (i = 1, 2), \end{aligned}$$

where

$$A_1 = -\alpha_1(t) \prod_{0 < t_k < t} (1 + \varrho_{1k}) - 2m_2\gamma_1(t) \prod_{0 < t_k < t} (1 + \varrho_{1k}) \prod_{0 < t_k < t} (1 + \varrho_{2k}) + \beta_2(t) \prod_{0 < t_k < t} (1 + \varrho_{1k}),$$

$$A_2 = -\alpha_2(t) \prod_{0 < t_k < t} (1 + \varrho_{2k}) - 2m_1\gamma_2(t) \prod_{0 < t_k < t} (1 + \varrho_{1k}) \prod_{0 < t_k < t} (1 + \varrho_{2k}) + \beta_1(t) \prod_{0 < t_k < t} (1 + \varrho_{2k}).$$

By hypothesis (H7) there exist constants α_i ($i = 1, 2$) and $T^* > T$ such that

$$(4.5) \quad A_i(t) \geq \alpha_i > 0 \quad (i = 1, 2) \quad \text{for } t \geq T^*.$$

Integrating both sides of (4.11) over the interval $[T^*, t]$ yields

$$(4.6) \quad V(t) + \sum_{i=1}^2 \int_{T^*}^t A_i(s) |z_i^*(s) - z_i(s)| ds \leq V(T^*).$$

It follows from (4.12) and (4.13) that

$$(4.7) \quad \sum_{i=1}^2 \int_{T^*}^t A_i(s) |z_i^*(s) - z_i(s)| ds \leq V(T^*) < \infty \quad \text{for } t \geq T^*.$$

Since $z_i^*(t)$ and $z_i(t)$ ($i = 1, 2$) are bounded for $t \geq T^*$, so $|z_i^*(t) - z_i(t)|$ ($i = 1, 2$) are uniformly continuous on $[T^*, \infty)$. By Barbalat's Lemma [1] we have

$$\lim_{t \rightarrow \infty} |z_i^*(t) - z_i(t)| = \lim_{t \rightarrow \infty} \left[\prod_{0 < t_k < t} (1 + \varrho_{ik})^{-1} |x_i^*(t) - x_i(t)| \right] = 0 \quad (i = 1, 2).$$

Thus

$$(4.8) \quad \lim_{t \rightarrow \infty} |x_i^*(t) - x_i(t)| = 0 \quad (i = 1, 2).$$

By Theorems 7.4 and 8.2 in [18] we know that the positive periodic solution $x^*(t) = (x_1^*(t), x_2^*(t))^T$ of equation (2.3) is uniformly asymptotically stable. The proof of Theorem 4.1 is complete. \square

References

- [1] *I. Barbălat*: Systèmes d'équations différentielles d'oscillations non linéaires. Acad. Républ. Popul. Roum., Rev. Math. Pur. Appl. 4 (1959), 267–270. (In French.)
- [2] *F. Chen*: Almost periodic solution of the non-autonomous two-species competitive model with stage structure. Appl. Math. Comput. 181 (2006), 685–693.

- [3] *S. Chen, T. Wang, J. Zhang*: Positive periodic solution for non-autonomous competition Lotka-Volterra patch system with time delay. *Nonlinear Anal., Real World Appl.* *5* (2004), 409–419.
- [4] *L. Dong, L. Chen, P. Shi*: Periodic solutions for a two-species nonautonomous competition system with diffusion and impulses. *Chaos Solitons Fractals* *32* (2007), 1916–1926.
- [5] *M. Fan, K. Wang, D. Jiang*: Existence and global attractivity of positive periodic solutions to periodic n -species Lotka-Volterra competition systems with several deviating arguments. *Math. Biosci.* *160* (1999), 47–61.
- [6] *R. E. Gaines, J. L. Mawhin*: Coincidence Degree, and Nonlinear Differential Equations, *Lecture Notes in Mathematics* 568. Springer, Berlin, 1977.
- [7] *Z. Li, F. Chen*: Extinction in periodic competitive stage-structured Lotka-Volterra model with the effects of toxic substances. *J. Comput. Appl. Math.* *231* (2009), 143–153.
- [8] *Z. Liu, L. Chen*: Periodic solution of a two-species competitive system with toxicant and birth pulse. *Chaos Solitons Fractals* *32* (2007), 1703–1712.
- [9] *Z. Liu, J. Hui, J. Wu*: Permanence and partial extinction in an impulsive delay competitive system with the effect of toxic substances. *J. Math. Chem.* *46* (2009), 1213–1231.
- [10] *L. Nie, J. Peng, Z. Teng*: Permanence and stability in multi-species non-autonomous Lotka-Volterra competitive systems with delays and feedback controls. *Math. Comput. Modelling* *49* (2009), 295–306.
- [11] *J. Shen, J. Li*: Existence and global attractivity of positive periodic solutions for impulsive predator-prey model with dispersion and time delays. *Nonlinear Anal., Real World Appl.* *10* (2009), 227–243.
- [12] *X. Song, L. Chen*: Periodic solution of a delay differential equation of plankton allelopathy. *Acta Math. Sci., Ser. A, Chin. Ed.* *23* (2003), 8–13. (In Chinese.)
- [13] *X. H. Tang, X. Zou*: Global attractivity of non-autonomous Lotka-Volterra competition system without instantaneous negative feedback. *J. Differ. Equations* *192* (2003), 502–535.
- [14] *X. Tang, D. Cao, X. Zou*: Global attractivity of positive periodic solution to periodic Lotka-Volterra competition systems with pure delay. *J. Differ. Equations* *228* (2006), 580–610.
- [15] *Y. Xia*: Positive periodic solutions for a neutral impulsive delayed Lotka-Volterra competition system with the effect of toxic substance. *Nonlinear Anal., Real World Appl.* *8* (2007), 204–221.
- [16] *R. Xu, M. A. J. Chaplain, F. A. Davidson*: Periodic solution of a Lotka-Volterra predator-prey model with dispersion and time delays. *Appl. Math. Comput.* *148* (2004), 537–560.
- [17] *J. Yan, A. Zhao*: Oscillation and stability of linear impulsive delay differential equations. *J. Math. Anal. Appl.* *227* (1998), 187–194.
- [18] *T. Yoshizawa*: *Stability Theory by Ljapunov's Second Method*, Publications of the Mathematical Society of Japan. Vol. 9. The Mathematical Society of Japan, Tokyo, 1966.

Authors' addresses: *Changjin Xu*, Guizhou Key Laboratory of Economics System Simulation, Guizhou University of Finance and Economics, Guiyang 550004, P. R. China, e-mail: xcj4030126.com; *Qianhong Zhang*, Guizhou Key Laboratory of Economics System Simulation, Guizhou University of Finance and Economics, Guiyang 550004, P. R. China; *Maoxin Liao*, School of Mathematics and Physics, Nanhua University, Hengyang 421001, P. R. China.