

Applications of Mathematics

Juan Liu; Xiaodi Li

Impulsive stabilization of high-order nonlinear retarded differential equations

Applications of Mathematics, Vol. 58 (2013), No. 3, 347--367

Persistent URL: <http://dml.cz/dmlcz/143282>

Terms of use:

© Institute of Mathematics AS CR, 2013

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

IMPULSIVE STABILIZATION OF HIGH-ORDER NONLINEAR
RETARDED DIFFERENTIAL EQUATIONS

JUAN LIU, Xinxiang, XIAODI LI, Ji'nan

(Received March 31, 2010)

Abstract. In this paper, impulsive stabilization of high-order nonlinear retarded differential equations is investigated by using Lyapunov functions and some analysis methods. Our results show that several non-impulsive unstable systems can be stabilized by imposition of impulsive controls. Some recent results are extended and improved. An example is given to demonstrate the effectiveness of the proposed control and stabilization methods.

Keywords: high-order nonlinear retarded differential equation, Lyapunov function, impulsive stabilization, exponential stability

MSC 2010: 34A37, 34K20

1. INTRODUCTION AND PRELIMINARIES

Over the last decade, impulsive control and impulsive stabilization for delay differential equations have attracted a great deal of attention due to its potential applications in many fields such as biological systems, chemical reactions, dosage supply in pharmacokinetics, ecosystems management, dynamic portfolio management and stabilization and synchronization in chaotic secure communication systems and other chaos systems [5], [6], [10], [13], [15], [16]. For example, in pest management, the impulse models the process of periodic release of infective pests at fixed moments to control pests population size, which can be described by an impulsive control system [16]. In such system, the equilibrium solution that can be effectively controlled by impulses denotes that the pests population size can be kept at acceptably low levels

This work was jointly supported by the National Natural Sciences Foundation of China: Tianyuan Foundation (No. 11226136), the Project of Shandong Province Higher Educational Science and Technology Program (No. J12LI04) and the Research Fund for Excellent Young and Middle-aged Scientists of Shandong Province (No. BS2012DX039).

in the long term. Moreover, in many cases, impulsive control can produce better performance than continuous control; even in some cases, only impulsive approaches can be used. A representative example is that a central bank cannot change its interest rate everyday in order to regulate the money supply in a financial market [15]. In recent years, various results for impulsive control systems have been reported, see [1], [2], [3], [4], [7], [8], [9], [10], [11], [12], [14]. Since impulses can make unstable systems stable, and stable systems can become unstable after impulse effects, it is reasonable to ask whether the solutions of high-order nonlinear retarded differential equations have similar properties. However, to the best of author's knowledge, there is almost no result on impulsive stabilization of high-order nonlinear retarded differential equations.

In this paper, we consider the high-order nonlinear retarded differential equations

$$(1.1) \quad x^{(n)}(t) + a(t)x^{(\gamma)}(t) + b(t)x^\varrho(t) + \sum_{i=1}^N f_i(t, x^\delta(g_i(t))) = 0, \quad t \geq t_0,$$

and the corresponding equations with impulses

$$(1.2) \quad \begin{cases} x^{(n)}(t) + a(t)x^{(\gamma)}(t) + b(t)x^\varrho(t) + \sum_{i=1}^N f_i(t, x^\delta(g_i(t))) = 0, & t \geq t_0, t \neq t_k, \\ x(t_k) = I_k(x(t_k^-)), x^{(j)}(t_k) = J_{jk}(x^{(j)}(t_k^-)), & j = 1, 2, \dots, n-1, t = t_k. \end{cases}$$

We also consider other high-order nonlinear retarded differential equations

$$(1.3) \quad x^{(n)}(t) + a(t)x^{(\gamma)}(t) + b(t)x^\varrho(t) + \sum_{i=1}^N \int_{g_i(t)}^t f_i(t-u, x^\delta(u)) du = 0, \quad t \geq t_0,$$

and the corresponding equations with impulses

$$(1.4) \quad \begin{cases} x^{(n)}(t) + a(t)x^{(\gamma)}(t) + b(t)x^\varrho(t) + \sum_{i=1}^N \int_{g_i(t)}^t f_i(t-u, x^\delta(u)) du = 0, & t \geq t_0, \\ x(t_k) = I_k(x(t_k^-)), x^{(j)}(t_k) = J_{jk}(x^{(j)}(t_k^-)), & j = 1, 2, \dots, n-1, t = t_k. \end{cases}$$

The following assumptions will be needed throughout the paper:

- (A₁) The impulsive sequence t_k satisfies $0 \leq t_0 < t_1 < \dots < t_k < \dots$, $\lim_{k \rightarrow +\infty} t_k = +\infty$;
- (A₂) $I_k, J_{jk} : \mathbb{R} \rightarrow \mathbb{R}$ are continuous and $I_k(0) = J_{jk}(0) = 0$, $k \in \mathbb{Z}_+$;
- (A₃) $a(t), b(t) : [t_0, \infty) \rightarrow \mathbb{R}$ are continuous functions;
- (A₄) $f_i : [t_0, \infty) \times \mathbb{R} \rightarrow \mathbb{R}$, $f_i(t, 0) = 0$, $t \geq t_0$. There exists a sequence of functions $p_i(t)$, where $p_i : [t_0, \infty) \rightarrow \mathbb{R}$ are continuous such that for all $t \geq t_0$, $|f_i(t, x)| \leq |p_i(t)||x|$, $i = 1, 2, \dots, N$, where $N \geq 1$;

(A₅) $g_i: [t_0, \infty) \times \mathbb{R} \rightarrow \mathbb{R}$ are continuous satisfying $0 \leq t - g_i(t) < \infty$ for all $t \geq t_0 \geq 0$, $i = 1, 2, \dots, N$;

(A₆) γ, ϱ, δ are constants and $1 \leq \gamma < n, \varrho \geq 1, \delta \geq 1, \gamma \in \mathbb{Z}_+$;

(A₇) $x'(t)$ denotes the right hand derivative of $x(t)$, i.e.,

$$x'(t_k) = x'(t_k^+) = \lim_{h \rightarrow 0^+} (x(t_k + h) - x(t_k^+))/h, \text{ and } x''(t) = [x'(t)]', \dots, \\ x^{(n)}(t) = [x^{(n-1)}(t)]'.$$

For any $\sigma \geq 0$, let $\tau_i^\sigma = \sup_{t \geq \sigma} \{t - g_i(t)\}$, $\tau^\sigma = \max_{1 \leq i \leq N} \tau_i^\sigma$ and let $\Phi(\sigma)$ denote the set of functions $\varphi: [\sigma - \tau^\sigma, \sigma] \rightarrow \mathbb{R}$ which have at most finitely many discontinuity points of the first kind and are right continuous at these points.

So for any $\sigma \geq 0$ and $\varphi \in \Phi$, we can define the initial value condition of (1.1), (1.2), (1.3), (1.4)

$$(1.5) \quad x(t) = \varphi(t), \quad t \in [\sigma - \tau^\sigma, \sigma]; \quad x^{(j)}(\sigma) = x_{j0}, \quad j = 1, 2, \dots, n - 1.$$

If we let $n = 2, \gamma = 1, \delta = \varrho = 1, N = 1, g_1(t) = t - \tau_1, f_1(t, x) = p(t)x$, then (1.1) and (1.3) reduce to the differential equations

$$\begin{cases} x''(t) + a(t)x'(t) + b(t)x(t) + p(t)x(t - \tau_1) = 0, & t \geq t_0, \\ x(t) = \varphi(t), & t_0 - \tau_1 \leq t \leq t_0, \quad x'(t_0) = x_0 \end{cases}$$

and

$$\begin{cases} x''(t) + a(t)x'(t) + b(t)x(t) + \int_{t-\tau_1}^t p(t-u)x(u) \, du = 0, & t \geq t_0, \\ x(t) = \varphi(t), & t_0 - \tau_1 \leq t \leq t_0, \quad x'(t_0) = x_0. \end{cases}$$

The existence of solutions and impulsive stabilization of these equations was extensively investigated in [14].

If we let $a(t) = 0, f_i = 0$, then (1.1) or (1.3) reduces to the differential equation

$$(1.6) \quad \begin{cases} x^{(n)}(t) + b(t)x^e(t) = 0, & t \geq t_0, \\ x(t_0) = x_0, \quad x^{(j)}(t_0) = x_{j0}, & j = 1, 2, \dots, n - 1. \end{cases}$$

The stabilization of the solutions of (1.6) with impulse has been investigated in [9].

In the paper [4], the authors study the impulsive stabilization for the following second order delay differential equations:

$$\begin{cases} x''(t) + f(t, x(t), x'(t)) + g(t, x(t), x(t - \tau)) = 0, & t \geq t_0, \\ x(t) = \varphi(t), & t_0 - \tau \leq t \leq t_0, \quad x'(t_0) = y_0 \end{cases}$$

with some necessary assumptions which include: $f, g: [t_0, \infty) \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ are continuous such that $f(t, 0, 0) = g(t, 0, 0) = 0$ and there exist constants $F > 0, G > 0$ such that for all $t \geq t_0$ and $u, v \in \mathbb{R}, |f(t, u, v)| \leq F|u|, |g(t, u, v)| \leq G|v|$.

In the present paper, we deal with the equations (1.1) and (1.3), which are more general than those in [14] and [9]. To some extent the systems studied in [4] are more general than the systems we study (when $n = 2$, $\gamma = 1$, $\delta = \sigma = 1$, $N = 1$, $g_1(t) = t - \tau_1$ in (1.1)), but there still exist some cases to which the methods cannot be applied, for example $f(t, x, x') = x^3 + x'$. The results we study can solve it.

In this paper, since (1.1) may reduce to first order impulsive differential equation, we can obtain a global existence result of the solution of system (1.1), see [11], [2], [1]. So we always assume the solution of (1.1) to exist globally in this paper.

Definition 1.1. For any $\sigma \geq 0$ and $\varphi \in \Phi$, a function $x: [\sigma - \tau^\sigma, \sigma + a) \rightarrow \mathbb{R}$, $a > 0$, is said to be a solution of (1.2) and (1.5) through $(\sigma, \varphi, \{x_{j0}\}_{j=1}^{n-1})$ if

- (i) $x(t)$ and $x^{(j)}(t)$, $j = 1, 2, \dots, n-1$, are continuous on $[\sigma - \tau^\sigma, \sigma + a) \setminus \{t_k; k \in \mathbb{Z}_+\}$ and are right continuous at t_k ;
- (ii) $x(t)$ satisfies (1.1) and (1.5);
- (iii) $x(t)$ and $x^{(j)}(t)$, $j = 1, 2, \dots, n-1$ fulfil (1.2) for each $k \in \mathbb{Z}_+$.

Remark 1.1. The definition of a solution of (1.4) and (1.5) is similar to Definition 1.1, we omit it.

Remark 1.2. In the present paper, for convenience we use τ instead of τ^σ .

Definition 1.2. The zero solution of (1.1) is said to be exponentially stabilized by impulses, if there exist $\alpha > 0$, a sequence $\{t_k\}_{k=1}^\infty$, I_k , J_{jk} satisfying (A₁) and (A₂) such that for all $\varepsilon > 0$ there exists a $\delta^* > 0$ such that, when the solution $x(t)$ of (1.1) through $(\sigma, \varphi, \{x_{j0}\}_{j=1}^{n-1})$ fulfils

$$(1.7) \quad \left(\|\varphi\|_{t_0}^2 + \sum_{j=1}^{n-1} (x_{j0})^2 \right)^{\frac{1}{2}} \leq \delta^*,$$

then

$$(1.8) \quad \left(x^2(t) + \sum_{j=1}^{n-1} (x^{(j)}(t))^2 \right)^{\frac{1}{2}} \leq \varepsilon \exp[-\alpha(t - t_0)], \quad t \geq t_0.$$

where $\|\varphi\|_t = \sup_{t-\tau \leq s \leq t} |\varphi(s)|$.

Definition 1.3. The zero solution of (1.1) is said to be exponentially stabilized by periodic impulses if there exist $\alpha > 0$, a sequence $\{t_k\}_{k=1}^\infty$ satisfying (A₁) and $t_k - t_{k-1} = d$ (> 0 constant), I_k , J_{jk} satisfying (A₂) and

$$I_k(x) = I(x), J_{jk}(x) = J(x), \quad x \in \mathbb{R},$$

such that for all $\varepsilon > 0$ there exists a $\delta > 0$ such that the solution $x(t)$ of (1.1) through $(\sigma, \varphi, \{x_{j0}\}_{j=1}^{n-1})$ fulfils (1.7) and (1.8).

2. MAIN RESULTS

Theorem 2.1. *Assume that conditions (A₃)–(A₇) hold. Moreover, suppose that*

(A₈) *there exists a constant $\beta > 0$ such that $g'_i(t) \geq \beta$, $t \geq 0$, $i = 1, 2, \dots, N$;*

(A₉) *there exist constants $a, b, p_i \geq 0$, $i = 1, 2, \dots, N$ such that $|a(t)| \leq a$, $|b(t)| \leq b$,
 $|p_i(t)| \leq p_i$;*

(A₁₀)

$$\frac{\tau}{\beta} \sum_{i=1}^N p_i < \exp \left[- \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \right) \tau \right].$$

Then the zero solution of (1.1) can be exponentially stabilized by impulses.

Proof. Since (A₁₀) holds, there exist $\alpha > 0$ and $\lambda \geq \tau$ such that

$$\frac{1}{\beta} \sum_{i=1}^N p_i \tau_i \leq \frac{\tau}{\beta} \sum_{i=1}^N p_i \leq \exp[-2\alpha(\lambda + \tau)] \exp \left[- \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \right) \lambda \right].$$

Then one may choose a sequence $\{t_k\}_{k=1}^\infty$ satisfying (A₁) and $\tau \leq t_{k+1} - t_k \leq \lambda$, $t_0 = \sigma$. Note that $g'_i(t) \geq \beta > 0$, hence g is nondecreasing in t for $t \geq 0$. Thus one may choose a sequence $\{\eta_k\}_{k=1}^\infty$, $\eta_k \in (0, 1)$ such that, when the solution $x(t) = x(t, t_k, x(t_k), \{x_{jk}\}_{j=1}^{n-1})$ through $(t_k, x(t_k), \{x_{jk}\}_{j=1}^{n-1})$ fulfils

$$\left(x^2(t_k) + \sum_{j=1}^{n-1} (x^{(j)}(t_k))^2 \right)^{\frac{1}{2}} \leq \eta_k,$$

then

$$(2.1) \quad \left(x^2(t) + \sum_{j=1}^{n-1} (x^{(j)}(t))^2 \right)^{\frac{1}{2}} < 1, \quad t \in [t_k, t_{k+1}).$$

Let

$$\begin{aligned} |I_k(u)| &= d_k |u|, \quad |J_{jk}(v)| = d_k |v|, \\ d_k &= \min \left\{ \eta_k \exp[\alpha(t_1 - \sigma)], \left(\Gamma_k - \sum_{i=1}^N \frac{p_i}{\beta} \tau_i \right)^{\frac{1}{2}} \right\}, \\ \Gamma_k &= \exp[-2\alpha(t_{k+1} - t_k + \tau)] \exp \left[- \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \right) (t_{k+1} - t_k) \right]. \end{aligned}$$

It is obvious that $d_k \geq 0$.

For any $\varepsilon \in (0, 1)$, let

$$\delta^* = \min \left\{ \eta_0, \varepsilon, \varepsilon \left(1 + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i \right)^{-1/4} \right. \\ \left. \times \exp[-\alpha(t_1 - \sigma)] \exp \left[-\frac{1}{2} \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \right) (t_1 - \sigma) \right] \right\}.$$

Next, we prove that each solution $x(t) = x(t, \sigma, \varphi, \{x_{j0}\}_{j=1}^{n-1})$ of (1.2) and (1.5) with

$$\left(\|\varphi\|_\sigma^2 + \sum_{j=1}^{n-1} (x_{j0})^2 \right)^{\frac{1}{2}} \leq \delta^*$$

satisfies

$$\left(x^2(t) + \sum_{j=1}^{n-1} (x^{(j)}(t))^2 \right)^{\frac{1}{2}} \leq \varepsilon \exp[-\alpha(t - t_0)], \quad t \geq t_0,$$

where $\|\varphi\|_t = \sup_{t-\tau \leq s \leq t} |\varphi(s)|$.

First, for $t \in [\sigma, t_1)$, we choose a Lyapunov function

$$V(t) = x^2(t) + \sum_{i=1}^{n-1} (x^{(i)}(t))^2 + \sum_{i=1}^N \frac{p_i}{\beta} \int_{g_i(t)}^t x^{2\delta}(s) ds.$$

It follows from conditions (A₈)–(A₁₀) and (2.1) that

$$(1) \quad V(t) \geq x^2(t) + \sum_{i=1}^{n-1} (x^{(i)}(t))^2;$$

(2)

$$\begin{aligned} V(t) &\leq x^2(t) + \sum_{i=1}^{n-1} (x^{(i)}(t))^2 + \sum_{i=1}^N \frac{p_i}{\beta} \int_{t-\tau_i}^t x^{2\delta}(s) ds \\ &\leq x^2(t) + \sum_{i=1}^{n-1} (x^{(i)}(t))^2 + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i \sup_{t-\tau_i \leq s \leq t} |x(s)|^{2\delta} \\ &\leq x^2(t) + \sum_{i=1}^{n-1} (x^{(i)}(t))^2 + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i \|x\|_t^{2\delta} \\ &\leq x^2(t) + \sum_{i=1}^{n-1} (x^{(i)}(t))^2 + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i \|x\|_t^2 \\ &\leq \left(1 + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i \right) \left(\|x\|_t^2 + \sum_{i=1}^{n-1} (x^{(i)}(t))^2 \right), \end{aligned}$$

where $\|x\|_t = \sup_{t-\tau \leq s \leq t} |x(s)|$.

(3) We denote by $V'(t)$ the right upper derivative of $V(t)$ along the solution of (1.1)–(1.3). Then

$$\begin{aligned}
V'(t) &= 2x(t)x'(t) + 2x'(t)x''(t) + \dots + 2x^{(n-1)}(t)x^{(n)}(t) \\
&\quad + \sum_{i=1}^N \frac{p_i}{\beta} x^{2\delta}(t) - \sum_{i=1}^N \frac{p_i}{\beta} x^{2\delta}(g_i(t))g'_i(t) \\
&= 2x(t)x'(t) + 2x'(t)x''(t) + \dots \\
&\quad + 2x^{(n-1)}(t) \left\{ -a(t)x^{(\gamma)}(t) - b(t)x^e(t) - \sum_{i=1}^N f_i(t, x^\delta(g_i(t))) \right\} \\
&\quad + \sum_{i=1}^N \frac{p_i}{\beta} x^{2\delta}(t) - \sum_{i=1}^N \frac{p_i}{\beta} x^{2\delta}(g_i(t))g'_i(t) \\
&\leq x^2(t) + [x'(t)]^2 + \dots + [x^{(n-2)}(t)]^2 + [x^{(n-1)}(t)]^2 \\
&\quad + a([x^{(n-1)}(t)]^2 + [x^{(\gamma)}(t)]^2) + b([x^{(n-1)}(t)]^2 + x^2(t)) \\
&\quad + 2|x^{(n-1)}(t)| \sum_{i=1}^N |p_i(t)| |x^{2\delta}(g_i(t))| + \sum_{i=1}^N \frac{p_i}{\beta} x^{2\delta}(t) - \sum_{i=1}^N \frac{p_i}{\beta} x^{2\delta}(g_i(t))g'_i(t) \\
&\leq x^2(t) + [x'(t)]^2 + \dots + [x^{(n-2)}(t)]^2 + [x^{(n-1)}(t)]^2 \\
&\quad + a([x^{(n-1)}(t)]^2 + [x^{(\gamma)}(t)]^2) + b([x^{(n-1)}(t)]^2 + x^2(t)) \\
&\quad + \sum_{i=1}^N p_i([x^{(n-1)}(t)]^2 + x^{2\delta}(g_i(t))) + \sum_{i=1}^N \frac{p_i}{\beta} x^{2\delta}(t) - \sum_{i=1}^N \frac{p_i}{\beta} x^{2\delta}(g_i(t))g'_i(t) \\
&\leq x^2(t) + [x'(t)]^2 + \dots + [x^{(n-2)}(t)]^2 + [x^{(n-1)}(t)]^2 \\
&\quad + a([x^{(n-1)}(t)]^2 + [x^{(\gamma)}(t)]^2) + b([x^{(n-1)}(t)]^2 + x^2(t)) \\
&\quad + \sum_{i=1}^N p_i[x^{(n-1)}(t)]^2 + \sum_{i=1}^N \frac{p_i}{\beta} x^2(t) + \sum_{i=1}^N p_i x^{2\delta}(g_i(t)) \left(1 - \frac{g'_i(t)}{\beta}\right) \\
&\leq \left(1 + b + \sum_{i=1}^N \frac{p_i}{\beta}\right) x^2(t) + 2[x'(t)]^2 + \dots + 2[x^{(\gamma-1)}(t)]^2 + (a+2)[x^{(\gamma)}(t)]^2 \\
&\quad + 2[x^{(\gamma+1)}(t)]^2 + \dots + \left(2 + a + \sum_{i=1}^N \frac{p_i}{\beta}\right) [x^{(n-1)}(t)]^2 \\
&\leq \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta}\right) V(t),
\end{aligned}$$

which implies that

$$V(t) \leq V(t_0) \exp \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \right) (t - \sigma), \quad t \in [\sigma, t_1].$$

Thus for $t \in [\sigma, t_1)$ we get

$$\begin{aligned}
x^2(t) + \sum_{i=1}^{n-1} (x^{(j)}(t))^2 &\leq V(t) \leq V(t_0) \exp\left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta}\right)(t - \sigma) \\
&< V(t_0) \exp\left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta}\right)(t_1 - \sigma) \\
&\leq \left(1 + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i\right) \left(\|x\|_\sigma^2 + \sum_{i=1}^{n-1} (x^{(j)}(\sigma))^2\right) \exp\left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta}\right)(t_1 - \sigma) \\
&= \left(1 + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i\right) \left(\|\varphi\|_\sigma^2 + \sum_{j=1}^{n-1} x_{j0}^2\right) \exp\left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta}\right)(t_1 - \sigma) \\
&\leq \left(1 + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i\right) \delta^{*2} \exp\left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta}\right)(t_1 - \sigma) \\
&\leq \varepsilon^2 \exp[-2\alpha(t_1 - \sigma)] < \varepsilon^2 \exp[-2\alpha(t - \sigma)],
\end{aligned}$$

which implies that

$$\left(x^2(t) + \sum_{i=1}^{n-1} (x^{(j)}(t))^2\right)^{\frac{1}{2}} < \varepsilon \exp[-\alpha(t - \sigma)], \quad t \in [\sigma, t_1).$$

Especially,

$$\left(x^2(t_1^-) + \sum_{i=1}^{n-1} (x^{(j)}(t_1^-))^2\right)^{\frac{1}{2}} \leq \varepsilon \exp[-\alpha(t_1 - \sigma)].$$

It then follows that

$$\begin{aligned}
\left(x^2(t_1) + \sum_{i=1}^{n-1} (x^{(j)}(t_1))^2\right)^{\frac{1}{2}} &= d_1 \left(x^2(t_1^-) + \sum_{i=1}^{n-1} (x^{(j)}(t_1^-))^2\right)^{\frac{1}{2}} \\
&\leq d_1 \varepsilon \exp[-\alpha(t_1 - \sigma)] < d_1 \exp[-\alpha(t_1 - \sigma)] \leq \eta_1,
\end{aligned}$$

which implies that, for $t \in [t_1, t_2)$,

$$(2.2) \quad \left(x^2(t) + \sum_{i=1}^{n-1} (x^{(j)}(t))^2\right)^{\frac{1}{2}} < 1.$$

For $t \in [t_1, t_2)$, we still choose a Lyapunov function

$$V(t) = x^2(t) + \sum_{i=1}^{n-1} (x^{(j)}(t))^2 + \sum_{i=1}^N \frac{p_i}{\beta} \int_{g_i(t)}^t x^{2\delta}(s) \, ds.$$

Then we have

$$\begin{aligned}
& x^2(t) + \sum_{i=1}^{n-1} (x^{(j)}(t))^2 \\
& \leq V(t) \\
& \leq V(t_1) \exp\left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta}\right)(t - t_1) \\
& < V(t_1) \exp\left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta}\right)(t_2 - t_1) \\
& = \left[x^2(t_1) + \sum_{i=1}^{n-1} (x^{(j)}(t_1))^2 + \sum_{i=1}^N \frac{p_i}{\beta} \int_{g_i(t_1)}^{t_1} x^{2\delta}(s) \, ds \right] \\
& \quad \times \exp\left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta}\right)(t_2 - t_1) \\
& \leq \left\{ d_1^2 \left[x^2(t_1^-) + \sum_{i=1}^{n-1} (x^{(j)}(t_1^-))^2 \right] + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i \sup_{t_1 - \tau_i \leq t \leq t_1} x^{2\delta}(t) \right\} \\
& \quad \times \exp\left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta}\right)(t_2 - t_1) \\
& \leq \left\{ d_1^2 \left[x^2(t_1^-) + \sum_{i=1}^{n-1} (x^{(j)}(t_1^-))^2 \right] + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i \sup_{t_1 - \tau \leq t \leq t_1} x^2(t) \right\} \\
& \quad \times \exp\left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta}\right)(t_2 - t_1) \\
& \leq \left\{ d_1^2 \sup_{t_1 - \tau \leq t \leq t_1} \left[x^2(t) + \sum_{i=1}^{n-1} (x^{(j)}(t))^2 \right] + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i \sup_{t_1 - \tau \leq t \leq t_1} x^2(t) \right\} \\
& \quad \times \exp\left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta}\right)(t_2 - t_1) \\
& \leq \left(d_1^2 + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i \right) \sup_{t_1 - \tau \leq t \leq t_1} \left[x^2(t) + \sum_{i=1}^{n-1} (x^{(j)}(t))^2 \right] \\
& \quad \times \exp\left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta}\right)(t_2 - t_1) \\
& \leq \Gamma_1 \varepsilon^2 \exp(-2\alpha(t_1 - \sigma - \tau)) \exp\left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta}\right)(t_2 - t_1) \\
& \leq \varepsilon^2 \exp[-2\alpha(t_2 - \sigma)] \\
& < \varepsilon^2 \exp[-2\alpha(t - \sigma)], \quad t \in [t_1, t_2].
\end{aligned}$$

Hence,

$$\left(x^2(t) + \sum_{i=1}^{n-1} (x^{(i)}(t))^2\right)^{\frac{1}{2}} < \varepsilon \exp[-\alpha(t - \sigma)], \quad t \in [t_1, t_2].$$

Arguing as before, by induction hypothesis we may prove, in general, that for $k \geq 1$,

$$\left(x^2(t) + \sum_{i=1}^{n-1} (x^{(i)}(t))^2\right)^{\frac{1}{2}} < \varepsilon \exp[-\alpha(t - \sigma)], \quad t \in [t_k, t_{k+1}).$$

Therefore, we finally obtain

$$\left(x^2(t) + \sum_{i=1}^{n-1} (x^{(i)}(t))^2\right)^{\frac{1}{2}} < \varepsilon \exp[-\alpha(t - \sigma)], \quad t \geq \sigma,$$

and the proof is therefore complete. \square

Remark 2.1. In Theorem 2.1, we choose linear functions $I_k(u) = d_k u$, $J_{jk}(v) = d_k v$. In fact, from the procedure in the proof of Theorem 2.1, it is not difficult to realize that we only need $I_k(u)$, $J_{jk}(v)$ to satisfy: $|I_k(u)| \leq d_k |u|$, $|J_{jk}(v)| \leq d_k |v|$.

Remark 2.2. If $\gamma = n - 1$ in Theorem 2.1, then the condition $|a(t)| \leq a$ can be replaced by $a(t) \geq 0$.

Remark 2.3. Suppose that all the conditions in Theorem 2.1 hold. Then the procedure in Theorem 2.1 can be used to prove the exponential stabilization for the high-order nonlinear retarded differential equations

$$x^{(n)}(t) + h(t, x(t), x^{(\gamma)}(t)) + \sum_{i=1}^N f_i(t, x^\delta(g_i(t))) = 0, \quad t \geq t_0,$$

and the corresponding equations with impulses

$$\begin{cases} x^{(n)}(t) + h(t, x(t), x^{(\gamma)}(t)) + \sum_{i=1}^N f_i(t, x^\delta(g_i(t))) = 0, & t \geq t_0, \quad t \neq t_k, \\ x(t_k) = I_k(t_k^-), \quad x^{(j)}(t_k) = J_{jk}(t_k^-), & j = 1, 2, \dots, n-1, \quad t = t_k, \end{cases}$$

where there exist two continuous functions h_1, h_2 such that $|h(t, x, y)| \leq |h_1(t)||x| + |h_2(t)||y|$.

Remark 2.4. In Remark 2.3, we can further consider the high-order nonlinear retarded differential equations

$$x^{(n)}(t) + h(t, x(t), x^{(\gamma)}(t)) + f(t, x(g_1(t)), x(g_2(t)), \dots, x(g_N(t))) = 0, \quad t \geq t_0,$$

and the corresponding equations with impulses

$$\begin{cases} x^{(n)}(t) + h(t, x(t), x^{(\gamma)}(t)) + f(t, x(g_1(t)), x(g_2(t)), \dots, x(g_N(t))) = 0, \\ t \geq t_0, t \neq t_k, \\ x(t_k) = I_k(t_k^-), x^{(j)}(t_k) = J_{jk}(t_k^-), \quad j = 1, 2, \dots, n-1, t = t_k, \end{cases}$$

where there exists a sequence of continuous functions ω_i , $i = 1, 2, \dots, N$ such that $|f(t, \mu_1, \mu_2, \dots, \mu_N)| \leq \sum_{i=1}^N |\omega_i(t)| |\mu_i|$.

Theorem 2.2. (Case: $\delta = \varrho = 1$.) Assume that the conditions in Theorem 2.1 hold. Then the zero solution of (1.1) can be exponentially stabilized by periodic impulses.

Proof. Here one may choose a sequence $\{t_k\}_{k=1}^\infty$ satisfying (A₁) and $t_{k+1} - t_k = \lambda \geq \tau$, $t_0 = \sigma$. Note that since $\delta = \varrho = 1$, we only need to choose

$$|I_k(u)| = d|u|, \quad |J_{jk}(v)| = d|v|, \quad d = \left(\Gamma - \sum_{i=1}^N \frac{p_i \tau_i}{\beta} \right)^{\frac{1}{2}},$$

$$\Gamma = \exp[-2\alpha(\lambda + \tau)] \exp \left[- \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \right) \lambda \right].$$

Since this proof is similar to the proof of Theorem 2.1, we omit it partly. Finally, it can be deduced that each solution $x(t) = x(t, \sigma, \varphi, \{x_{j0}\}_{j=1}^{n-1})$ of (1.2) and (1.5) through $(\sigma, \varphi, \{x_{j0}\}_{j=1}^{n-1})$ fulfils

$$\left(\|\varphi\|_\sigma^2 + \sum_{j=1}^{n-1} (x_{j0})^2 \right)^{\frac{1}{2}} \leq \delta^*,$$

hence

$$\left(x^2(t) + \sum_{j=1}^{n-1} (x^{(j)}(t))^2 \right)^{\frac{1}{2}} \leq \varepsilon \exp[-\alpha(t - t_0)], \quad t \geq t_0$$

where $\|\varphi\|_t = \sup_{t-\tau \leq s \leq t} |\varphi(s)|$. □

Theorem 2.3. (Case: either δ or $\rho > 1$.) Assume that the conditions in Theorem 2.1 still hold. Moreover, suppose that

(A₁₁) there exists a nonnegative constant θ such that $g_i(t + \theta) = g_i(t) + \theta$;

(A₁₂) there exists a nonnegative constant μ such that $a(t + \mu) = a(t)$, $b(t + \mu) = b(t)$, $f(t + \mu, x) = f(t, x)$.

Then the zero solution of (1.1) can be exponentially stabilized by periodic impulses.

Proof. Suppose that $\tau < T$, $T = \theta\mu$. Or else, we can choose kT to replace T such that $kT > \tau$, $k \in \mathbb{Z}_+$. Then let $t_{k+1} - t_k = T$, $t_0 = \sigma$. Similarly to the proof of Theorem 2.1, for $t \in [t_1, t_2)$ we choose

$$d_1 = \min \left\{ \eta_1 \exp[\alpha T], \left(\Gamma_1 - \sum_{i=1}^N \frac{p_i \tau_i}{\beta} \right)^{\frac{1}{2}} \right\},$$

$$\Gamma_1 = \exp[-\alpha(T + \tau)] \exp \left[- \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \right) T \right].$$

Since η_1 depends only on system (1.1), d_1 also depends only on system (1.1). It then follows from Theorem 2.1 that the solution $x(t) = x(t, t_1, x(t_1), \{x_{j1}\}_{j=1}^{n-1})$ through $(t_1, x(t_1), \{x_{j1}\}_{j=1}^{n-1})$ fulfils

$$\left(x^2(t_1) + \sum_{j=1}^{n-1} (x^{(j)}(t_1))^2 \right)^{\frac{1}{2}} \leq \eta_1,$$

hence

$$\left(x^2(t) + \sum_{j=1}^{n-1} (x^{(j)}(t))^2 \right)^{\frac{1}{2}} < 1, \quad t \in [t_1, t_2).$$

Next we prove that, for $t \in [t_k, t_{k+1})$, the solution $x(t) = x(t, t_k, x(t_k), \{x_{jk}\}_{j=1}^{n-1})$ through $(t_k, x(t_k), \{x_{jk}\}_{j=1}^{n-1})$ fulfils

$$\left(x^2(t_k) + \sum_{j=1}^{n-1} (x^{(j)}(t_k))^2 \right)^{\frac{1}{2}} \leq \eta_1,$$

hence

$$\left(x^2(t) + \sum_{j=1}^{n-1} (x^{(j)}(t))^2 \right)^{\frac{1}{2}} < 1, \quad t \in [t_k, t_{k+1}).$$

Since $t_{k+1} - t_k = T$, $[t_k, t_{k+1}) = [t_1 + (k-1)T, t_1 + kT)$. We define $\Omega(t) = x(t + (k-1)T)$ where $x(t + (k-1)T) = x(t + (k-1)T, t_k, x(t_k), \{x_{jk}\}_{j=1}^{n-1})$. Hence, we

note that the solution $x(t + (k - 1)T)$ of (1.1) begins at $t = t_k$ and ends at $t = t_{k+1}$, which means that the function $\Omega(t)$ begins at $t = t_1$ and ends at $t = t_2$. On the other hand, in view of (A₁₁) and (A₁₂) we get

$$\begin{aligned} \Omega^{(n)}(t) &= x^{(n)}(t + (k - 1)T) \\ &= -a(t + (k - 1)T)x^{(\gamma)}(t + (k - 1)T) - b(t + (k - 1)T)x^\varrho(t + (k - 1)T) \\ &\quad - \sum_{i=1}^N f_i(t + (k - 1)T, x^\delta(g_i(t + (k - 1)T))) \\ &= -a(t)x^{(\gamma)}(t + (k - 1)T) - b(t)x^\varrho(t + (k - 1)T) - \sum_{i=1}^N f_i(t, x^\delta(g_i(t) + (k - 1)T)) \\ &= -a(t)\Omega^{(\gamma)}(t) - b(t)\Omega^\varrho(t) - \sum_{i=1}^N f_i(t, \Omega^\delta(g_i(t))). \end{aligned}$$

Hence, $\Omega(t)$ is a solution of (1.1) which begins at $t = t_1$ and ends at $t = t_2$. As mentioned above, we obtain

$$\begin{aligned} \left(\Omega^2(t_1) + \sum_{j=1}^{n-1} (\Omega^{(j)}(t_1))^2 \right)^{\frac{1}{2}} &= \left(x^2(t_1 + (k - 1)T) + \sum_{j=1}^{n-1} x^{(j)}(t_1 + (k - 1)T)^2 \right)^{\frac{1}{2}} \\ &= \left(x^2(t_k) + \sum_{j=1}^{n-1} (x^{(j)}(t_k))^2 \right)^{\frac{1}{2}} \leq \eta_1. \end{aligned}$$

Consequently, for $t \in [t_1, t_2)$,

$$\left(x^2(t + (k - 1)T) + \sum_{j=1}^{n-1} (x^{(j)}(t + (k - 1)T))^2 \right)^{\frac{1}{2}} = \left(\Omega^2(t) + \sum_{j=1}^{n-1} (\Omega^{(j)}(t))^2 \right)^{\frac{1}{2}} < 1, \quad t \in [t_1, t_2),$$

which implies that

$$\left(x^2(t) + \sum_{j=1}^{n-1} (x^{(j)}(t))^2 \right)^{\frac{1}{2}} < 1, \quad t \in [t_k, t_{k+1}).$$

Then we only need to choose

$$\begin{aligned} d_k &= d = \min \left\{ \eta_1 \exp[\alpha T], \left(\Gamma - \sum_{i=1}^N \frac{p_i \tau_i}{\beta} \right)^{\frac{1}{2}} \right\}, \\ \Gamma &= \exp[-\alpha(T + \tau)] \exp \left[- \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \right) T \right]. \end{aligned}$$

We see that d_k depends only on system (1.1). The rest of the proof is similar to Theorem 2.1 and we omit it here. The proof is complete. \square

Remark 2.5. Let $f_i = a = 0$ in Theorem 2.3. The corresponding results have been investigated in [9]. If $n = 2$, $\gamma = 1$, $\varrho = 1$, $f_i(t, x) = p(t)x$, $g_i(t) = t - \tau$, the corresponding results have been given in [14].

Next we consider high-order nonlinear retarded differential equations (1.3) and the corresponding equations with impulses (1.4).

Theorem 2.4. Assume that conditions (A₃)–(A₉) hold. Moreover, suppose that (A₁₁)

$$\frac{\tau^2}{2\beta} \sum_{i=1}^N p_i < \exp \left[- \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i \right) \tau \right].$$

Then the zero solution of (1.3) can be exponentially stabilized by impulses.

Proof. Since (A₁₁) holds, there exist $\alpha > 0$ and $\lambda \geq \tau$ such that

$$\frac{1}{2\beta} \sum_{i=1}^N p_i \tau_i^2 \leq \tau \sum_{i=1}^N p_i \leq \exp[-\alpha(\lambda + \tau)] \exp \left[- \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i \right) \lambda \right].$$

Similarly to Theorem 2.1, we choose a sequence $\{t_k\}_{k=1}^\infty$ satisfying (A₁) and $\tau \leq t_{k+1} - t_k \leq \lambda$, $t_0 = \sigma$. Considering condition (A₈), we choose a sequence $\{\eta_k\}_{k=1}^\infty$, $\eta_k \in (0, 1)$ such that, when the solution $x(t) = x(t, t_k, x(t_k), \{x_{jk}\}_{j=1}^{n-1})$ through $(t_k, x(t_k), \{x_{jk}\}_{j=1}^{n-1})$ fulfils

$$\left(x^2(t_k) + \sum_{j=1}^{n-1} (x^{(j)}(t_k))^2 \right)^{\frac{1}{2}} \leq \eta_k,$$

then

$$\left(x^2(t) + \sum_{j=1}^{n-1} (x^{(j)}(t))^2 \right)^{\frac{1}{2}} < 1, \quad t \in [t_k, t_{k+1}).$$

Let

$$\begin{aligned} |I_k(u)| &= d_k|u|, \quad |J_{jk}(v)| = d_k|v|, \\ d_k &= \min \left\{ \eta_k \exp[\alpha(t_1 - \sigma)], \left(\Gamma_k - \sum_{i=1}^N \frac{p_i}{2\beta} \tau_i^2 \right)^{\frac{1}{2}} \right\}, \\ \Gamma_k &= \exp[-2\alpha(t_{k+1} - t_k + \tau)] \exp \left[- \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i \right) (t_{k+1} - t_k) \right], \end{aligned}$$

which implies that $d_k \geq 0$.

For any $\varepsilon \in (0, 1)$, let

$$\delta^* = \min \left\{ \eta_0, \varepsilon, \varepsilon \left(1 + \sum_{i=1}^N \frac{p_i}{2\beta} \tau_i^2 \right)^{-1/4} \right. \\ \left. \times \exp[-\alpha(t_1 - \sigma)] \exp \left[-\frac{1}{2} \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \right) (t_1 - \sigma) \right] \right\}.$$

Next, we prove that each solution $x(t) = x(t, \sigma, \varphi, \{x_{j0}\}_{j=1}^{n-1})$ of (1.4) and (1.5) with

$$\left(\|\varphi\|_\sigma^2 + \sum_{j=1}^{n-1} (x_{j0})^2 \right)^{\frac{1}{2}} \leq \delta^*$$

satisfies

$$\left(x^2(t) + \sum_{j=1}^{n-1} (x^{(j)}(t))^2 \right)^{\frac{1}{2}} \leq \varepsilon \exp[-\alpha(t - t_0)], \quad t \geq t_0,$$

where $\|\varphi\|_t = \sup_{t-\tau \leq s \leq t} |\varphi(s)|$.

First, for $t \in [\sigma, t_1)$ we choose a Lyapunov function

$$V(t) = x^2(t) + \sum_{i=1}^{n-1} (x^{(i)}(t))^2 + \sum_{i=1}^N \frac{p_i}{\beta} \int_{g_i(t)}^t \int_u^t x^{2\delta}(s) \, ds \, du.$$

Then $V(t)$ satisfies:

(1) $V(t) \geq x^2(t) + \sum_{i=1}^{n-1} (x^{(i)}(t))^2;$

(2)

$$\begin{aligned} V(t) &\leq x^2(t) + \sum_{i=1}^{n-1} (x^{(i)}(t))^2 + \sum_{i=1}^N \frac{p_i}{\beta} \sup_{g_i(t) \leq s \leq t} |x(s)|^{2\delta} \int_{g_i(t)}^t \int_u^t \, ds \, du \\ &\leq x^2(t) + \sum_{i=1}^{n-1} (x^{(i)}(t))^2 + \sum_{i=1}^N \frac{p_i}{\beta} \sup_{g_i(t) \leq s \leq t} |x(s)|^{2\delta} \frac{(t - g_i(t))^2}{2} \\ &\leq x^2(t) + \sum_{i=1}^{n-1} (x^{(i)}(t))^2 + \sum_{i=1}^N \frac{p_i}{\beta} \sup_{t-\tau_i \leq s \leq t} |x(s)|^{2\delta} \frac{\tau_i^2}{2} \\ &\leq x^2(t) + \sum_{i=1}^{n-1} (x^{(i)}(t))^2 + \sum_{i=1}^N \frac{p_i}{2\beta} \tau_i^2 \|x\|_t^2 \\ &\leq \left(1 + \sum_{i=1}^N \frac{p_i}{2\beta} \tau_i^2 \right) \left(\|x\|_t^2 + \sum_{i=1}^{n-1} (x^{(i)}(t))^2 \right), \end{aligned}$$

where $\|x\|_t = \sup_{t-\tau \leq s \leq t} |x(s)|$.

(3) We denote by $V'(t)$ the right upper derivative of $V(t)$ along the solution of (1.4) and (1.5). Then

$$\begin{aligned}
V'(t) &= 2x(t)x'(t) + 2x'(t)x''(t) + \dots + 2x^{(n-1)}(t)x^{(n)}(t) \\
&\quad + \sum_{i=1}^N \frac{p_i}{\beta} x^{2\delta}(t)(t - g_i(t)) - \sum_{i=1}^N \frac{p_i}{\beta} g_i'(t) \int_{g_i(t)}^t x^{2\delta}(s) \, ds \\
&\leq 2x(t)x'(t) + 2x'(t)x''(t) + \dots + 2x^{(n-1)}(t) \left\{ -a(t)x^{(\gamma)}(t) - b(t)x^\varrho(t) \right. \\
&\quad \left. - \sum_{i=1}^N \int_{g_i(t)}^t f_i(t-u, x^\delta(u)) \, du \right\} + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i x^{2\delta}(t) - \sum_{i=1}^N \frac{p_i}{\beta} g_i'(t) \int_{g_i(t)}^t x^{2\delta}(s) \, ds \\
&\leq x^2(t) + [x'(t)]^2 + \dots + [x^{(n-2)}(t)]^2 + [x^{(n-1)}(t)]^2 + a([x^{(n-1)}(t)]^2 \\
&\quad + [x^{(\gamma)}(t)]^2) + b([x^{(n-1)}(t)]^2 + x^{2\varrho}(t)) \\
&\quad + 2|x^{(n-1)}(t)| \sum_{i=1}^N \int_{g_i(t)}^t |p_i(t-u)| |x^{2\delta}(u)| \, du \\
&\quad + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i x^{2\delta}(t) - \sum_{i=1}^N \frac{p_i}{\beta} g_i'(t) \int_{g_i(t)}^t x^{2\delta}(s) \, ds \\
&\leq x^2(t) + [x'(t)]^2 + \dots + [x^{(n-2)}(t)]^2 + [x^{(n-1)}(t)]^2 + a([x^{(n-1)}(t)]^2 \\
&\quad + [x^{(\gamma)}(t)]^2) + b([x^{(n-1)}(t)]^2 + x^2(t)) + \sum_{i=1}^N p_i \int_{g_i(t)}^t ([x^{(n-1)}(t)]^2 + x^{2\delta}(u)) \, du \\
&\quad + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i x^{2\delta}(t) - \sum_{i=1}^N \frac{p_i}{\beta} g_i'(t) \int_{g_i(t)}^t x^{2\delta}(s) \, ds \\
&\leq x^2(t) + [x'(t)]^2 + \dots + [x^{(n-2)}(t)]^2 + [x^{(n-1)}(t)]^2 + a([x^{(n-1)}(t)]^2 \\
&\quad + [x^{(\gamma)}(t)]^2) + b([x^{(n-1)}(t)]^2 + x^2(t)) + \sum_{i=1}^N p_i \tau_i [x^{(n-1)}(t)]^2 \\
&\quad + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i x^2(t) + \sum_{i=1}^N p_i \int_{g_i(t)}^t x^{2\delta}(u) \, du \left(1 - \frac{g_i'(t)}{\beta}\right) \\
&\leq \left(1 + b + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i\right) x^2(t) + 2[x'(t)]^2 + \dots + 2[x^{(\gamma-1)}(t)]^2 + (a+2)[x^{(\gamma)}(t)]^2 \\
&\quad + 2[x^{(\gamma+1)}(t)]^2 + \dots + \left(2 + a + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i\right) [x^{(n-1)}(t)]^2 \\
&\leq \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i\right) V(t),
\end{aligned}$$

which implies that

$$V(t) \leq V(t_0) \exp \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i \right) (t - \sigma), \quad t \in [\sigma, t_1].$$

Thus for $t \in [\sigma, t_1)$ we get

$$\begin{aligned} x^2(t) + \sum_{i=1}^{n-1} (x^{(j)}(t))^2 &\leq V(t) \leq V(\sigma) \exp \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i \right) (t - \sigma) \\ &< V(\sigma) \exp \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i \right) (t_1 - \sigma) \\ &\leq \left(1 + \sum_{i=1}^N \frac{p_i}{2\beta} \tau_i^2 \right) \left(\|x\|_{\sigma}^2 + \sum_{i=1}^{n-1} (x^{(j)}(\sigma))^2 \right) \exp \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i \right) (t_1 - \sigma) \\ &= \left(1 + \sum_{i=1}^N \frac{p_i}{2\beta} \tau_i^2 \right) \left(\|\varphi\|_{\sigma}^2 + \sum_{j=1}^{n-1} x_{j0}^2 \right) \exp \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i \right) (t_1 - \sigma) \\ &\leq \left(1 + \sum_{i=1}^N \frac{p_i}{2\beta} \tau_i^2 \right) \delta^{*2} \exp \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i \right) (t_1 - \sigma) \\ &= \varepsilon^2 \exp[-2\alpha(t_1 - \sigma)] < \varepsilon^2 \exp[-2\alpha(t - \sigma)], \end{aligned}$$

which implies that

$$\left(x^2(t) + \sum_{i=1}^{n-1} (x^{(j)}(t))^2 \right)^{\frac{1}{2}} < \varepsilon \exp[-\alpha(t - \sigma)], \quad t \in [\sigma, t_1].$$

Especially,

$$\left(x^2(t_1^-) + \sum_{i=1}^{n-1} (x^{(j)}(t_1^-))^2 \right)^{\frac{1}{2}} < \varepsilon \exp[-\alpha(t_1 - \sigma)].$$

It then follows that

$$\begin{aligned} \left(x^2(t_1) + \sum_{i=1}^{n-1} (x^{(j)}(t_1))^2 \right)^{\frac{1}{2}} &= d_1 \left(x^2(t_1^-) + \sum_{i=1}^{n-1} (x^{(j)}(t_1^-))^2 \right)^{\frac{1}{2}} \\ &< d_1 \varepsilon \exp[-\alpha(t_1 - \sigma)] < d_1 \exp[-\alpha(t_1 - \sigma)] = \eta_1. \end{aligned}$$

Thus we obtain that for $t \in [t_1, t_2)$,

$$\left(x^2(t) + \sum_{i=1}^{n-1} (x^{(j)}(t))^2 \right)^{\frac{1}{2}} < 1.$$

For $t \in [t_1, t_2)$, we choose a Lyapunov function

$$V(t) = x^2(t) + \sum_{i=1}^{n-1} (x^{(j)}(t))^2 + \sum_{i=1}^N \frac{p_i}{\beta} \int_{g_i(t)}^t \int_u^t x^{2\delta}(s) ds du.$$

Then we have

$$\begin{aligned} x^2(t) + \sum_{i=1}^{n-1} (x^{(j)}(t))^2 &\leq V(t) \leq V(t_1) \exp\left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i\right) (t - t_1) \\ &< V(t_1) \exp\left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i\right) (t_2 - t_1) \\ &= \left[x^2(t_1) + \sum_{i=1}^{n-1} (x^{(j)}(t_1))^2 + \sum_{i=1}^N \frac{p_i}{\beta} \int_{g_i(t_1)}^{t_1} \int_u^{t_1} x^{2\delta}(s) ds du \right] \\ &\quad \times \exp\left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i\right) (t_2 - t_1) \\ &\leq \left\{ d_1^2 \left[x^2(t_1^-) + \sum_{i=1}^{n-1} (x^{(j)}(t_1^-))^2 \right] + \sum_{i=1}^N \frac{p_i}{2\beta} \tau_i^2 \sup_{t_1 - \tau_i \leq t \leq t_1} x^{2\delta}(t) \right\} \\ &\quad \times \exp\left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i\right) (t_2 - t_1) \\ &\leq \left\{ d_1^2 \left[x^2(t_1^-) + \sum_{i=1}^{n-1} (x^{(j)}(t_1^-))^2 \right] + \sum_{i=1}^N \frac{p_i}{2\beta} \tau_i^2 \sup_{t_1 - \tau \leq t \leq t_1} x^2(t) \right\} \\ &\quad \times \exp\left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i\right) (t_2 - t_1) \\ &\leq \left\{ d_1^2 \sup_{t_1 - \tau \leq t \leq t_1} \left[x^2(t) + \sum_{i=1}^{n-1} (x^{(j)}(t))^2 \right] + \sum_{i=1}^N \frac{p_i}{2\beta} \tau_i^2 \sup_{t_1 - \tau \leq t \leq t_1} x^2(t) \right\} \\ &\quad \times \exp\left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i\right) (t_2 - t_1) \\ &\leq \left(d_1^2 + \sum_{i=1}^N \frac{p_i}{2\beta} \tau_i^2 \right) \sup_{t_1 - \tau \leq t \leq t_1} \left[x^2(t) + \sum_{i=1}^{n-1} (x^{(j)}(t))^2 \right] \\ &\quad \times \exp\left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i\right) (t_2 - t_1) \\ &\leq \Gamma_1 \varepsilon^2 \exp[-2\alpha(t_1 - \sigma - \tau)] \exp\left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \tau_i\right) (t_2 - t_1) \\ &\leq \varepsilon^2 \exp[-2\alpha(t_2 - \sigma)] < \varepsilon^2 \exp[-2\alpha(t - \sigma)], \quad t \in [t_1, t_2). \end{aligned}$$

Hence,

$$\left(x^2(t) + \sum_{i=1}^{n-1} (x^{(i)}(t))^2\right)^{\frac{1}{2}} < \varepsilon \exp[-\alpha(t - \sigma)], \quad t \in [t_1, t_2].$$

By the induction hypothesis, we may prove that, for $k \geq 1$,

$$\left(x^2(t) + \sum_{i=1}^{n-1} (x^{(i)}(t))^2\right)^{\frac{1}{2}} < \varepsilon \exp[-\alpha(t - \sigma)], \quad t \in [t_k, t_{k+1}).$$

Therefore, we arrive at

$$\left(x^2(t) + \sum_{i=1}^{n-1} (x^{(i)}(t))^2\right)^{\frac{1}{2}} < \varepsilon \exp[-\alpha(t - \sigma)], \quad t \geq \sigma,$$

and the proof is complete. □

Remark 2.6. Remarks 2.1–2.3 are still valid for Theorem 2.4.

3. AN EXAMPLE

Example. Consider the equation

$$(3.1) \quad \begin{cases} x^{(6)}(t) + x^{(3)}(t) + 0.9x(t) - 0.5x(t - 0.01) - 0.5x(t - 0.02) \\ \quad + a(t)x^3(t - 0.015) = 0, \quad t \geq 0, \\ x(t) = \varphi(t), \quad -0.02 \leq t \leq 0, \quad x^{(j)}(0) = x_{j0}, \quad j = 1, \dots, 5, \end{cases}$$

where $a(t) \in \Gamma$, $\Gamma = \{s(t) \in [0, \infty) : |s(t)| \leq 0.5\}$.

When $a(t) = 0 \in \Gamma$, then its characteristic equation is

$$\lambda^6 + \lambda^3 - 0.5e^{-0.01\lambda} - 0.5e^{-0.02\lambda} + 0.9 = 0.$$

Using the software Mathematica, we obtain a characteristic root with positive real part. Hence the non-impulsive equation (4.1) is unstable for some $a(t) \in \Gamma$. But if we take $\beta = 1$, $\tau = \lambda = 0.02$, $\alpha = \frac{1}{2}$, $p_i = 0.5$, $i = 1, 2, 3$, $N = 3$, then it is easy to check that

$$\begin{aligned} \frac{\tau}{\beta} \sum_{i=1}^N p_i &= 0.03 < e^{-0.02} e^{-0.108} = \exp[-\alpha(\lambda + \tau)] \exp \left[- \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \right) \lambda \right] \\ &< \exp \left[- \left(2 + a + b + \sum_{i=1}^N \frac{p_i}{\beta} \right) \tau \right]. \end{aligned}$$

We choose a sequence $\{t_k\}_{k=1}^{\infty}$ satisfying (A₁) and $t_{k+1} - t_k = 0.02$, $t_0 = 0$, $k \in \mathbb{Z}_+$, such that

$$\begin{aligned}x(t_k) &= dx(t_k^-), \\x^{(j)}(t_k) &= dx^{(j)}(t_k^-), \quad j = 1, 2, \dots, 5,\end{aligned}$$

where $d = \min\{\eta_1 \exp[0.01], \sqrt{e^{-0.128} - 0.0225}\}$ and η_1 depends only on the first equation of (4.1) on $[0.02, 0.04]$. Then the hypotheses in Theorem 2.3 are satisfied and hence the unstable equation (4.2) can be exponentially stabilized by periodic impulses for all $a(t) \in \Gamma$.

References

- [1] *D. D. Bainov, P. S. Simeonov*: Systems with Impulsive Effect: Stability, Theory and Applications. Ellis Horwood, New York, 1989.
- [2] *X. Fu, B. Yan, Y. Liu*: Introduction of Impulsive Differential Systems. Science Press, Beijing, 2005.
- [3] *L. P. Gimenes, M. Federson*: Existence and impulsive stability for second order retarded differential equations. *Appl. Math. Comput.* *177* (2006), 44–62.
- [4] *L. P. Gimenes, M. Federson, P. Táboas*: Impulsive stability for systems of second order retarded differential equations. *Nonlinear Anal., Theory Methods Appl.* *67* (2007), 545–553.
- [5] *A. Khadra, X. Liu, X. Shen*: Application of impulsive synchronization to communication security. *IEEE Trans. Circuits Systems I Fund. Theory Appl.* *50* (2003), 341–351.
- [6] *V. Lakshmikantham, D. D. Bajnov, P. S. Simeonov*: Theory of Impulsive Differential Equations. World Scientific, Singapore, 1989.
- [7] *X. Li*: Exponential stability of Hopfield neural networks with time-varying delays via impulsive control. *Math. Methods Appl. Sci.* *33* (2010), 1596–1604.
- [8] *X. Li*: New results on global exponential stabilization of impulsive functional differential equations with infinite delays or finite delays. *Nonlinear Anal., Real World Appl.* *11* (2010), 4194–4201.
- [9] *X. Li, P. Weng*: Impulsive stabilization of a class of nonlinear differential equations. *Appl. Math., Ser. A (Chin. Ed.)* *18* (2003), 408–416.
- [10] *X. Liu*: Impulsive stabilization and applications to population growth models. *Rocky Mt. J. Math.* *25* (1995), 381–395.
- [11] *X. Liu, G. Ballinger*: Existence and continuability of solutions for differential equations with delays and state-dependent impulses. *Nonlinear Anal., Theory Methods Appl.* *51* (2002), 633–647.
- [12] *Z. Luo, J. Shen*: Impulsive stabilization of functional differential equations with infinite delays. *Appl. Math. Lett.* *16* (2003), 695–701.
- [13] *J. E. Prussing, L. J. Wellnitz, W. G. Heckathorn*: Optimal impulsive time-fixed direct-ascent interception. *J. Guidance Control Dynam.* *12* (1989), 487–494.
- [14] *A. Weng, J. Sun*: Impulsive stabilization of second-order delay differential equations. *Nonlinear Anal., Real World Appl.* *8* (2007), 1410–1420.
- [15] *T. Yang*: Impulsive Systems and Control: Theory and Applications. Nova Science Publishers, New York, 2001.

- [16] *H. Zhang, J. Jiao, L. Chen*: Pest management through continuous and impulsive control strategies. *BioSystems* 90 (2007), 350–361.

Authors' addresses: *Juan Liu*, Department of Mathematics, Henan Institute of Science and Technology, Xinxiang 453003, P. R. China; *Xiaodi Li* (corresponding author), School of Mathematical Sciences, Shandong Normal University, Ji'nan 250014, P. R. China, e-mail: sodymath@163.com.