

# Applications of Mathematics

---

Sanjoy Ghosal

Statistical convergence of a sequence of random variables and limit theorems

*Applications of Mathematics*, Vol. 58 (2013), No. 4, 423--437

Persistent URL: <http://dml.cz/dmlcz/143339>

## Terms of use:

© Institute of Mathematics AS CR, 2013

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

STATISTICAL CONVERGENCE OF A SEQUENCE OF  
RANDOM VARIABLES AND LIMIT THEOREMS

SANJOY GHOSAL, Kalyani

(Received July 29, 2011)

*Abstract.* In this paper the ideas of three types of statistical convergence of a sequence of random variables, namely, statistical convergence in probability, statistical convergence in mean of order  $r$  and statistical convergence in distribution are introduced and the inter-relation among them is investigated. Also their certain basic properties are studied.

*Keywords:* asymptotic density, random variable, statistical convergence, statistical convergence in probability, statistical convergence in mean of order  $r$ , statistical convergence in distribution

*MSC 2010:* 40Axx, 40Cxx, 60Fxx, 60Gxx

1. INTRODUCTION AND BACKGROUND

The idea of statistical convergence was known to A. Zygmund as early as 1935 and in particular after 1951 when Steinhaus [22] and Fast [4] reintroduced statistical convergence for sequences of real numbers, several generalizations and applications of this notion have been investigated (see [1], [5], [6], [8], [10], [12], [19], [21] where many more references can be found). Recall that a subset  $A$  of the set  $\mathbb{N}$  of natural numbers is said to have ‘asymptotic density’  $d(A)$  if

$$d(A) = \lim_{n \rightarrow \infty} \frac{1}{n} |\{k \leq n : k \in A\}|,$$

where the vertical bars denote the cardinality of the enclosed set. The sequence  $\{x_n\}_{n \in \mathbb{N}}$  is said to be statistically convergent to a real number  $x$  if for each  $\varepsilon > 0$ ,

$$\lim_{n \rightarrow \infty} \frac{1}{n} |\{k \leq n : |x_k - x| \geq \varepsilon\}| = 0;$$

and we write  $x_n \xrightarrow{\text{st}} x$  or  $\text{st-}\lim_{n \rightarrow \infty} x_n = x$ .

Now statistical convergence has turned out to be one of the most active areas of research in summability theory and has several applications in different fields of mathematics: number theory [3], trigonometric series [23], probability theory [7], measure theory [15], optimization [16], approximation theory [9], Hopfield neural network [14] (in this paper a geometrical vision of the dynamic states is explained). Statistical convergence has also been discussed in more general abstract spaces such as the fuzzy number space [20], locally convex space [13] and Banach space [11].

On the other hand, if for each positive integer  $n$  a random variable  $X_n$  is defined on a given event space  $S$  (the same for all  $n$ ) with respect to a given class of events  $\Delta$  and a probability function  $P: \Delta \rightarrow \mathbb{R}$  (where  $\mathbb{R}$  denotes the set of all real numbers) then we say that  $X_1, X_2, X_3, \dots, X_n, \dots$  is a sequence of random variables and as in analysis we denoted the sequence by  $\{X_n\}_{n \in \mathbb{N}}$ .

From the practical point of view the discussion of a random variable  $X$  will be highly significant if it is known that there exists a real constant  $c$  for which  $P(|X - c| < \varepsilon) \simeq 1$ , where  $\varepsilon > 0$  is sufficiently small, that is, it is nearly certain that values of  $X$  lie in a very small neighbourhood of  $c$ .

For a sequence of random variables  $\{X_n\}_{n \in \mathbb{N}}$ , each  $X_n$  may not have the above property but it may happen that the aforesaid property (with respect to a real constant  $c$ ) becomes more and more distinguished as  $n$  gradually increases and the question of existence of such a real constant  $c$  will be answered by a concept of convergence in probability of the sequence  $\{X_n\}_{n \in \mathbb{N}}$ .

Again the sequence  $\{X_n\}_{n \in \mathbb{N}}$  may be such that as  $n$  gradually increases the distribution function  $F_n(x)$  of  $X_n$  may more and more resemble the distribution function of a particular random variable and the question of existence of such a distribution function is related to the concept of 'convergence in distribution' of the sequence  $\{X_n\}_{n \in \mathbb{N}}$ .

Besides the above mentioned two modes of convergence, there are other modes of convergence of the sequence  $\{X_n\}_{n \in \mathbb{N}}$ . In this paper, we will limit our discussion to three types of statistical convergence of a sequence of random variables, namely,

- (a) statistical convergence in probability,
- (b) statistical convergence in mean of order  $r$ ,
- (c) statistical convergence in distribution.

We will conclude the paper discussing some fundamental limit theorems related to the modes of convergence (a), (b), (c) which effectively extend and improve all the existing results in this direction [18].

Now probability convergence has several applications in different fields of mathematics: graph theory (minimal spanning tree) [17], measure theory [15], Lorentz gas [2].

## 2. STATISTICAL CONVERGENCE IN PROBABILITY

Let  $\{X_n\}_{n \in \mathbb{N}}$  be a sequence of random variables, where each  $X_n$  is defined on the same event space  $S$  with respect to a given class of subsets (of  $S$ ) as the class  $\Delta$  of events and a given probability function  $P: \Delta \rightarrow \mathbb{R}$ . The sequence  $\{X_n\}_{n \in \mathbb{N}}$  is said to be statistically convergent in probability to a random variable  $X$  (where  $X: S \rightarrow \mathbb{R}$ ) if for any  $\varepsilon, \delta > 0$

$$\lim_{n \rightarrow \infty} \frac{1}{n} |\{k \leq n: P(|X_k - X| \geq \varepsilon) \geq \delta\}| = 0,$$

$$\text{or equivalently, } \lim_{n \rightarrow \infty} \frac{1}{n} |\{k \leq n: 1 - P(|X_k - X| < \varepsilon) \geq \delta\}| = 0$$

and we write  $\text{st-}\lim_{n \rightarrow \infty} P(|X_n - X| \geq \varepsilon) = 0$  or  $\text{st-}\lim_{n \rightarrow \infty} P(|X_n - X| < \varepsilon) = 1$  or  $X_n \xrightarrow{\text{st}_p} X$ .

**Theorem 2.1.** *If a sequence of constants  $x_n \xrightarrow{\text{st}} x$  then regarding a constant as a random variable having a one-point distribution at that point, we may also write  $x_n \xrightarrow{\text{st}_p} x$ .*

**Proof.** Let  $\varepsilon$  be any positive real number. Then  $d(M) = 1$ , where  $M = \{n \in \mathbb{N}: |x_n - x| < \varepsilon\}$ . Thus for any  $\delta > 0$ ,  $B = \{n \in \mathbb{N}: 1 - P(|x_n - x| < \varepsilon) \geq \delta\} \subseteq \mathbb{N} \setminus M$  implies  $d(B) = 0$ . This implies  $x_n \xrightarrow{\text{st}_p} x$ .  $\square$

The following example shows that in general the converse of Theorem 2.1 is not true.

**Example 2.1.1.** Let the probability density function  $X_n$  be

$$f_n(x) = \begin{cases} 1, & \text{where } 0 < x < 1; \\ \frac{nx^{n-1}}{2^n}, & \text{where } 0 < x < 2; \end{cases} \quad \begin{cases} 0 \text{ otherwise, if } n = 2^m \text{ where } \forall m \in \mathbb{N}; \\ 0 \text{ otherwise, if } n \neq 2^m \text{ where } \forall m \in \mathbb{N}. \end{cases}$$

Let  $0 < \varepsilon, \delta < 1$ . Then

$$\begin{aligned} P(|X_n - 2| \geq \varepsilon) &= \begin{cases} 1 & \text{if } n = 2^m \text{ where } \forall m \in \mathbb{N}, \\ 1 - P(|X_n - 2| < \varepsilon) = 1 - \left\{1 - \left(\frac{2-\varepsilon}{2}\right)^n\right\} = \left(1 - \frac{\varepsilon}{2}\right)^n & \text{if } n \neq 2^m \text{ where } \forall m \in \mathbb{N} \end{cases} \\ \Rightarrow \lim_{n \rightarrow \infty} \frac{1}{n} |\{k \leq n: P(|X_k - 2| \geq \varepsilon) \geq \delta\}| &\leq \lim_{n \rightarrow \infty} \frac{1}{n} |\{2^0, 2^1, 2^2, \dots\} \cup A| = 0, \\ &\quad (\text{where } A \text{ is a finite subset of } \mathbb{N}) \\ \Rightarrow \lim_{n \rightarrow \infty} \frac{1}{n} |\{k \leq n: P(|X_k - 2| \geq \varepsilon) \geq \delta\}| &= 0. \end{aligned}$$

However, it is not ordinary statistical convergence of a sequence of numbers.

Although the concept of statistical convergence in probability is basically different from that of ordinary statistical convergence of sequence of numbers, the following simple results hold for statistical convergence in probability as well.

**Theorem 2.2** (Elementary Properties).

- (i)  $X_n \xrightarrow{\text{st}_p} X$  and  $X_n \xrightarrow{\text{st}_p} Y \Rightarrow P\{X = Y\} = 1$ ,
- (ii)  $X_n \xrightarrow{\text{st}_p} X \Leftrightarrow X_n - X \xrightarrow{\text{st}_p} 0$ ,
- (iii)  $X_n \xrightarrow{\text{st}_p} X \Rightarrow cX_n \xrightarrow{\text{st}_p} cX$  where  $c \in \mathbb{R}$ ,
- (iv)  $X_n \xrightarrow{\text{st}_p} X$  and  $Y_n \xrightarrow{\text{st}_p} Y \Rightarrow X_n + Y_n \xrightarrow{\text{st}_p} X + Y$ ,
- (v)  $X_n \xrightarrow{\text{st}_p} X$  and  $Y_n \xrightarrow{\text{st}_p} Y \Rightarrow X_n - Y_n \xrightarrow{\text{st}_p} X - Y$ ,
- (vi)  $X_n \xrightarrow{\text{st}_p} x \Rightarrow X_n^2 \xrightarrow{\text{st}_p} x^2$ ,
- (vii)  $X_n \xrightarrow{\text{st}_p} x$  and  $Y_n \xrightarrow{\text{st}_p} y \Rightarrow X_n \cdot Y_n \xrightarrow{\text{st}_p} x \cdot y$ ,
- (viii)  $X_n \xrightarrow{\text{st}_p} x$  and  $Y_n \xrightarrow{\text{st}_p} y \Rightarrow X_n/Y_n \xrightarrow{\text{st}_p} x/y$  provided  $y \neq 0$ ,
- (ix)  $X_n \xrightarrow{\text{st}_p} X$  and  $Y_n \xrightarrow{\text{st}_p} Y \Rightarrow X_n \cdot Y_n \xrightarrow{\text{st}_p} X \cdot Y$ ,
- (x) if  $0 \leq X_n \leq Y_n$  and  $Y_n \xrightarrow{\text{st}_p} 0 \Rightarrow X_n \xrightarrow{\text{st}_p} 0$ ,
- (xi) if  $X_n \xrightarrow{\text{st}_p} X$ , then for each  $\varepsilon, \delta > 0$  there exists  $k \in \mathbb{N}$  such that

$$d(\{n \in \mathbb{N} : P(|X_n - X_k| \geq \varepsilon) \geq \delta\}) = 0$$

(this is called the statistical Cauchy condition in probability).

*Proof.* Let  $\varepsilon, \delta$  be any positive real numbers.

(i) Let  $k \in \{n \in \mathbb{N} : P(|X_n - X| \geq \frac{1}{2}\varepsilon) < \frac{1}{2}\delta\} \cap \{n \in \mathbb{N} : P(|X_n - Y| \geq \frac{1}{2}\varepsilon) < \frac{1}{2}\delta\}$  (the existence of  $k$  is granted, since the asymptotic density of both the sets is 1). Then  $P(|X - Y| \geq \varepsilon) \leq P(|X_k - X| \geq \frac{1}{2}\varepsilon) + P(|X_k - Y| \geq \frac{1}{2}\varepsilon) < \delta$ . This implies  $P\{X = Y\} = 1$ .

For (ii), (iii), (iv), (v) the proofs are straightforward and hence omitted.

(vi) If  $Z_n \xrightarrow{\text{st}_p} 0$  then  $Z_n^2 \xrightarrow{\text{st}_p} 0$  for  $\{n \in \mathbb{N} : P(|Z_n^2 - 0| \geq \varepsilon) \geq \delta\} = \{n \in \mathbb{N} : P(|Z_n - 0| \geq \sqrt{\varepsilon}) \geq \delta\}$ . Now  $X_n^2 = (X_n - x)^2 + 2x(X_n - x) + x^2 \xrightarrow{\text{st}_p} x^2$ .

(vii) We have  $X_n Y_n = \frac{1}{4}\{(X_n + Y_n)^2 - (X_n - Y_n)^2\} \xrightarrow{\text{st}_p} \frac{1}{4}\{(x + y)^2 - (x - y)^2\} = xy$ .

(viii) Let  $A, B$  represent the events ' $|Y_n - y| < |y|$ ', ' $|1/Y_n - 1/y| \geq \varepsilon$ ', respectively.

Now

$$\left| \frac{1}{Y_n} - \frac{1}{y} \right| = \frac{|Y_n - y|}{|yY_n|} = \frac{|Y_n - y|}{|y| \cdot |y + (Y_n - y)|} \leq \frac{|Y_n - y|}{|y| \cdot (|y| - |Y_n - y|)}.$$

If  $A, B$  occur simultaneously, then

$$|Y_n - y| \geq \frac{\varepsilon|y|^2}{1 + \varepsilon|y|} \quad (\text{by the above inequality}).$$

Let  $\varepsilon_0 = \varepsilon|y|^2/(1 + \varepsilon|y|)$  and let  $C$  be the event ' $|Y_n - y| \geq \varepsilon_0$ '. This implies  $AB \subseteq C \Rightarrow P(B) \leq P(C) + P(\bar{A})$ , where bar denotes the set complement. This implies  $\{n \in \mathbb{N} : P(|1/Y_n - 1/y| \geq \varepsilon) \geq \delta\} \subseteq \{n \in \mathbb{N} : P(|Y_n - y| \geq \varepsilon_0) \geq \frac{1}{2}\delta\} \cup \{n \in \mathbb{N} : P(|Y_n - y| \geq |y|) \geq \frac{1}{2}\delta\}$ . Hence,  $1/Y_n \xrightarrow{\text{stp}} 1/y$  provided  $y \neq 0$ . Finally,  $X_n/Y_n \xrightarrow{\text{stp}} x/y$  provided  $y \neq 0$  (by Theorem 2.2 (vii)).

(ix) First prove that if  $X_n \xrightarrow{\text{stp}} X$  and  $Z$  is a random variable then  $X_n Z \xrightarrow{\text{stp}} XZ$ . Since  $Z$  is a random variable, so given  $\delta > 0$ , there exists an  $\alpha > 0$  such that  $P(|Z| > \alpha) \leq \frac{1}{2}\delta$ . Then for any  $\varepsilon > 0$ ,  $P(|X_n Z - XZ| \geq \varepsilon) = P(|X_n - X||Z| \geq \varepsilon, |Z| > \alpha) + P(|X_n - X||Z| \geq \varepsilon, |Z| \leq \alpha) \leq \frac{1}{2}\delta + P(|X_n - X| \geq \varepsilon/\alpha)$ . This implies  $\{n \in \mathbb{N} : P(|X_n Z - XZ| \geq \varepsilon) \geq \delta\} \subseteq \{n \in \mathbb{N} : P(|X_n - X| \geq \varepsilon/\alpha) \geq \frac{1}{2}\delta\}$ . Next  $(X_n - X)(Y_n - Y) \xrightarrow{\text{stp}} 0$ . This implies  $X_n Y_n \xrightarrow{\text{stp}} XY$ .

(x) Proof is straightforward and hence omitted.

(xi) Choose  $k \in \mathbb{N}$  such that  $P(|X_k - X| \geq \frac{1}{2}\varepsilon) < \frac{1}{2}\delta$  ( $k$  exists since  $d(\{n \in \mathbb{N} : P(|X_n - X| \geq \frac{1}{2}\varepsilon) < \frac{1}{2}\delta\}) = 1$ ). Then  $\{n \in \mathbb{N} : P(|X_n - X_k| \geq \varepsilon) \geq \delta\} \subseteq \{n \in \mathbb{N} : P(|X_n - X| \geq \frac{1}{2}\varepsilon) \geq \frac{1}{2}\delta\}$ .  $\square$

**Theorem 2.3.** Let  $\{X_n\}_{n \in \mathbb{N}}$  be a sequence of random variables such that there exists a sequence of constants  $\{x_n\}_{n \in \mathbb{N}}$  with the property that

$$X_n - x_n \xrightarrow{\text{stp}} 0.$$

If  $m(X_n)$  is a median of  $X_n$  then

$$X_n - m(X_n) \xrightarrow{\text{stp}} 0 \quad \text{and} \quad x_n - m(X_n) \xrightarrow{\text{st}} 0.$$

**Proof.** Proof is straightforward and hence omitted.  $\square$

**Theorem 2.4.** Let  $X_n \xrightarrow{\text{stp}} X$  and let  $g: \mathbb{R} \rightarrow \mathbb{R}$  be a continuous on  $\mathbb{R}$ . Then  $g(X_n) \xrightarrow{\text{stp}} g(X)$ .

**Proof.** Since  $X$  is a random variable, hence for each  $\delta > 0$  there exists  $\alpha$  such that  $P(|X| > \alpha) \leq \frac{1}{2}\delta$ . Since  $g$  is uniformly continuous on  $[-\alpha, \alpha]$ , hence for each  $\varepsilon > 0$  there exists  $\delta_0$  such that

$$|g(x_n) - g(x)| < \varepsilon \quad \text{whenever} \quad |x| \leq \alpha \quad \text{and} \quad |x_n - x| < \delta_0.$$

It follows that

$$P(|g(X_n) - g(X)| \geq \varepsilon) \leq P(|X_n - X| \geq \delta_0) + P(|X| > \alpha) \leq P(|X_n - X| \geq \delta_0) + \frac{1}{2}\delta.$$

This implies  $\{n \in \mathbb{N} : P(|g(X_n) - g(X)| \geq \varepsilon) \geq \delta\} \subseteq \{n \in \mathbb{N} : P(|X_n - X| \geq \delta_0) \geq \frac{1}{2}\delta\}$ . Hence the result.  $\square$

**Corollary 2.4.1.** If  $X_n \xrightarrow{\text{stp}} x$  and  $g: \mathbb{R} \rightarrow \mathbb{R}$  is a continuous function, then  $g(X_n) \xrightarrow{\text{stp}} g(x)$ .

**Proposition 2.1.** Let  $\{a_n\}_{n \in \mathbb{N}}$ ,  $\{b_n\}_{n \in \mathbb{N}}$ ,  $\{c_n\}_{n \in \mathbb{N}}$  be three sequences of real numbers such that  $d(\{n \in \mathbb{N}: a_n \leq b_n \leq c_n\}) = 1$  and  $\text{st-lim}_{n \rightarrow \infty} a_n = \text{st-lim}_{n \rightarrow \infty} c_n = x$ . Then  $\text{st-lim}_{n \rightarrow \infty} b_n = x$ .

The next theorem is the generalization of Tchebycheff's theorem.

**Theorem 2.5.** If  $\{X_n\}_{n \in \mathbb{N}}$  is a sequence of random variables such that for any  $n$ ,  $X_n$  has a finite mean  $m_n$ , then

$$X_n - m_n \xrightarrow{\text{stp}} 0 \quad \text{provided } \text{st-lim}_{n \rightarrow \infty} \sigma_n = 0.$$

*Proof.* The proof is straightforward and hence omitted. □

**Corollary 2.5.1.** If moreover  $\text{st-lim}_{n \rightarrow \infty} m_n = m$ , then by Theorem 2.2, (v) together with the preceding remark  $X_n \xrightarrow{\text{stp}} m$ .

**Example 2.5.1.** (i) The following example shows that if  $\{X_n\}_{n \in \mathbb{N}}$  is a sequence of random variables such that for any  $n$ ,  $X_n$  has a finite mean  $m_n$  and finite standard deviation  $\sigma_n$  then

$$X_n - m_n \xrightarrow{\text{stp}} 0, \quad \text{but} \quad \lim_{n \rightarrow \infty} \sigma_n \neq 0.$$

Let  $X_n = \begin{cases} \pm n & \text{if } n = m^2 \quad \forall m \in \mathbb{N} \text{ with probability } \frac{1}{2}, \\ 0 & \text{if } n \neq m^2 \quad \forall m \in \mathbb{N} \text{ with probability } 1. \end{cases}$

Then  $m_n = E(X_n) = 0 \quad \forall n \in \mathbb{N}$  and

$$\sigma_n = \begin{cases} n & \text{if } n = m^2 \quad \forall m \in \mathbb{N}, \\ 0 & \text{if } n \neq m^2 \quad \forall m \in \mathbb{N} \end{cases}$$

$\Rightarrow \lim_{n \rightarrow \infty} \sigma_n \neq 0$  (in fact it is an unbounded sequence) but  $\text{st-lim}_{n \rightarrow \infty} \sigma_n = 0$ . Then by Tchebycheff's inequality we get

$$P(|X_n - 0| \geq \varepsilon) \leq \frac{\sigma_n^2}{\varepsilon^2} \quad \forall n \in \mathbb{N} \text{ and } \varepsilon > 0.$$

Since  $\text{st-lim}_{n \rightarrow \infty} \sigma_n = 0$ , by Proposition 2.1 we get  $\text{st-lim}_{n \rightarrow \infty} P(|X_n - 0| \geq \varepsilon) = 0$ , i.e.,  $X_n - 0 \xrightarrow{\text{stp}} 0$ .

(ii) Again let

$$X_n = \begin{cases} \pm 1 & \text{if } n = 2^m \quad \forall m \in \mathbb{N} \text{ with probability } \frac{1}{2}, \\ 0 & \text{if } n \neq 2^m \quad \forall m \in \mathbb{N} \text{ with probability } 1. \end{cases}$$

Then  $m_n = E(X_n) = 0 \quad \forall n \in \mathbb{N}$  and

$$\sigma_n = \begin{cases} 1 & \text{if } n = 2^m \quad \forall m \in \mathbb{N}, \\ 0 & \text{if } n \neq 2^m \quad \forall m \in \mathbb{N} \end{cases}$$

$\Rightarrow \lim_{n \rightarrow \infty} \sigma_n \neq 0$  but  $\sigma_n \xrightarrow{\text{stp}} 0$ . Then by Tchebycheff's inequality we get

$$P(|X_n - 0| \geq \varepsilon) \leq \frac{\sigma_n^2}{\varepsilon^2} \quad \forall n \in \mathbb{N} \text{ and } \varepsilon > 0.$$

Since  $\sigma_n \xrightarrow{\text{stp}} 0$ , by Proposition 2.1 we get  $\text{st-}\lim_{n \rightarrow \infty} P(|X_n - 0| \geq \varepsilon) = 0$ , i.e.,  $X_n - 0 \xrightarrow{\text{stp}} 0$ .

**Theorem 2.6.** Let  $g: \mathbb{R} \rightarrow \mathbb{R}$  be a non-decreasing function such that  $g(x) > 0$  for all  $x \in \mathbb{R}$ . If  $\{X_n\}_{n \in \mathbb{N}}$  is a sequence of random variables such that for any  $n$ ,  $X_n$  has a finite mean  $m_n$ , then

$$X_n - m_n \xrightarrow{\text{stp}} 0, \text{ provided } \text{st-}\lim_{n \rightarrow \infty} E\{g(|X_n - m_n|)\} = 0.$$

**Proof.** Proof is easily done by the inequality  $P(|X - m| \geq \varepsilon) \leq E\{g(|X - m|)\} / g(\varepsilon)$ .  $\square$

**Theorem 2.7.** Let  $\{X_n\}_{n \in \mathbb{N}}$  be a sequence of random variables such that  $S_n = X_1 + X_2 + \dots + X_n$  has a finite mean  $M_n$  and a finite variance  $\Sigma_n$  for all  $n$ . Then  $(S_n - M_n)/n \xrightarrow{\text{stp}} 0$  provided  $\text{st-}\lim_{n \rightarrow \infty} \Sigma_n/n^2 = 0$ .

**Proof.** The proof is straightforward and hence omitted.  $\square$

**Theorem 2.8.** Let  $\{X_n\}_{n \in \mathbb{N}}$  be a sequence of identically and independently distributed random variables and  $Y_n = (S_n - E(S_n))/n$ , where  $S_n = X_1 + X_2 + \dots + X_n$ . Then  $Y_n \xrightarrow{\text{stp}} 0$  if and only if  $\text{st-}\lim_{n \rightarrow \infty} E\{Y_n^2/(1 + Y_n^2)\} = 0$ .

**Proof.** Let  $\text{st-}\lim_{n \rightarrow \infty} E\{Y_n^2/(1 + Y_n^2)\} = 0$  and  $\varepsilon > 0$ , then  $|Y_n| \geq \varepsilon \Rightarrow |Y_n|^2 \geq \varepsilon^2 \Rightarrow Y_n^2 + Y_n^2 \varepsilon^2 \geq \varepsilon^2 + Y_n^2 \varepsilon^2 \Rightarrow (Y_n^2/(1 + Y_n^2))/(\varepsilon^2/(1 + \varepsilon^2)) \geq 1$ . This implies

$$P(|Y_n| \geq \varepsilon) \leq P\left\{\frac{Y_n^2/(1 + Y_n^2)}{\varepsilon^2/(1 + \varepsilon^2)}\right\} \leq E\left\{\frac{Y_n^2/(1 + Y_n^2)}{\varepsilon^2/(1 + \varepsilon^2)}\right\} \quad (\text{by Markov's Inequality})$$

$\Rightarrow Y_n \xrightarrow{\text{stp}} 0$ .



Conversely, let us assume that  $X_i$ 's are continuous and let  $Y_n$  have pdf  $f_n(y)$ . Then

$$\begin{aligned}
 E\left\{\frac{Y_n^2}{1+Y_n^2}\right\} &= \int_{-\infty}^{\infty} \frac{y_n^2}{1+y_n^2} f_n(y) \, dy \\
 \Rightarrow E\left\{\frac{Y_n^2}{1+Y_n^2}\right\} &= \int_{|Y_n| \geq \varepsilon} \frac{y_n^2}{1+y_n^2} f_n(y) \, dy + \int_{|Y_n| < \varepsilon} \frac{y_n^2}{1+y_n^2} f_n(y) \, dy \\
 &\leq \int_{|Y_n| \geq \varepsilon} f_n(y) \, dy + \int_{|Y_n| < \varepsilon} y_n^2 f_n(y) \, dy \\
 &\quad \left(\text{since } \frac{y_n^2}{1+y_n^2} < 1 \text{ and } \frac{y_n^2}{1+y_n^2} < y^2\right) \\
 &\leq P(|Y_n| \geq \varepsilon) + \varepsilon^2 \int_{|Y_n| < \varepsilon} f_n(y) \, dy = P(|Y_n| \geq \varepsilon) + \varepsilon^2 P(|Y_n| < \varepsilon) \\
 &\leq P(|Y_n| \geq \varepsilon) + \varepsilon^2 \quad (\text{since } P(|Y_n| < \varepsilon) \leq 1).
 \end{aligned}$$

Since  $Y_n \xrightarrow{\text{st}_p} 0$  and  $\varepsilon^2$  is an arbitrarily small positive real number, we get  $\text{st-}\lim_{n \rightarrow \infty} E\left\{\frac{Y_n^2}{1+Y_n^2}\right\} = 0$ .  $\square$

Note 2.8.1. The result of Theorem 2.8 holds even if  $E(X_i)$  does not exist. In this case we simply define  $Y_n = S_n/n$  rather than  $(S_n - E(S_n))/n$ .

A slightly stronger concept of statistical convergence in probability is defined by statistical convergence in mean of order  $r$ , which is shown by Theorem 3.1.

### 3. STATISTICAL CONVERGENCE IN MEAN OF ORDER $r$

A sequence of random variables  $\{X_n\}_{n \in \mathbb{N}}$  is said to be statistically convergent in the  $r^{\text{th}}$ -mean (where  $r > 0$ ) to a random variable  $X$  (where  $X: S \rightarrow \mathbb{R}$ ) if for any  $\delta > 0$

$$d(\{n \in \mathbb{N}: E(|X_n - X|^r) \geq \delta\}) = 0$$

provided  $E(|X_n|^r)$  exists for every  $n \in \mathbb{N}$  and  $E(|X|^r)$  exists and we write  $\text{st-}\lim_{n \rightarrow \infty} E(|X_n - X|^r) = 0$  or  $X_n \xrightarrow{\text{st}_{\text{rm}}} X$ .

Statistical convergences in mean of orders one and two are called statistical convergence in mean and quadratic mean (or mean square), respectively.

In this section we observe the above form of statistical convergence and a notion of distance, which is widely used in statistics and time series analysis. It is based on the  $L_p$  ( $p \geq 1$ ) metric  $\rho$  which is defined by  $\rho(X, Y) = E\{|X - Y|^p\}^{1/p}$ .

**Theorem 3.1.** Let  $X_n \xrightarrow{\text{st}_{1^m}} X$  (any  $r > 0$ ). Then  $X_n \xrightarrow{\text{st}_p} X$ , i.e., statistical convergence in  $r^{\text{th}}$ -mean implies statistical convergence in probability.

**Proof.** Proof is easily done by Bienayme-Tchebycheff's inequality, i.e.,  $P(|X - m|^r \geq \varepsilon) \leq E\{|X - m|^r\}/\varepsilon^r$ .  $\square$

The following example shows that in general the converse of Theorem 3.1 is not true.

**Example 3.1.1.** We consider the sequence of random variables  $\{X_n\}_{n \in \mathbb{N}}$  defined by

$$X_n \in \{0, n\} \quad \text{with } P(X_n = 0) = 1 - \frac{1}{n^r}$$

and

$$P(X_n = n) = \frac{1}{n^r}, \quad \text{where } r > 0, n \in \mathbb{N}.$$

For any  $\varepsilon > 0$ ,  $P(|X_n - 0| \geq \varepsilon) = P(X_n = n)$  if  $0 < \varepsilon \leq n$  and  $P(|X_n - 0| \geq \varepsilon) = 0$  if  $\varepsilon > n$ , hence for any  $\delta > 0$ ,  $\{n \in \mathbb{N} : P(|X_n - 0| \geq \varepsilon) \geq \delta\} = \text{finite set} \Rightarrow \text{st-lim}_{n \rightarrow \infty} P(|X_n - 0| \geq \varepsilon) = 0$ . But  $E(|X_n|^r) = 1$  for all  $n \in \mathbb{N} \Rightarrow \{n \in \mathbb{N} : E(|X_n - 0|^r) \geq \frac{1}{2}\} = \mathbb{N}$ . This implies  $\{E(|X_n - 0|^r)\}_{n \in \mathbb{N}}$  is not statistically convergent to 0. Hence the result.

**Theorem 3.2.** Let  $\{X_n\}_{n \in \mathbb{N}}$  be a sequence of random variables such that  $P(|X_n| \leq M) = 1$  for all  $n$  and some constant  $M > 0$ . Suppose that  $X_n \xrightarrow{\text{st}_p} X$ . Then  $X_n \xrightarrow{\text{st}_{1^m}} X$  for any  $r > 0$ .

**Proof.** Proof is obvious and hence omitted.  $\square$

**Theorem 3.3.** Let  $X_n \xrightarrow{\text{st}_{1^m}} X$  and  $Y_n \xrightarrow{\text{st}_{1^m}} Y$  (any  $r > 0$ ) be such that  $X_n - X, Y_n - Y \geq 0$ . Then  $X_n + Y_n \xrightarrow{\text{st}_{1^m}} X + Y$ .

**Proof.** Proof is easily done by the inequality  $E(X + Y)^r \leq 2^r[E(X)^r + E(Y)^r]$ , where  $X$  and  $Y$  are non-negative random variables and  $r > 0$ .  $\square$

**Theorem 3.4.** (i)  $X_n \xrightarrow{\text{st}_{1^m}} X \Leftrightarrow \sup_{A \in \Delta} \left| \int_A X_n dP - \int_A X dP \right| \xrightarrow{\text{st}} 0$ .

(ii) If  $X_n \xrightarrow{\text{st}_{1^m}} X$  then  $E(X_n) \xrightarrow{\text{st}} E(X)$ .

**Proof.** Proof is easily done by 'Scheff's lemma'.  $\square$

**Proposition 3.1.** *Let  $\{a_n\}_{n \in \mathbb{N}}$  be a sequence of non negative real numbers such that  $\text{st-lim}_{n \rightarrow \infty} a_n = a$  and  $a \geq 0$ . Then  $\text{st-lim}_{n \rightarrow \infty} (a_n)^q = a^q$ , where  $q \in \mathbb{Q}^+$ .*

*Proof.* Let  $q = m/r \in \mathbb{Q}^+$ . If  $a = 0$  then the result is obvious. So choose  $a > 0$ . It is sufficient to prove that  $\text{st-lim}_{n \rightarrow \infty} a_n = a$  implies  $\text{st-lim}_{n \rightarrow \infty} (a_n)^{1/r} = a^{1/r}$ , where  $r \in \mathbb{N} \setminus \{1\}$ . Then  $\{n \in \mathbb{N} : |a_n - a| < \frac{1}{2}a\} \subseteq \{n \in \mathbb{N} : a_n > \frac{1}{2}a\} = M$  (say, then  $d(M) = 1$ ). If  $n \in M$  then

$$\begin{aligned} |a_n - a| &= |(a_n)^{1/r} - a^{1/r}| |\{(a_n)^{(r-1)/r} + (a_n)^{(r-2)/r} a^{1/r} \\ &\quad + \dots + (a_n)^{1/r} a^{(r-2)/r} + a^{(r-1)/r}\}| \\ &\geq L |(a_n)^{1/r} - a^{1/r}|, \text{ where } L = \frac{a^{(r-1)/r}}{2(1 - 1/\sqrt[r]{2})}, \end{aligned}$$

i.e.,  $M \subseteq \{n \in \mathbb{N} : 0 \leq |(a_n)^{1/r} - a^{1/r}| \leq L^{-1}|a_n - a|\}$ . Then by Proposition 2.1,  $\text{st-lim}_{n \rightarrow \infty} \sqrt[r]{a_n} = \sqrt[r]{a}$ . □

**Theorem 3.5.** *Let  $\{X_n\}_{n \in \mathbb{N}}$  be a sequence of random variables such that  $X_n \xrightarrow{\text{st}_{2^n}} X$ . Then  $\text{st-lim}_{n \rightarrow \infty} E(X_n) = E(X)$  and  $\text{st-lim}_{n \rightarrow \infty} E(X_n^2) = E(X^2)$ .*

*Proof.* The proof is parallel to that of Theorem 8 in [18] (and using the result of the Proposition 3.1) and so it is omitted. □

**Theorem 3.6.** *Let  $\{X_n\}_{n \in \mathbb{N}}$  be a sequence of random variables and  $r \in \mathbb{Q}^+$  such that  $X_n \xrightarrow{\text{st}_{\text{rm}}} X$ . Then  $\text{st-lim}_{n \rightarrow \infty} E(|X_n|^s) = E(|X|^s)$  for  $s \leq r$  and  $s \in \mathbb{Q}^+$ .*

*Proof.* The proof is parallel to that of Theorem 9, 10 in [18] (and using the result of the Proposition 3.1) and so it is omitted. □

#### 4. STATISTICAL CONVERGENCE IN DISTRIBUTION

Let  $\{X_n\}_{n \in \mathbb{N}}$  be a sequence of random variables, where  $F_n(x)$  is the distribution function of  $X_n$  for  $n \in \mathbb{N}$ . If there exists a random variable  $X$  whose distribution function is  $F(x)$  such that  $\text{st-lim}_{n \rightarrow \infty} F_n(x) = F(x)$  at every point of continuity  $x$  of  $F(x)$ , then  $\{X_n\}_{n \in \mathbb{N}}$  is said to be statistically convergent in distribution to  $X$  and we write  $X_n \xrightarrow{\text{st}_d} X$ .

**Theorem 4.1.** Let  $\{X_n\}_{n \in \mathbb{N}}$  be a sequence of random variables. Also, let  $f_n(x_i) = P(X_n = x_i)$ , for all  $i \in \mathbb{N}$ , be the probability mass function of  $X_n$ , for all  $n \in \mathbb{N}$ , and  $f(x_i) = P(X = x_i)$ , for all  $i \in \mathbb{N}$  be the probability mass function of  $X$ . Then

$$f_n(x) \xrightarrow{\text{st}} f(x) \text{ for all } x \Leftrightarrow X_n \xrightarrow{\text{st}_d} X.$$

**Proof.** Proof is straightforward and hence omitted.  $\square$

**Proposition 4.1.** Let  $\{a_n\}_{n \in \mathbb{N}}$ ,  $\{b_n\}_{n \in \mathbb{N}}$  be two sequences of real numbers such that  $a_n \leq b_n$  for all  $n \in \mathbb{N}$ . Then

$$\text{st-}\underline{\lim} a_n \leq \text{st-}\underline{\lim} b_n \text{ and } \text{st-}\overline{\lim} a_n \leq \text{st-}\overline{\lim} b_n.$$

**Proof.** The proof is straightforward and hence omitted.  $\square$

**Theorem 4.2.** Let  $X_n \xrightarrow{\text{st}_p} X$ . Then  $X_n \xrightarrow{\text{st}_d} X$ , i.e., statistical convergence in probability implies statistical convergence in distribution.

**Proof.** Let  $F_n(x)$  and  $F(x)$  be the distribution functions of  $X_n$  and  $X$ , respectively. Now for any two real numbers  $x$  and  $y$  with  $x < y$ , we have

$$(X \leq x) = (X_n \leq y, X \leq x) + (X_n > y, X \leq x).$$

Since  $(X_n \leq y, X \leq x) \subseteq (X_n \leq y)$ , we have

$$(X \leq x) \subseteq (X_n \leq y) + (X_n > y, X \leq x).$$

Therefore,

$$\begin{aligned} (1) \quad P(X \leq x) &\leq P\{(X_n \leq y) + (X_n > y, X \leq x)\} \\ &\leq P(X_n \leq y) + P(X_n > y, X \leq x) \\ &\Rightarrow F_n(y) \geq F(x) - P(X_n > y, X \leq x) \end{aligned}$$

Now if  $X_n > y, X \leq x$  occur simultaneously, then  $X_n > y, -X \geq -x$  and so  $X_n - X > y - x$ , i.e.,  $(X_n > y, X \leq x) \subseteq (X_n - X > y - x) \subseteq (|X_n - X| > y - x)$  which implies

$$P(X_n > y, X \leq x) \leq P(|X_n - X| > y - x).$$

Since  $x < y$  and  $X_n \xrightarrow{\text{st}_p} X$ , we get

$$\text{st-}\lim_{n \rightarrow \infty} P(X_n > y, X \leq x) = 0.$$

Now from (1) we get  $\text{st-}\underline{\lim} F_n(y) \geq F(x)$  (by the above Proposition 4.1).

Similarly, if  $y < z$  then

$$(X_n \leq y) = (X \leq z, X_n \leq y) + (X > z, X_n \leq y),$$

then  $F_n(y) \leq F(z) + P(X > z, X_n \leq y)$  and  $\text{st-lim}_{n \rightarrow \infty} P(X > z, X_n \leq y) = 0$ .

Finally, we get  $\text{st-lim}_{n \rightarrow \infty} F_n(y) \leq F(z)$ .

Thus, for  $x < y < z$  we have

$$F(x) \leq \text{st-lim}_{n \rightarrow \infty} F_n(y) \leq \text{st-lim}_{n \rightarrow \infty} F_n(y) \leq F(z).$$

If  $F$  is continuous at  $y$  then

$$F(y) = \lim_{x \rightarrow y^-} F(x) \leq \text{st-lim}_{n \rightarrow \infty} F_n(y) \leq \text{st-lim}_{n \rightarrow \infty} F_n(y) \leq \lim_{z \rightarrow y^+} F(z) = F(y).$$

This implies  $\text{st-lim}_{n \rightarrow \infty} F_n(y) = \text{st-lim}_{n \rightarrow \infty} F_n(y) = F(y)$ , i.e.,  $X_n \xrightarrow{\text{st-d}} X$ . Hence the result.  $\square$

**Note 4.2.1.** The converse of Theorem 4.2 is not true in general, i.e.,  $X_n \xrightarrow{I_d} X$  does not imply  $X_n \xrightarrow{I_p} X$  in general. For, let us consider random variables  $X, X_1, X_2, \dots$  having identical distribution. Let the spectrum of the two dimensional random variable  $(X_n, X)$  be  $(0, 0), (0, 1), (1, 0), (1, 1)$  and

$$\begin{aligned} P(X_n = 0, X = 0) &= 0 = P(X_n = 1, X = 1), \\ P(X_n = 0, X = 1) &= \frac{1}{2} = P(X_n = 1, X = 0). \end{aligned}$$

Hence, the marginal distribution of  $X_n$  is given by  $X_n = i$  ( $i = 0, 1$ ), with p.m.f  $p_{x_n i} = P(X_n = i)$ , where  $p_{x_n 0} = \frac{1}{2} = p_{x_n 1}$  and that of  $X = j$  ( $j = 0, 1$ ), with p.m.f  $p_{x j} = P(X = j)$ , where  $p_{x 0} = \frac{1}{2} = p_{x 1}$ .

If  $F_n(x)$  and  $F(x)$  are the distribution functions of  $X_n$  and  $X$ , respectively, then

$$F(x) = F_n(x) = \begin{cases} 0, & x < 0, \\ \frac{1}{2}, & 0 \leq x < 1, \\ 1, & x \geq 1. \end{cases}$$

Therefore,  $\text{st-lim}_{n \rightarrow \infty} F_n(x) = F(x)$  for all  $x \in \mathbb{R}$ , i.e.,  $X_n \xrightarrow{\text{st-d}} X$  as  $n \rightarrow \infty$ . But  $P(|X_n - X| \geq \frac{1}{2}) = P(|X_n - X| = 1) = P(X_n = 0, X = 1) + P(X_n = 1, X = 0) = 1$ . Therefore,  $\text{st-lim}_{n \rightarrow \infty} P(|X_n - X| \geq \frac{1}{2}) \neq 0$ . Then  $\text{st-lim}_{n \rightarrow \infty} P(|X_n - X| \geq \varepsilon) \neq 0$ . Hence, the result follows.

**Theorem 4.3.** Let  $\{X_n\}_{n \in \mathbb{N}}$  be a sequence of random variables such that  $X_n \xrightarrow{\text{st}_d} X$ . Let  $\{\zeta_n\}_{n \in \mathbb{N}}$  be a sequence of positive constants such that  $\zeta_n \xrightarrow{\text{st}} 0$ . Then  $\zeta_n X_n \xrightarrow{\text{st}_p} 0$ .

*Proof.* The proof is straightforward and hence omitted. □

The following result is easy to establish.

**Theorem 4.4.** Let  $\{X_n\}_{n \in \mathbb{N}}$  be a sequence of random variables such that  $X_n \xrightarrow{\text{st}_d} X$ , and  $c$  a constant. Then

- (a)  $X_n + c \xrightarrow{\text{st}_d} X + c$ , and
- (b)  $cX_n \xrightarrow{\text{st}_d} cX$ ,  $c \neq 0$ .

**Theorem 4.5.** Let  $c$  be a constant, then  $X_n \xrightarrow{\text{st}_d} c \Rightarrow X_n \xrightarrow{\text{st}_p} c$ .

*Proof.* The proof is straightforward and hence omitted. □

**Corollary 4.5.1.** Let  $c$  be a constant, then  $X_n \xrightarrow{\text{st}_d} c \Leftrightarrow X_n \xrightarrow{\text{st}_p} c$ .

**Theorem 4.6.** If  $\{X_n\}_{n \in \mathbb{N}}$  and  $\{Y_n\}_{n \in \mathbb{N}}$  are sequences of random variables on some probability space with  $X_n - Y_n \xrightarrow{\text{st}_p} 0$ , and  $Y_n \xrightarrow{\text{st}_d} X$ , then  $X_n \xrightarrow{\text{st}_d} X$ .

*Proof.* Let  $x, x \pm \varepsilon$  be a point of continuity of the distribution function  $F$  corresponding to the random variable  $X$ , where  $\varepsilon > 0$ . Then  $P(X_n \leq x) = P(Y_n \leq x + Y_n - X_n) = P(Y_n \leq x + Y_n - X_n; Y_n - X_n \leq \varepsilon) + P(Y_n \leq x + Y_n - X_n; Y_n - X_n > \varepsilon) \leq P(Y_n \leq x + \varepsilon) + P(Y_n - X_n > \varepsilon)$ . This implies  $\text{st-}\overline{\lim} F_n(x) \leq F(x + \varepsilon)$  and analogously  $F(x - \varepsilon) \leq \text{st-}\underline{\lim} F_n(x)$ . Since  $\varepsilon$  is arbitrary, so  $F(x) = \text{st-lim} F_n(x)$ . □

**Theorem 4.7.** If  $\{X_n\}_{n \in \mathbb{N}}$  and  $\{Y_n\}_{n \in \mathbb{N}}$  are sequences of random variables on some probability space and  $c$  is a constant, then

- (a)  $X_n \xrightarrow{\text{st}_d} X, Y_n \xrightarrow{\text{st}_p} c \Rightarrow X_n + Y_n \xrightarrow{\text{st}_d} X + c$ ,
- (b)  $X_n \xrightarrow{\text{st}_d} X, Y_n \xrightarrow{\text{st}_p} 0 \Rightarrow X_n Y_n \xrightarrow{\text{st}_p} 0$ ,
- (c)  $X_n \xrightarrow{\text{st}_d} X, Y_n \xrightarrow{\text{st}_p} c \Rightarrow X_n Y_n \xrightarrow{\text{st}_d} cX$  if  $c \neq 0$ ,
- (d)  $X_n \xrightarrow{\text{st}_d} X, Y_n \xrightarrow{\text{st}_p} c \Rightarrow X_n / Y_n \xrightarrow{\text{st}_d} X/c$  if  $c \neq 0$ .

*Proof.* (a) Proof is parallel to that of Theorem 15 (Slutsky's theorem) in [18] and so omitted.

(b) For any  $\delta > 0$ , choose  $\pm\alpha \in$  set of points of continuity of the distribution function  $F$  of  $X$  such that  $F(\alpha) - F(-\alpha) \geq 1 - \delta$ . Any  $\varepsilon > 0$ ,  $P(|X_n Y_n| \geq \varepsilon) = P(|X_n Y_n| \geq \varepsilon, |Y_n| < \varepsilon/\alpha) + P(|X_n Y_n| \geq \varepsilon, |Y_n| \geq \varepsilon/\alpha) \leq P(|X_n| > \alpha) + P(|Y_n| \geq \varepsilon/\alpha)$ . So  $\text{st-}\overline{\lim} P(|X_n Y_n| \geq \varepsilon) < \delta$ . This implies  $X_n Y_n \xrightarrow{\text{st}_p} 0$ .

Parts (c) and (d) are obvious and hence omitted. □

Applications. (i) Much of classical probability theory and its applications to statistics concerns limit theorems; i.e., the asymptotic behaviour of a sequence of random variables. The sequence could consist of sample averages, cumulative sums, extremes, sample quantiles, sample correlations and so on. Whereas probability theory discusses limit theorems, the theory of statistics is concerned with large sample properties of statistics, where a statistic is just a function of the sample.

(ii) A classical significant statistical application of ‘statistical convergence in probability’ is to quantile estimation and geometric probability (by using the theorem Weak laws of large numbers).

(iii) Classical application of ‘statistical convergence in mean of order  $r$ ’ is to time series analysis.

(iv) In statistical estimation the delta method allows us to take a basic ‘statistical convergence in distribution’, for instance to a limiting normal distribution, apply smooth functions and conclude that the functions are asymptotically normal as well.

#### References

- [1] *M. Balcerzak, K. Dems, A. KomisarSKI*: Statistical convergence and ideal convergence for sequences of functions. *J. Math. Anal. Appl.* *328* (2007), 715–729.
- [2] *L. A. Bunimovich, Ya. G. Sinai*: Statistical properties of Lorentz gas with periodic configuration of scatterers. *Commun. Math. Phys.* *78* (1981), 479–497.
- [3] *P. Erdős, G. Tenenbaum*: On densities of certain sequences of integers. *Proc. Lond. Math. Soc., III. Ser.* *59* (1989), 417–438. (In French.)
- [4] *H. Fast*: Sur la convergence statistique. *Colloq. Math.* *2* (1951), 241–244. (In French.)
- [5] *J. A. Fridy*: On statistical convergence. *Analysis* *5* (1985), 301–313.
- [6] *J. A. Fridy*: Statistical limit points. *Proc. Am. Math. Soc.* *118* (1993), 1187–1192.
- [7] *J. A. Fridy, M. K. Khan*: Tauberian theorems via statistical convergence. *J. Math. Anal. Appl.* *228* (1998), 73–95.
- [8] *J. A. Fridy, C. Orhan*: Statistical limit superior and limit inferior. *Proc. Am. Math. Soc.* *125* (1997), 3625–3631.
- [9] *A. D. Gadjiev, C. Orhan*: Some approximation theorems via statistical convergence. *Rocky Mt. J. Math.* *32* (2002), 129–138.
- [10] *M. Katětov*: Products of filters. *Commentat. Math. Univ. Carol.* *9* (1968), 173–189.
- [11] *E. Kolk*: The statistical convergence in Banach spaces. *Tartu ÜI. Toimetised* *928* (1991), 41–52.
- [12] *P. Kostyrko, M. Macaj, T. Šalát, O. Strauch*: On statistical limit points. *Proc. Am. Math. Soc.* *129* (2001), 2647–2654.
- [13] *I. J. Maddox*: Statistical convergence in a locally convex space. *Math. Proc. Camb. Philos. Soc.* *104* (1988), 141–145.
- [14] *V. G. Martínez, G. S. Torrubia, C. T. Blanc*: A statistical convergence application for the Hopfield networks. *Information Theory and Applications* *15* (2008), 84–88.
- [15] *H. I. Miller*: A measure theoretical subsequence characterization of statistical convergence. *Trans. Am. Math. Soc.* *347* (1995), 1811–1819.
- [16] *S. Pehlivan, M. A. Mamedov*: Statistical cluster points and turnpike. *Optimization* *48* (2000), 93–106.

- [17] *M. D. Penrose, J. E. Yukich*: Weak laws of large numbers in geometric probability. *Ann. Appl. Probab.* 13 (2003), 277–303.
- [18] *V. K. Rohatgi*: *An Introduction to Probability Theory and Mathematical Statistics*. Wiley Series in Probability and Mathematical Statistics, John Wiley & Sons, New York, 1976.
- [19] *T. Šalát*: On statistically convergent sequences of real numbers. *Math. Slovaca* 30 (1980), 139–150.
- [20] *E. Savaş*: On statistically convergent sequences of fuzzy numbers. *Inf. Sci.* 137 (2001), 277–282.
- [21] *I. J. Schoenberg*: The integrability of certain functions and related summability methods. *Am. Math. Mon.* 66 (1959), 361–375.
- [22] *H. Steinhaus*: Sur la convergence ordinaire et la convergence asymptotique. *Colloq. Math.* 2 (1951), 73–74.
- [23] *A. Zygmund*: *Trigonometric Series*. Cambridge University Press, 1979.

*Author's address: Sanjoy Ghosal, Faculty of Science, Department of Mathematics, Kalyani Government Engineering College, Kalyani, Nadia-741235, West Bengal, India, e-mail: sanjoykrghosal@yahoo.co.in.*