

Jiantao Li; Xiankun Du

A note on the kernels of higher derivations

Czechoslovak Mathematical Journal, Vol. 63 (2013), No. 3, 583–588

Persistent URL: <http://dml.cz/dmlcz/143478>

Terms of use:

© Institute of Mathematics AS CR, 2013

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A NOTE ON THE KERNELS OF HIGHER DERIVATIONS

JIANTAO LI, XIANKUN DU, Changchun

(Received February 21, 2012)

Abstract. Let $k \subseteq k'$ be a field extension. We give relations between the kernels of higher derivations on $k[X]$ and $k'[X]$, where $k[X] := k[x_1, \dots, x_n]$ denotes the polynomial ring in n variables over the field k . More precisely, let $D = \{D_n\}_{n=0}^{\infty}$ a higher k -derivation on $k[X]$ and $D' = \{D'_n\}_{n=0}^{\infty}$ a higher k' -derivation on $k'[X]$ such that $D'_m(x_i) = D_m(x_i)$ for all $m \geq 0$ and $i = 1, 2, \dots, n$. Then (1) $k[X]^D = k$ if and only if $k'[X]^{D'} = k'$; (2) $k[X]^D$ is a finitely generated k -algebra if and only if $k'[X]^{D'}$ is a finitely generated k' -algebra. Furthermore, we also show that the kernel $k[X]^D$ of a higher derivation D of $k[X]$ can be generated by a set of closed polynomials.

Keywords: higher derivation; field extension; closed polynomial

MSC 2010: 13A50

1. INTRODUCTION

Throughout this paper, $k[X] = k[x_1, x_2, \dots, x_n]$ is the polynomial ring in n variables over a field k .

Let R be a commutative ring and A an R -algebra. Recall that a set of R -linear endomorphisms $D = \{D_n\}_{n=0}^{\infty}$ is called a *higher R -derivation* on A if D satisfies the following conditions:

- (1) D_0 is the identity map of A ;
- (2) $D_n(ab) = \sum_{i+j=n} D_i(a)D_j(b)$ for all $n \geq 0$ and for any $a, b \in A$.

A higher R -derivation D is called locally finite if for any $a \in A$, there exists $n \in \mathbb{N}$ such that $D_m(a) = 0$ for any integer $m \geq n$. And D is called iterative if for any

This research was supported by NSF of China (No. 11071097, No. 11101176) and “211 Project” and “985 Project” of Jilin University.

$i, j \in \mathbb{N}$, $D_i \circ D_j = \binom{i+j}{i} D_{i+j}$. For a higher R -derivation D on A , A^D is the kernel of D defined by $A^D = \bigcap_{n \geq 1} \text{Ker} D_n$.

Higher derivations and their kernels play an important role when we deal with some curves and affine surfaces. For example, a G_a -action on an affine scheme $X = \text{Spec}(A)$ can be interpreted in terms of a locally finite iterative higher derivation on A , and many things become easier to treat by observing the locally finite iterative higher derivation on A associated with the G_a -action [4]. Recently, in [2], [7], [8] the kernels of higher derivations have been studied. In [3], the structure of higher derivations is studied. In this short note, we prove the following results.

(1) Let $k \subseteq k'$ be a field extension. Let $D = \{D_n\}_{n=0}^\infty$ be a higher k -derivation on the polynomial ring $k[X]$ over a field k and let $D' = \{D'_n\}_{n=0}^\infty$ be a higher k' -derivation on $k'[X]$ such that $D'_m(x_i) = D_m(x_i)$ for all $m \geq 0$ and $i = 1, 2, \dots, n$. Then

- (i) $k[X]^D = k$ if and only if $k'[X]^{D'} = k'$;
- (ii) $k[X]^D$ is a finitely generated k -algebra if and only if $k'[X]^{D'}$ is a finitely generated k' -algebra.

(2) Let $D = \{D_n\}_{n=0}^\infty$ be a higher k -derivation of $k[X]$. Then there exists a set S of closed polynomials in $k[X]$ such that $k[X]^D = k[S]$.

In the case of derivations on $k[X]$, similar results can be found in [5].

2. HIGHER DERIVATIONS UNDER EXTENSION OF FIELDS

Let $k \subseteq k'$ be a field extension and A a k -algebra. Let $D = \{D_n\}_{n=0}^\infty$ be a higher k -derivation on A . Consider the tensor product $A \otimes_k k'$, then $A \otimes_k k'$ is a k' -algebra. Consider the set of k' -linear mappings $D \otimes 1$ of $A \otimes_k k'$ given by

$$D \otimes 1 = \{D_n \otimes 1: A \otimes_k k' \rightarrow A \otimes_k k'\}_{n=0}^\infty.$$

Proposition 2.1. *The set of k' -linear mappings $D \otimes 1$ is a higher k' -derivation on the k' -algebra $A \otimes_k k'$. Moreover, if D is locally finite or iterative, then $D \otimes 1$ is also locally finite or iterative, respectively.*

Proof. It is obvious that $D_0 \otimes 1$ is the identity map of $A \otimes_k k'$.

For any $a \otimes p, b \otimes q \in A \otimes_k k'$,

$$\begin{aligned} (D_n \otimes 1)((a \otimes p) \cdot (b \otimes q)) &= (D_n \otimes 1)(ab \otimes pq) = D_n(ab) \otimes pq \\ &= \left(\sum_{i+j=n} D_i(a)D_j(b) \right) \otimes pq = \sum_{i+j=n} D_i(a)D_j(b) \otimes pq \\ &= \sum_{i+j=n} ((D_i(a) \otimes p) \cdot (D_j(b) \otimes q)) \\ &= \sum_{i+j=n} (D_i \otimes 1)(a \otimes p) \cdot (D_j \otimes 1)(b \otimes q). \end{aligned}$$

Since $D_i \otimes 1$ are k' -linear mappings of $A \otimes_k k'$, $i = 0, 1, 2, \dots$, it follows from the above equation that $(D_n \otimes 1)(xy) = \sum_{i+j=n} (D_i \otimes 1)(x) \cdot (D_j \otimes 1)(y)$ for any $x = \sum_i a_i \otimes p_i$, $y = \sum_j b_j \otimes q_j \in A \otimes_k k'$, where $a_i, b_j \in A$ and $p_i, q_j \in k'$. Thus, $D \otimes 1$ is a higher k' -derivation on $A \otimes_k k'$.

Moreover, D is locally finite, that is, for any $a \in A$ there exists $n \in \mathbb{N}$ such that $D_m(a) = 0$ for any $m \geq n$. Define $\nu_D(a) = n$ where n is the least integer such that $D_{n-1}(a) \neq 0$ and $D_m(a) = 0$ for any $m \geq n$. Note that any element of $A \otimes_k k'$ is a finite sum of the form $\sum_i a_i \otimes p_i$, where $a_i \in A$, $p_i \in k'$. Set $N = \max_i \{\nu_D(a_i)\}$. Then for any $m \geq N$,

$$(D_m \otimes 1) \left(\sum_i a_i \otimes p_i \right) = \sum_i (D_m \otimes 1)(a_i \otimes p_i) = \sum_i D_m(a_i) \otimes p_i = \sum_i 0 \otimes p_i = 0.$$

Thus, $D \otimes 1$ is a locally finite higher k' -derivation.

If D is iterative, that is, $D_i \circ D_j = \binom{i+j}{i} D_{i+j}$ for any $i, j \in \mathbb{N}$, then

$$(D_i \otimes 1) \circ (D_j \otimes 1) = (D_i \circ D_j) \otimes 1 = \left(\binom{i+j}{i} D_{i+j} \right) \otimes 1 = \binom{i+j}{i} D_{i+j} \otimes 1.$$

Thus, $D \otimes 1$ is an iterative higher k' -derivation. □

The kernel of $D \otimes 1$ is closely related to the kernel of D .

Proposition 2.2. *The k' -algebras $A^D \otimes_k k'$ and $(A \otimes_k k')^{D \otimes 1}$ are isomorphic.*

Proof. It is obvious that k' is flat as a k -algebra. Then the exact sequences

$$0 \longrightarrow \text{Ker} D_i \xrightarrow{\text{id}} A \xrightarrow{D_i} A, \quad i = 1, 2, \dots$$

of k -vector spaces induce the following exact sequences of k' -vector spaces:

$$0 \longrightarrow \text{Ker} D_i \otimes_k k' \xrightarrow{\text{id} \otimes 1} A \otimes_k k' \xrightarrow{D_i \otimes 1} A \otimes_k k', \quad i = 1, 2, \dots$$

Thus,

$$\text{Ker}D_i \otimes_k k' \cong \text{Im id} \otimes 1 \cong \text{Ker}D_i \otimes 1.$$

Note that if U, V are vector spaces and U_1, U_2 are subspaces of U , then it is known (see for instance [6, Chapter 14: Tensor Products]) that

$$(U_1 \otimes V) \cap (U_2 \otimes V) \cong (U_1 \cap U_2) \otimes V.$$

Therefore,

$$\begin{aligned} (A \otimes_k k')^{D \otimes 1} &= \bigcap_{i \geq 1} \text{Ker}D_i \otimes 1 \cong \bigcap_{i \geq 1} \text{Ker}D_i \otimes_k k' \cong \left(\bigcap_{i \geq 1} \text{Ker}D_i \right) \otimes_k k' \\ &= A^D \otimes_k k'. \end{aligned}$$

□

Lemma 2.3 ([5]). *A is a finitely generated algebra over k if and only if $A \otimes_k k'$ is a finitely generated algebra over k' .*

As a direct corollary of Proposition 2.2 and Lemma 2.3, we have

Corollary 2.4. *Let $k \subseteq k'$ be a field extension and let $D = \{D_n\}_{n=0}^\infty$ and $D \otimes 1 = \{D_n \otimes 1\}_{n=0}^\infty$ be higher derivations on A and $A \otimes_k k'$, respectively. Then*

- (1) $A^D = k$ if and only if $(A \otimes_k k')^{D \otimes 1} = k'$;
- (2) A^D is a finitely generated k -algebra if and only if $(A \otimes_k k')^{D \otimes 1}$ is a finitely generated k' -algebra.

In most cases, we deal with higher derivations on polynomial rings. In that case, we can get the following theorem.

Theorem 2.5. *Let $D = \{D_n\}_{n=0}^\infty$ be a higher k -derivation on the polynomial ring $k[X]$ over a field k . Let $k \subseteq k'$ be a field extension and $D' = \{D'_n\}_{n=0}^\infty$ a higher k' -derivation on $k'[X]$ such that $D'_m(x_i) = D_m(x_i)$ for all $m \geq 0$ and $i = 1, 2, \dots, n$. Then*

- (1) $k[X]^D = k$ if and only if $k'[X]^{D'} = k'$;
- (2) $k[X]^D$ is a finitely generated k -algebra if and only if $k'[X]^{D'}$ is a finitely generated k' -algebra.

3. KERNELS OF HIGHER k -DERIVATIONS ON $k[X]$

Recall that a polynomial $f \in k[X] \setminus k$ is called closed if the subalgebra $k[f]$ is a maximal element in the family \mathcal{L} of subalgebras defined by $\mathcal{L} = \{k[f] : f \in k[X] \setminus k\}$. It is known that f is a closed polynomial if and only if $k[f]$ is integrally closed in $k[X]$. Define a generative polynomial h_f of a polynomial $f \in k[X] \setminus k$ as a closed polynomial such that $f = F(h_f)$ for some $F \in k[t]$. Such a generative polynomial h_f is unique up to affine transformations [1]. In [5], it is shown that if d is a derivation of $k[X]$ and $f \in k[X]^d$ is a nonconstant polynomial, then the generative polynomial h_f of f also belongs to $k[X]^d$, and hence, there is a set S of closed polynomials such that $k[X]^d = k[S]$. Next we generalize such results to the case of higher derivations.

For a higher R -derivation $D = \{D_n\}_{n=0}^\infty$ on an R -algebra A we can define a mapping φ_D from A to $A[[t]]$ (here, $A[[t]]$ is the formal power series ring in one variable t over A) associated with D by

$$\varphi_D(a) = \sum_{i \geq 0} D_i(a)t^i \quad \text{for any } a \in A.$$

Lemma 3.1 ([2]). *Let $D = \{D_n\}_{n=0}^\infty$ be a set of R -endomorphisms of the R -algebra A where D_0 is the identity map. Then the following statements are equivalent.*

- (1) D is a higher R -derivation on A ;
- (2) φ_D is a homomorphism of the R -algebras A and $A[[t]]$.

Moreover, D is a locally finite higher R -derivation on A if and only if φ_D is a homomorphism of the R -algebras A and $A[[t]]$.

It is clear that for any $a \in A$, $a \in A^D$ if and only if $\varphi_D(a) = a$.

We are now in a position to show the main result of this section.

Theorem 3.2. *Let $D = \{D_n\}_{n=0}^\infty$ be a higher k -derivation of $k[X]$. Then there exists a set S of closed polynomials in $k[X]$ such that $k[X]^D = k[S]$.*

Proof. If $k[X]^D = k$, then put $S = \emptyset$. Now assume that $k[X]^D \neq k$. Choose $f \in k[X]^D \setminus k$ and let h_f be the generative polynomial of f . Then h_f is a closed polynomial and there exists $F(x) = a_n x^n + \dots + a_1 x + a_0 \in k[x]$ such that $f = F(h_f)$, where $a_i \in k$ and $a_n \neq 0$, and we can assume that F has the minimal degree. Thus,

$$f = a_n h_f^n + \dots + a_1 h_f + a_0.$$

Applying φ_D to the above equation, we obtain

$$\varphi_D(f) = a_n \varphi_D(h_f)^n + \dots + a_1 \varphi_D(h_f) + a_0.$$

Since $f \in k[X]^D$, it follows that $\varphi_D(f) = f$. Thus,

$$f = a_n \varphi_D(h_f)^n + \dots + a_1 \varphi_D(h_f) + a_0.$$

If $\varphi_D(h_f) \neq h_f$, then $\varphi_D(h_f) = \sum_{i \geq 0} D_i(h_f)t^i$ is algebraically independent over $k[X]$, even over the field of rational functions $k(X)$. Hence $\varphi_D(h_f)$ does not satisfy a non-trivial algebraic equation over $k[X]$. This contradiction forces $\varphi_D(h_f) = h_f$. Thus, $h_f \in k[X]^D$. Therefore, it is proved that if $f \in k[X]^d$, then the generative polynomial h_f of f also belongs to $k[X]^d$.

Set $S = \{h_f : f \in k[X]^D \setminus k\}$. Then S is clearly a set of closed polynomials in $k[X]$ and $k[X]^D = k[S]$. \square

References

- [1] *I. V. Arzhantsev, A. P. Petravchuk*: Closed polynomials and saturated subalgebras of polynomial algebras. *Ukr. Math. J.* 59 (2007), 1783–1790.
- [2] *H. Kojima, N. Wada*: Kernels of higher derivations in $R[x, y]$. *Commun. Algebra* 39 (2011), 1577–1582.
- [3] *M. Mirzavaziri*: Characterization of higher derivations on algebras. *Commun. Algebra* 38 (2010), 981–987.
- [4] *M. Miyanishi*: Lectures on Curves on Rational and Unirational Surfaces. Tata Institute of Fundamental Research Lectures on Mathematics and Physics, Berlin, Springer, 1978.
- [5] *A. Nowicki*: Polynomial Derivations and their Rings of Constants. N. Copernicus Univ. Press, Toruń, 1994.
- [6] *S. Roman*: Advanced Linear Algebra. 3rd edition, Graduate Texts in Mathematics 135, New York, Springer, 2008.
- [7] *R. Tanimoto*: An algorithm for computing the kernel of a locally finite iterative higher derivation. *J. Pure Appl. Algebra* 212 (2008), 2284–2297.
- [8] *N. Wada*: Some results on the kernels of higher derivations on $k[x, y]$ and $k(x, y)$. *Colloq. Math.* 122 (2011), 185–189.

Authors' address: J i a n t a o L i, X i a n k u n D u, School of Mathematics, Jilin University, 130012, Changchun, China, e-mail: jtlimath@gmail.com, duxk@jlu.edu.cn.