

Guanghai Cheng

New bounds for the minimum eigenvalue of the Fan product of two M -matrices

Czechoslovak Mathematical Journal, Vol. 64 (2014), No. 1, 63–68

Persistent URL: <http://dml.cz/dmlcz/143949>

Terms of use:

© Institute of Mathematics AS CR, 2014

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

NEW BOUNDS FOR THE MINIMUM EIGENVALUE OF THE
FAN PRODUCT OF TWO M -MATRICES

GUANGHUI CHENG, Chengdu

(Received September 17, 2012)

Abstract. In this paper, we mainly use the properties of the minimum eigenvalue of the Fan product of M -matrices and Cauchy-Schwarz inequality, and propose some new bounds for the minimum eigenvalue of the Fan product of two M -matrices. These results involve the maximum absolute value of off-diagonal entries of each row. Hence, the lower bounds for the minimum eigenvalue are easily calculated in the practical examples. In theory, a comparison is given in this paper. Finally, to illustrate our results, a simple example is also considered.

Keywords: Fan product; minimum eigenvalue; M -matrix

MSC 2010: 15A18, 15A42

1. INTRODUCTION

For convenience, the set $\{1, 2, \dots, n\}$ is denoted by \mathbb{N} , where n is any positive integer. A matrix $A = (a_{ij}) \in \mathbb{R}^{n \times n}$ is called a nonnegative (positive) matrix if $a_{ij} \geq 0$ ($a_{ij} > 0$). A matrix $A \in \mathbb{R}^{n \times n}$ is called a nonsingular M -matrix [1] if there exists $P \geq 0$ and $\alpha > 0$ such that

$$A = \alpha I - P \quad \text{and} \quad \alpha > \varrho(P),$$

where $\varrho(P)$ is the spectral radius (Perron root) of the nonnegative matrix P and I is the $n \times n$ identity matrix. Denote by \mathcal{M}_n the set of all $n \times n$ nonsingular M -matrices. Denote

$$\tau(A) = \min\{\operatorname{Re} \lambda : \lambda \in \sigma(A)\},$$

where $\sigma(A)$ denotes the spectrum of A . If $A \in \mathcal{M}_n$, then

$$\tau(A) = \frac{1}{\varrho(A^{-1})}$$

is a positive real eigenvalue, and the corresponding eigenvector is nonnegative [3].

A matrix A is irreducible if there does not exist a permutation matrix P such that

$$PAP^T = \begin{bmatrix} A_{1,1} & A_{1,2} \\ 0 & A_{2,2} \end{bmatrix},$$

where $A_{1,1}$ and $A_{2,2}$ are square matrices.

Let $A, B \in \mathbb{C}^{n \times n}$. The Fan product of A and B is denoted by $A \star B \equiv C = (c_{ij}) \in \mathbb{C}^{n \times n}$ and is defined by

$$c_{ij} = \begin{cases} -a_{ij}b_{ij}, & i \neq j, \\ a_{ii}b_{ii}, & i = j. \end{cases}$$

If $A, B \in \mathcal{M}_n$, then $A \star B$ is a M -matrix. Let $A, B \in \mathcal{M}_n$. In [2], Fang gave a lower bound for $\tau(A \star B)$ as follows:

$$(1.1) \quad \tau(A \star B) \geq \min_{1 \leq i \leq n} \{a_{ii}\tau(B) + b_{ii}\tau(A) - \tau(A)\tau(B)\}.$$

In [4], Liu and Chen gave a sharper lower bound for $\tau(A \star B)$ as follows:

$$(1.2) \quad \tau(A \star B) \geq \frac{1}{2} \min_{i \neq j} \{a_{ii}b_{ii} + a_{jj}b_{jj} - [(a_{ii}b_{ii} - a_{jj}b_{jj})^2 + 4(a_{ii} - \tau(A))(b_{ii} - \tau(B))(a_{jj} - \tau(A))(b_{jj} - \tau(B))]^{1/2}\}.$$

In this paper, our aim is to propose some new lower bounds for the minimum eigenvalue of the Fan product of two M -matrices.

2. SOME LOWER BOUNDS FOR THE MINIMUM EIGENVALUE OF THE FAN PRODUCT OF M -MATRICES

Lemma 2.1 ([1]). *If $A \in \mathcal{M}_n$ is irreducible, $Az \geq kz$ for a nonnegative nonzero vector z , then $k \leq \tau(A)$.*

Lemma 2.2 (Cauchy-Schwarz inequality). *For any vectors $u = (u_1, u_2, \dots, u_n) \in \mathbb{R}^n$ and $v = (v_1, v_2, \dots, v_n) \in \mathbb{R}^n$, it holds that*

$$\left| \sum_{i=1}^n u_i v_i \right| \leq \left(\sum_{i=1}^n u_i^2 \right) \left(\sum_{i=1}^n v_i^2 \right).$$

Theorem 2.1. *If $A = (a_{ij}) \in \mathcal{M}_n$ and $B = (b_{ij}) \in \mathcal{M}_n$, then*

$$(1.1) \quad \tau(A \star B) \geq \min_{1 \leq i \leq n} \{(a_{ii} - \alpha_i)b_{ii} + \alpha_i\tau(B)\},$$

where $\alpha_i = \max_{k \neq i} \{a_{ik}\}$, for all $i \in \mathbb{N}$.

Proof. It is easy to see that (2.1) holds with equality for $n = 1$. Next, we assume that $n \geq 2$. Two cases will be discussed in the following.

Case 1. If $A \star B$ is irreducible, then A and B are irreducible. Hence, there exists a positive vector $v = (v_1, v_2, \dots, v_n)^T$ such that

$$Bv = \tau(B)v.$$

So, we have

$$b_{ii}v_i - \sum_{j \neq i} |b_{ij}|v_j = \tau(B)v_i, \quad \forall i \in \mathbb{N},$$

i.e.,

$$(2.2) \quad \sum_{j \neq i} |b_{ij}|v_j = [b_{ii} - \tau(B)]v_i, \quad \forall i \in \mathbb{N}.$$

Let $\alpha_i = \max_{k \neq i} \{|a_{ik}|\}$, for all $i \in \mathbb{N}$. Denote $C = A \star B$. For all $i \in \mathbb{N}$, by (2.2), we have

$$\begin{aligned} (Cv)_i &= a_{ii}b_{ii}v_i - \sum_{j \neq i} |a_{ij}||b_{ij}|v_j = a_{ii}b_{ii}v_i - \sum_{j \neq i} |b_{ij}||a_{ij}|v_j \\ &\geq a_{ii}b_{ii}v_i - \alpha_i \sum_{j \neq i} |b_{ij}|v_j = a_{ii}b_{ii}v_i - \alpha_i [b_{ii} - \tau(B)]v_i \\ &= [(a_{ii} - \alpha_i)b_{ii} + \alpha_i\tau(B)]v_i. \end{aligned}$$

By Lemma 2.1, we obtain

$$\tau(A \star B) \geq \min_{1 \leq i \leq n} \{(a_{ii} - \alpha_i)b_{ii} + \alpha_i\tau(B)\}.$$

Case 2. If $A \star B$ is reducible, let $T = (t_{ij})$ be the permutation matrix such that $t_{12} = t_{23} = \dots = t_{n-1,n} = t_{n,1} = 1$ and the remaining $t_{ij} = 0$. Then there exists a positive real number ε such that $A - \varepsilon T$ and $B - \varepsilon T$ are two irreducible M -matrices, i.e., $(A - \varepsilon T) \star (B - \varepsilon T)$ is irreducible. Apply Case 1 and then use the continuity argument to complete the proof. \square

Since the Fan product is commutative, the inequality (2.1) remains correct if A and B are switched. Moreover, the following result can be immediately obtained.

Theorem 2.2. *If $A = (a_{ij}) \in \mathcal{M}_n$ and $B = (b_{ij}) \in \mathcal{M}_n$, then*

$$(2.3) \quad \tau(A \star B) \geq \min_{1 \leq i \leq n} \{(b_{ii} - \beta_i)a_{ii} + \beta_i\tau(A)\},$$

where $\beta_i = \max_{k \neq i} \{|b_{ik}|\}$, for all $i \in \mathbb{N}$.

From Theorem 2.1 and Theorem 2.2 we can obtain the following result.

Theorem 2.3. If $A = (a_{ij}) \in \mathcal{M}_n$ and $B = (b_{ij}) \in \mathcal{M}_n$, then

$$\tau(A \star B) \geq \max\left\{\min_{1 \leq i \leq n} \{(a_{ii} - \alpha_i)b_{ii} + \alpha_i \tau(B)\}, \min_{1 \leq i \leq n} \{(b_{ii} - \beta_i)a_{ii} + \beta_i \tau(A)\}\right\},$$

where $\alpha_i = \max_{k \neq i} \{|a_{ik}|\}$ and $\beta_i = \max_{k \neq i} \{|b_{ik}|\}$, for all $i \in \mathbb{N}$.

Theorem 2.4. If $A = (a_{ij}) \in \mathcal{M}_n$ and $B = (b_{ij}) \in \mathcal{M}_n$, then

$$(2.4) \quad \tau(A \star B) \geq \min_{1 \leq i \leq n} \{a_{ii}b_{ii} - \alpha_i^{1/2}\beta_i^{1/2}[a_{ii} - \tau(A)]^{1/2}[b_{ii} - \tau(B)]^{1/2}\},$$

where $\alpha_i = \max_{k \neq i} \{|a_{ik}|\}$ and $\beta_i = \max_{k \neq i} \{|b_{ik}|\}$, for all $i \in \mathbb{N}$.

Proof. It is easy to see that (2.4) holds with equality for $n = 1$. Next, we assume that $n \geq 2$. Two cases will be discussed in the following.

Case 1. If $A \star B$ is irreducible, then A and B are irreducible. There exist two positive vectors $u = (u_1^2, u_2^2, \dots, u_n^2)^T$ and $v = (v_1^2, v_2^2, \dots, v_n^2)^T$ such that

$$Au = \tau(A)u,$$

and

$$Bv = \tau(B)v.$$

In order to prove the following, let $u_i > 0$ and $v_i > 0$ for all $i \in \mathbb{N}$. Hence, we have

$$a_{ii}u_i^2 - \sum_{j \neq i} |a_{ij}|u_j^2 = \tau(A)u_i^2, \quad \forall i \in \mathbb{N},$$

and

$$b_{ii}v_i^2 - \sum_{j \neq i} |b_{ij}|v_j^2 = \tau(B)v_i^2, \quad \forall i \in \mathbb{N},$$

i.e.,

$$(2.5) \quad \sum_{j \neq i} |a_{ij}|u_j^2 = [a_{ii} - \tau(A)]u_i^2, \quad \forall i \in \mathbb{N},$$

and

$$(2.6) \quad \sum_{j \neq i} |b_{ij}|v_j^2 = [b_{ii} - \tau(B)]v_i^2, \quad \forall i \in \mathbb{N}.$$

Let $\alpha_i = \max_{k \neq i} \{|a_{ik}|\}$ and $\beta_i = \max_{k \neq i} \{|b_{ik}|\}$, for all $i \in \mathbb{N}$. Define a positive vector $z = (z_1, z_2, \dots, z_n)^T$, where

$$z_i = u_i v_i, \quad \forall i \in \mathbb{N}.$$

Denote $C = A \star B$. For all $i \in \mathbb{N}$, by Lemma 2.2 and equalities (2.5), (2.6), we have

$$\begin{aligned}
(Cz)_i &= a_{ii}b_{ii}z_i - \sum_{j \neq i} |a_{ij}| |b_{ij}| z_j = a_{ii}b_{ii}z_i - \sum_{j \neq i} |b_{ij}| |a_{ij}| u_j v_j \\
&\geq a_{ii}b_{ii}z_i - \left(\sum_{j \neq i} |a_{ij}|^2 u_j^2 \right)^{1/2} \left(\sum_{j \neq i} |b_{ij}|^2 v_j^2 \right)^{1/2} \\
&\geq a_{ii}b_{ii}z_i - \alpha_i^{1/2} \beta_i^{1/2} \left(\sum_{j \neq i} |a_{ij}| u_j^2 \right)^{1/2} \left(\sum_{j \neq i} |b_{ij}| v_j^2 \right)^{1/2} \\
&= a_{ii}b_{ii}z_i - \alpha_i^{1/2} \beta_i^{1/2} [a_{ii} - \tau(A)]^{1/2} [b_{ii} - \tau(B)]^{1/2} u_i v_i \\
&= (a_{ii}b_{ii} - \alpha_i^{1/2} \beta_i^{1/2} [a_{ii} - \tau(A)]^{1/2} [b_{ii} - \tau(B)]^{1/2}) z_i.
\end{aligned}$$

By Lemma 2.1, we get

$$\tau(A \star B) \geq \min_{1 \leq i \leq n} \{a_{ii}b_{ii} - \alpha_i^{1/2} \beta_i^{1/2} [a_{ii} - \tau(A)]^{1/2} [b_{ii} - \tau(B)]^{1/2}\}.$$

Case 2. If $A \star B$ is reducible, the proof is similar to the one of Theorem 2.1. \square

3. EXAMPLE

In this section, we will show an example to illustrate our results.

Example 3.1 ([4]). Consider two 3×3 M -matrices as follows.

$$A = \begin{bmatrix} 2 & -1 & 0 \\ 0 & 1 & -0.5 \\ -0.5 & -1 & 2 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & -0.25 & -0.25 \\ -0.5 & 1 & -0.25 \\ -0.25 & -0.5 & 1 \end{bmatrix}$$

By direct calculation, $\tau(A) = 0.5402$, $\tau(B) = 0.3432$ and $\tau(A \star B) = 0.9377$. According to inequalities (1.1) and (1.2), we have

$$\tau(A \star B) \geq \min_{1 \leq i \leq 3} \{a_{ii}\tau(B) + b_{ii}\tau(A) - \tau(A)\tau(B)\} = 0.6980$$

and

$$\begin{aligned}
\tau(A \star B) &\geq \frac{1}{2} \min_{i \neq j} \{a_{ii}b_{ii} + a_{jj}b_{jj} - [(a_{ii}b_{ii} - a_{jj}b_{jj})^2 \\
&\quad + 4(a_{ii} - \tau(A))(b_{ii} - \tau(B))(a_{jj} - \tau(A))(b_{jj} - \tau(B))]^{1/2}\} = 0.7654.
\end{aligned}$$

According to inequalities (2.1), (2.3) and (2.4), we have

$$\tau(A \star B) \geq \min_{1 \leq i \leq 3} \{(a_{ii} - \alpha_i)b_{ii} + \alpha_i\tau(B)\} = 0.6716,$$

$$\tau(A \star B) \geq \min_{1 \leq i \leq 3} \{(b_{ii} - \beta_i)a_{ii} + \beta_i\tau(A)\} = 0.7701,$$

and

$$\tau(A \star B) \geq \min_{1 \leq i \leq 3} \{a_{ii}b_{ii} - \alpha_i^{1/2}\beta_i^{1/2}[a_{ii} - \tau(A)]^{1/2}[b_{ii} - \tau(B)]^{1/2}\} = 0.7252,$$

respectively.

Although we can not prove that our results are sharper than the ones of [2], [4] in theory, we can see that our results are sharper than the ones of [2], [4] for some matrices from Example 3.1.

Addendum. After this paper was accepted, I learned that Theorem 2.3 is the same as Theorem 2 in the paper H. Li: New estimation of the eigenvalue bounds of the Hadamard product and the Fan product of matrices, Henan Science, 30 (2012), 680–683; but my results are independent and obtained by a different method.

References

- [1] *A. Berman, R. J. Plemmons*: Nonnegative Matrices in the Mathematical Sciences. Classics in Applied Mathematics 9, SIAM, Philadelphia, 1994.
- [2] *M. Z. Fang*: Bounds on eigenvalues of the Hadamard product and the Fan product of matrices. Linear Algebra. Appl. 425 (2007), 7–15.
- [3] *R. A. Horn, C. R. Johnson*: Topics in Matrix Analysis. Cambridge University Press, Cambridge, 1991.
- [4] *Q. B. Liu, G. L. Chen*: On two inequalities for the Hadamard product and the Fan product of matrices. Linear Algebra Appl. 431 (2009), 974–984.

Author's address: Guanghui Cheng, School of Mathematical Sciences, University of Electronic Science and Technology of China, Chengdu, Sichuan, 611731, P.R. China, e-mail: ghcheng@126.com.