

Igor Milovanović; Emina Milovanović

Remark on inequalities for the Laplacian spread of graphs

*Czechoslovak Mathematical Journal*, Vol. 64 (2014), No. 1, 285–287

Persistent URL: <http://dml.cz/dmlcz/143966>

## Terms of use:

© Institute of Mathematics AS CR, 2014

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

REMARK ON INEQUALITIES FOR THE LAPLACIAN  
SPREAD OF GRAPHS

IGOR MILOVANOVIĆ, EMINA MILOVANOVIĆ, Niš

(Received September 21, 2013)

*Abstract.* Two inequalities for the Laplacian spread of graphs are proved in this note. These inequalities are reverse to those obtained by Z. You, B. Liu: The Laplacian spread of graphs, Czech. Math. J. 62 (2012), 155–168.

*Keywords:* Laplacian eigenvalue; spread of a graph

*MSC 2010:* 15A18, 05C50

## 1. INTRODUCTION

Let  $G = (V, E)$  be an undirected connected graph with  $m$  edges and  $n$ ,  $n \geq 3$  vertices,  $V = \{x_1, x_2, \dots, x_n\}$ . Denote by  $d_i = d(x_i)$ ,  $i = 1, 2, \dots, n$  the degree of each vertex, and by  $M_1 = \sum_{i=1}^n d_i^2$  the first Zafreb index (see [1]). The Laplacian spectrum of  $G$  are the eigenvalues  $\mu_1 \geq \mu_2 \geq \dots \geq \mu_{n-1} > \mu_n = 0$ , whereas  $P(\mu) = \mu(\mu^{n-1} + c_1\mu^{n-2} + \dots + c_{n-1})$  is the Laplacian characteristic polynomial. The Laplacian spread of a graph is defined as

$$\text{LS}(G) = \mu_1 - \mu_{n-1}.$$

In [4] Z. You and B. Liu proved several inequalities for  $\text{LS}(G)$ . Here, we are interested in the following two of them:

$$(1.1) \quad \text{LS}(G) \geq \frac{2}{n-1} \sqrt{(n-1)(M_1 + 2m) - 4m^2}$$

---

The research has been supported by the Serbian Ministry of Education, Science and Technological development, under grants No. TR32012 and TR32009.

and

$$(1.2) \quad \text{LS}(G) \geq \mu_1 - \sqrt{\frac{M_1 + 2m - \mu_1^2}{n - 2}}.$$

In this paper we are going to prove two inequalities which are reverse to (1.1) and (1.2).

## 2. MAIN RESULT

**Theorem 2.1.** *For  $\text{LS}(G)$  of a connected undirected graph  $G$  with  $n$ ,  $n \geq 3$ , vertices and  $m$  edges, the inequality*

$$(2.1) \quad \text{LS}(G) \leq \sqrt{\frac{2}{n-1}} \sqrt{(n-1)(M_1 + 2m) - 4m^2}$$

is valid. Equality in (2.1) holds if and only if  $G \cong K_n$ .

*Proof.* Since the Laplacian eigenvalues of  $G$ ,  $\mu_1, \mu_2, \dots, \mu_{n-1}$  are positive and form a decreasing sequence, according to the identity

$$T = (n-1)(M_1 + 2m) - 4m^2 = (n-1) \sum_{i=1}^{n-1} \mu_i^2 - \left( \sum_{i=1}^{n-1} \mu_i \right)^2 = \sum_{1 \leq i < j \leq n-1} (\mu_i - \mu_j)^2$$

we have that

$$(2.2) \quad T \geq \sum_{k=2}^{n-2} ((\mu_1 - \mu_k)^2 + (\mu_k - \mu_{n-1})^2) + (\mu_1 - \mu_{n-1})^2.$$

Now, if in (2.2) we apply Jensen's discrete inequality for convex functions (see for example [2], [3]), we obtain

$$T \geq \frac{n-3}{2} (\mu_1 - \mu_{n-1})^2 + (\mu_1 - \mu_{n-1})^2 = \frac{n-1}{2} (\mu_1 - \mu_{n-1})^2.$$

Since  $T \geq 0$  and  $\mu_1 - \mu_{n-1} \geq 0$ , the above inequality directly yields the inequality (2.1).

Equality in (2.2) and in Jensen's inequality holds if and only if  $\mu_1 = \mu_2 = \dots = \mu_{n-1}$ . Accordingly, we conclude that equality in (2.1) holds if and only if  $G \cong K_n$ . □

**Corollary 2.1.** *If the observed graph  $G = (V, E)$  is  $k$ -regular, then the inequality*

$$\text{LS}(G) \leq \sqrt{\frac{2nk(n-k-1)}{n-1}}$$

holds.

**Remark 2.1.** Since

$$(n-1) \sum_{i=1}^{n-1} \mu_i^2 - \left( \sum_{i=1}^{n-1} \mu_i \right)^2 = (n-2)C_1^2 - 2(n-1)C_2,$$

where  $C_1$  and  $C_2$  are the coefficients of the Laplacian characteristic polynomial, the inequalities (1.1) and (2.1) can be represented as

$$\frac{2}{n-1} \sqrt{(n-2)C_1^2 - 2(n-1)C_2} \leq \text{LS}(G) \leq \sqrt{\frac{2}{n-1}} \sqrt{(n-2)C_1^2 - 2(n-1)C_2}.$$

**Theorem 2.2.** *For  $\text{LS}(G)$  of a connected undirected graph  $G$  with  $n$  vertices and  $m$  edges, the inequality*

$$(2.3) \quad \text{LS}(G) \leq \sqrt{M_1 + 2m - (n-2)\mu_{n-1}^2 - \mu_{n-1}}$$

is valid. Equality in (2.3) holds if and only if  $G \cong K_n$ , or  $G \cong K_{1,n}$ , or  $G \cong K_{n/2, n/2}$ .

**Proof.** Inequality (2.3) can be easily obtained by using the inequality

$$M_1 + 2m = \mu_1^2 + \dots + \mu_{n-1}^2 \geq \mu_1^2 + (n-2)\mu_{n-1}^2.$$

□

#### References

- [1] *I. Gutman, K. C. Das:* The first Zagreb index 30 years after. *MATCH Commun. Math. Comput. Chem.* 50 (2004), 83–92.
- [2] *D. S. Mitrinović:* Analytic Inequalities. In Cooperation with P. M. Vasić. *Die Grundlehren der mathematischen Wissenschaften* 165, Springer, Berlin, 1970.
- [3] *D. S. Mitrinović, J. E. Pečarić, A. M. Fink:* Classical and New Inequalities in Analysis. *Mathematics and Its Applications (East European Series)* 61, Kluwer Academic Publishers, Dordrecht, 1993.
- [4] *Z. You, B. Liu:* The Laplacian spread of graphs. *Czech. Math. J.* 62 (2012), 155–168.

*Authors' address:* Igor Ž. Milovanović, Emina I. Milovanović, Faculty of Electronic Engineering, Beogradska 14, P.O.Box 73, 18000 Niš, Serbia, e-mail: igor@elfak.ni.ac.rs, ema@elfak.ni.ac.rs.