

Carlos Escudero; Filippo Gazzola; Robert Hakl; Ireneo Peral; Pedro José Torres
Existence results for a fourth order partial differential equation arising in condensed
matter physics

Mathematica Bohemica, Vol. 140 (2015), No. 4, 385–393

Persistent URL: <http://dml.cz/dmlcz/144457>

Terms of use:

© Institute of Mathematics AS CR, 2015

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

EXISTENCE RESULTS FOR A FOURTH ORDER
PARTIAL DIFFERENTIAL EQUATION
ARISING IN CONDENSED MATTER PHYSICS

CARLOS ESCUDERO, Madrid, FILIPPO GAZZOLA, Milano, ROBERT HAKL, Brno,
IRENEO PERAL, Madrid, PEDRO JOSÉ TORRES, Granada

(Received September 17, 2013)

Abstract. We study a higher order parabolic partial differential equation that arises in the context of condensed matter physics. It is a fourth order semilinear equation which nonlinearity is the determinant of the Hessian matrix of the solution. We consider this model in a bounded domain of the real plane and study its stationary solutions both when the geometry of this domain is arbitrary and when it is the unit ball and the solution is radially symmetric. We also consider the initial-boundary value problem for the full parabolic equation. We summarize our results on existence of solutions in these cases and propose an open problem related to the existence of self-similar solutions.

Keywords: higher order parabolic equation; existence of solution; blow-up in finite time; higher order elliptic equation; variational method; strongly singular boundary value problem

MSC 2010: 34B16, 35G20, 35J50, 35J60, 35K25, 35K91

1. INTRODUCTION

We are interested in the initial-boundary problem for $u = u(x, y, t)$ solving the parabolic equation

$$(1.1) \quad \partial_t u + \Delta^2 u = \det(D^2 u) + \lambda h,$$

subject to the initial condition $u(x, y, 0) = u_0(x, y)$ and where h is a function depending in general on both the space and time coordinates and belonging to a suitable Lebesgue space, $\lambda \in \mathbb{R}$. This equation is to be solved for $(x, y) \in \Omega \subset \mathbb{R}^2$, where Ω is

The research has been supported by projects MTM2010-18128, RYC-2011-09025 and SEV-2011-0087, MINECO, Spain, and project RVO 67985840, Czech Republic.

open, bounded and provided with a smooth boundary, and $t > 0$. We will consider two different sets of boundary conditions: Dirichlet and Navier.

This partial differential equation is a model that arises in the coarse-grained description of epitaxial growth processes in the field of condensed matter physics. The function $u = u(x, y, t)$ models the height of the growing film at the spatial point (x, y) at time t . Note that the evolution is dictated by the competition between the determinant of the Hessian matrix of the solution and the bilaplacian. These terms model the dynamics at the solid surface. The function h models the introduction of new mass on the surface, and the parameter λ measures the intensity of this income of new mass. A formal derivation of this model in terms of geometric quantities can be found in [3] and references therein.

We have partially analysed this model in a series of recent works [1]–[4]. We have shown the existence of stationary solutions to this partial differential equation, in the case h is time independent, for two different sets of boundary conditions: Dirichlet and Navier. The proofs are different for the two sets of boundary conditions because the variational structure that is present in the Dirichlet problem is absent in the Navier one [4]. For solutions that are radially symmetric we recover the variational structure in both cases, and the proofs of existence of solutions are built making explicit use of this fact [3]. Furthermore, in the case of radially symmetric solutions it is possible to prove non-existence of solutions for large enough data [2]. For the evolution problem it is possible to prove local existence of solutions for arbitrary data and global existence of solutions for small (but otherwise arbitrary) data. Depending on the boundary conditions and the concomitant presence of a variational structure in the equation as well as on the size of the data it is possible to prove blow-up of the solution in finite time and convergence to a stationary solution in the long time limit [1].

A summary of these results will be exposed in the next section. Together with these proven facts, there is a number of questions that remain open for both the partial differential equation (1.1) and its stationary counterpart. One of them is the existence of self-similar solutions and its possible role in the blow-up structure. We describe an open question related to the existence of self-similar solutions in Section 3.

2. SUMMARY OF PREVIOUS RESULTS CONCERNING EXISTENCE OF SOLUTIONS

In this section we list without proof some results we have recently obtained. The proofs can be found in the references quoted in the text next to the statement in question. We are going to consider the following two sets of boundary conditions $u = \partial_n u = 0$ on Ω , which we refer to as Dirichlet boundary conditions, and $u = \Delta u = 0$

on Ω , which we refer to as Navier boundary conditions. Following [4] we find that the Dirichlet problem for the stationary version of (1.1)

$$(2.1) \quad \Delta^2 u = \det(D^2 u) + \lambda h,$$

where of course h is assumed to be time independent, has a variational structure:

Theorem 2.1. *The critical points of the functional*

$$(2.2) \quad \begin{aligned} J_\lambda: W_0^{2,2}(\Omega) &\rightarrow \mathbb{R}, \\ u \rightarrow J_\lambda(u) &= \frac{1}{2} \int_\Omega |\Delta u|^2 \, dx \, dy - \int_\Omega u_x u_y u_{xy} \, dx \, dy - \lambda \int_\Omega h u \, dx \, dy \end{aligned}$$

are weak solutions to the Dirichlet problem for (2.1).

This variational structure can be used to prove the existence of at least two solutions to the corresponding boundary value problem.

Theorem 2.2. *Let $h \in L^1(\Omega)$. Then there exists a $\lambda_0 > 0$ such that for $0 \leq \lambda < \lambda_0$, the Dirichlet problem for equation (2.1) has at least two solutions in $W_0^{2,2}(\Omega)$.*

The proof can be found in [4] and makes use of the mountain pass geometry of functional (2.2). For the Navier problem for equation (2.1) we cannot use the above mentioned variational methods because we do not know of any suitable functional in this case. Instead we have the following result:

Theorem 2.3. *Let $h \in L^1(\Omega)$. Then there exists a $\lambda_0 > 0$ such that for $0 \leq \lambda < \lambda_0$, the Navier problem for equation (2.1) has at least one solution in $W_0^{1,2}(\Omega) \cap W^{2,2}(\Omega)$.*

The proof makes use of the Banach fixed point theorem, see [4]. Note also that this proof can be immediately adapted for the case of Dirichlet boundary conditions.

The radial problem corresponding to (2.1) reads

$$\frac{1}{r} \left\{ r \left[\frac{1}{r} (ru')' \right]' \right\}' = \frac{1}{r} u' u'' + \lambda h(r),$$

where $r = \sqrt{x^2 + y^2}$. In this case, for both sets of boundary conditions, the problem admits a variational formulation. And in both cases, due to the mountain pass geometry of the associated functional, it is possible to prove the existence of at least two solutions for small enough λ , see [3]. Furthermore, it is also possible to prove the non-existence of solutions for large enough λ as well as rigorous bounds for the values

of λ that separate existence from non-existence, see [2]. Note that these bounds are rather precise in certain cases when compared to the numerical estimations of the critical values of λ calculated in [3]. With respect to the full evolution problem (1.1), we can prove the following theorem:

Theorem 2.4. *Let $T > 0$; for any $u_0 \in W_0^{2,2}(\Omega)$, any $h \in L^2(0, T; L^2(\Omega))$ and any $\lambda \in \mathbb{R}$ the Dirichlet problem has a unique solution in*

$$\mathcal{X}_T := C(0, T; W_0^{2,2}(\Omega)) \cap L^2(0, T; W^{4,2}(\Omega)) \cap W^{1,2}(0, T; L^2(\Omega)),$$

provided T is sufficiently small. Furthermore, for any $T \in (0, \infty]$ there exists a unique solution to this problem in the same space provided $\|u_0\|_{W^{2,2}}$ and λ are small enough. Moreover, if $[0, T^)$ denotes the maximal interval of continuation of u and if $T^* < \infty$ then $\|u(t)\|_{W^{2,2}} \rightarrow \infty$ as $t \rightarrow T^*$.*

An analogous result holds for the Navier problem. Note that for this last statement, unlike in the previous results concerning the stationary problems, we have allowed the datum h to depend on both space and time coordinates. The proof makes use of the Banach fixed point theorem and of the potential well techniques and can be found in [1].

We have proved more results concerning the parabolic problem (1.1) but we are not going to expose them because due to their more technical nature they require more preparatory results. They concern the asymptotic properties of the solutions to the Dirichlet problem that converge to zero in the long time provided $\lambda = 0$ and the initial datum is small enough. Also, the blow-up in finite time that takes place for both the Navier and Dirichlet problems in case the initial datum is large enough. These results can be found in [1]. Precisely these results, and in particular those regarding the blow-up, motivate in part the study of self-similar solutions to the problems at hand. In the next section we outline some facts about this sort of solutions.

3. THE SEARCH FOR SELF-SIMILAR SOLUTIONS

In this section we describe an open problem related to the existence of self-similar solutions to partial differential equation (1.1). For the first time we are going to consider this equation on all the plane instead of on a bounded domain Ω . Our first step is setting $\lambda = 0$. Next we look for solutions with the form $u(r, t) = t^{-\beta} f(r/t^\alpha)$, where α and β are two real parameters to be fitted in order to find a closed ordinary differential equation involving the self-similar variable $\eta = r/t^\alpha$ and the solution to equation (1.1) expressed as a function of this variable only $f(\eta)$.

Substituting our self-similar ansatz in (1.1) we find that this equation adopts the closed form

$$4f'(\eta) - \eta^4 f'(\eta) - 4\eta f''(\eta) - 4\eta^2 f'(\eta)f''(\eta) + 8\eta^2 f'''(\eta) + 4\eta^3 f''''(\eta) = 0$$

only if $\alpha = 1/4$ and $\beta = 0$. Note that the simplicity of this fact possibly makes it one of the simplest ways of looking for self-similar solutions. Now we have to provide suitable boundary conditions for this ordinary differential equation. Note that by its very nature the self-similar variable describes a rotationally invariant solution. Therefore we assume the following symmetry conditions on the solution to equation (1.1): $\partial_r u(0, 0, t) = 0$ and $\partial_r \Delta_r u(0, 0, t) = 0$, for a rotationally invariant u , where $\Delta_r(\cdot) = (1/r)\partial_r[r\partial_r(\cdot)]$. Finally, we impose that both u and Δu decrease to zero as $r \rightarrow \infty$. So this leads to the boundary value problem

$$(3.1) \quad 4f'(\eta) - \eta^4 f'(\eta) - 4\eta f''(\eta) - 4\eta^2 f'(\eta)f''(\eta) + 8\eta^2 f'''(\eta) + 4\eta^3 f''''(\eta) = 0, \\ f'(0) = f'''(0) = 0, \quad f(\eta), f''(\eta) \rightarrow 0 \quad \text{as } \eta \rightarrow \infty.$$

The existence of solutions to this problem automatically implies the existence of solutions of the form $u(x, y, t) = f(r/t^{1/4})$ to partial differential equation (1.1). The obvious fact that $u \equiv 0$ solves (1.1) with $\lambda = 0$ and with the assumed boundary conditions directly translates into the fact that $f \equiv 0$ solves boundary value problem (3.1).

Noting that equation (3.1) does not depend on f but on its derivatives we can obtain a third order ordinary differential equation for $g(\eta) = f'(\eta)$. The corresponding boundary value problem reads

$$(3.2) \quad 4g - \eta^4 g - 4\eta g' - 4\eta^2 g g' + 8\eta^2 g'' + 4\eta^3 g''' = 0, \\ g(0) = g''(0) = 0, \quad g'(\eta) \rightarrow 0 \quad \text{as } \eta \rightarrow \infty.$$

Note that this problem is strongly singular and is to be solved for $\eta \in [0, \infty)$. Obviously $g \equiv 0$ solves this boundary value problem. Therefore we will be interested in nontrivial solutions. Since one expects the solutions of (3.1), if any, to behave like a Gaussian, one should first try to exclude the cases where $f''(0) > 0$. We do so in the next statement.

Proposition 3.1. *A local solution of the problem*

$$(3.3) \quad 4g - \eta^4 g - 4\eta g' - 4\eta^2 g g' + 8\eta^2 g'' + 4\eta^3 g''' = 0, \\ g(0) = g''(0) = 0, \quad g'(0) > 0,$$

blows up in finite time. More precisely, there exists $\bar{\eta} > 0$ such that $g'(\eta) > 0$ for $\eta \in (0, \bar{\eta})$ and $\lim_{\eta \rightarrow \bar{\eta}} g(\eta) = \infty$.

Proof. Let g be a local solution of (3.3). For contradiction, assume that there exists the first $\eta_0 > 0$ such that $g'(\eta_0) = 0$. Then $g(\eta) > 0$ and $g'(\eta) > 0$ for all $\eta \in (0, \eta_0)$. In turn, if we rewrite the equation appearing in (3.3) as

$$4(\eta^3 g'' - \eta^2 g' + \eta g)' = \eta^2 g(4g' + \eta^2),$$

this readily shows that $(\eta^3 g'' - \eta^2 g' + \eta g)' > 0$ in $(0, \eta_0]$. Since the term inside the bracket vanishes at $\eta = 0$ this yields $\eta^3 g''(\eta) - \eta^2 g'(\eta) + \eta g(\eta) > 0$ in $(0, \eta_0]$, which we may also rewrite as

$$g''(\eta) - \frac{g'(\eta)}{\eta} + \frac{g(\eta)}{\eta^2} > 0 \quad \text{for } \eta \in (0, \eta_0].$$

This proves that $(g' - g/\eta)' > 0$ and since the term inside the bracket vanishes as $\eta \rightarrow 0$, this also gives $g'(\eta_0) > g(\eta_0)/\eta_0 > 0$, contradicting the characterization of η_0 . We have so proved that

$$(3.4) \quad g'(\eta) > 0 \quad \text{for all } \eta \in [0, \bar{\eta})$$

where g is a local solution to (3.3) and $\bar{\eta}$ is the endpoint of its interval of continuation, possibly infinite. Moreover, what we have seen also proves that

$$(3.5) \quad \lim_{\eta \rightarrow \bar{\eta}} g(\eta) = \infty$$

in both the cases $\bar{\eta} < \infty$ (blow up in finite time) and $\bar{\eta} = \infty$ (global solution).

Now put $\eta = e^r$, $g(\eta) = h(\log \eta)$, $h(r) = g(e^r)$. Then the equation in (3.3) reads

$$(3.6) \quad 4(h'''(r) - h''(r) - h'(r)) = 4e^r h'(r)h(r) + (e^{4r} - 4)h(r), \quad r \in (-\infty, \infty),$$

while conditions (3.4) and (3.5) become

$$(3.7) \quad h'(r) > 0 \quad \text{for all } r < \bar{r}, \quad \lim_{r \rightarrow \bar{r}} h(r) = \infty, \quad \bar{r} = \log \bar{\eta}.$$

For contradiction, assume that $\bar{r} = \infty$. From (3.6) and (3.7) we infer that

$$(h''(r) - h'(r) - h(r))' \geq e^r h'(r)h(r) \geq 2h'(r)h(r) \quad \text{for all } r \geq \log 2.$$

By integrating this inequality over $[\log 2, r]$ we get

$$h''(r) - h'(r) - h(r) \geq h(r)^2 + \gamma \quad \text{for all } r \geq \log 2$$

where $\gamma = h''(\log 2) - h'(\log 2) - h(\log 2) - h(\log 2)^2$. By (3.7) we may multiply this inequality by $h'(r)$ and maintain its sense:

$$\begin{aligned} h''(r)h'(r) &\geq h'(r)^2 + h(r)h'(r) + h(r)^2h'(r) + \gamma h'(r) \\ &\geq h(r)h'(r) + h(r)^2h'(r) + \gamma h'(r) \end{aligned}$$

for all $r \geq \log 2$. Let us now integrate this inequality over $[\log 2, r]$; we obtain

$$\frac{1}{2}h'(r)^2 \geq \frac{1}{2}h(r)^2 + \frac{1}{3}h(r)^3 + \gamma h(r) + \delta$$

where $\delta = \frac{1}{2}h'(\log 2)^2 - \frac{1}{2}h(\log 2)^2 - \frac{1}{3}h(\log 2)^3 - \gamma h(\log 2)$. By (3.7) we know that there exists $R > \log 2$ such that the last inequality yields $h'(r)^2 \geq \frac{1}{4}h(r)^3$ for all $r \geq R$. By taking the square root of this inequality we obtain $h'(r)/h(r)^{3/2} \geq 1/2$ which, upon integration over $[R, r]$ yields

$$\frac{1}{h(R)^{1/2}} \geq \frac{r-R}{4} + \frac{1}{h(r)^{1/2}} \quad \text{for all } r \geq R.$$

By letting $r \rightarrow \infty$ we reach a contradiction. □

In the case $g'(0) < 0$, we have numerically integrated this differential equation, for an example see Figure 1. Based on Proposition 3.1 and on these preliminary numerical experiments we conjecture the following result:

Conjecture 3.1. *There exist no nontrivial solutions to boundary value problem (3.2).*

It is interesting to highlight one property of the solutions to (3.3) for any $g'(0)$ used in the numerical integrations shown in Figure 1. If we assume that $g \in C^3[0, \infty)$, we have

$$\begin{aligned} g(\eta) &= g'(0)\eta + g'''(0)\frac{\eta^3}{6} + o(\eta^3), & g'(\eta) &= g'(0) + g'''(0)\frac{\eta^2}{2} + o(\eta^2), \\ g''(\eta) &= g'''(0)\eta + o(\eta), & g'''(\eta) &= g'''(0) + o(1) \quad \text{as } \eta \rightarrow 0. \end{aligned}$$

Substituting this into the equation and simplifying we arrive at

$$\left[\frac{8}{3}g'''(0) - g'(0)^2 \right] \eta^3 + o(\eta^3) = 0,$$

which implies $g'''(0) = 3g'(0)^2/8 > 0$. Consequently $g''(\eta) > 0$ in a right neighborhood of $\eta = 0$ and thus $g(\eta)$ is convex in a right neighborhood of $\eta = 0$. Note that this feature can be observed in the numerical solutions in Figure 1 (a).




Figure 1. Numerical solutions of the ordinary differential equation corresponding to boundary value problem (3.2). The boundary conditions $g(0) = g''(0) = 0$ were employed in this numerical integration. The third condition is arbitrary and chosen to be: (a) $g'(0) = -1$ (solid line), $g'(0) = -10$ (dashed line) and $g'(0) = -10^2$ (dotted line); (b) $g'(0) = -1$ (solid line), $g'(0) = -10^2$ (dashed line) and $g'(0) = -10^4$ (dotted line); (c) $g'(0) = -10^4$ (dash-dotted line), $g'(0) = -10^5$ (solid line), $g'(0) = -10^6$ (dashed line) and $g'(0) = -10^7$ (dotted line); (d) $g'(0) = -10^4$ (dash-dotted line), $g'(0) = -10^5$ (solid line), $g'(0) = -8 \times 10^5$ (dashed line) and $g'(0) = -2.5 \times 10^6$ (dotted line).

Let us now make some final remarks. Note first that a proof of Conjecture 3.1 would not imply the nonexistence of (any kind of) self-similar solutions to partial differential equation (1.1). The existence of these is actually a question that remains open. Also, the trivial solution $g \equiv 0$ corresponds to a constant f , but the only constant f that solves boundary value problem (3.1) is $f \equiv 0$ due to the extra boundary condition. As we have already mentioned, this corresponds to the solution $u \equiv 0$ of the partial differential equation. In connection to this, any nontrivial solution to problem (3.2) gives rise to a nontrivial solution f to (3.1) only if it satisfies an extra cancelation condition.

References

- [1] *C. Escudero, F. Gazzola, I. Peral*: Global existence versus blow-up results for a fourth order parabolic PDE involving the Hessian. *J. Math. Pures Appl.* (9) *103* (2015), 924–957.
- [2] *C. Escudero, R. Hakl, I. Peral, P. J. Torres*: Existence and nonexistence results for a singular boundary value problem arising in the theory of epitaxial growth. *Math. Methods Appl. Sci.* *37* (2014), 793–807.
- [3] *C. Escudero, R. Hakl, I. Peral, P. J. Torres*: On radial stationary solutions to a model of non-equilibrium growth. *Eur. J. Appl. Math.* *24* (2013), 437–453.
- [4] *C. Escudero, I. Peral*: Some fourth order nonlinear elliptic problems related to epitaxial growth. *J. Differ. Equations* *254* (2013), 2515–2531.

Authors' addresses: *Carlos Escudero*, Departamento de Matemáticas, Facultad de Ciencias, Universidad Autónoma de Madrid, C/Francisco Tomás y Valiente 7, E-28049 Madrid, Spain, e-mail: carlos.escudero@uam.es; *Filippo Gazzola*, Politecnico di Milano, Sede Milano Leonardo, Edificio Nave, 5° Piano, Milan, Italy, e-mail: filippo.gazzola@polimi.it; *Robert Hakl*, Institute of Mathematics of The Czech Academy of Sciences, Žitkova 22, 616 62 Brno, Czech Republic, e-mail: hakl@ipm.cz; *Ireneo Peral*, Departamento Matemáticas, Facultad de Ciencias, Universidad Autónoma de Madrid, C/Francisco Tomás y Valiente 7, E-28049 Madrid, Spain, e-mail: ireneo.peral@uam.es; *Pedro José Torres*, Departamento de Matemática Aplicada, Facultad de Ciencias, Universidad de Granada, Campus de Fuentenueva, 180 71 Granada, Spain, e-mail: ptorres@ugr.es.