

Zhiyong Chen; Haibin Wang; Xuejun Wang; Shuhe Hu

Strong Convergence for weighed sums of negatively superadditive dependent random variables

Kybernetika, Vol. 52 (2016), No. 1, 52–65

Persistent URL: <http://dml.cz/dmlcz/144862>

Terms of use:

© Institute of Information Theory and Automation AS CR, 2016

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

STRONG CONVERGENCE FOR WEIGHED SUMS OF NEGATIVELY SUPERADDITIVE DEPENDENT RANDOM VARIABLES

ZHIYONG CHEN, HAIBIN WANG, XUEJUN WANG AND SHUHE HU

In this paper, the strong law of large numbers for weighted sums of negatively superadditive dependent (NSD, in short) random variables is obtained, which generalizes and improves the corresponding one of Bai and Cheng ([2]) for independent and identically distributed random variables to the case of NSD random variables.

Keywords: NSD random variables, weighted sums, strong law of large numbers

Classification: 60F15

1. INTRODUCTION

Let $\{X_n, n \geq 1\}$ be a sequence of random variables defined on a fixed probability space (Ω, \mathcal{F}, P) . We say that the sequence $\{X_n, n \geq 1\}$ satisfies the strong law of large numbers if there exist some increasing sequence $\{a_n, n \geq 1\}$ and some sequence $\{c_n, n \geq 1\}$ such that

$$\frac{1}{a_n} \sum_{i=1}^n (X_i - c_i) \rightarrow 0 \quad \text{a.s. as } n \rightarrow \infty.$$

Many authors have studied the strong law of large numbers for sequences of independent and identically distributed random variables. The following Theorems A is due to Bai and Cheng ([2]).

Theorem A. Suppose that $1 < \alpha, \beta < \infty$, $1 \leq p < 2$, and $1/p = \frac{1}{\alpha} + \frac{1}{\beta}$. Let $\{X_n, n \geq 1\}$ be a sequence of independent and identically distributed random variables satisfying $EX_1 = 0$, and let $\{a_{nk}, 1 \leq k \leq n, n \geq 1\}$ be an array of real numbers satisfying

$$\limsup_{n \rightarrow \infty} \left(\frac{1}{n} \sum_{k=1}^n |a_{nk}|^\alpha \right)^{1/\alpha} < \infty.$$

If $E|X_1|^\beta < \infty$, then

$$n^{-1/p} \sum_{k=1}^n a_{nk} X_k \rightarrow 0 \text{ a.s. as } n \rightarrow \infty.$$

The result of Theorem A for independent and identically distributed random variables has been extended to the case of dependent random variables. See for example, Jing and Liang ([12]) established Marcinkiewicz–Zygmund strong law of large numbers for weighted sums of negatively associated random variables. Meng and Lin ([14]) obtained the Marcinkiewicz–Zygmund strong law of large numbers for $\tilde{\rho}$ -mixing random variables. Moreover, Shen ([17]) discussed the strong limit theorem for weighted sums of sequences of negatively dependent random variables. Sung ([22]) gave some sufficient conditions to prove the strong law of large numbers for weighted sums of random variables. Recently, Hu et al. ([9]) established the strong law of large numbers of partial sums for pairwise of negatively quadrant dependent sequences. Shen and Wu ([16]) investigated strong and weak convergence for asymptotically almost negatively associated random variables. Shen ([18]) established a general result on strong convergence for weighted sums of a class of random variables. Inspired by the literatures above, we will extend and improve the result of Theorem A to the case of weighted sums of negatively superadditive dependent random variables.

The definitions of negatively associated random variables and negatively superadditive dependent random variables are as follows.

Definition 1.1. A finite collection of random variables X_1, X_2, \dots, X_n is said to be negatively associated (NA) if for every pair of disjoint subsets A_1, A_2 of $\{1, 2, \dots, n\}$,

$$Cov\{f(X_i : i \in A_1), g(X_j : j \in A_2)\} \leq 0, \tag{1.1}$$

whenever f and g are coordinatewise nondecreasing such that this covariance exists. An infinite sequence $\{X_n, n \geq 1\}$ is NA if every finite subcollection is NA.

The notion of NA was first introduced by Alam and Lai Saxena ([1]) and carefully studied by Joag-Dev and Proschan ([11]). Joag-Dev and Proschan ([11]) showed that many well known multivariate distributions possess the NA property. For more details about NA random variables, one can refer to Block et al. ([3]), Matuala ([13]), Budsaba et al. ([4]), Wu and Jiang ([28, 29]), Wang et al. ([23]), Yang et al. ([30]), Gerasimov et al. ([8]), and so on.

Definition 1.2. A function $\phi : R^n \rightarrow R$ is called superadditive if $\phi(\mathbf{x} \vee \mathbf{y}) + \phi(\mathbf{x} \wedge \mathbf{y}) \geq \phi(\mathbf{x}) + \phi(\mathbf{y})$ for all $\mathbf{x}, \mathbf{y} \in R^n$, where \vee is for componentwise maximum and \wedge is for componentwise minimum.

Next, we provide the concept of negatively superadditive dependent random variables as follows.

Definition 1.3. A random vector $\mathbf{X} = (X_1, X_2, \dots, X_n)$ is said to be negatively superadditive dependent (NSD) if

$$E\phi(X_1, X_2, \dots, X_n) \leq E\phi(X_1^*, X_2^*, \dots, X_n^*), \tag{1.2}$$

where $X_1^*, X_2^*, \dots, X_n^*$ are independent such that X_i^* and X_i have the same distribution for each i , and ϕ is a superadditive function such that the expectations in (1.2) exist.

A sequence $\{X_n, n \geq 1\}$ of random variables is said to be NSD if for all $n \geq 1$, (X_1, X_2, \dots, X_n) is NSD.

The concept of NSD random variables was introduced by Hu ([10]). An example of an NSD sequence which is not NA was constructed by Hu ([10]), and illustrated that NSD implies NOD (negatively orthant dependent). Christofides and Vaggelatou ([5]) indicated that the family of NSD sequence contains NA. So we can see that NSD is weaker than NA. Negatively superadditive dependent structure is an extension of negatively associated structure and sometimes more useful than it and can be used to get many important probability inequalities. A number of limit theorems and applications for NSD random variables have been found by many authors. See for example, Eghbal et al. ([6]) derived two maximal inequalities and strong law of large numbers of quadratic forms of NSD random variables, and Eghbal et al. ([7]) provided some Kolmogorov inequality for quadratic forms of nonnegative NSD uniformly bounded random variables, Shen et al. ([20]) obtained Kolmogorov-type inequality and the almost sure convergence for NSD sequences, Shen et al. ([21]) established some inequalities for NSD random variables, Shen et al. ([19]) gave some applications of the Rosenthal-type inequality for NSD random variables, Wang et al. ([26]) presented some results on complete convergence for weighted sums of NSD random variables and gave its application in the EV regression model, and so forth.

Finally, we will present the concept of stochastic domination, which will be used frequently in this paper.

Definition 1.4. A sequence $\{X_n, n \geq 1\}$ of random variables is said to be stochastically dominated by a random variable X if there exists a positive constant C , such that

$$P(|X_n| > x) \leq CP(|X| > x) \quad (1.3)$$

for all $x \geq 0$ and $n \geq 1$.

The main purpose of the paper is to study the strong law of large numbers for weighted sums of NSD random variables, which generalizes the corresponding one of Theorem A for independent and identically distributed random variables. The techniques used in the paper are the truncation method and the moment inequality for NSD random variables.

Throughout this paper, let $a \wedge b = \min(a, b)$, $a \vee b = \max(a, b)$ and C denotes a positive constant which may be different in various places. Let $I(A)$ be the indicator function of the set A and $a_n = O(b_n)$ stands for $a_n \leq Cb_n$.

2. PRELIMINARY LEMMAS

The main results of this paper are dependent on the following lemmas.

Lemma 2.1. (cf. Hu, [10]) Let $\{X_n, n \geq 1\}$ be a sequence of NSD random variables, and let $\{f_n, n \geq 1\}$ be a sequence of nondecreasing functions, then $\{f_n(X_n), n \geq 1\}$ is still a sequence of NSD random variables.

Lemma 2.2. (cf. Hu, [10]) Let $\mathbf{X} = (X_1, X_2, \dots, X_n)$ be an NSD random vector, and let $\mathbf{X}^* = (X_1^*, X_2^*, \dots, X_n^*)$ be independent vector such that X_i^* and X_i have the same distribution for each i . Then for any nondecreasing convex function f ,

$$Ef \left(\max_{1 \leq k \leq n} \sum_{i=1}^k X_i \right) \leq Ef \left(\max_{1 \leq k \leq n} \sum_{i=1}^k X_i^* \right). \tag{2.1}$$

Lemma 2.2 is the so called comparison theorem on moments between the NSD and independent random variables. Similarly to the proof of Theorem 2 of Shao ([15]) and by using Lemma 2.2, Wang et al. ([25]) got the following Rosenthal-type maximal inequality for NSD random variables.

Lemma 2.3. (Rosenthal-type maximal inequality) Let $\{X_n, n \geq 1\}$ be a sequence of NSD random variables with $EX_n = 0$ and $E|X_n|^p < \infty$ for some $p > 2$. Then there exists a positive constant C_p depending only on p such that

$$E \left(\max_{1 \leq i \leq n} \left| \sum_{i=1}^n X_i \right|^p \right) \leq C_p \left\{ \sum_{i=1}^n E|X_i|^p + \left(\sum_{i=1}^n EX_i^2 \right)^{p/2} \right\}, \quad n \geq 1. \tag{2.2}$$

Lemma 2.4. (cf. Shen et al., [20]) Let $\{X_n, n \geq 1\}$ be a sequence of NSD random variables. If

$$\sum_{n=1}^{\infty} \text{Var}(X_n) < \infty, \tag{2.3}$$

then $\sum_{n=1}^{\infty} (X_n - EX_n)$ converges almost surely.

Lemma 2.5. (cf. Wu, [27]) Let $\{X_n, n \geq 1\}$ be a sequence of random variables which is stochastically dominated by a random variable X . For any $\alpha > 0$ and $b > 0$, the following two statements hold:

$$\begin{aligned} E|X_n|^\alpha I(|X_n| \leq b) &\leq C_1 [E|X|^\alpha I(|X| \leq b) + b^\alpha P(|X| > b)], \\ E|X_n|^\alpha I(|X_n| > b) &\leq C_2 E|X|^\alpha I(|X| > b), \end{aligned} \tag{2.4}$$

where C_1 and C_2 are positive constants.

3. MAIN RESULTS AND THEIR PROOFS

In this section, we will provide some results on strong convergence for weighted sums of NSD random variables.

Theorem 3.1. Let $0 < p < 2$, $0 < \alpha, \beta < \infty$, and $1/p = \frac{1}{\alpha} + \frac{1}{\beta}$. Assume that $\{X_n, n \geq 1\}$ is a sequence of NSD random variables stochastically dominated by a random variable X such that $E|X|^\beta < \infty$. Let $EX_n = 0, n \geq 1$, if $\beta > 1$ and $\{a_{ni}, i \geq 1, n \geq 1\}$ be an array of real numbers satisfying

$$\sum_{i=1}^n |a_{ni}|^\alpha = O(n). \tag{3.1}$$

Then

$$n^{-1/p} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} X_i \right| \rightarrow 0 \text{ a.s. as } n \rightarrow \infty. \quad (3.2)$$

Proof. For fixed $n \geq 1$ and $1 \leq i \leq n$, denote

$$Y_i = X_i I(|X_i| \leq n^{1/\beta}) + n^{1/\beta} I(X_i > n^{1/\beta}) - n^{1/\beta} I(X_i < -n^{1/\beta}),$$

$$Z_i = (X_i + n^{1/\beta}) I(X_i < -n^{1/\beta}) + (X_i - n^{1/\beta}) I(X_i > n^{1/\beta}).$$

Meanwhile, one can see that $a_{ni} = a_n^+ - a_{ni}^-$. Without loss of generality, we can assume that $a_{ni} > 0$. Hence, $X_i = Y_i + Z_i$, which implies that

$$\begin{aligned} n^{-1/p} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} X_i \right| &\leq n^{-1/p} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} Y_i \right| + n^{-1/p} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} Z_i \right| \\ &\leq n^{-1/p} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} (Y_i - EY_i) \right| + n^{-1/p} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} EY_i \right| \\ &\quad + n^{-1/p} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} Z_i \right| \\ &\doteq H + I + J. \end{aligned} \quad (3.3)$$

To prove (3.2) we need to prove $H \rightarrow 0$ a.s., $I \rightarrow 0$ and $J \rightarrow 0$ a.s. as $n \rightarrow \infty$.

Firstly, we will show that $H \rightarrow 0$ a.s.

Combining (3.1) with Hölder's inequality, we have for $1 \leq \gamma < \alpha$ that

$$\sum_{i=1}^n |a_{ni}|^\gamma \leq \left(\sum_{i=1}^n |a_{ni}|^\alpha \right)^{\gamma/\alpha} \left(\sum_{i=1}^n 1 \right)^{1-\gamma/\alpha} \leq Cn. \quad (3.4)$$

Jensen's inequality implies that for any $0 < \alpha \leq \gamma$,

$$\sum_{i=1}^n |a_{ni}|^\gamma \leq \left(\sum_{i=1}^n |a_{ni}|^\alpha \right)^{\gamma/\alpha} \leq Cn^{\gamma/\alpha}. \quad (3.5)$$

It follows from (3.4) and (3.5) that

$$\sum_{i=1}^n |a_{ni}|^\gamma \leq Cn^{(1 \vee \gamma/\alpha)}. \quad (3.6)$$

By Borel–Cantelli lemma, we only need to show that for any $\varepsilon > 0$,

$$\sum_{n=1}^{\infty} P \left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} (Y_i - EY_i) \right| > \varepsilon n^{1/p} \right) < \infty. \quad (3.7)$$

For fixed $n \geq 1$, it easily seen that $\{a_{ni}(Y_i - EY_i), 1 \leq i \leq n\}$ are still NSD random variables by Lemma 2.1. Taking $r > 1/\min\{1/\alpha, 1/\beta, 1/2, 1/p - 1/2\}$, which implies that $r > \alpha$, $r > \beta$ and $r > 2$. It follows from Markov's inequality and Lemma 2.3 that

$$\begin{aligned}
& \sum_{n=1}^{\infty} P\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni}(Y_i - EY_i) \right| > \varepsilon n^{1/p}\right) \\
& \leq C \sum_{n=1}^{\infty} n^{-r/p} E\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni}(Y_i - EY_i) \right|^r\right) \\
& \leq C \sum_{n=1}^{\infty} n^{-r/p} \sum_{i=1}^n E|a_{ni}(Y_i - EY_i)|^r + C \sum_{i=1}^{\infty} n^{-r/p} \left(\sum_{i=1}^n E|a_{ni}(Y_i - EY_i)|^2\right)^{r/2} \\
& \doteq H_1 + H_2.
\end{aligned}$$

For H_1 , we have by C_r inequality, Jensen's inequality, (3.5) and Lemma 2.5 that

$$\begin{aligned}
H_1 & \leq C \sum_{n=1}^{\infty} n^{-r/p} \sum_{i=1}^n |a_{ni}|^r E|Y_i|^r \\
& \leq C \sum_{n=1}^{\infty} n^{-r/p} \sum_{i=1}^n |a_{ni}|^r [E|X_i|^r I(|X_i| \leq n^{1/\beta}) + n^{r/\beta} P(|X_i| > n^{1/\beta})] \\
& \leq C \sum_{n=1}^{\infty} n^{-r/p} \sum_{i=1}^n |a_{ni}|^r [E|X|^r I(|X| \leq n^{1/\beta}) + n^{r/\beta} P(|X| > n^{1/\beta})] \\
& \leq C \sum_{n=1}^{\infty} n^{-r/\beta} E|X|^r I(|X| \leq n^{1/\beta}) + C \sum_{n=1}^{\infty} P(|X| > n^{1/\beta}) \\
& \leq C \sum_{n=1}^{\infty} n^{-r/\beta} \sum_{i=1}^n E|X|^r I((i-1)^{1/\beta} < |X| \leq i^{1/\beta}) \\
& \quad + C \sum_{n=1}^{\infty} \sum_{i=n}^{\infty} P(i^{1/\beta} < |X| \leq (i+1)^{1/\beta}) \\
& \leq C \sum_{i=1}^{\infty} E|X|^r I((i-1)^{1/\beta} < |X| \leq i^{1/\beta}) \sum_{n=i}^{\infty} n^{-r/\beta} \\
& \quad + C \sum_{i=1}^{\infty} EI(i^{1/\beta} < |X| \leq (i+1)^{1/\beta})i \\
& \leq C \sum_{i=1}^{\infty} i^{(r-\beta)/\beta} E|X|^\beta I((i-1)^{1/\beta} < |X| \leq i^{1/\beta}) i^{-r/\beta+1} \\
& \quad + CE|X|^\beta \sum_{i=1}^{\infty} I(i^{1/\beta} < |X| \leq (i+1)^{1/\beta}) \\
& \leq CE|X|^\beta < \infty.
\end{aligned} \tag{3.8}$$

Next, we prove that $H_2 < \infty$. By C_r inequality, Jensen's inequality, (3.6) and Lemma 2.5 again, we can get that

$$\begin{aligned}
\sum_{i=1}^n E|a_{ni}(Y_i - EY_i)|^2 &\leq \sum_{i=1}^n a_{ni}^2 EY_i^2 \\
&\leq C \sum_{i=1}^n a_{ni}^2 [EX_i^2 I(|X_i| \leq n^{1/\beta}) + n^{2/\beta} P(|X_i| > n^{1/\beta})] \\
&\leq C \sum_{i=1}^n a_{ni}^2 [EX^2 I(|X| \leq n^{1/\beta}) + n^{2/\beta} P(|X| > n^{1/\beta})] \\
&\leq Cn^{(1+2/\alpha)} [EX^2 I(|X| \leq n^{1/\beta}) + n^{2/\beta} P(|X| > n^{1/\beta})]. \quad (3.9)
\end{aligned}$$

It follows by Markov's inequality and the fact $E|X|^\beta < \infty$ that

$$\begin{aligned}
&EX^2 I(|X| \leq n^{1/\beta}) + n^{2/\beta} P(|X| > n^{1/\beta}) \\
&\leq \begin{cases} CEX^2 I(|X| \leq n^{1/\beta}) + EX^2 I(|X| > n^{1/\beta}), & \text{for } \beta \geq 2, \\ Cn^{(2-\beta)/\beta} E|X|^\beta I(|X| \leq n^{1/\beta}) + n^{-1+2/\beta} E|X|^\beta I(|X| > n^{1/\beta}), & \text{for } \beta < 2, \end{cases} \\
&\leq \begin{cases} CEX^2, & \text{for } \beta \geq 2, \\ Cn^{-1+2/\beta} EX^\beta, & \text{for } \beta < 2. \end{cases} \quad (3.10)
\end{aligned}$$

If we denote $\delta = \max\{(-1 + 2/p), 2/\alpha, 2/\beta, 1\}$, then we can get by (3.9) and (3.10) that

$$\sum_{i=1}^n E|a_{ni}(Y_i - EY_i)|^2 \leq Cn^\delta. \quad (3.11)$$

It is easily seen that

$$\begin{aligned}
\left(-\frac{1}{p} + \frac{\delta}{2}\right)r &= \max\left\{-\frac{1}{2}, -\frac{1}{\beta}, -\frac{1}{\alpha}, -\frac{1}{p} + \frac{1}{2}\right\}r \\
&= -\min\left\{\frac{1}{2}, \frac{1}{\beta}, \frac{1}{\alpha}, \frac{1}{p} - \frac{1}{2}\right\}r < -1. \quad (3.12)
\end{aligned}$$

Therefore, we have by (3.11) and (3.12) that

$$H_2 \leq C \sum_{i=1}^{\infty} n^{(-1/p + \delta/2)r} < \infty, \quad (3.13)$$

which together with $H_1 < \infty$ yields (3.7).

On the other hand, we will prove that

$$I \doteq n^{-1/p} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} EY_i \right| \rightarrow 0, \quad \text{as } n \rightarrow \infty. \quad (3.14)$$

If $0 < \beta \leq 1$, then we have by Lemma 2.5 and (3.6) that

$$\begin{aligned}
I &\leq n^{-1/p} \sum_{i=1}^n |a_{ni} E Y_i| \\
&\leq n^{-1/p} \sum_{i=1}^n |a_{ni}| [E|X_i| I(|X_i| \leq n^{1/\beta}) + n^{1/\beta} P(|X_i| > n^{1/\beta})] \\
&\leq C n^{-1/p} \sum_{i=1}^n |a_{ni}| [E|X| I(|X| \leq n^{1/\beta}) + n^{1/\beta} P(|X| > n^{1/\beta})] \\
&\leq C n^{-1/p} \sum_{i=1}^n |a_{ni}| [n^{(1-\beta)/\beta} E|X|^\beta I(|X| \leq n^{1/\beta}) + n^{1/\beta-1} E|X|^\beta I(|X| > n^{1/\beta})] \\
&= C n^{-1/\alpha-1} E|X|^\beta \sum_{i=1}^n |a_{ni}| \\
&\leq C n^{-1/\alpha-1+(1\vee 1/\alpha)} \rightarrow 0, \quad \text{as } n \rightarrow \infty.
\end{aligned} \tag{3.15}$$

If $\beta > 1$, then we have by $E X_n = 0$, Lemma 2.5 and (3.6) that

$$\begin{aligned}
I &\leq n^{-1/p} \sum_{i=1}^n |a_{ni} E Y_i| \\
&\leq n^{-1/p} \sum_{i=1}^n |a_{ni}| [E|X_i| I(|X_i| > n^{1/\beta}) + n^{1/\beta} P(|X_i| > n^{1/\beta})] \\
&\leq C n^{-1/p} \sum_{i=1}^n |a_{ni}| E|X| I(|X| > n^{1/\beta}) \\
&\leq C n^{-1/p} \sum_{i=1}^n |a_{ni}| n^{1/\beta-1} E|X|^\beta I(|X| > n^{1/\beta}) \\
&\leq C n^{-1/\alpha-1} E|X|^\beta \sum_{i=1}^n |a_{ni}| \\
&\leq C n^{-1/\alpha-1+(1\vee 1/\alpha)} \rightarrow 0, \quad \text{as } n \rightarrow \infty.
\end{aligned} \tag{3.16}$$

Hence, (3.14) follows from (3.15) and (3.16) immediately.

Finally, we prove $J \rightarrow 0$ a.s. as $n \rightarrow \infty$.

The condition $E|X|^\beta < \infty$ yields that

$$\sum_{n=1}^{\infty} P(Z_n \neq 0) = \sum_{n=1}^{\infty} P(|X_n| > n^{1/\beta}) \leq C \sum_{n=1}^{\infty} P(|X| > n^{1/\beta}) \leq C E|X|^\beta < \infty, \tag{3.17}$$

which implies that $P(Z_n \neq 0, \text{i.o.}) = 0$ by Borel–Cantelli lemma. Hence, we have by (3.1) that

$$\begin{aligned}
J &\doteq n^{-1/p} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} Z_i \right| \leq n^{-1/p} \sum_{i=1}^n |a_{ni} Z_i| \\
&\leq n^{-1/p} \left(\max_{1 \leq i \leq n} |a_{ni}|^\alpha \right)^{1/\alpha} \sum_{i=1}^n |Z_i| \leq n^{-1/p} \left(\sum_{i=1}^n |a_{ni}|^\alpha \right)^{1/\alpha} \sum_{i=1}^n |Z_i| \\
&\leq C n^{-1/\beta} \sum_{i=1}^n |Z_i| \rightarrow 0, \quad \text{a.s., as } n \rightarrow \infty.
\end{aligned} \tag{3.18}$$

Therefore, the desired result (3.2) follows from (3.7), (3.14) and (3.18) immediately. This completes the proof of the Theorem. \square

Taking $a_{ni} \equiv 1$ in Theorem 3.1, then (3.1) is always valid for any $\alpha > 0$. Hence, for any $0 < p < \min(\beta, 2)$, letting $\alpha = p\beta/(\beta-p) > 0$, we can obtain the following corollary.

Corollary 3.2. Let $\{X_n, n \geq 1\}$ be a sequence of NSD identically distributed random variables with $E|X_1|^\beta < \infty$. If $\beta > 1$, further assume that $EX_1 = 0$, then for any $0 < p < \min(\beta, 2)$,

$$n^{-1/p} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} X_i \right| \rightarrow 0 \quad \text{a.s. as } n \rightarrow \infty. \tag{3.19}$$

Remark 3.3. Theorem 3.1 generalizes and improves Theorem A of Bai and Cheng ([2]) for independent and identically distributed random variables to the case of NSD random variables, since Theorem 3.1 removes the identically distributed condition and expands the ranges of α , β , and p , respectively.

Remark 3.4. Comparing Theorem 3.1 with Theorem 2.1 of Wang et al. ([24]), we have the following improvements:

- (i) the moment condition $E|X|^\beta < \infty$ in Theorem 3.1 is weaker than (2.1) in Wang et al ([24]);
- (ii) the condition (3.1) in Theorem 3.1 is weaker than $\sum_{i=1}^n |a_{ni}|^\alpha = O(n^\delta)$ for some $0 < \alpha < 2$ and $0 < \delta < 1$ in Wang et al ([24]).

Theorem 3.5. Let $1 < r < 2$ and $\{X_n, n \geq 1\}$ be a sequence of mean zero NSD random variables, which is stochastically dominated by a random variable X . Let $\{a_n, n \geq 1\}$ be a sequence of positive constants satisfying $A_n \doteq \sum_{k=1}^n a_k \uparrow \infty$. Denote $c_n = A_n/a_n$ for each $n \geq 1$. Assume that

$$E|X|^r < \infty, \tag{3.20}$$

$$N(n) \doteq \text{Card}\{i : c_i \leq n\} = O(n^r), \quad n \geq 1, \tag{3.21}$$

then

$$A_n^{-1} \sum_{k=1}^n a_k X_k \rightarrow 0 \quad \text{a.s., as } n \rightarrow \infty. \tag{3.22}$$

Proof. Let $N(0) = 0$ and denote

$$X_n^{(c_n)} = -c_n I(X_n < -c_n) + X_n I(|X_n| \leq c_n) + c_n I(X_n > c_n), \quad n \geq 1.$$

It follows from (3.20) and (3.21) that

$$\begin{aligned} \sum_{i=1}^{\infty} P\left(X_i \neq X_i^{(c_i)}\right) &= \sum_{i=1}^{\infty} P(|X_i| > c_i) = \sum_{j=1}^{\infty} \sum_{c_i \leq j < c_{i+1}} P(|X_i| > c_i) \\ &\leq C \sum_{j=1}^{\infty} \sum_{j-1 < c_i \leq j} P(|X| > j-1) \\ &= C \sum_{j=1}^{\infty} (N(j) - N(j-1)) P(|X| > j-1) \\ &= C \sum_{j=1}^{\infty} (N(j) - N(j-1)) \sum_{n=j}^{\infty} P(n-1 < |X| \leq n) \\ &= C \sum_{n=1}^{\infty} \sum_{j=1}^n (N(j) - N(j-1)) P(n-1 < |X| \leq n) \\ &\leq C \sum_{n=1}^{\infty} n^r P(n-1 < |X| \leq n) \leq CE|X|^r < \infty. \end{aligned}$$

By the inequality above and Borel–Cantelli lemma, we can get $P(X_i \neq X_i^{(c_i)}, i.o.) = 0$. Therefore, in order to prove (3.22), we only need to prove

$$A_n^{-1} \sum_{i=1}^n a_i X_i^{(c_i)} \rightarrow 0 \quad \text{a.s.}, \quad n \rightarrow \infty. \quad (3.23)$$

By C_r inequality, Lemma 2.5, (3.20) and (3.21) again,

$$\begin{aligned} \sum_{k=1}^{\infty} \text{Var} \left(\frac{a_k X_k^{(c_k)}}{A_k} \right) &\leq \sum_{k=1}^{\infty} c_k^{-2} E(X_k^{(c_k)})^2 \\ &\leq 3 \sum_{k=1}^{\infty} c_k^{-2} E [c_k^2 I(|X_k| > c_k) + X_k^2 I(|X_k| \leq c_k)] \\ &\leq C \sum_{k=1}^{\infty} P(|X| > c_k) + C \sum_{k=1}^{\infty} c_k^{-2} E X^2 I(|X| \leq c_k) \\ &\leq C + C \sum_{j=1}^{\infty} \sum_{j-1 < c_k \leq j} c_k^{-2} E X^2 I(|X| \leq c_k) \\ &\leq C + C \sum_{j=1}^{\infty} \sum_{j-1 < c_k \leq j} c_k^{-2} E X^2 I(|X| \leq j) \end{aligned}$$

$$\begin{aligned}
&\leq C + C \sum_{j=2}^{\infty} (N(j) - N(j-1))(j-1)^{-2} \sum_{k=1}^j EX^2 I(k-1 < |X| \leq k) \\
&\leq C + C \sum_{k=2}^{\infty} \sum_{j=k}^{\infty} (N(j) - N(j-1))(j-1)^{-2} EX^2 I(k-1 < |X| \leq k) \\
&\leq C + C \sum_{k=2}^{\infty} \sum_{j=k}^{\infty} N(j)((j-1)^{-2} - j^{-2}) EX^2 I(k-1 < |X| \leq k) \\
&\leq C + C \sum_{k=2}^{\infty} \sum_{j=k}^{\infty} j^{r-3} EX^2 I(k-1 < |X| \leq k) \\
&\leq C + C \sum_{k=2}^{\infty} k^{r-2} E|X|^r k^{2-r} I(k-1 < |X| \leq k) \\
&= C + C \sum_{k=2}^{\infty} E|X|^r I(k-1 < |X| \leq k) \\
&\leq C + CE|X|^r < \infty.
\end{aligned}$$

Hence, by the inequality above, Lemma 2.4 and Kronecker's lemma, we have

$$A_n^{-1} \sum_{i=1}^n a_i \left(X_i^{(c_i)} - EX_i^{(c_i)} \right) \rightarrow 0 \text{ a.s., } n \rightarrow \infty. \quad (3.24)$$

In order to prove (3.22), it suffices to prove that

$$A_n^{-1} \sum_{i=1}^n a_i EX_i^{(c_i)} \rightarrow 0, \quad n \rightarrow \infty. \quad (3.25)$$

By (3.21), it easily seen that $c_n \rightarrow \infty$ as $n \rightarrow \infty$. Notice that $EX_n = 0$ for each $n \geq 1$, we have

$$|EX_n I(|X_n| \leq c_n)| = |EX_n I(|X_n| > c_n)|.$$

It follows by Lemma 2.5, (3.20) and (3.21) that,

$$\begin{aligned}
\sum_{k=1}^{\infty} \left| \frac{a_k EX_k^{(c_k)}}{A_k} \right| &\leq \sum_{k=1}^{\infty} [P(|X_k| > c_k) + c_k^{-1} |EX_k I(|X_k| \leq c_k)|] \\
&= \sum_{k=1}^{\infty} [P(|X_k| > c_k) + c_k^{-1} |EX_k I(|X_k| > c_k)|] \\
&\leq \sum_{k=1}^{\infty} P(|X_k| > c_k) + \sum_{k=1}^{\infty} c_k^{-1} E|X_k| I(|X_k| > c_k) \\
&\leq C \sum_{k=1}^{\infty} P(|X| > c_k) + C \sum_{k=1}^{\infty} c_k^{-1} E|X| I(|X| > c_k)
\end{aligned}$$

$$\begin{aligned}
&\leq C + C \sum_{k=1}^{\infty} \sum_{c_k \leq j < c_{k+1}} c_k^{-1} E|X| I(|X| > c_k) \\
&\leq C + C \sum_{k=1}^{\infty} \sum_{j-1 < c_k \leq j} c_k^{-1} E|X| I(|X| > j-1) \\
&\leq C + C \sum_{j=2}^{\infty} (N(j) - N(j-1))(j-1)^{-1} \sum_{k=j-1}^{\infty} EXI(k < |X| \leq k+1) \\
&\leq C + C \sum_{k=1}^{\infty} \sum_{j=2}^{k+1} N(j)((j-1)^{-1} - j^{-1}) EXI(k < |X| \leq k+1) \\
&\leq C + C \sum_{k=1}^{\infty} \sum_{j=2}^{k+1} j^{r-2} EXI(k < |X| \leq k+1) \\
&\leq C + C \sum_{k=1}^{\infty} k^{r-1} E|X| I(k < |X| \leq k+1) \\
&\leq C + C \sum_{k=1}^{\infty} E|X|^r I(k < |X| \leq k+1) \leq C + CE|X|^r < \infty.
\end{aligned}$$

By Kronecker's lemma, we can get(3.24) immediately. The proof is complete. \square

ACKNOWLEDGEMENT

The authors are most grateful to the Editor Lucie Fajfrová and two anonymous referees for careful reading of the manuscript and valuable suggestions which helped significantly improving an earlier version of this paper. This work is supported by the National Natural Science Foundation of China (11471272, 11171001, 11201001, 11501004, 11501005), the Natural Science Foundation of Fujian Province (2013J01019) and Anhui Province (1508085J06).

(Received June 5, 2015)

REFERENCES

-
- [1] K. Alam and K.M.L. Saxena: Positive dependence in multivariate distributions. *Commun. Statist. – Theory and Methods* 10 (1981), 12, 1183–1196. DOI:10.1080/03610928108828102
 - [2] Z. D. Bai and P. E. Cheng: Marcinkiewicz strong laws for linear statistics. *Statist. Probab. Lett.* 46 (2000), 2, 105–112. DOI:10.1016/s0167-7152(99)00093-0
 - [3] H. M. Block, T. H. Savits, and M. Shaked: Some concepts of negative dependence. *Ann. Probab.* 10 (1982), 3, 765–772. DOI:10.1214/aop/1176993784
 - [4] K. Budsaba, P. Chen, K. Panishkan, and A. Volodin: Strong laws for weighted sums and certain types U-statistics based on negatively associated random variables. *Siberian Advances in Mathematics* 19 (2009), 4, 225–232. DOI:10.3103/s1055134409040014
 - [5] T.C. Christofides and E. Vaggelatou: A connection between supermodular ordering and positive/negative association. *J. Multivariate Anal.* 88 (2004), 1, 138–151. DOI:10.1016/s0047-259x(03)00064-2

- [6] N. Eghbal, M. Amini, and A. Bozorgnia: Some maximal inequalities for quadratic forms of negative superadditive dependence random variables. *Statist. Probab. Lett.* *80* (2010), 7, 587–591. DOI:10.1016/j.spl.2009.12.014
- [7] N. Eghbal, M. Amini, and A. Bozorgnia: On the Kolmogorov inequalities for quadratic forms of dependent uniformly bounded random variables. *Statist. Probab. Lett.* *81* (2011), 8, 1112–1120. DOI:10.1016/j.spl.2011.03.005
- [8] M. Gerasimov, V. Kruglov, and A. Volodin: On negatively associated random variables. *Lobachevskii J. Math.* *33* (2012), 1, 47–55. DOI:10.1134/s1995080212010052
- [9] S. H. Hu, X. T. Liu, X. H. Wang, and X. T. Li: Strong law of large numbers of partial sums for pairwise NQD sequences. *J. Math. Res. Appl.* *33* (2013), 1, 111–116.
- [10] T. Z. Hu: Negatively superadditive dependence of random variables with applications. *Chinese J. App. Probab. Statist.* *16* (2000), 133–144.
- [11] K. Joag-Dev and F. Proschan: Negative association of random variables with applications. *Ann. Statist.* *11* (1983), 1, 286–295. DOI:10.1214/aos/1176346079
- [12] B. Y. Jing and H. Y. Liang: Strong limit theorems for weighted sums of negatively associated random variables. *J. Theoret. Probab.* *21* (2008), 4, 890–909. DOI:10.1007/s10959-007-0128-4
- [13] P. Matula: A note on the almost sure convergence of sums of negatively dependent random variables. *Statistics and Probability Letters*, *15* (1992), 209–213. DOI:10.1016/0167-7152(92)90191-7
- [14] Y. J. Meng and Z. Y. Lin: Strong laws of large numbers for $\tilde{\rho}$ -mixing random variables. *J. Math. Anal. Appl.* *365* (2010), 711–717. DOI:10.1016/j.jmaa.2009.12.009
- [15] Q. M. Shao: A comparison theorem on moment inequalities between negatively associated and independent random variables. *J. Theoret. Probab.* *13* (2000), 2, 343–356.
- [16] A. T. Shen and R. C. Wu: Strong and weak convergence for asymptotically almost negatively associated random variables. *Discrete Dynamics in Nature and Society* *2013* (2013), 1–7. DOI:10.1155/2013/235012
- [17] A. T. Shen: On the strong convergence rate for weighted sums of arrays of rowwise negatively orthant dependent random variables. *RACSAM* *107* (2013), 2, 257–271. DOI:10.1007/s13398-012-0067-5
- [18] A. T. Shen: On strong convergence for weighted sums of a class of random variables. *Abstract Appl. Anal.* *2013* (2013), 1–7. DOI:10.1155/2013/216236
- [19] A. T. Shen, Y. Zhang, and A. Volodin: Applications of the Rosenthal-type inequality for negatively superadditive dependent random variables. *Metrika* *78* (2015), 295–311. DOI:10.1007/s00184-014-0503-y
- [20] Y. Shen, X. J. Wang, W. Z. Yang, and S. H. Hu: Almost sure convergence theorem and strong stability for weighted sums of NSD random variables. *Acta Mathematica Sinica, English Series*, *29* (2013), 4, 743–756. DOI:10.1007/s10114-012-1723-6
- [21] Y. Shen, X. J. Wang, and S. H. Hu: On the strong convergence and some inequalities for negatively superadditive dependent sequences. *J. Inequalities Appl.* *2013* (2013), 1, 448. DOI:10.1186/1029-242x-2013-448
- [22] S. H. Sung: On the strong law of large numbers for weighted sums of random variables. *Computers Math. Appl.* *62* (11) (2011), 4277–4287. DOI:10.1016/j.camwa.2011.10.018

- [23] X. J. Wang, X. Q. Li, S. H. Hu, and W. Z. Yang: Strong limit theorems for weighted sums of negatively associated random variables. *Stochast. Anal. Appl.* *29* (2011), 1, 1–14. DOI:10.1080/07362994.2010.515484
- [24] X. J. Wang, S. H. Hu, and W. Z. Yang: Complete convergence for arrays of rowwise negatively orthant dependent random variables. *RACSAM* *106* (2012), 2, 235–245. DOI:10.1007/s13398-011-0048-0
- [25] X. J. Wang, X. Deng, L. L. Zheng, and S. H. Hu: Complete convergence for arrays of rowwise negatively superadditive-dependent random variables and its applications. *Statistics* *48* (4) (2014), 834–850. DOI:10.1080/02331888.2013.800066
- [26] X. J. Wang, A. T. Shen, Z. Y. Chen, and S. H. Hu: Complete convergence for weighted sums of NSD random variables and its application in the EV regression model. *Test* *24* (2015), 166–184. DOI:10.1007/s11749-014-0402-6
- [27] Q. Y. Wu: *Probability Limit Theory for Mixing Sequence*. Science Press of China, Beijing 2006.
- [28] Q. Y. Wu and Y. Y. Jiang: A law of the iterated logarithm of partial sums for NA random variables. *J. Korean Statist. Soc.* *39* (2010), 199–206. DOI:10.1016/j.jkss.2009.06.001
- [29] Q. Y. Wu and Y. Y. Jiang: Chover’s law of the iterated logarithm for negatively associated sequences. *J. Systems Sci. Complex.* *23* (2010), 293–302. DOI:10.1007/s11424-010-7258-y
- [30] W. Z. Yang, S. H. Hu, X. J. Wang, and Q. C. Zhang: Berry-Esséen bound of sample quantiles for negatively associated sequence. *J. Inequalities Appl.* *2011* (2011), 1, 83. DOI:10.1186/1029-242x-2011-83

Zhiyong Chen, School of Mathematical Sciences, Xiamen University, Xiamen 361005. P. R. China.

e-mail: zyichen1024@163.com

Haibin Wang, School of Mathematical Sciences, Xiamen University, Xiamen 361005. P. R. China.

e-mail: whb@xmu.edu.cn

Xuejun Wang, Corresponding author. School of Mathematical Sciences, Anhui University, Hefei, 230601. P. R. China.

e-mail: wxjahdx2000@126.com

Shuhe Hu, School of Mathematical Sciences, Anhui University, Hefei, 230601. P. R. China.

e-mail: hushuhe@263.net