

# Applications of Mathematics

---

Lam Quoc Anh; Dinh Vinh Hien

On well-posedness for parametric vector quasiequilibrium problems with moving cones

*Applications of Mathematics*, Vol. 61 (2016), No. 6, 651–668

Persistent URL: <http://dml.cz/dmlcz/145914>

## Terms of use:

© Institute of Mathematics AS CR, 2016

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON WELL-POSEDNESS FOR PARAMETRIC VECTOR  
QUASIEQUILIBRIUM PROBLEMS WITH MOVING CONES

LAM QUOC ANH, Cantho, DINH VINH HIEN, Hochiminh

(Received August 31, 2015)

*Abstract.* In this paper we consider weak and strong quasiequilibrium problems with moving cones in Hausdorff topological vector spaces. Sufficient conditions for well-posedness of these problems are established under relaxed continuity assumptions. All kinds of well-posedness are studied: (generalized) Hadamard well-posedness, (unique) well-posedness under perturbations. Many examples are provided to illustrate the essentialness of the imposed assumptions. As applications of the main results, sufficient conditions for lower and upper bounded equilibrium problems and elastic traffic network problems to be well-posed are derived.

*Keywords:* quasiequilibrium problem; lower bounded equilibrium problem; upper bounded equilibrium problem; network traffic problem; well-posedness;  $C$ -upper semicontinuity;  $C$ -lower semicontinuity

*MSC 2010:* 49K40, 90C31, 91B50

## 1. INTRODUCTION

The equilibrium problem was introduced by Blum and Oettli [16] in 1994. The mathematical formulation of the problem unifies various important problems related to optimization, namely, constrained minimization, variational inequality, complementarity problem, Nash equilibria, minimax problem, fixed-point and coincidence-point problems, traffic network problem, etc. Due to the important role of the problem, it has been intensively studied for all of the main topics, such as existence theory [10], [14], [18], [22], [25], [28], [31], [35], [38], [40], stability and sensitivity analysis theory [1], [3], [4], [5], [6], [13], [15], and solution methods [17], [32], [37], [42], etc.

---

This research is funded by Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number 101.01-2014.44.

Well-posedness may be understood in two ways. The first concept is the well-posedness in the sense of Hadamard [27]. This property includes the existence, uniqueness and continuous dependence of the optimal solution and optimal value on the perturbed data of the problem. The second concept was introduced by Tikhonov in 1966 in [43], where the author proposed it as the existence and uniqueness of the solution and convergence of each minimizing sequence to the solution. Well-posedness under perturbations (termed also parametric well-posedness) is a generalized notion of those of Hadamard and Tikhonov. Recently, the study of generalized well-posed concepts was extended to mathematical programming [30], optimization problem (see e.g., [39], [47]–[49]), variational inequality [24], Nash equilibria [34]. Well-posedness for equilibrium problems is a theme of great importance and has received increasing attention of many researchers recently. The sufficient and necessary conditions and metric characterizations of some types of generalized well-posedness for the class of equilibrium problems were considered. For more details, we refer the reader to [7], [8], [9], [12], [23], [29], [33], and the references therein.

Optimization-related problems with moving cones play an important role in many practical situations. For example, when we want to choose a product from the market, of course, the characteristics of the chosen product have to satisfy our requirements, the so-called requirement domain. Naturally, the higher quality of the product is the better choice, and hence the requirement domain is a cone which is called the requirement cone. Since our requirement cones may be different from those of the other ones, the requirement cone will depend on some parameters.

Motivated and inspired by the above observations, in this paper we consider the weak and strong quasiequilibrium problems with moving cones. We introduce the concepts of moving cone semicontinuity and investigate their properties. Under the moving cone semicontinuity assumptions, we establish sufficient and/or necessary conditions for such problems to be (uniquely) well-posed under perturbations and (generalized) Hadamard well-posed. Moreover, since we consider the problems with moving cones, the main results in [7], [8], [9], [12], [23], [29], [33] cannot be applied to such problems. Therefore, our results are new and different from the existing ones in the literature.

The rest of the paper is organized as follows. Section 2 is devoted to problem statements and preliminary facts. The concepts of moving cone semicontinuity and their properties are also introduced and discussed in this section. We study, in Section 3, the necessary and/or sufficient conditions for the weak and strong vector quasiequilibrium problems with moving cones to be (uniquely) well-posed under perturbations and (generalized) Hadamard well-posed. The applications of the main results to some special cases of quasiequilibrium problems are presented in Section 4.

## 2. PRELIMINARIES

Let  $X, \Lambda$  be two metric spaces, and  $Y$  a Hausdorff topological vector space. Let  $K: X \times \Lambda \rightrightarrows X$  and  $C: \Lambda \rightrightarrows Y$  be two set-valued mappings such that, for each  $\lambda \in \Lambda$ ,  $C(\lambda)$  is a closed, convex, pointed and solid cone. Let  $e: \Lambda \rightarrow Y$  be a continuous mapping such that, for each  $\lambda \in \Lambda$ ,  $e(\lambda) \in \text{int } C(\lambda)$ . Let  $f: X \times X \times \Lambda \rightarrow Y$  be a vector valued mapping. For each  $\lambda \in \Lambda$ , we consider the following parametric vector quasiequilibrium problems:

(WQEP $_{\lambda}$ ) Find  $\bar{x} \in K(\bar{x}, \lambda)$  such that, for each  $y \in K(\bar{x}, \lambda)$ ,

$$f(\bar{x}, y, \lambda) \in (Y \setminus -\text{int } C(\lambda)).$$

(SQEP $_{\lambda}$ ) Find  $\bar{x} \in K(\bar{x}, \lambda)$  such that, for each  $y \in K(\bar{x}, \lambda)$ ,

$$f(\bar{x}, y, \lambda) \in C(\lambda).$$

Instead of writing  $\{(WQEP_{\lambda}): \lambda \in \Lambda\}$  and  $\{(SQEP_{\lambda}): \lambda \in \Lambda\}$  for the sets of such problems, we will simply write (WQEP) and (SQEP), respectively, in the sequel. For each  $\lambda \in \Lambda$ , the solution sets of (WQEP $_{\lambda}$ ) and (SQEP $_{\lambda}$ ) are denoted by  $S^w(\lambda)$  and  $S^s(\lambda)$ , respectively, i.e.,

$$S^w(\lambda) = \{x \in K(x, \lambda): f(x, y, \lambda) \in (Y \setminus -\text{int } C(\lambda)) \quad \forall y \in K(x, \lambda)\},$$

$$S^s(\lambda) = \{x \in K(x, \lambda): f(x, y, \lambda) \in C(\lambda) \quad \forall y \in K(x, \lambda)\}.$$

**Definition 2.1.** Let  $Q$  be a set-valued mapping from  $X$  into  $Y$ .

- (i)  $Q$  is said to be upper semicontinuous (usc, shortly) at  $x_0$  if for any open superset  $U$  of  $Q(x_0)$ , there is a neighborhood  $N$  of  $x_0$  such that  $Q(N) \subset U$ .
- (ii)  $Q$  is said to be lower semicontinuous (lsc, shortly) at  $x_0$  if for any open subset  $U$  of  $Y$  with  $Q(x_0) \cap U \neq \emptyset$ , there is a neighborhood  $N$  of  $x_0$  such that for all  $x \in N$ ,  $Q(x) \cap U \neq \emptyset$ .

The mapping  $Q$  is said to be continuous at  $x_0$  if it is both usc and lsc at  $x_0$ . We say that  $Q$  satisfies a certain property on a subset  $A \subset X$  if  $Q$  satisfies it at every points of  $A$ . If  $A = X$  we omit “on  $X$ ” in the statement.

**Lemma 2.1** ([11]).

- (i)  $Q$  is usc at  $x_0$ , if for each superset  $U$  of  $Q(x_0)$  and for every sequence  $\{x_n\}$  in  $X$  converging to  $x_0$ , there is  $n_0$  such that for all  $n \geq n_0$ ,  $Q(x_n) \subset U$ .
- (ii)  $Q$  is lsc at  $x_0$  if, for each sequence  $x_n \rightarrow x_0$  and  $y_0 \in Q(x_0)$ , there exists  $y_n \in Q(x_n)$  such that  $y_n \rightarrow y_0$ .

**Definition 2.2.** Let  $Q$  be a set-valued mapping from  $X$  into  $Y$ .

- (i)  $Q$  is said to be Hausdorff upper semicontinuous ( $H$ -usc, shortly) at  $x_0$  if, for each neighborhood  $B$  of the origin  $\theta_Y$  in  $Y$ , there exists a neighborhood  $N$  of  $x_0$  such that,  $Q(x) \subset Q(x_0) + B$  for every  $x \in N$ .
- (ii)  $Q$  is said to be Hausdorff lower semicontinuous ( $H$ -lsc, shortly) at  $x_0$  if, for each neighborhood  $B$  of the origin  $\theta_Y$  in  $Y$ , there exists a neighborhood  $N$  of  $x_0$  such that,  $Q(x_0) \subset Q(x) + B$  for every  $x \in N$ .

$Q$  is said to be Hausdorff continuous at  $x_0$  if it is both  $H$ -usc and  $H$ -lsc at  $x_0$ .

Picking up the main ideas from [18], we introduce the concepts of moving cone upper and lower semicontinuity as follows.

**Definition 2.3.** Let  $X, \Lambda$  be two topological spaces,  $Y$  a topological vector space, and  $C: \Lambda \rightrightarrows Y$  a set-valued mapping with pointed solid convex cone values. Let  $g: X \times \Lambda \rightarrow Y$  be a vector valued mapping.

- (i)  $g$  is said to be  $C$ -upper semicontinuous ( $C$ -usc, for short) at  $(x_0, \lambda_0)$  if, for any neighborhood  $V$  of  $\theta_Y$  in  $Y$ , there exists a neighborhood  $U$  of  $(x_0, \lambda_0)$  such that, for all  $(x, \lambda) \in U$ ,

$$g(x, \lambda) \in g(x_0, \lambda_0) + V - C(\lambda_0).$$

- (ii)  $g$  is said to be  $C$ -lower semicontinuous ( $C$ -lsc, for short) if  $-g$  is  $C$ -upper semicontinuous.

The mapping  $g$  is said to be  $C$ -continuous at  $(x_0, \lambda_0)$  if it is both  $C$ -usc and  $C$ -lsc at  $(x_0, \lambda_0)$ . We say that  $g$  satisfies a certain property on a subset  $A \subset X \times \Lambda$  if  $g$  satisfies it at every point of  $A$ . If  $A = X \times \Lambda$  we omit "on  $X \times \Lambda$ " in the statement.

**Proposition 2.1.** Let  $X, \Lambda, C, g$  be as in Definition 2.3. The following conditions are equivalent to each other.

- (i)  $g$  is  $C$ -upper semicontinuous.
- (ii) For each  $(x_0, \lambda_0) \in X \times \Lambda$  and  $d \in \text{int } C(\lambda_0)$ , there is a neighborhood  $U$  of  $(x_0, \lambda_0)$  such that  $g(x, \lambda) \in g(x_0, \lambda_0) + d - \text{int } C(\lambda_0)$ , for all  $(x, \lambda) \in U$ .
- (iii) For each  $(x_0, \lambda_0) \in X \times \Lambda$  and  $a \in Y$ ,  $g^{-1}(a - \text{int } C(\lambda_0))$  is open.

**Proof.** First, we show that (i) and (ii) are equivalent to each other. Let  $(x_0, \lambda_0) \in X \times \Lambda$  and  $d \in \text{int } C(\lambda_0)$ . Let  $V = d - \text{int } C(\lambda_0)$ . Then  $V$  is a neighborhood of  $\theta_Y$  in  $Y$ . If (i) is satisfied, then there is a neighborhood  $U$  of  $(x_0, \lambda_0)$  such that

$$g(x, \lambda) \in g(x_0, \lambda_0) + V - C(\lambda_0) \quad \forall (x, \lambda) \in U.$$

Since  $C(\lambda_0)$  is a convex cone,  $-\text{int } C(\lambda_0) - C(\lambda_0) = -\text{int } C(\lambda_0)$ , we conclude that

$$g(x, \lambda) \in g(x_0, \lambda_0) + d - \text{int } C(\lambda_0) \quad \forall (x, \lambda) \in U,$$

i.e., (ii) holds. Conversely, let  $(x_0, \lambda_0) \in X \times \Lambda$  and let  $V$  be a neighborhood of  $\theta_Y$ . Since  $C(\lambda_0)$  is a pointed solid cone, there is  $d \in \text{int } C(\lambda_0)$  such that  $d \in V$ . If (ii) is satisfied, there exists a neighborhood  $U$  of  $(x_0, \lambda_0)$  such that

$$g(x, \lambda) \in g(x_0, \lambda_0) + d - \text{int } C(\lambda_0) \quad \forall (x, \lambda) \in U,$$

and hence,  $g(x, \lambda) \in g(x_0, \lambda_0) + V - C(\lambda_0)$  for all  $(x, \lambda) \in U$ , i.e., (i) holds.

Next, we prove that (ii) and (iii) are equivalent to each other. Assume that, for each  $(x_0, \lambda_0) \in X \times \Lambda$  and  $a \in Y$ ,  $g^{-1}(a - \text{int } C(\lambda_0))$  is open. Let  $d \in \text{int } C(\lambda_0)$ , we have  $(x_0, \lambda_0) \in g^{-1}(g(x_0, \lambda_0) + d - \text{int } C(\lambda_0))$ , which is open. Thus, there is a neighborhood  $U$  of  $(x_0, \lambda_0)$  such that  $U \subset g^{-1}(g(x_0, \lambda_0) + d - \text{int } C(\lambda_0))$ , i.e.,

$$g(x, \lambda) \in g(x_0, \lambda_0) + d - \text{int } C(\lambda_0) \quad \forall (x, \lambda) \in U.$$

Conversely, for each  $a \in Y$  and  $(x_0, \lambda_0) \in g^{-1}(a - \text{int } C(\lambda_0))$ , let  $d = a - g(x_0, \lambda_0)$ . Since  $d \in \text{int } C(\lambda_0)$ , there is a neighborhood  $U$  of  $(x_0, \lambda_0)$  such that

$$g(x, \lambda) \in g(x_0, \lambda_0) + d - \text{int } C(\lambda_0) \quad \forall (x, \lambda) \in U,$$

i.e.,  $U \subset g^{-1}(g(x_0, \lambda_0) + d - \text{int } C(\lambda_0))$ . Hence,  $g^{-1}(a - \text{int } C(\lambda_0))$  is open.  $\square$

Next, we propose other properties of the  $C$ -upper semicontinuity.

**Proposition 2.2.** *Let  $X, \Lambda, C$  be as in Definition 2.3 and let  $f, g: X \times \Lambda \rightarrow Y$  be vector valued mappings. If  $f$  and  $g$  are  $C$ -upper semicontinuous, then*

- (i)  $f + g$  is  $C$ -upper semicontinuous;
- (ii)  $kf$  is  $C$ -upper semicontinuous, for each  $k \in (0, \infty)$ .

*Proof.* (i) For each  $(x_0, \lambda_0) \in X \times \Lambda$  and  $d \in \text{int } C(\lambda_0)$ , since  $f$  and  $g$  are  $C$ -usc, there are two neighborhoods  $U_1$  and  $U_2$  of  $(x_0, \lambda_0)$  such that

$$\begin{aligned} f(x, \lambda) &\in f(x_0, \lambda_0) + \frac{1}{2}d - \text{int } C(\lambda_0) \quad \forall (x, \lambda) \in U_1, \\ g(x, \lambda) &\in g(x_0, \lambda_0) + \frac{1}{2}d - \text{int } C(\lambda_0) \quad \forall (x, \lambda) \in U_2. \end{aligned}$$

Therefore,

$$(f + g)(x, \lambda) \in (f + g)(x_0, \lambda_0) + d - \text{int } C(\lambda_0) \quad \forall (x, \lambda) \in U = U_1 \cap U_2.$$

Applying Proposition 2.2, we conclude that  $f + g$  is  $C$ -upper semicontinuous.

- (ii) The  $C$ -upper semicontinuity of  $kf$  can be proved similarly.  $\square$

Passing to the  $C$ -lower semicontinuity, we also establish conclusions similar to those in Propositions 2.1 and 2.2.

**Proposition 2.3.** Let  $X, \Lambda, C, g$  be as in Definition 2.3. The following conditions are equivalent to each other.

- (i)  $g$  is  $C$ -lower semicontinuous.
- (ii) For each  $(x_0, \lambda_0) \in X \times \Lambda$  and  $d \in \text{int } C(\lambda_0)$ , there is a neighborhood  $U$  of  $(x_0, \lambda_0)$  such that  $g(x, \lambda) \in g(x_0, \lambda_0) - d + \text{int } C(\lambda_0)$  for all  $(x, \lambda) \in U$ .
- (iii) For each  $(x_0, \lambda_0) \in X \times \Lambda$  and  $a \in Y$ ,  $g^{-1}(a + \text{int } C(\lambda_0))$  is open.

**Proposition 2.4.** Let  $X, \Lambda, C, f, g$  be as in Proposition 2.2. If  $f$  and  $g$  are  $C$ -lower semicontinuous, then

- (i)  $f + g$  is  $C$ -lower semicontinuous;
- (ii)  $kf$  is  $C$ -lower semicontinuous for each  $k \in (0, \infty)$ .

If  $X = \Lambda = Y = \mathbb{R}$ ,  $C(\lambda) \equiv \mathbb{R}_+$ , and  $g(x, \lambda) \equiv g(x)$ , then Propositions 2.1 and 2.3 imply that the  $C$ -upper (lower) semicontinuity of  $g$  reduces to the ordinary upper (lower, respectively) semicontinuity of such function.

**Definition 2.4.** Let  $\lambda \in \Lambda$  and let  $\{\lambda_n\} \subset \Lambda$  be a sequence converging to  $\lambda$ . A sequence  $\{x_n\}, x_n \in K(x_n, \lambda_n)$ , is said to be an approximating sequence for  $(\text{WQEP}_\lambda)$  or  $(\text{SQEP}_\lambda)$  corresponding to  $\{\lambda_n\}$ , if there exists a sequence  $\{\varepsilon_n\} \downarrow 0$  such that, for each  $n \in \mathbb{N}$  and  $y \in K(x_n, \lambda_n)$ ,

$$f(x_n, y, \lambda_n) + \varepsilon_n e(\lambda_n) \in (Y \setminus -\text{int } C(\lambda_n)),$$

or  $f(x_n, y, \lambda_n) + \varepsilon_n e(\lambda_n) \in C(\lambda_n)$ , respectively.

**Definition 2.5.** The problem  $(\text{WQEP})$  or  $(\text{SQEP})$  is said to be well-posed under perturbations (well-posed, shortly), if for each  $\lambda \in \Lambda$ ,

- (i)  $(\text{WQEP}_\lambda)$  or  $(\text{SQEP}_\lambda)$  has solutions;
- (ii) for any sequence  $\{\lambda_n\} \subset \Lambda$  converging to  $\lambda$ , every approximating sequence  $\{x_n\}$  for  $(\text{WQEP}_\lambda)$  or  $(\text{SQEP}_\lambda)$  corresponding to  $\{\lambda_n\}$  has a subsequence converging to an element in  $S^w(\lambda)$  or  $S^s(\lambda)$ , respectively.

The problem  $(\text{WQEP})$  or  $(\text{SQEP})$  is said to be uniquely well-posed under perturbations (uniquely well-posed, shortly) if for each  $\lambda \in \Lambda$  the solution set  $S^w(\lambda)$  or  $S^s(\lambda)$  is a singleton, and every approximating sequence for  $(\text{WQEP}_\lambda)$  or  $(\text{SQEP}_\lambda)$  tends to the unique solution, respectively.

**Definition 2.6.** The problem  $(\text{WQEP})$  or  $(\text{SQEP})$  is said to be generalized Hadamard well-posed if, for each  $\lambda \in \Lambda$ ,

- (i)  $(\text{WQEP}_\lambda)$  or  $(\text{SQEP}_\lambda)$  has solutions;
- (ii) for any sequence  $\{\lambda_n\} \subset \Lambda$  converging to  $\lambda$  and  $x_n \in S^w(\lambda_n)$  or  $x_n \in S^s(\lambda_n)$ ,  $\{x_n\}$  has a subsequence converging to some point of  $S^w(\lambda)$  or  $S^s(\lambda)$ , respectively.

The problem (WQEP) or (SQEP) is said to be Hadamard well-posed if for each  $\lambda \in \Lambda$  the solution set  $S^w(\lambda)$  or  $S^s(\lambda)$  is a singleton, and every  $x_n \in S^w(\lambda_n)$  or  $x_n \in S^s(\lambda_n)$ ,  $\{x_n\}$  converges to the unique solution, respectively.

For each  $\lambda \in \Lambda$ ,  $\varepsilon \geq 0$ , we denote the  $\varepsilon$ -solution sets of (WQEP $_\lambda$ ) and (SQEP $_\lambda$ ) by  $\widetilde{S}^w(\lambda, \varepsilon)$  and  $\widetilde{S}^s(\lambda, \varepsilon)$ , respectively, defined as follows:

$$\begin{aligned} \widetilde{S}^w(\lambda, \varepsilon) &= \{x \in K(x, \lambda) : f(x, y, \lambda) + \varepsilon e(\lambda) \in (Y \setminus -\text{int } C(\lambda)) \quad \forall y \in K(x, \lambda)\}, \\ \text{and } \widetilde{S}^s(\lambda, \varepsilon) &= \{x \in K(x, \lambda) : f(x, y, \lambda) + \varepsilon e(\lambda) \in C(\lambda) \quad \forall y \in K(x, \lambda)\}. \end{aligned}$$

**Lemma 2.2** ([9]). *Let  $S: X \rightrightarrows Y$  be a set-valued mapping between two topological spaces. Then the following assertions hold.*

- (i) *If  $S(\bar{x})$  is compact, then  $S$  is usc at  $\bar{x}$  if and only if for any sequence  $\{x_n\}$  convergent to  $\bar{x}$  and  $y_n \in S(x_n)$ , there is a subsequence  $\{y_{n_k}\}$  convergent to some  $y \in S(\bar{x})$ .*
- (ii) *If, in addition,  $S(\bar{x}) = \{\bar{y}\}$  is a singleton, then the above limit point  $y$  must be  $\bar{y}$  and the whole  $\{y_n\}$  converges to  $\bar{y}$ .*

### 3. MAIN RESULTS

Since the existence theory of the equilibrium problems has been intensively studied, in this paper we only focus on the well-posedness of such problems, and hence we always assume that the solutions of the problems considered exist in a neighborhood of the reference point.

**Theorem 3.1.** *Assume that, for given  $\lambda \in \Lambda$ , the following assumptions hold:*

- (i)  *$f(\cdot, \cdot, \lambda)$  is  $C$ -upper semicontinuous;*
- (ii)  *$K(\cdot, \lambda)$  is continuous with compact values.*

*Then the solution sets  $S^w(\lambda)$  and  $S^s(\lambda)$  are closed in  $X$ .*

*Proof.* Since the techniques of the proofs are similar, we discuss only the closedness of  $S^w(\lambda)$ . Let an arbitrary sequence  $\{x_n\} \subset S^w(\lambda)$  such that  $x_n \rightarrow \bar{x} \in X$ . We need to show that  $\bar{x}$  belongs to  $S^w(\lambda)$ . Since  $x_n$  is a solution of (WQEP $_\lambda$ ) for each  $n \in \mathbb{N}$ ,  $x_n \in K(x_n, \lambda)$ , and

$$(3.1) \quad f(x_n, y, \lambda) \in (Y \setminus -\text{int } C(\lambda)) \quad \forall y \in K(x_n, \lambda).$$

Since  $K(\cdot, \lambda)$  is usc with compact values,  $\bar{x} \in K(\bar{x}, \lambda)$ . If  $\bar{x} \notin S^w(\lambda)$ , then there exists  $y_0 \in K(\bar{x}, \lambda)$  such that

$$f(\bar{x}, y_0, \lambda) \in -\text{int } C(\lambda).$$



Due to the openness of  $-\text{int } C(\lambda)$ , there is a neighborhood  $V$  of  $\theta_Y$  such that  $(f(\bar{x}, y_0, \lambda) + V) \subset -\text{int } C(\lambda)$ . Since  $f(\cdot, \cdot, \lambda)$  is  $C$ -usc at  $(\bar{x}, y_0)$ , and  $x_n \rightarrow \bar{x}$ , there exist a neighborhood  $U$  of  $y_0$  and  $n_0 \in \mathbb{N}$  such that for all  $y \in U$  and  $n \geq n_0$  we have

$$(3.2) \quad f(x_n, y, \lambda) \in f(\bar{x}, y_0, \lambda) + V - C(\lambda) \subset -\text{int } C(\lambda) - C(\lambda) = -\text{int } C(\lambda).$$

Due to  $y_0 \in K(\bar{x}, \lambda)$ ,  $K(\bar{x}, \lambda) \cap U \neq \emptyset$ . Since  $K(\cdot, \lambda)$  is lsc at  $\bar{x}$ , there exists  $n_1 \in \mathbb{N}$  such that

$$(3.3) \quad K(x_n, \lambda) \cap U \neq \emptyset \quad \forall n \geq n_1.$$

Choose up  $n \geq \max\{n_0, n_1\}$ . Combining (3.2) with (3.3), we infer the existence of  $\bar{y} \in K(x_n, \lambda) \cap U$  such that

$$f(x_n, \bar{y}, \lambda) \in -\text{int } C(\lambda),$$

which is in contradiction with (3.1). Therefore,  $\bar{x} \in S^w(\lambda)$ . □

Now we pass to the semicontinuity results of the approximating solution sets, which play an important role in the well-posedness for the corresponding problems.

**Theorem 3.2.** *Assume that  $X$  is compact,  $f$  is  $C$ -upper semicontinuous, and  $K$  is continuous with compact values. Then*

- (a)  $\widetilde{S}^w$  is upper semicontinuous with compact values on  $\Lambda \times \mathbb{R}_+$  if for given  $\lambda \in \Lambda$  and an arbitrary sequence  $\{\lambda_n\} \subset \Lambda$  converging to  $\lambda$  there exists  $\lambda_{n_0}$  such that  $C(\lambda) \subset C(\lambda_{n_0})$ .
- (b)  $\widetilde{S}^s$  is upper semicontinuous with compact values on  $\Lambda \times \mathbb{R}_+$  if for given  $\lambda \in \Lambda$  and an arbitrary sequence  $\{\lambda_n\} \subset \Lambda$  converging to  $\lambda$  there exists  $\lambda_{n_0}$  such that  $C(\lambda_{n_0}) \subset C(\lambda)$ .

*Proof.* As an example we present only the proof for (b). We first prove the upper semicontinuity with compact values of  $S^s(\cdot) = \widetilde{S}^s(\cdot, 0)$  on  $\Lambda$ . For each  $\lambda \in \Lambda$ , Theorem 3.1 yields that  $S^s(\lambda)$  is a closed subset of  $X$ , so  $S^s(\lambda)$  is compact. We show that  $S^s$  is usc on  $\Lambda$ . Assume, to arrive at a contradiction, that there exists  $\lambda \in \Lambda$  such that  $S^s$  is not usc at  $\lambda$ . Then there are an open superset  $V$  of  $S^s(\lambda)$  and a sequence  $\{\lambda_n\} \subset \Lambda$ ,  $\lambda_n \rightarrow \lambda$  such that, for each  $n \in \mathbb{N}$ , there exists  $x_n \in S^s(\lambda_n) \setminus V$ . Since  $x_n \in S^s(\lambda_n)$ ,  $x_n \in K(x_n, \lambda_n)$  and

$$f(x_n, y, \lambda_n) \in C(\lambda_n) \quad \forall y \in K(x_n, \lambda_n).$$

By the compactness of  $X$ , we can assume that  $x_n$  converges to  $\bar{x}$  for some  $\bar{x} \in X$ . Since  $K$  is usc with compact values,  $K$  is closed, and hence  $\bar{x} \in K(\bar{x}, \lambda)$ . Since  $x_n \notin V$  for all  $n$ ,  $\bar{x}$  does not belong to  $S^s(\lambda) \subset V$ , i.e., there exists  $y_0 \in K(\bar{x}, \lambda)$  such that

$$f(\bar{x}, y_0, \lambda) \in (Y \setminus C(\lambda)),$$

and hence, there is a neighborhood  $B$  of  $\theta_Y$  such that

$$(f(\bar{x}, y_0, \lambda) + B) \subset (Y \setminus C(\lambda)).$$

Since  $f$  is  $C$ -usc at  $(\bar{x}, y_0, \lambda)$ , there is a neighborhood  $U$  of  $y_0$  and  $n_0 \in \mathbb{N}$  such that for all  $y \in U$  and  $n \geq n_0$ ,

$$f(x_n, y, \lambda_n) \in f(\bar{x}, y_0, \lambda) + B - C(\lambda) \subset (Y \setminus C(\lambda)) - C(\lambda) \subset (Y \setminus C(\lambda)).$$

Since  $y_0 \in K(\bar{x}, \lambda)$ , we have  $K(\bar{x}, \lambda) \cap U \neq \emptyset$ . By the lower semicontinuity of  $K$  at  $(\bar{x}, \lambda)$ , there exists  $n_1 \in \mathbb{N}$  such that for all  $n \geq n_1$ ,

$$K(x_n, \lambda_n) \cap U \neq \emptyset.$$

Applying the assumption in (b), we obtain the existence of  $n_2$ , where  $n_2 \geq \max\{n_0, n_1\}$ ,  $C(\lambda_{n_2}) \subset C(\lambda)$ , and hence

$$(Y \setminus C(\lambda)) \subset (Y \setminus C(\lambda_{n_2})).$$

Therefore,  $f(x_{n_2}, y, \lambda_{n_2}) \notin C(\lambda_{n_2})$  for some  $y \in K(x_{n_2}, \lambda_{n_2}) \cap U$ , which is impossible as  $x_{n_2} \in S^s(\lambda_{n_2})$ . Hence,  $S^s$  is usc at  $\lambda$ .

Now we show that  $\widetilde{S^s}$  is usc with compact values on  $\Lambda \times \mathbb{R}_+$ . Let  $g: X \times X \times \mathbb{R}_+ \times \Lambda \rightarrow Y$  be defined by

$$g(x, y, \varepsilon, \lambda) = f(x, y, \lambda) + \varepsilon e(\lambda).$$

Since  $f$  is  $C$ -usc and  $e$  is continuous, Proposition 2.2 implies that  $g$  is  $C$ -usc. Using the techniques as in the proof for  $S^s$ , where  $f$  is replaced by  $g$ , we conclude that  $\widetilde{S^s}$  is upper semicontinuous with compact values on  $\Lambda \times \mathbb{R}_+$ .  $\square$

The essentialness of the assumptions imposed in (a) and (b) of Theorem 3.2 is shown by the following examples.

**Example 3.1.** Let  $X = [0, \frac{1}{2}\pi]$ ;  $\Lambda = [0, 2]$ ;  $Y = \mathbb{R}^2$ ,  $K(x, \lambda) = [\frac{1}{2}\lambda x, x]$ ,  $e(\lambda) \equiv (1; 1)$ ,  $f(x, y, \lambda) = (x + \cos x; \lambda - y)$ ,

$$C(\lambda) = \begin{cases} \mathbb{R}_+^2, & \text{if } \lambda \neq 1, \\ \mathbb{R} \times \mathbb{R}_+, & \text{if } \lambda = 1. \end{cases}$$

Then  $X$  is compact,  $f$  and  $K$  are continuous with compact values. Direct calculations give

$$S^w(\lambda) = \begin{cases} [0, \frac{1}{2}\pi], & \text{if } \lambda \neq 1, \\ [0, 1], & \text{if } \lambda = 1, \end{cases}$$

and

$$\widetilde{S}^w(\lambda, \varepsilon) = \begin{cases} [0, \frac{1}{2}\pi], & \text{if } \lambda \neq 1, \\ [0, \varepsilon + 1] \cap [0, \frac{1}{2}\pi], & \text{if } \lambda = 1. \end{cases}$$

It is obvious that  $S^w$  and  $\widetilde{S}^w$  are not usc at 1 and  $(1, 0)$ , respectively. The reason is that the imposed assumption in (a) is violated.

**Example 3.2.** Let  $X = [0, 2]$ ;  $\Lambda = [0, 1]$ ;  $Y = \mathbb{R}^2$ ,  $K(x, \lambda) = [\lambda x, x]$ ,  $e(\lambda) \equiv (1; 1)$ ,  $f(x, y, \lambda) = (e^{x+y}; \lambda - y)$ ,

$$C(\lambda) = \begin{cases} \mathbb{R}_+^2, & \text{if } \lambda = \frac{1}{2}, \\ \mathbb{R}_+ \times \mathbb{R}, & \text{if } \lambda \neq \frac{1}{2}. \end{cases}$$

Then  $X$  is compact,  $f$  and  $K$  are continuous with compact values. Easy calculations yield

$$S^s(\lambda) = \begin{cases} [0, \frac{1}{2}], & \text{if } \lambda = \frac{1}{2}, \\ [0, 2], & \text{if } \lambda \neq \frac{1}{2}, \end{cases}$$

and

$$\widetilde{S}^s(\lambda, \varepsilon) = \begin{cases} [0, \frac{1}{2} + \varepsilon] \cap [0, 2], & \text{if } \lambda = \frac{1}{2}, \\ [0, 2], & \text{if } \lambda \neq \frac{1}{2}. \end{cases}$$

It is clear that  $S^s$  and  $\widetilde{S}^s$  are not usc at  $\frac{1}{2}$  and  $(\frac{1}{2}, 0)$ , respectively. The reason is that the imposed assumption in (b) is violated.

Although the imposed assumptions in (a) and (b) of Theorem 3.2 cannot be dispensed with in the statement, we can replace such assumptions together with the imposed condition on  $f$  as follows.

**Theorem 3.3.** Assume that  $X$  is compact and  $K$  is continuous with compact values. Then

- (a)  $\widetilde{S}^w$  is upper semicontinuous with compact values on  $\Lambda \times \mathbb{R}_+$  if one of the following two conditions holds:
- (i)  $f$  is continuous and  $Y \setminus -\text{int } C(\cdot)$  is closed.
  - (ii)  $f$  is  $C$ -upper semicontinuous and  $Y \setminus -\text{int } C(\cdot)$  is  $H$ -upper semicontinuous.
- (b)  $\widetilde{S}^s$  is upper semicontinuous with compact values on  $\Lambda \times \mathbb{R}_+$  if one of the following two conditions holds:
- (i)  $f$  is continuous and  $C$  is closed in  $\Lambda$ .
  - (ii)  $f$  is  $C$ -upper semicontinuous and  $C$  is  $H$ -upper semicontinuous.

*Proof.* We verify only (a) as an example. The proof of part (b) is similar.

(i) For given  $\lambda \in \Lambda$ , we first prove that  $S^w$  is usc at  $\lambda$ . Suppose to the contrary that  $S^w$  is not usc at  $\lambda$ . Then we can pick a superset  $U$  of  $S^w(\lambda)$  and a sequence  $\{\lambda_n\}$  in  $\Lambda$  converging to  $\lambda$  such that there is  $x_n \in S^w(\lambda_n) \setminus U$ , for all  $n$ . Since  $X$  is compact, we can assume that  $x_n$  tends to  $\bar{x}$ , for some  $\bar{x} \in X$ . By the upper semicontinuity with compact values of  $K$  at  $(\bar{x}, \lambda)$ , we have  $\bar{x} \in K(\bar{x}, \lambda)$ . For each  $y \in K(\bar{x}, \lambda)$ , since  $K$  is lsc at  $(\bar{x}, \lambda)$ , there is  $y_n \in K(x_n, \lambda_n)$ ,  $y_n \rightarrow y$ . As  $x_n \in S^w(\lambda_n)$ , we have

$$f(x_n, y_n, \lambda_n) \in (Y \setminus -\text{int } C(\lambda_n)).$$

Combining the closedness of  $Y \setminus -\text{int } C(\cdot)$  with the continuity of  $f$ , we conclude that

$$f(\bar{x}, y, \lambda) \in (Y \setminus -\text{int } C(\lambda)),$$

i.e.,  $\bar{x} \in S^w(\lambda)$ , which is impossible as  $x_n \notin U$  for all  $n$ . Hence  $S^w$  is usc at  $\lambda$ .

Theorem 3.1 yields that  $S^w(\lambda)$  is a closed subset of  $X$ , and hence it is compact.

For given  $(\lambda, \varepsilon) \in \Lambda \times \mathbb{R}_+$ , it is clear that  $g(x, y, \varepsilon, \lambda) = f(x, y, \lambda) + \varepsilon e(\lambda)$  is continuous as  $f$  and  $e$  are continuous. Arguing as above, we conclude that  $\widetilde{S}^w$  is upper semicontinuous with compact values at  $(\lambda, \varepsilon)$ .

(ii) Similarly to the first part, if  $S^w$  is not usc at  $\lambda$ , then there exist a superset  $U$  of  $S^w(\lambda)$  and a sequence  $\{\lambda_n\} \rightarrow \lambda$  such that there is  $x_n \in S^w(\lambda_n) \setminus U$  for all  $n$ ,  $x_n \rightarrow \bar{x} \in K(\bar{x}, \lambda)$ . For each  $y \in K(\bar{x}, \lambda)$  there is  $y_n \in K(x_n, \lambda_n)$ ,  $y_n \rightarrow y$ , and

$$f(x_n, y_n, \lambda_n) \in (Y \setminus -\text{int } C(\lambda_n)).$$

For each neighborhood  $B$  of  $\theta_Y$ , there is a balanced neighborhood  $B_1$  of  $\theta_Y$ , i.e.,  $-B_1 = B_1$ , such that  $B_1 + B_1 \subset B$ . Since  $Y \setminus -\text{int } C(\cdot)$  is  $H$ -usc at  $\lambda$  and  $f$  is  $C$ -usc at  $(\bar{x}, y, \lambda)$ , there is  $n_0 \in \mathbb{N}$  such that, for each  $n \geq n_0$ , we have

$$(Y \setminus -\text{int } C(\lambda_n)) \subset (Y \setminus -\text{int } C(\lambda)) + B_1, \text{ and} \\ f(x_n, y_n, \lambda_n) \in f(\bar{x}, y, \lambda) + B_1 - C(\lambda).$$

By the balance of  $B_1$ , we have

$$f(\bar{x}, y, \lambda) \in f(x_n, y_n, \lambda_n) - B_1 + C(\lambda) = f(x_n, y_n, \lambda_n) + B_1 + C(\lambda),$$

so,

$$f(\bar{x}, y, \lambda) \in (Y \setminus -\text{int } C(\lambda_n)) + B_1 + C(\lambda).$$

Since  $C(\lambda)$  is a convex cone,  $(Y \setminus -\text{int } C(\lambda)) + C(\lambda) \subset (Y \setminus -\text{int } C(\lambda))$ . Therefore,

$$f(\bar{x}, y, \lambda) \in (Y \setminus -\text{int } C(\lambda)) + B_1 + B_1 + C(\lambda) \subset (Y \setminus -\text{int } C(\lambda)) + B.$$

Since  $B$  is arbitrary and  $Y \setminus -\text{int } C(\lambda)$  is closed, we conclude that  $f(\bar{x}, y, \lambda) \in (Y \setminus -\text{int } C(\lambda))$ , i.e.,  $\bar{x} \in S^w(\lambda)$ , which is impossible as  $x_n \notin U$  for all  $n$ . Hence  $S^w$  is usc at  $\lambda$ . Theorem 3.1 yields the closedness of  $S^w(\lambda)$ , and hence  $S^w(\lambda)$  is compact as  $X$  is compact.

Using the same arguments as above, the upper semicontinuity with compact values of  $\widetilde{S}^w$  is also established.  $\square$

Now let us present an important relationship between well-posedness and stability of the problems considered.

**Theorem 3.4.**

- (i) *The problems (WQEP) and (SQEP) are well-posed under perturbations if and only if, for each  $\lambda \in \Lambda$ ,  $\widetilde{S}^w$  and  $\widetilde{S}^s$  are upper semicontinuous with compact values at  $(\lambda, 0)$ , respectively.*
- (ii) *The problems (WQEP) and (SQEP) are generalized Hadamard well-posed if and only if, for each  $\lambda \in \Lambda$ ,  $S^w$  and  $S^s$  are upper semicontinuous with compact values at  $\lambda$ , respectively.*

**Proof.** We show only the well-posedness under perturbations for the problem (SQEP), since the other cases are similar. Suppose that for given  $\lambda \in \Lambda$ ,  $\widetilde{S}^s$  is usc at  $(\lambda, 0)$  and  $S^s(\lambda)$  is compact. Let  $\{\lambda_n\} \subset \Lambda$  an arbitrary sequence converging to  $\lambda$  and  $\{x_n\}$  an approximating sequence for (SQEP) $_{\lambda}$  corresponding to  $\{\lambda_n\}$ . Then there exists a sequence  $\{\varepsilon_n\} \downarrow 0$  such that, for each  $n \in \mathbb{N}$ ,  $x_n \in K(x_n, \lambda_n)$  and

$$f(x_n, y, \lambda_n) + \varepsilon_n e(\lambda_n) \in C(\lambda_n) \quad \forall y \in K(x_n, \lambda_n),$$

i.e.,  $x_n \in \widetilde{S}^s(\lambda_n, \varepsilon_n)$ . Since  $\widetilde{S}^s$  is upper semicontinuous with compact values at  $(\lambda, 0)$ , Lemma 2.2 implies that there is a subsequence  $\{x_{n_k}\}$  of  $\{x_n\}$ ,  $x_{n_k} \rightarrow \bar{x}$ , for some  $\bar{x} \in S^s(\lambda)$ . Therefore, the problem (SQEP) is well-posed.

Conversely, suppose that the problem (SQEP) is well-posed. Let  $\lambda \in \Lambda$ ,  $\{(\lambda_n, \varepsilon_n)\} \subset \Lambda \times \mathbb{R}_+$ , with  $(\lambda_n, \varepsilon_n) \rightarrow (\lambda, 0)$  and  $x_n \in \widetilde{S^s}(\lambda_n, \varepsilon_n)$ . Then, for each  $n \in \mathbb{N}$ ,  $x_n \in K(x_n, \lambda_n)$ , and

$$f(x_n, y, \lambda_n) + \varepsilon_n e(\lambda_n) \in C(\lambda_n) \quad \forall y \in K(x_n, \lambda_n),$$

i.e.,  $\{x_n\}$  is an approximating sequence of  $(\text{SQEP}_\lambda)$  corresponding to  $\{\lambda_n\}$ . Using the well-posedness of the problem (SQEP), we can find a subsequence of  $\{x_n\}$  converging to some point of  $S^s(\lambda) = \widetilde{S^s}(\lambda, 0)$ . Using Lemma 2.2, we obtain that  $\widetilde{S^s}$  is upper semicontinuous with compact values at  $(\lambda, 0)$ .  $\square$

Combining Theorems 3.2 and 3.3, we establish the results of well-posedness under perturbations and generalized Hadamard well-posedness for the problems (WQEP) and (SQEP).

**Corollary 3.1.**

- (a) *If the conditions of Theorem 3.2(a) or those of Theorem 3.3(a) are satisfied, then the problem (WQEP) is well-posed under perturbations and generalized Hadamard well-posed.*
- (b) *If the conditions of Theorem 3.2(b) or those of Theorem 3.3(b) are satisfied, then the problem (SQEP) is well-posed under perturbations and generalized Hadamard well-posed.*

Passing to uniquely well-posedness under perturbations and Hadamard well-posedness, we obtain results similar to those of Theorem 3.4 and Corollary 3.1.

**Theorem 3.5.** *Suppose that, for each  $\lambda \in \Lambda$ ,  $(\text{WQEP}_\lambda)$  and  $(\text{SQEP}_\lambda)$  have a unique solution. Then*

- (i)  *$(\text{WQEP})$  and  $(\text{SQEP})$  are uniquely well-posed under perturbations if and only if, for each  $\lambda \in \Lambda$ ,  $\widetilde{S^w}$  and  $\widetilde{S^s}$  are upper semicontinuous with compact values at  $(\lambda, 0)$ , respectively;*
- (ii)  *$(\text{WQEP})$  and  $(\text{SQEP})$  are Hadamard well-posed if and only if, for each  $\lambda \in \Lambda$ ,  $S^w$  and  $S^s$  are upper semicontinuous with compact values at  $\lambda$ , respectively.*

**Corollary 3.2.**

- (a) *If the conditions of Theorem 3.2(a) or those of Theorem 3.3(a) are satisfied and assuming further that, for each  $\lambda \in \Lambda$ , the problem  $(\text{WQEP}_\lambda)$  has a unique solution, then  $(\text{WQEP})$  is uniquely well-posed under perturbations and Hadamard well-posed.*

- (b) *If the conditions of Theorem 3.2(b) or those of Theorem 3.3(b) are satisfied and assuming further that, for each  $\lambda \in \Lambda$ , the problem (SQEP $_{\lambda}$ ) has a unique solution, then (SQEP) is uniquely well-posed under perturbations and Hadamard well-posed.*

#### 4. APPLICATIONS

**4.1. Lower and upper bounded equilibrium problems.** Let  $X$  and  $\Lambda$  be as in Section 2. Let  $g: X \times X \times \Lambda \rightarrow \mathbb{R}$  be a vector valued mapping and let  $\alpha, \beta \in \mathbb{R}_+$ ,  $\alpha < \beta$ , be fixed. For each  $\lambda \in \Lambda$ , we consider the lower and upper bounded equilibrium problem studied in [19], [45]:

(BEP $_{\lambda}$ ) Find  $\bar{x} \in K(\lambda)$  such that, for each  $y \in K(\lambda)$ ,

$$\alpha \leq g(\bar{x}, y, \lambda) \leq \beta.$$

Setting  $Y = \mathbb{R}^2$ ,  $C(\lambda) \equiv \mathbb{R}_+^2$  and  $f(x, y, \lambda) = (g(x, y, \lambda) - \alpha; \beta - g(x, y, \lambda))$ , (SQEP $_{\lambda}$ ) reduces to (BEP $_{\lambda}$ ).

Combining Theorems 3.2 and 3.4, we establish sufficient conditions for the well-posedness of the lower and upper bounded equilibrium problem as follows.

**Corollary 4.1.** *Assume that*

- (i)  *$g$  is continuous;*
- (ii)  *$K$  is continuous and with compact values.*

*Then the family  $\{(\text{BEP}_{\lambda}) : \lambda \in \Lambda\}$  is well-posed under perturbations and generalized Hadamard well-posed. Moreover, if the solution set is a singleton, then the problem is uniquely and Hadamard well-posed.*

Recently, there have been many works devoted to the lower and upper bounded equilibrium problems, such as existence conditions [19], [45], [21], stability conditions [46], and the references therein. To the best of our knowledge, there have not been any papers on the well-posedness for the lower and upper bounded equilibrium problems.

**4.2. Traffic network problems.** The Wardrop equilibrium flows for the transportation network problem was first introduced in 1952 by Wardrop [44] together with a basic traffic network principle. Since then, traffic network problems have been intensively studied from both the theory and methodology view points. Contributing to the development of traffic network problems, Smith [41] derived a spotlight point by claiming that the Wardrop equilibrium flows of a network are expressed as solutions of the corresponding variational inequality. Because of diverse practical situations, these demands may depend on the equilibrium vector flows which

were considered in [20], [36]; the so-called elastic network problems, and then the Wardrop equilibrium flows of the elastic network problem are obtained as solutions of the corresponding quasivariational inequality.

Let us recall the mathematical model of the elastic traffic network problem studied in [2], [6], [20], [36]. Let  $N$  the set of nodes and  $L$  that of arcs (or links). We denote the collection of origin-destination pairs (O/D pairs for short) by  $W = (W_1, \dots, W_l)$ . Suppose that the pair  $W_j$ ,  $j = 1, \dots, l$ , is connected by a set  $P_j$  of paths and  $P_j$  contains  $r_j$  paths,  $r_j \geq 1$ . Let us denote the path vector flow by  $F = (F_1, \dots, F_m)$ , where  $m = r_1 + \dots + r_l$ . The capacity of these paths must be taken into account in practice (see, [26]), and hence we assume that the compact capacity restriction is defined by

$$F \in X := \{F \in \mathbb{R}^m : 0 \leq \gamma_s \leq F_s \leq \Gamma_s, s = 1, \dots, m\}.$$

Suppose that the travel cost on the path flow  $F_s$ ,  $s = 1, \dots, m$ , depends on the whole path vector flow  $F$  and is  $T_s(F) \geq 0$ . Then we derive the path cost vector  $T(F) = (T_1(F), \dots, T_m(F))$ .

A path vector flow  $H$  is named an equilibrium vector flow as defined in [44], if for each  $W_j$  and  $p, s \in P_j$ ,

$$[T_p(H) < T_s(H)] \implies [H_s = \gamma_s \text{ or } H_p = \Gamma_p].$$

Now assume that the perturbation on the traffic is expressed by a parameter  $\lambda$  of a metric space  $A$ . Suppose that the travel demands  $g_j$  of the O/D pair  $W_j$  depend on  $\lambda \in A$  and also on the equilibrium vector flow  $H$  as interpreted in [20], [36]. We denote the travel vector demand by  $g = (g_1, \dots, g_l)$  and set

$$\varphi_{js} = \begin{cases} 1, & \text{if } s \in P_j, \\ 0, & \text{if } s \notin P_j, \end{cases}$$

$$\varphi = \{\varphi_{js}\}, j = 1, \dots, l; s = 1, \dots, m.$$

Then the path vector flows meeting the travel demands are termed the feasible path vector flows and form the constraint set

$$K(H, \lambda) = \{F \in X : \varphi F = g(H, \lambda)\}.$$

The matrix  $\varphi$  is called the O/D pair—path incidence matrix. Assume further that the path costs also depend on the parameter  $\lambda$ ,  $T_s(F, \lambda)$ ,  $s = 1, \dots, m$ .

Our traffic network problem is equivalent to a quasivariational inequality as follows.



**Lemma 4.1** ([41]). *A path vector flow  $H \in K(H, \lambda)$  is an equilibrium flow if and only if it is a solution of the following quasivariational inequality:*

(QVI $_{\lambda}$ ) *find  $H \in K(H, \lambda)$  such that, for each  $F \in K(H, \lambda)$ ,*

$$\langle T(H, \lambda), F - H \rangle \geq 0.$$

Setting  $f(H, F, \lambda) = \langle T(H, \lambda), F - H \rangle$ , (WQEP $_{\lambda}$ ) and (SQEP $_{\lambda}$ ) become (QVI $_{\lambda}$ ).

**Lemma 4.2** ([6]). *If  $g$  is continuous at  $(H, \lambda)$ , then  $K$  is continuous with convex and compact values at  $(H, \lambda)$ .*

The following result is derived from Theorems 3.2, 3.4 and Lemmas 4.1, 4.2.

**Corollary 4.2.** *If  $g$  and  $T$  are continuous, then the family of parametric elastic network problems is well-posed under perturbations and generalized Hadamard well-posed. Furthermore, if (QVI $_{\lambda}$ ) has a unique solution, then this problem is uniquely well-posed under perturbations and Hadamard well-posed.*

**Acknowledgement.** The authors would like to thank the referees for their valuable remarks and suggestions, which helped to improve the paper.

#### References

- [1] *M. Ait Mansour, H. Riahi:* Sensitivity analysis for abstract equilibrium problems. *J. Math. Anal. Appl.* *306* (2005), 684–691.
- [2] *M. Ait Mansour, L. Scrimali:* Hölder continuity of solutions to elastic traffic network models. *J. Glob. Optim.* *40* (2008), 175–184.
- [3] *L. Q. Anh, P. Q. Khanh:* Semicontinuity of the solution set of parametric multivalued vector quasiequilibrium problems. *J. Math. Anal. Appl.* *294* (2004), 699–711.
- [4] *L. Q. Anh, P. Q. Khanh:* On the Hölder continuity of solutions to parametric multivalued vector equilibrium problems. *J. Math. Anal. Appl.* *321* (2006), 308–315.
- [5] *L. Q. Anh, P. Q. Khanh:* Semicontinuity of the approximate solution sets of multivalued quasiequilibrium problems. *Numer. Funct. Anal. Optim.* *29* (2008), 24–42.
- [6] *L. Q. Anh, P. Q. Khanh:* Continuity of solution maps of parametric quasiequilibrium problems. *J. Glob. Optim.* *46* (2010), 247–259.
- [7] *L. Q. Anh, P. Q. Khanh, D. T. M. Van:* Well-posedness without semicontinuity for parametric quasiequilibria and quasioptimization. *Comput. Math. Appl.* *62* (2011), 2045–2057.
- [8] *L. Q. Anh, P. Q. Khanh, D. T. M. Van:* Well-posedness under relaxed semicontinuity for bilevel equilibrium and optimization problems with equilibrium constraints. *J. Optim. Theory Appl.* *153* (2012), 42–59.
- [9] *L. Q. Anh, P. Q. Khanh, D. T. M. Van, J.-C. Yao:* Well-posedness for vector quasiequilibria. *Taiwanese J. Math.* *13* (2009), 713–737.
- [10] *Q. H. Ansari, F. Flores-Bazán:* Generalized vector quasi-equilibrium problems with applications. *J. Math. Anal. Appl.* *277* (2003), 246–256.

- [11] *J.-P. Aubin, H. Frankowska*: Set-Valued Analysis. Modern Birkhäuser Classics, Birkhäuser, Boston, 2009.
- [12] *M. Bianchi, G. Kassay, R. Pini*: Well-posed equilibrium problems. *Nonlinear Anal., Theory Methods Appl., Ser. A, Theory Methods* 72 (2010), 460–468.
- [13] *M. Bianchi, R. Pini*: A note on stability for parametric equilibrium problems. *Oper. Rest. Lett.* 31 (2003), 445–450.
- [14] *M. Bianchi, R. Pini*: Coercivity conditions for equilibrium problems. *J. Optimization Theory Appl.* 124 (2005), 79–92.
- [15] *M. Bianchi, R. Pini*: Sensitivity for parametric vector equilibria. *Optimization* 55 (2006), 221–230.
- [16] *E. Blum, W. Oettli*: From optimization and variational inequalities to equilibrium problems. *Math. Stud.* 63 (1994), 123–145.
- [17] *R. Burachik, G. Kassay*: On a generalized proximal point method for solving equilibrium problems in Banach spaces. *Nonlinear Anal., Theory Methods Appl., Ser. A, Theory Methods* 75 (2012), 6456–6464.
- [18] *A. Capătă, G. Kassay*: On vector equilibrium problems and applications. *Taiwanese J. Math.* 15 (2011), 365–380.
- [19] *O. Chadli, Y. Chiang, J. C. Yao*: Equilibrium problems with lower and upper bounds. *Appl. Math. Lett.* 15 (2002), 327–331.
- [20] *M. De Luca*: Generalized quasi-variational inequalities and traffic equilibrium problem. *Variational Inequalities and Network Equilibrium Problems* (F. Giannessi, eds.). Proc. Conf., Erice, 1994, Plenum, New York, 1995, pp. 45–54.
- [21] *X. Ding*: Equilibrium problems with lower and upper bounds in topological spaces. *Acta Math. Sci., Ser. B, Engl. Ed.* 25 (2005), 658–662.
- [22] *B. Djafari Rouhani, E. Tarafdar, P. J. Watson*: Existence of solutions to some equilibrium problems. *J. Optimization Theory Appl.* 126 (2005), 97–107.
- [23] *Y.-P. Fang, R. Hu, N.-J. Huang*: Well-posedness for equilibrium problems and for optimization problems with equilibrium constraints. *Comput. Math. Appl.* 55 (2008), 89–100.
- [24] *Y.-P. Fang, N.-J. Huang, J.-C. Yao*: Well-posedness of mixed variational inequalities, inclusion problems and fixed point problems. *J. Global Optim.* 41 (2008), 117–133.
- [25] *F. Flores-Bazán*: Existence theorems for generalized noncoercive equilibrium problems: the quasi-convex case. *SIAM J. Optim.* 11 (2001), 675–690.
- [26] *F. Giannessi*: Theorems of alternative, quadratic programs and complementarity problems. *Variational Inequalities and Complementarity Problems. Proc. Int. School Math., Erice, 1978, Wiley, Chichester, 1980*, pp. 151–186.
- [27] *J. Hadamard*: Sur le problèmes aux dérivées partielles et leur signification physique. *Bull. Univ. Princeton* 13 (1902), 49–52. (In French.)
- [28] *N. X. Hai, P. Q. Khanh*: Existence of solutions to general quasiequilibrium problems and applications. *J. Optim. Theory Appl.* 133 (2007), 317–327.
- [29] *N.-J. Huang, X.-J. Long, C.-W. Zhao*: Well-posedness for vector quasi-equilibrium problems with applications. *J. Ind. Manag. Optim.* 5 (2009), 341–349.
- [30] *A. Ioffe, R. E. Lucchetti*: Typical convex program is very well posed. *Math. Program.* 104 (2005), 483–499.
- [31] *A. N. Iusem, G. Kassay, W. Sosa*: On certain conditions for the existence of solutions of equilibrium problems. *Math. Program.* 116 (2009), 259–273.
- [32] *A. N. Iusem, W. Sosa*: Iterative algorithms for equilibrium problems. *Optimization* 52 (2003), 301–316.
- [33] *K. Kimura, Y.-C. Liou, S.-Y. Wu, J.-C. Yao*: Well-posedness for parametric vector equilibrium problems with applications. *J. Ind. Manag. Optim.* 4 (2008), 313–327.

- [34] *M. B. Lignola, J. Morgan*:  $\alpha$ -well-posedness for Nash equilibria and for optimization problems with Nash equilibrium constraints. *J. Glob. Optim.* *36* (2006), 439–459.
- [35] *X.-J. Long, N.-J. Huang, K.-L. Teo*: Existence and stability of solutions for generalized strong vector quasi-equilibrium problem. *Math. Comput. Modelling* *47* (2008), 445–451.
- [36] *A. Maugeri*: Variational and quasi-variational inequalities in network flow models. Recent developments in theory and algorithms. *Variational Inequalities and Network Equilibrium Problems. Proc. Conf., Erice, 1994, Plenum, New York, 1995*, pp. 195–211.
- [37] *L. D. Muu, W. Oettli*: Convergence of an adaptive penalty scheme for finding constrained equilibria. *Nonlinear Anal., Theory Methods Appl.* *18* (1992), 1159–1166.
- [38] *M. A. Noor, K. I. Noor*: Equilibrium problems and variational inequalities. *Mathematica* *47(70)* (2005), 89–100.
- [39] *J. P. Revalski, N. V. Zhikov*: Well-posed constrained optimization problems in metric spaces. *J. Optimization Theory Appl.* *76* (1993), 145–163.
- [40] *I. Sadeqi, C. G. Alizadeh*: Existence of solutions of generalized vector equilibrium problems in reflexive Banach spaces. *Nonlinear Anal., Theory Methods Appl., Ser. A, Theory Methods* *74* (2011), 2226–2234.
- [41] *M. J. Smith*: The existence, uniqueness and stability of traffic equilibria. *Transportation Res. Part B* *13* (1979), 295–304.
- [42] *J. J. Strodiot, T. T. V. Nguyen, V. H. Nguyen*: A new class of hybrid extragradient algorithms for solving quasi-equilibrium problems. *J. Glob. Optim.* *56* (2013), 373–397.
- [43] *A. N. Tikhonov*: On the stability of the functional optimization problem. *U.S.S.R. Comput. Math. Math. Phys.* *6* (1966), 28–33; translation from *Zh. Vychisl. Mat. Mat. Fiz.* *6* (1966), 631–634. (In Russian.)
- [44] *J. G. Wardrop*: Some theoretical aspects of road traffic research. *Proceedings of the Institute of Civil Engineers, Part II* (1952), 325–378.
- [45] *C. Zhang*: A class of equilibrium problems with lower and upper bound. *Nonlinear Anal., Theory Methods Appl., Ser. A, Theory Methods (electronic only)* *63* (2005), e2377–e2385.
- [46] *C. Zhang, J. Li, Z. Feng*: The existence and the stability of solutions for equilibrium problems with lower and upper bounds. *J. Nonlinear Anal. Appl.* *2012* (2012), Article ID jnaa-00135, 13 pages.
- [47] *T. Zolezzi*: Well-posedness criteria in optimization with application to the calculus of variations. *Nonlinear Anal., Theory Methods Appl.* *25* (1995), 437–453.
- [48] *T. Zolezzi*: Well-posedness and optimization under perturbations. *Ann. Oper. Res.* *101* (2001), 351–361.
- [49] *T. Zolezzi*: On well-posedness and conditioning in optimization. *ZAMM, Z. Angew. Math. Mech.* *84* (2004), 435–443.

*Authors' addresses:* *Lam Quoc Anh*, Department of Mathematics, Teacher College, Cantho University, Cantho City, 92000, Vietnam, e-mail: [quocanh@ctu.edu.vn](mailto:quocanh@ctu.edu.vn); *Dinh Vinh Hien*, Department of Fundamental Sciences, Hochiminh City University of Food Industry, Hochiminh City, 70000, Vietnam, e-mail: [hiedinhvinh@gmail.com](mailto:hiedinhvinh@gmail.com).