

Haosui Duanmu; William Weiss

Finitely-additive, countably-additive and internal probability measures

Commentationes Mathematicae Universitatis Carolinae, Vol. 59 (2018), No. 4, 467–485

Persistent URL: <http://dml.cz/dmlcz/147551>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2018

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

Finitely-additive, countably-additive and internal probability measures

HAOSUI DUANMU, WILLIAM WEISS

To the memory of Bohuslav Balcar

Abstract. We discuss two ways to construct standard probability measures, called push-down measures, from internal probability measures. We show that the Wasserstein distance between an internal probability measure and its push-down measure is infinitesimal. As an application to standard probability theory, we show that every finitely-additive Borel probability measure P on a separable metric space is a limit of a sequence of countably-additive Borel probability measures $\{P_n\}_{n \in \mathbb{N}}$ in the sense that $\int f dP = \lim_{n \rightarrow \infty} \int f dP_n$ for all bounded uniformly continuous real-valued function f if and only if the space is totally bounded.

Keywords: nonstandard model in mathematics; nonstandard analysis; nonstandard measure theory; convergence of probability measures

Classification: 03H05, 26E35, 28E05, 60B10

1. Introduction

One of the foundational problems in probability theory is to study the connection between finitely-additive measures and countably additive measures (see, e.g., [19]). In contrast to Prokhorov's theorem and the Vitali–Hahn–Saks theorem which state that a sequence of countably additive measures converge to a countably additive measure under regularity conditions, we prove that for every finitely-additive probability measure P on a totally bounded separable metric space, there is a sequence of countably additive probability measures $\{P_n\}_{n \in \mathbb{N}}$ such that $\int f dP_n \rightarrow \int f dP$ for every bounded uniformly continuous real-valued function. On the other hand, unlike the Portmanteau lemma, such convergence fails for merely bounded continuous functions, showing that the hypothesis of the Portmanteau theorem is sharp.

In Section 2, we give a gentle introduction to nonstandard analysis as well as nonstandard measure theory. Nonstandard measure theory provides powerful machinery to study this problem. On one hand internal measures have the same first-order logical properties as finitely-additive measures. On the other hand internal measures can be easily extended to countably additive measures (Loeb measure), using Loeb's construction in [13]. We can also reverse the procedure to construct

finitely-additive probability measures from internal probability measures or Loeb measures. There is a rich literature on constructing standard measures using the standard part map and Loeb measures in very general settings (see, e.g., [3], [12], [15], [1], [17], [2] and [18]). On the other hand, by the transfer principle, we can construct finitely-additive probability measures from internal probability measures. In this paper, we establish some connections between finitely-additive measures and countably-additive measures by studying the relation between these two forms of push-down, and in doing so, we prove the theorem mentioned in the first paragraph, which has no known standard proof.

In Section 3, we define the Wasserstein distance between finitely-additive measures and show that the Wasserstein distance, ${}^*W(\nu, {}^*P)$, between a finitely additive measure P and an internal probability measure ν is infinitesimal provided that the underlying space is a bounded σ -compact metric space and ${}^*P({}^*A) \approx \nu({}^*A)$ for all Borel sets A . In contrast to this, we give an example of an internal probability measure ν with $\nu({}^*A) \approx \lambda(A)$ for $A \in \mathcal{B}[[0, 1]]$ where λ denote the Lebesgue measure on $[0, 1]$. Meanwhile, we have

$$(1.1) \quad \sup_{B \in {}^*\mathcal{B}[[0, 1]]} |\nu(B) - {}^*\lambda(B)| = 1.$$

Given an internal probability measure ν on $({}^*X, {}^*\mathcal{B}[X])$, the internal push-down measure ν^p is a finitely-additive measure on $(X, \mathcal{B}[X])$ defined as $\nu^p(A) = \text{st}(\nu({}^*A))$ and the external push-down measure ν_p is a countably additive measure on $(X, \mathcal{B}[X])$ defined as $\nu_p(A) = \overline{\nu}(\text{st}^{-1}(A))$ where $\overline{\nu}$ denote the Loeb extension of ν . H. Duanmu, D. M. Roy and A. Smith (in paper that is in preparation) showed that ${}^*W(\nu, {}^*\nu_p) \approx 0$ if the underlying space is compact. In Section 4, we generalize this result to bounded σ -compact spaces. We also show that ${}^*W(\nu, \nu^p) \approx 0$. Thus, the Wasserstein distance is only a pseudometric on the space of all finitely-additive probability measures.

There exists a rich literature on studying the relationship between finitely-additive probability measures and countably additive probability measures (e.g., see [19] and [10]). For an uncountable σ -algebra \mathcal{F} , the set of finitely-additive probability measures can be viewed as a subset of the compact product space $[0, 1]^\mathcal{F}$. T. Seidenfeld pointed out that the set of finitely-additive probability measures with finite support forms a dense subset of the set of all finitely-additive probability measures. Thus, every finitely-additive probability measure is an accumulation point of a set of countably additive probability measures. However, no point in $[0, 1]^\mathcal{F}$ has a countable local base hence we cannot conclude that every finitely-additive probability measure is the limit of a countable sequence of countably additive probability measures under pointwise convergence.

In Section 5 we show that for every finitely-additive probability measure P , there is a sequence of countably-additive probability measures $\{P_n\}_{n \in \mathbb{N}}$ such that $\int f dP_n$ converges to $\int f dP$ for every bounded uniformly continuous real-valued function f . We denote such convergence by weak convergence. In Example 5.14,

we show that our theorem fails if we replace bounded uniformly continuous real-valued function by merely bounded continuous real-valued function, hence our result is sharp. We conclude with a nonstandard characterization of weak convergence to finitely-additive probability measures, which is similar in spirit to Theorem 4 in [4].

2. Preliminaries

In this section, we give a short introduction to nonstandard analysis. A large part of this introduction is taken from the preliminary section in [11]. For a detailed introduction to nonstandard models, we recommend the first four chapters of [5].

Given any set S containing \mathbb{R} as a subset, the superstructure $\mathbb{V}(S)$ over S is defined as

- (1) $\mathbb{V}_1(S) = S$;
- (2) $\mathbb{V}_{n+1}(S) = \mathbb{V}_n(S) \cup \mathcal{P}(S)$;
- (3) $\mathbb{V}(S) = \bigcup_{n \in \mathbb{N}} \mathbb{V}_n(S)$.

The starting point of nonstandard analysis is to construct a set ${}^*\mathbb{R} \supset \mathbb{R}$ and a mapping $*$: $\mathbb{V}(\mathbb{R}) \mapsto \mathbb{V}({}^*\mathbb{R})$ with three basic properties. We first state the following two basic notions from mathematical logic. A *formula* is a statement φ built up from equality and \in relations $x = y, x \in y$, the connectives \wedge, \vee, \neg and bounded quantifiers $(\forall x \in y), (\exists x \in y)$. An *internal object* is an element of the set $\bigcup \{ {}^*A : A \in \mathbb{V}(S) \}$. A set in $\mathbb{V}(S)$ which is not internal is called *external*. We now state the three basic properties.

- (1) *Extension principle*: *S is a proper extension of S and ${}^*s = s$ for all $s \in S$.
- (2) *Transfer principle*: Let $S_1, \dots, S_n \in \mathbb{V}(S)$. Any formula which is true of S_1, \dots, S_n is true of ${}^*S_1, \dots, {}^*S_n$.
- (3) *κ -saturation principle*: Let κ be a cardinal number and let \mathcal{F} be a collection of internal sets. If \mathcal{F} has the finite intersection property with cardinality no more than κ , then the total intersection of \mathcal{F} is nonempty.

An internal set A is a *hyperfinite set* if there exists an internal bijection f between A and $\{n \in {}^*\mathbb{N} : n \leq N_0\}$ for some $N_0 \in {}^*\mathbb{N}$.

In this paper, the nonstandard model is as saturated as it needs to be. Let (X, \mathcal{T}) be a topological space. The monad of a point $x \in X$ is the set $\bigcap_{x \in U \in \mathcal{T}} {}^*U$. An element $x \in {}^*X$ is *near-standard* if it is in the monad of some $y \in X$. We say y is the standard part of x and write $y = \text{st}(x)$. We use $\text{NS}({}^*X)$ to denote the collection of near-standard elements of *X and we say $\text{NS}({}^*X)$ is the *near-standard part* of *X . The standard part map st is a function from $\text{NS}({}^*X)$ to X . For a metric space X , two elements $x, y \in {}^*X$ are *infinitely close* if ${}^*d(x, y) \approx 0$. For two elements $a, b \in {}^*\mathbb{R}$, we write $a \lesssim b$ to mean $a < b$ or $a \approx b$.

Let X be a topological space. The σ -algebra on X is always taken to be the Borel σ -algebra and is denoted by $\mathcal{B}[X]$. We use $\mathcal{M}_1(X)$ to denote the collection

of all countably additive probability measures on $(X, \mathcal{B}[X])$ and let $\mathcal{M}(X)$ denote the collection of all *charges*, that is, finitely-additive probability measures, on $(X, \mathcal{B}[X])$. An internal probability measure μ on $(^*X, ^*\mathcal{B}[X])$ is an element of $^*\mathcal{M}(X)$. Namely, an internal probability measure μ on $(^*X, ^*\mathcal{B}[X])$ is an internal function from $^*\mathcal{B}[X] \rightarrow ^*[0, 1]$ such that

- (1) $\mu(\emptyset) = 0$;
- (2) $\mu(^*X) = 1$; and
- (3) for $A, B \in ^*\mathcal{B}[X]$ with $A \cap B = \emptyset$, $\mu(A \cup B) = \mu(A) + \mu(B)$.

We use $(^*X, \overline{^*\mathcal{B}[X]}, \overline{\mu})$ to denote the Loeb extension of the internal probability space $(^*X, ^*\mathcal{B}[X], \mu)$.

3. Wasserstein metric

Let P be a charge on $(X, \mathcal{B}[X])$ and let ν be an internal probability measure on $(^*X, ^*\mathcal{B}[X])$. Suppose $^*P(^*A) \approx \nu(^*A)$ for all $A \in \mathcal{B}[X]$. We investigate the relation between *P and ν . Note that if $\nu = ^*P_1$ for some charge P_1 on $(X, \mathcal{B}[X])$, by the transfer principle, it is easy to see that $P_1 = P$. So we are interested in the case where ν is not the nonstandard extension of any standard charge.

Integration with respect to charges is similar to integration with respect to countably additive probability measures except that we only have finite additivity. However, we need to be careful about what functions are integrable. We quote the following result regarding integrability of charges.

Lemma 3.1 ([6, Corollary 4.5.9]). *Let P be a charge on $(X, \mathcal{B}[X])$. Let f be a bounded real-valued measurable function on X . Then f is integrable with respect to P .*

The Wasserstein distance is usually defined for countably additive probability measures. In this paper we extend the definition of Wasserstein distance to charges.

Definition 3.2. Let μ and ν be two charges on some bounded metric space (Y, d) with Borel σ -algebra $\mathcal{B}[Y]$. The Wasserstein distance between μ and ν is given by

$$(3.1) \quad W(\mu, \nu) = \sup \left\{ \left| \int f \, d\mu - \int f \, d\nu \right| : f \in \mathcal{L}_1(Y) \right\}$$

where $\mathcal{L}_1(Y)$ denote the set of 1-Lipschitz functions from Y to \mathbb{R} , i.e. those functions f such that $|f(x) - f(y)| \leq d(x, y)$ for all $x, y \in Y$.

As Y is bounded, every $f \in \mathcal{L}_1(Y)$ is bounded measurable. Thus, by Lemma 3.1, the Wasserstein metric on charges is well-defined. The following two lemmas provide a sufficient criterion to establish that the Wasserstein distance between two given internal probability measures is infinitesimal.

Lemma 3.3. *Let (X, d) be a bounded σ -compact metric space. Let P be a countably additive probability measure on $(X, \mathcal{B}[X])$ and let ν be an internal probability measure on $({}^*X, {}^*\mathcal{B}[X])$. Suppose for every $n \in \mathbb{N}$, there is a countable partition $\{V_i^n : i \in \mathbb{N}\}$ of X consisting of nonempty Borel sets with diameters no greater than $1/n$ such that ${}^*P({}^*V_i^n) \approx \nu({}^*V_i^n)$. Then ${}^*W(\nu, {}^*P) \approx 0$.*

PROOF: Fix $n \in \mathbb{N}$. Let $\{V_i^n : i \in \mathbb{N}\}$ be a countable partition of X consisting of nonempty Borel sets with diameters no greater than $1/n$ such that ${}^*P({}^*V_i^n) \approx \nu({}^*V_i^n)$. For every $i \in \mathbb{N}$, let B_i denote the set of all internal functions $g : {}^*\mathbb{N} \mapsto {}^*\mathcal{B}[X]$ such that

- (1) $g(i) = {}^*V_i^n$;
- (2) $\forall k \in {}^*\mathbb{N} \quad \sup\{d(x, y) : x, y \in g(k)\} \leq 1/n$;
- (3) $\forall k \in {}^*\mathbb{N} \quad |\nu(g(k)) - {}^*P(g(k))| \leq 1/i$.

Let \mathcal{B} be the collection of all B_i . Then \mathcal{B} has countable cardinality and the finite intersection property. By the saturation principle, there is an internal function g_0 which is an element of B_i for all $i \in \mathbb{N}$. Note that $g_0(i) = {}^*V_i^n$ for all $i \in \mathbb{N}$ and $\nu(g_0(k)) \approx {}^*P(g_0(k))$ for all $k \in {}^*\mathbb{N}$. By overspill, there is a $K \in {}^*\mathbb{N} \setminus \mathbb{N}$ such that $g_0(i) \neq \emptyset$ for all $i \leq K + 1$. For any $i \leq K$, pick $x_i \in g_0(i)$. Pick some $F \in {}^*\mathcal{L}_1(X)$. As $F \in {}^*\mathcal{L}_1(X)$ and the diameter of every $g_0(i)$ is no greater than $1/n$, we have

$$(3.2) \quad \forall i \leq K \quad \forall x \in g_0(i) \quad |F(x_i) - F(x)| \leq \frac{1}{n}.$$

Finally, we pick some $x_{K+1} \in {}^*X \setminus \bigcup_{i \leq K} g_0(i)$. As X is bounded and $F \in {}^*\mathcal{L}_1(X)$, we know that the function $|F(x) - F(x_{K+1})|$ is bounded by a standard real number. Define $g_1 : \{1, 2, \dots, K + 1\} \rightarrow {}^*\mathcal{B}[X]$ to be the internal function such that $g_1(i) = g_0(i)$ for all $i \leq K$ and $g_1(K + 1) = {}^*X \setminus \bigcup_{i \leq K} g_0(i)$. Hence we have

$$(3.3) \quad \left| \int_{{}^*X} F(x)\nu(dx) - \sum_{i \leq K+1} \int_{g_1(i)} F(x_i)\nu(dx) \right|$$

$$(3.4) \quad \leq \sum_{i \leq K} \int_{g_1(i)} |F(x) - F(x_i)|\nu(dx) + \int_{g_1(K+1)} |F(x) - F(x_{K+1})|\nu(dx)$$

$$(3.5) \quad \lesssim \frac{1}{n}.$$

Similarly, we have

$$(3.6) \quad \left| \int_{{}^*X} F(x){}^*P(dx) - \sum_{i \leq K+1} \int_{g_1(i)} F(x_i){}^*P(dx) \right| \lesssim \frac{1}{n}.$$

We now compare $\sum_{i \leq K+1} \int_{g_1(i)} F(x_i) \nu(dx)$ and $\sum_{i \leq K+1} \int_{g_1(i)} F(x_i) {}^*P(dx)$. Note that

$$(3.7) \quad \left| \sum_{i \leq K+1} \int_{g_1(i)} F(x_i) \nu(dx) - \sum_{i \leq K+1} \int_{g_1(i)} F(x_i) {}^*P(dx) \right|$$

$$(3.8) \quad = \left| \sum_{i \leq K+1} F(x_i) (\nu(g_1(i)) - {}^*P(g_1(i))) \right|$$

$$(3.9) \quad = \left| \sum_{i \leq K+1} (F(x_1) + k_i) (\nu(g_1(i)) - {}^*P(g_1(i))) \right|$$

$$(3.10) \quad = \left| \sum_{i \leq K+1} F(x_1) (\nu(g_1(i)) - {}^*P(g_1(i))) + \sum_{i \leq K+1} k_i (\nu(g_1(i)) - {}^*P(g_1(i))) \right|$$

$$(3.11) \quad = \left| \sum_{i \leq K+1} k_i (\nu(g_1(i)) - {}^*P(g_1(i))) \right|,$$

where k_i is the difference between $F(x_i)$ and $F(x_1)$.

As X is bounded and $F \in {}^*\mathcal{L}_1(X)$, we know that $k_i \in \text{NS}({}^*\mathbb{R})$ for all $i \leq K + 1$. Suppose $\left| \sum_{i \leq K+1} k_i (\nu(g_1(i)) - {}^*P(g_1(i))) \right| \approx 0$ then we have

$$(3.12) \quad \left| \int_{*X} F(x) \nu(dx) - \int_{*X} F(x) {}^*P(dx) \right| \lesssim \frac{2}{n}.$$

As n is arbitrary, we know that $\int_{*X} F(x) \nu(dx) \approx \int_{*X} F(x) {}^*P(dx)$ hence we have ${}^*W(\nu, {}^*P) \approx 0$ by Definition 3.2. Thus, in order to finish the proof, it is sufficient to prove $\left| \sum_{i \leq K+1} k_i (\nu(g_1(i)) - {}^*P(g_1(i))) \right| \approx 0$.

Claim 3.4. $\sum_{i \leq K+1} k_i (\nu(g_1(i)) - {}^*P(g_1(i))) \approx 0$.

PROOF: Pick some $k \in \mathbb{N}$. As P is countably additive, there exists $m \in \mathbb{N}$ such that $\sum_{i \leq m} \nu(g_1(i)) \geq 1 - 1/k$ and $\sum_{i \leq m} {}^*P(g_1(i)) \geq 1 - 1/k$. Thus we have

$$(3.13) \quad \sum_{i \leq K+1} k_i (\nu(g_1(i)) - {}^*P(g_1(i)))$$

$$(3.14) \quad = \sum_{i \leq m} k_i (\nu(g_1(i)) - {}^*P(g_1(i))) + \sum_{m < i \leq K+1} k_i (\nu(g_1(i)) - {}^*P(g_1(i)))$$

$$(3.15) \quad \approx \sum_{m < i \leq K+1} k_i (\nu(g_1(i)) - {}^*P(g_1(i))) \leq \frac{\max\{k_i : i \leq K + 1\}}{k}.$$

As $\max\{k_i : i \leq K + 1\}$ is near-standard and k is arbitrary, we have the desired result. □

Hence we have completed the proof of Lemma 3.3. □

By using a similar argument as in Lemma 3.3, we obtain the following result for charges.

Lemma 3.5. *Let (X, d) be a bounded σ -compact metric space. Let P be a charge on $(X, \mathcal{B}[X])$ and let ν be an internal probability measure on $({}^*X, {}^*\mathcal{B}[X])$. Suppose for every $n \in \mathbb{N}$, there is a finite partition $\{V_i^n : i \leq N\}$ of X consisting of nonempty Borel sets with diameters no greater than $1/n$ such that ${}^*P({}^*V_i^n) \approx \nu({}^*V_i^n)$. Then ${}^*W(\nu, {}^*P) \approx 0$.*

The following example shows that Lemma 3.3 and Lemma 3.5 are sharp.

Example 3.6. Let $X = (0, 1)$ endowed with the standard metric and Borel σ -algebra $\mathcal{B}[X]$. Let ν be an internal probability measure concentrating on some infinitesimal ε . Let P be a charge with $P((0, 1 - 1/n)) = 0$ for all $n \in \mathbb{N}$ and $P((0, 1)) = 1$. Pick $n \in \mathbb{N}$. We can pick $m \geq n$ such that $[1/m, 1 - 1/m]$ is a nonempty subset of $(0, 1)$. We can partition $[1/m, 1 - 1/m]$ into k Borel sets with diameter no greater than $1/n$ for some $k \in \mathbb{N}$. We denote these sets by V_i^n for $i \leq k$. Let $V_{k+j}^n = [1/(2^j m), 1/(2^{j-1} m)] \cup (1 - 1/(2^{j-1} m), 1 - 1/(2^j m)]$. Thus, $\{V_i^n : i \in \mathbb{N}\}$ forms a countable partition of $(0, 1)$ consisting of Borel sets with diameter no greater than $1/n$. Note that $\nu({}^*V_i^n) = {}^*P({}^*V_i^n) = 0$.

On the other hand, let f be the identity function from $(0, 1) \rightarrow (0, 1)$. Then we have $\int {}^*f(x)\nu(dx) \approx 0$ while $\int {}^*f(x){}^*P(dx) \approx 1$. Hence, we have ${}^*W(\nu, {}^*P) \not\approx 1$.

We conclude this section with the following theorem, which is a direct consequence of Lemma 3.5.

Theorem 3.7. *Let (X, d) be a bounded σ -compact metric space. Let P be a charge on $(X, \mathcal{B}[X])$ and let ν be an internal probability measure on $({}^*X, {}^*\mathcal{B}[X])$. Suppose $\nu({}^*B) \approx {}^*P({}^*B)$ for all $B \in \mathcal{B}[X]$. Then ${}^*W(\nu, {}^*P) \approx 0$.*

4. Construction of standard charges

To construct exotic standard objects, in this section, we discuss two procedures on constructing standard measures/charges on metric spaces. We also establish some connections between standard objects obtained from these two different approaches.

We begin with an internal probability measure and then use the transfer principle to push it down to get a charge.

Definition 4.1. Let $(X, \mathcal{B}[X])$ be a measurable space and let ν be an internal probability measure on $({}^*X, {}^*\mathcal{B}[X])$. Its *internal push-down* is a function $\nu^p : \mathcal{B}[X] \mapsto [0, 1]$ defined by $\nu^p(A) = \text{st}(\nu({}^*A))$.

The following lemma follows immediately from Definition 4.1.

Lemma 4.2. *Let ν be an internal probability measure on $({}^*X, {}^*\mathcal{B}[X])$. Then its internal push-down measure ν^p is a charge on $(X, \mathcal{B}[X])$.*

In most cases, ν^p is not a countably additive probability measure. Moreover, the nonstandard extension of ν^p is usually not the same as ν .

The following theorem is a direct consequence of Theorem 3.7.

Theorem 4.3. *Let (X, d) be a bounded σ -compact metric space. Let ν be an internal probability measure on $(*X, *\mathcal{B}[X])$. Let ν^p be the internal push-down of ν . Then $*W(\nu, *\nu^p) \approx 0$.*

Although the internal push-down of an internal probability measure always exists, it is merely finitely-additive in most cases. The properties of internal push-down are closely related to the internal probability measure via transfer principle. To get a countably additive probability measure, we shall use the standard part map to push down the Loeb measure.

Definition 4.4. Let X be a Hausdorff space with Borel σ -algebra $\mathcal{B}[X]$, let ν be an internal probability measure defined on $(*X, *\mathcal{B}[X])$, and let

$$(4.1) \quad \mathcal{C} = \{C \subset X : \text{st}^{-1}(C) \in \overline{*}\mathcal{B}[X]\}.$$

The *external push-down measure* ν_p is defined on the set \mathcal{C} by $\nu_p(C) = \overline{\nu}(\text{st}^{-1}(C))$.

The following two theorems guarantee that $\text{st}^{-1}(C) \in \overline{*}\mathcal{B}[X]$ for all $C \in \mathcal{B}[X]$ under moderate assumptions.

Theorem 4.5 ([5, Theorem 4.3.2]). *Let X be a regular topological space and let P be an internal probability measure on $(*X, *\mathcal{B}[X])$. Suppose $\text{NS}(*X) \in \overline{*}\mathcal{B}[X]$. Then $\text{st}^{-1}(A) \in \overline{*}\mathcal{B}[X]$ for all $A \in \mathcal{B}[X]$ (i.e., st is Borel measurable).*

Theorem 4.6 ([8, Theorem 5.6]). *Let X be a Čech-complete Tychonoff space with Borel σ -algebra $\mathcal{B}[X]$. Then $\text{NS}(*X) \in \overline{*}\mathcal{B}[X]$.*

In particular, we have $\text{NS}(*X) \in \overline{*}\mathcal{B}[X]$ for regular locally compact spaces; for complete metric spaces; and for regular σ -compact spaces.

For general Hausdorff Borel measurable space $(Y, \mathcal{B}[Y])$, the external push-down measure ν_p may not be a countably additive probability measure. In fact, if $\overline{\nu}(\text{NS}(*X)) = 0$ then ν_p is a null measure on $(X, \mathcal{B}[X])$. However, when Y is compact, the following theorem guarantees that ν_p is a countably additive probability measure on $(Y, \mathcal{B}[Y])$.

Theorem 4.7 ([7, Theorem 13.4.1]). *Let X be a Hausdorff space equipped with Borel σ -algebra $\mathcal{B}[X]$, and let ν be an internal probability measure defined on $(*X, *\mathcal{B}[X])$ with $\overline{\nu}(\text{NS}(*X)) = 1$. Then the external push-down measure ν_p of ν is the completion of a countably additive regular Borel probability measure.*

Given an internal probability measure ν on $(*X, *\mathcal{B}[X])$, it is easy to see that the total variation distance between ν and $*\nu_p$ may be large. For example, if ν is an internal probability measure concentrating on some infinitesimal ε then ν_p is a degenerate probability measure at point 0. The total variation distance between ν and $*\nu_p$ is 1 in this case. However, we show that ν and $*\nu_p$ are close in Wasserstein metric. We start by stating the following well-known definition.

Definition 4.8. Let X be a topological space and let $(X, \mathcal{B}[X], P)$ be a Borel probability space. A set $A \subset X$ is a P -continuity set if the boundary ∂A is contained in a measure 0 set.

Recall that a countably additive probability measure P on a Borel measurable space $(X, \mathcal{B}[X])$ is Radon provided that $P(E) = \sup\{P(K) : K \text{ compact and } K \subset E\}$. The following result, due to R. Anderson, is the first major result on representing standard measures using nonstandard measures via the standard part map.

Lemma 4.9 ([5, Theorem 4.1]). *Let $(X, \mathcal{B}[X], P)$ be a countably additive Radon probability measure. Then st is measure-preserving from $({}^*X, {}^*\mathcal{B}[X], {}^*P)$ to $(X, \mathcal{B}[X], P)$, i.e. $P(A) = {}^*P(\text{st}^{-1}(A))$ for all $A \in \mathcal{B}[X]$.*

The following result gives a nonstandard characterization of compact sets.

Theorem 4.10 ([5, Theorem 3.5.1]). *A set $A \subset X$ is compact if and only if for each $y \in {}^*A$, there is an $x \in A$ such that y is in the monad of x .*

Lemma 4.11 ([5, Exercise 4.27]). *If X is a Hausdorff regular space and A is an internal subset of $\text{NS}({}^*X)$, then $E = \text{st}(A)$ is compact.*

The following result is a partial converse of Lemma 4.9. It follows immediately from the fact that every Borel probability measure on a Polish space is Radon.

Lemma 4.12. *Let X be a σ -compact metric space with Borel σ -algebra $\mathcal{B}[X]$. Let ν be an internal probability measure on $(X, \mathcal{B}[X])$. Then the push-down measure ν_p is a Radon measure on $(X, \mathcal{B}[X])$.*

Lemma 4.13. *Let X be a σ -compact metric space with Borel σ -algebra $\mathcal{B}[X]$. Let ν be an internal probability measure on $({}^*X, {}^*\mathcal{B}[X])$. Suppose ν_p is a countably additive probability measure on $(X, \mathcal{B}[X])$. For every $n \in \mathbb{N}$, there exists a countable partition $\{A_i : i \in \mathbb{N}\}$ of X consisting of Borel sets with diameter no greater than $1/n$ such that $\nu({}^*A_i) \approx {}^*\nu_p({}^*A_i)$ for all $i \in \mathbb{N}$.*

PROOF: Pick $n \in \mathbb{N}$. For every $x \in X$, there are uncountably many open balls containing x with diameter no greater than $1/n$. The boundaries of these open balls form an uncountable collection of disjoint sets. Thus, for every $x \in X$, we can pick an open ball U_x containing x such that its diameter is no greater than $1/n$ and it is a ν_p -continuity set. As X is a Polish space, there is a countable subcollection of $\{U_x : x \in X\}$ that covers X . Denote this countable subcollection by $\mathcal{K}_n = \{U_{x_i} : i \in \mathbb{N}\}$.

Pick $i, j \leq m$. Note that $\partial(U_{x_i} \cap U_{x_j}) \subset \partial U_{x_i} \cup \partial U_{x_j}$ and $\partial(U_{x_i} \cup U_{x_j}) \subset \partial U_{x_i} \cup \partial U_{x_j}$. Hence any finite intersection (union) of elements from \mathcal{K}_n is a bounded open ν_p -continuity set. For any $i \in \mathbb{N}$, let

$$(4.2) \quad V_i = U_{x_i} \setminus \bigcup_{j < i} (U_{x_j} \cap U_{x_i}).$$

The following claim shows that $\{V_i : i \in \mathbb{N}\}$ is the desired partition.

Claim 4.14. $\nu(*U) \approx * \nu_p(*U)$ for any open ν_p -continuity set U .

PROOF: Pick any open ν_p -continuity set U . By Lemma 4.12, ν_p is a countably additive Radon probability measure. As U is a ν_p -continuity set, by Lemma 4.9, we have

$$(4.3) \quad \overline{* \nu_p}(\text{st}^{-1}(U)) = \nu_p(U) = \nu_p(\overline{U}) = \overline{* \nu_p}(\text{st}^{-1}(\overline{U})).$$

By the construction of ν_p , we also have $\overline{\nu}(\text{st}^{-1}(U)) = \nu_p(U)$ and $\overline{\nu}(\text{st}^{-1}(\overline{U})) = \nu_p(\overline{U})$. Thus, we have $\overline{\nu}(\text{st}^{-1}(U)) = \overline{* \nu_p}(\text{st}^{-1}(U))$ and $\overline{\nu}(\text{st}^{-1}(\overline{U})) = \overline{* \nu_p}(\text{st}^{-1}(\overline{U}))$. As U is an open set, we have $\text{st}^{-1}(U) \subset *U \cap \text{NS}(*X) \subset \text{st}^{-1}(\overline{U})$. As ν_p is a countably additive probability measure we have $\overline{\nu}(\text{NS}(*X)) = 1$. Hence $\overline{\nu}(*U) = \overline{\nu}(*U \cap \text{NS}(*X))$. Hence

$$(4.4) \quad \nu(*U) \approx \overline{\nu}(*U \cap \text{NS}(*X)) = * \nu_p(*U)$$

for all open ν_p -continuity set U . □

Hence we have completed the proof. □

We obtain the following result from Lemma 3.3 and Lemma 4.13.

Theorem 4.15. *Suppose X is a bounded σ -compact metric space. Let ν be an internal probability measure on $(*X, * \mathcal{B}[X])$. Suppose ν_p is a countably additive probability measure on $(X, \mathcal{B}[X])$. Then $*W(\nu, \nu_p) \approx 0$.*

The compact version of Lemma 4.13 and Theorem 4.15 will be proved in the paper which is in preparation. The structure of the proofs are similar.

By Theorems 4.3 and 4.15, we immediately obtain the following result.

Theorem 4.16. *Suppose X is a bounded σ -compact metric space. Let ν be an internal probability measure on $(*X, * \mathcal{B}[X])$. Let ν_p and ν^p denote the external push-down and internal push-down of ν , respectively. Suppose ν_p is a countably additive probability measure on $(X, \mathcal{B}[X])$. Then $W(\nu_p, \nu^p) = 0$.*

As ν_p and ν^p are different objects, the Wasserstein distance is only a pseudo-metric on $\mathcal{M}(X)$. In summary, we have the following result.

Theorem 4.17. *Suppose X is a bounded σ -compact metric space. Let ν and μ be two internal probability measures on $(*X, * \mathcal{B}[X])$. Suppose both ν_p and μ_p are countably additive probability measures on $(X, \mathcal{B}[X])$. Then the following statements are equivalent:*

- (1) $\nu_p = \mu_p$;
- (2) $*W(\nu, \mu) \approx 0$;
- (3) $W(\nu^p, \mu^p) = 0$;
- (4) $W(\nu_p, \mu_p) = 0$.

PROOF: As ν_p and μ_p are countably additive probability measures on $(X, \mathcal{B}[X])$, we have $\nu_p = \mu_p$ if and only if $W(\nu_p, \mu_p) = 0$. By Theorem 4.15 we know that

$W(\nu_p, \mu_p) = 0$ if and only if $*W(\nu, \mu) \approx 0$. By Theorem 4.3 we have $*W(\nu, \mu) \approx 0$ if and only if $W(\nu^p, \mu^p) = 0$. By Theorems 4.15 and 4.16, we have $*W(\nu, \mu) \approx 0$ if and only if $W(\nu_p, \mu^p) = 0$ hence finishing the proof. \square

5. Weak convergence of charges

In this section, we define a notion of weak convergence for charges and show that every charge is a weak limit of a countable sequence of countably additive probability measures under moderate conditions.

We start by proving the result when the underlying space is a compact metric space.

Lemma 5.1. *Let X be a compact metric space equipped with Borel σ -algebra $\mathcal{B}[X]$. For every charge P on $(X, \mathcal{B}[X])$, there is a countably-additive probability measure μ on $(X, \mathcal{B}[X])$ such that $W(P, \mu) = 0$.*

PROOF: Fix a charge P on $(X, \mathcal{B}[X])$. Let $\mu = (*P)_p$. As X is compact, μ defines a countably additive probability measure on $(X, \mathcal{B}[X])$. We always have $P = (*P)^p$, so by Theorem 4.16, we have the desired result. \square

To generalize Lemma 5.1 to non-compact spaces, we need several results on nonstandard integration theory. We start by quoting the following lemma.

Lemma 5.2 ([9, Lemma 6.5]). *Let X be a compact Hausdorff space equipped with Borel σ -algebra $\mathcal{B}[X]$, let ν be an internal probability measure on $(*X, *\mathcal{B}[X])$, and let $f: X \rightarrow \mathbb{R}$ be a bounded Borel measurable function. Define $g: *X \rightarrow \mathbb{R}$ by $g(s) = f(\text{st}(s))$. Then we have $\int f \, d\nu_p = \int g \, d\bar{\nu}$.*

Theorem 5.3. *Let X be a compact Hausdorff space equipped with Borel σ -algebra $\mathcal{B}[X]$, let ν be an internal probability measure on $(*X, *\mathcal{B}[X])$, and let $f: X \rightarrow \mathbb{R}$ be a bounded continuous function. Then we have $\int f \, d\nu_p = \int *f \, d\bar{\nu}$.*

PROOF: Since X is compact, we can define $g: *X \rightarrow \mathbb{R}$ by $g(s) = f(\text{st}(s))$. By Lemma 5.2, we have $\int f \, d\nu_p = \int g \, d\bar{\nu}$. As f is continuous, we have $*f(x) \approx g(x)$ for all $x \in *X$. Thus we have $\int g \, d\bar{\nu} = \int *f \, d\bar{\nu}$, completing the proof. \square

We now consider the relation between internal integration and integration with respect to internal push-down measures.

Theorem 5.4. *Let X be a metric space equipped with Borel σ -algebra $\mathcal{B}[X]$. Let ν be an internal probability measure on $(*X, *\mathcal{B}[X])$ and let $f: X \rightarrow \mathbb{R}$ be a bounded Borel measurable function. Then we have*

$$(5.1) \quad \int_{*X} *f(x)\nu(dx) \approx \int_X f(x)\nu^p(dx).$$

PROOF: Fix $\varepsilon \in \mathbb{R}_{>0}$. Let $\{K_1, K_2, \dots, K_n\}$ be a partition of a large enough interval of \mathbb{R} containing the range of f such that every $K_i \in \{K_1, K_2, \dots, K_n\}$ is

an interval with diameter not greater than ε . For $i \leq n$, let $F_i = f^{-1}(K_i)$. Then $\{F_1, \dots, F_n\} \subset \mathcal{B}[X]$ is a partition of X such that $|f(x) - f(x')| < \varepsilon$ for every $x, x' \in F_i$ for every $i = 1, 2, \dots, n$. Pick $x_i \in F_i$ for every $i = 1, 2, \dots, n$. Define $g: X \rightarrow \mathbb{R}$ by letting $g(x) = f(x_i)$ if $x \in F_i$ for every $i = 1, 2, \dots, n$. Then g is a simple bounded measurable real valued function on X . Thus, by Lemma 3.1, both g and f are integrable with respect to ν^p .

We now have

$$\begin{aligned} \left| \int_{*X} *f(x)\nu(dx) - \int_X f(x)\nu^p(dx) \right| &\leq \left| \int_{*X} *f(x)\nu(dx) - \int_{*X} *g(x)\nu(dx) \right| \\ &\quad + \left| \int_{*X} *g(x)\nu(dx) - \int_X g(x)\nu^p(dx) \right| \\ &\quad + \left| \int_X g(x)\nu^p(dx) - \int_X f(x)\nu^p(dx) \right| \end{aligned}$$

where all internal integrals are over $*X$ and all standard integrals are over X .

By the transfer principle, we have $|*f(x) - *f(x')| < \varepsilon$ for every $x, x' \in *F_i$ and for every $i = 1, 2, \dots, n$. Thus, we have $|*f(x) - *g(x)| < \varepsilon$ for all $x \in *X$. Thus, we have $|\int_{*X} *f(x)\nu(dx) - \int_{*X} *g(x)\nu(dx)| \leq \int_{*X} |*f(x) - *g(x)|\nu(dx) < \varepsilon$. Similarly, we have $|\int_X g(x)\nu^p(dx) - \int_X f(x)\nu^p(dx)| < \varepsilon$. For the term $|\int_{*X} *g(x)\nu(dx) - \int_X g(x)\nu^p(dx)|$, we have:

$$(5.2) \quad \int_{*X} *g(x)\nu(dx) = \sum_{i=1}^n \int_{*F_i} *g(x)\nu(dx)$$

$$(5.3) \quad = \sum_{i=1}^n *f(x_i)\nu(*F_i) = \sum_{i=1}^n f(x_i)\nu(*F_i) \approx \sum_{i=1}^n f(x_i)\nu^p(F_i)$$

$$(5.4) \quad = \sum_{i=1}^n \int_{F_i} g(x)\nu^p(dx) = \int_X g(x)\nu^p(dx).$$

Thus, we have $|\int_{*X} *f(x)\nu(dx) - \int_X f(x)\nu^p(dx)| \lesssim 2\varepsilon$. As ε is arbitrary, we have $\int_{*X} *f(x)\nu(dx) \approx \int_X f(x)\nu^p(dx)$. □

The following corollary is a direct consequence of Theorems 5.3 and 5.4:

Corollary 5.5. *Let X be a compact Hausdorff space equipped with Borel σ -algebra $\mathcal{B}[X]$, let ν be an internal probability measure on $(*X, *\mathcal{B}[X])$, and let $f: X \rightarrow \mathbb{R}$ be a bounded continuous function. Then we have $\int f d\nu_p = \int f d\nu^p$.*

Before we establish the main result of this section, we introduce the following definition.

Definition 5.6. A sequence of charges P_n is said to converge weakly to a charge P if $\int f dP_n \rightarrow \int f dP$ for all bounded uniformly continuous real-valued function f .

If $\{P_n\}_{n \in \mathbb{N}}$ and P are countably additive probability measures, we have the following well-known result.

Theorem 5.7 (The Portmanteau theorem [16, Theorem 10.1.1]). *Suppose that $\{P_n\}_{n \in \mathbb{N}}$ is a sequence of countably additive probability measures and P is a countably additive measure. Then the following are equivalent:*

- (1) $\int f \, dP_n \rightarrow \int f \, dP$ for all bounded continuous functions f ;
- (2) $\int f \, dP_n \rightarrow \int f \, dP$ for all bounded uniformly continuous functions f ;
- (3) $P_n(A) \rightarrow P(A)$ for all P -continuity sets A .

We also need the following theorem from point-set topology.

Theorem 5.8 ([14, Theorem 1]). *Let f be a real-valued function defined on some closed subset E of a metric space S . Suppose f is M -Lipschitz continuous for some $M \in \mathbb{R}$. Then f can be extended to an M -Lipschitz continuous function on S .*

We are now able to establish the main result of this section.

Theorem 5.9. *Let X be a separable metric space equipped with Borel σ -algebra $\mathcal{B}[X]$. Let P be a charge on $(X, \mathcal{B}[X])$. There is a sequence $\{P_n\}_{n \in \mathbb{N}}$ of finitely supported probability measures that converges to P weakly if and only if X is totally bounded.*

PROOF: Suppose X is totally bounded. In the case that X is compact, the result follows immediately from Corollary 5.5 by letting $\nu = *P$.

In the case that X is not compact, let \hat{X} denote the completion of X . Then \hat{X} is a compact space. We can extend P to a charge on $(\hat{X}, \mathcal{B}[\hat{X}])$ by letting $P(A) = P(A \cap \hat{X})$ for $A \in \mathcal{B}[\hat{X}]$. As \hat{X} is compact, by Theorem 4.7, $(*P)_p$ is a countably additive probability measure on $(\hat{X}, \mathcal{B}[\hat{X}])$. Pick $n \in \mathbb{N}$. As X is totally bounded and dense in \hat{X} , there is a finite collection of open balls such that their closure covers \hat{X} . Thus, we can decompose \hat{X} into finitely many mutually disjoint Borel sets $\{B_i^n : i \leq m\}$ with diameters no greater than $1/n$ and each B_i contain at least one element x_i from X . Define a finitely supported probability measure P_n on $(\hat{X}, \mathcal{B}[\hat{X}])$ by letting $P_n(\{x_i\}) = *P_p(B_i)$ for every $i \leq m$.

Claim 5.10. *The sequence $\{P_n\}_{n \in \mathbb{N}}$ converges to $*P_p$ weakly.*

PROOF: Pick a uniformly continuous function $f: \hat{X} \rightarrow \mathbb{R}$ and a positive $\varepsilon \in \mathbb{R}$. There exists $j \in \mathbb{N}$ such that $|f(x) - f(y)| < \varepsilon$ if $d(x, y) < 1/j$ for all $x, y \in \hat{X}$. Then, for every $n \geq j$ we have

$$\begin{aligned}
 (5.5) \quad & \left| \int_{\hat{X}} f(x) \, d*P_p - \int_{\hat{X}} f(x) \, dP_n \right| \\
 (5.6) \quad & = \left| \sum \int_{B_i^n} f(x) \, d*P_p - \sum \int_{B_i^n} f(x) \, dP_k \right|
 \end{aligned}$$

$$(5.7) \quad = \left| \sum \int_{B_i^n} f(x) d^*P_p - \sum \int_{B_i^n} f(x_i) d^*P_p \right|$$

$$(5.8) \quad \leq \int_{B_i^n} |f(x) - f(x_i)| d^*P_p < \varepsilon.$$

By Theorem 5.7 we have the desired result. □

We now show that the sequence $(P_n)_{n \in \mathbb{N}}$ converges to P weakly. Pick a bounded uniformly continuous function $f: X \rightarrow \mathbb{R}$. We can extend f to a bounded continuous function $\hat{f}: \hat{X} \rightarrow \mathbb{R}$. By Corollary 5.5 and Claim 5.10, we have $\int_{\hat{X}} \hat{f} dP_n \rightarrow \int_{\hat{X}} \hat{f} d(^*P)_p = \int_{\hat{X}} \hat{f} dP$. As the supports of P_n and P are subsets of X , we have $\int_{\hat{X}} \hat{f} dP_n = \int_X f dP_n$ for all $n \in \mathbb{N}$ and $\int_{\hat{X}} \hat{f} dP = \int_X f dP$. Thus the sequence $\{P_n\}_{n \in \mathbb{N}}$ converges to P weakly.

Now suppose X is not totally bounded.

Claim 5.11. *There exists a closed countably infinite discrete subset Y of X such that for every pair of distinct points $y_1, y_2 \in Y$, we have $d(y_1, y_2) > \varepsilon$ for some fixed $\varepsilon > 0$.*

PROOF: We explicitly construct a countably infinite discrete set. We pick any element $x_1 \in X$ in the first step. Suppose we have picked n distinct points $\{x_1, x_2, \dots, x_n\}$ up to step n . As X is not totally bounded, there exists $\varepsilon \in \mathbb{R}$ such that there is no finite cover of X by ε -open balls. Let U_i denote the ε -open ball centered at x_i for $i \leq n$. Thus, the set $X \setminus \bigcup_{i=1}^n U_i$ is nonempty. Pick any element in $X \setminus \bigcup_{i=1}^n U_i$ to be x_{n+1} . Note that $d(x_{n+1}, x_i) > \varepsilon$ for all $i \leq n$. Thus, we have constructed a countable discrete set $Y \subset X$ such that $d(y_1, y_2) > \varepsilon$ for every pair of distinct points $y_1, y_2 \in Y$. Hence, Y must be closed. □

Let \mathcal{A} be a non-principal ultrafilter and let P be a finitely additive probability measure concentrated on Y such that $P(A) = 1$ if and only if $A \in \mathcal{A}$. Suppose there is a sequence $\{P_n\}_{n \in \mathbb{N}}$ of countably additive probability measures converging to P weakly. As P concentrates on Y , without loss of generality, we can assume P_n concentrates on Y for every $n \in \mathbb{N}$ (replace P_n by $P_n/P_n(Y)$ if necessary).

Claim 5.12. *The sequence $\{P_n(B)\}_{n \in \mathbb{N}}$ converges to $P(B)$ on every subset $B \subset Y$.*

PROOF: Fix a set $B \subset Y$. Let $f: Y \rightarrow \mathbb{R}$ be the indicator function on B . By Claim 5.11, f is a Lipschitz continuous function. By Claim 5.11 and Theorem 5.8, f can be extended to a Lipschitz continuous function from X to \mathbb{R} . As $\{P_n\}_{n \in \mathbb{N}}$ converges to P weakly, we have $P_n(B) \rightarrow P(B)$ for every $B \subset Y$. □

As P_1 is countably additive, there is a finite set $B_1 \subset Y$ such that $P_1(B_1) > 3/4$. As $P(B_1) = 0$, there exists $n_2 \in \mathbb{N}$ such that $P_{n_2}(B_1) < 1/4$. We can pick a finite set $B_2 \subset Y$ such that $B_2 \cap B_1 = \emptyset$ and $P_{n_2}(B_2) > 3/4$. Following this procedure, we can extract a subsequence $\{P_{n_i} : i \in \mathbb{N}\}$ from $\{P_n\}_{n \in \mathbb{N}}$ and construct a sequence of finite sets $\{B_i : i \in \mathbb{N}\}$ such that

- (1) $B_i \cap \bigcup_{j < i} B_j = \emptyset$ for all $i \geq 2$;
- (2) $P_{n_i}(B_i) > 3/4$ for all $i \in \mathbb{N}$;
- (3) $P_{n_i}(\bigcup_{j < i} B_j) < 1/4$ for all $i \geq 2$.

Let $A = \bigcup_{i \in \mathbb{N}} B_{2i}$. Then $P(A)$ is either 0 or 1 but $\{P_{n_i}(A)\}_{i \in \mathbb{N}}$ is oscillating. \square

The Portmanteau theorem (Theorem 5.7) gives three equivalent statements for countably additive probability measures. It is natural to ask whether the same is true for charges. The following example shows that it is not the case.

Example 5.13. Consider the unit interval $(0, 1]$ equipped with Borel σ -algebra $\mathcal{B}[(0, 1]]$, let P be the internal push-down of the internal probability measure concentrating on $1/(2N)$ where $N \in {}^*\mathbb{N} \setminus \mathbb{N}$. For $n \in \mathbb{N}$, let P_n be the degenerate measure at $1/n$. Note that the sequence $(P_n)_{n \in \mathbb{N}}$ converges weakly to $({}^*P)_p$. As $({}^*P)_p(\{0\}) = 1$, using the same proof after Claim 5.10, we know that $\int f dP_n \rightarrow \int f dP$ for bounded uniformly continuous f . On the other hand, the sequence $\int \sin(1/x) dP_n$ does not converge although $\sin(1/x)$ is bounded continuous on $(0, 1]$. Let $A = \{1/(2n) : n \in \mathbb{N}\}$. As $P(\{0\}) = 0$, we know that A is a continuity set. Note that $P_n(A) = 1$ if n is even and $P_n(A) = 0$ if n is odd. So the sequence $(P_n(A))_{n \in \mathbb{N}}$ does not converge.

In light of Theorem 5.9 and Example 5.13, it is natural to ask the following two questions. Suppose P is a charge on a locally compact separable metric space X with Borel σ -algebra $\mathcal{B}[X]$, does there exist a sequence $\{P_n\}_{n \in \mathbb{N}}$ of countably additive probability measures such that $\int f dP_n \rightarrow \int f dP$ for every bounded continuous function f ? Does there exist a sequence $\{P_n\}_{n \in \mathbb{N}}$ of countably additive probability measures such that $P_n(A) \rightarrow P(A)$ for every P -continuity set A ? These questions are answered by Miklos Laczkovich, who communicated the following example.

Example 5.14 (Communicated by Miklos Laczkovich). Let P be a charge on all subsets of \mathbb{N} such that $P(\{k\}) = 0$ for all $k \in \mathbb{N}$ and $P(\{\mathbb{N}\}) = 1$ (such P can be constructed via an ultrafilter on the set of natural numbers). We extend P to all subsets of \mathbb{R} by letting $P(A) = P(A \cap \mathbb{N})$. We claim that there is no countable sequence $\{P_n\}_{n \in \mathbb{N}}$ of countably additive measures on \mathbb{R} such that $\int f dP_n \rightarrow \int f dP$ for every bounded continuous function f .

PROOF: Suppose there exists a sequence of P_n of countably additive probability measures on \mathbb{R} such that $\int f dP_n \rightarrow \int f dP$ for any bounded function f . We first show that if $k \geq 1$ is an integer and $0 < d < 1$, then $P_n([k - d, k + d]) \rightarrow 0$ as $n \rightarrow \infty$. Indeed, let $d < e < 1$ and let $f \geq 0$ be a continuous function which equals to 1 in $[k - d, k + d]$ and equals to 0 outside $(k - e, k + e)$. Then $P_n([k - d, k + d]) \leq \int f dP_n$, which converges to $\int f dP = 0$.

We now show that $P_n(\mathbb{N}) \rightarrow 1$ as $n \rightarrow \infty$. Assume that this is not valid. After passing to a subsequence, we have $P_n(\mathbb{N}) < 1 - a$ for all $n \in \mathbb{N}$, where $a > 0$. Then there exists $0 < d_1 < 1$ such that $P_1(U(\mathbb{N}, d_1)) < 1 - a/2$, where $U(\mathbb{N}, d_1) = \bigcup_{n \in \mathbb{N}} [n - d_1, n + d_1]$. Let $n_1 = 1$. As $P_{n_1}([1 - d_1, 1 + d_1])$, there exists

$n_2 > 1$ such that $P_{n_2}([1 - d_1, 1 + d_1]) < a/4$. As $P_{n_2}(\mathbb{N}) < 1 - a$, there exists $0 < d_2 < d_1$ such that

$$(5.9) \quad P_{n_2}([1 - d_1, 1 + d_1] \cup U(\mathbb{N}, d_2)) < 1 - \frac{a}{4}.$$

As $P_n([1 - d_1, 1 + d_1]) \rightarrow 0$ and $P_n([2 - d_2, 2 + d_2]) \rightarrow 0$, there exists an $n_3 > n_2$ such that

$$(5.10) \quad P_{n_3}([1 - d_1, 1 + d_1] \cup [2 - d_2, 2 + d_2]) < \frac{a}{4}.$$

Since $P_{n_3}(\mathbb{N}) < 1 - a$, there exists $0 < d_3 < d_2$ such that

$$(5.11) \quad P_{n_3}([1 - d_1, 1 + d_1] \cup [2 - d_2, 2 + d_2] \cup U(\mathbb{N}, d_3)) < 1 - \frac{a}{4}.$$

Continuing this procedure, we get positive numbers d_k and indices n_k such that $P_{n_k}(A) < 1 - a/4$ for all k , where $A = \bigcup_{k=1}^{\infty} [k - d_k, k + d_k]$. As there exists a bounded continuous function f which equals to 1 on \mathbb{N} and 0 outside of A , we have $\int f(x)P_{n_k}(dx) \leq P_{n_k}(A) < 1 - a/4$ for all k , contradicting with the fact that $\int f(x)P_{n_k}(dx) \rightarrow \int f(x)P(dx) = 1$.

Thus $P_n(\mathbb{N}) \rightarrow 1$ as $n \rightarrow \infty$. If $E \subset \mathbb{N}$, then $\limsup_{n \rightarrow \infty} P_n(E) \leq P(E)$ for all n . If $0 \leq f \leq 1$ is a continuous function which equals to 1 on E and 0 on $\mathbb{N} \setminus E$, then

$$(5.12) \quad P_n(E) \leq \int f(x)P_n(dx) \rightarrow \int f(x)P(dx) = P(E).$$

This implies that $P_n(E) \rightarrow P(E)$. If $\liminf_{n \rightarrow \infty} P_n(E) < P(E)$, then from

$$(5.13) \quad \limsup_{n \rightarrow \infty} P_n(\mathbb{N} \setminus E) \leq P(\mathbb{N} \setminus E)$$

and from additivity it follows that $\liminf_{n \rightarrow \infty} P_n(E) < 1$, which is impossible.

Thus, we have $P(E) = \lim_{n \rightarrow \infty} P_n(E)$ for all $E \subset \mathbb{N}$. By Vitali–Hahn–Saks theorem, P is countably additive, a contradiction. □

By using a similar argument, we can show that there is no countable sequence $\{P_n\}_{n \in \mathbb{N}}$ of countably additive probability measures on \mathbb{R} such that $P_n(A) \rightarrow P(A)$ for every P -continuity set A .

Example 5.14 shows that Theorem 5.9 is sharp. This implies that, when we talk about weak convergence to a charge, it is necessary to restrict ourselves to bounded uniformly continuous real-valued functions.

We conclude this paper by giving a nonstandard characterization of weak convergence defined in Definition 5.6. The work [4] gave a nonstandard characterization of weak convergence of countably additive probability measures. The main result of [4] is the following:

Theorem 5.15 ([4, Theorem 4]). *Let X be a metric space equipped with Borel σ -algebra $\mathcal{B}[X]$. Let $\{P_n\}_{n \in \mathbb{N}}$ be a sequence of countably additive probability measures on $(X, \mathcal{B}[X])$. Then the following are equivalent:*

- (1) *The measure $\{P_n\}_{n \in \mathbb{N}}$ converges weakly to some countably additive probability measure on $(X, \mathcal{B}[X])$.*
- (2) *For all infinite $N_1, N_2 \in {}^*\mathbb{N}$, we have $({}^*P_{N_1})_p(X) = 1$ and $({}^*P_{N_1})_p = ({}^*P_{N_2})_p$.*

By using Theorem 5.15, we present the following result.

Theorem 5.16. *Let X be a locally compact separable metric space equipped with Borel σ -algebra $\mathcal{B}[X]$. Let $\{P_n\}_{n \in \mathbb{N}}$ be a sequence of countably additive probability measures on $(X, \mathcal{B}[X])$. Then the following are equivalent:*

- (1) *$\{P_n\}_{n \in \mathbb{N}}$ converges weakly to some charge on $(X, \mathcal{B}[X])$.*
- (2) *For all infinite $N_1, N_2 \in {}^*\mathbb{N}$, we have $({}^*P_{N_1})_p = ({}^*P_{N_2})_p$.*

PROOF: Suppose that $\{P_n\}_{n \in \mathbb{N}}$ converges weakly to some charge P on $(X, \mathcal{B}[X])$. Let $\widehat{X} = X \cup \{a_0\}$ denote the metric one-point compactification of X . We extend each P_n and P to \widehat{X} by defining $P(\{a_0\}) = 0$ and $P_n(\{a_0\}) = 0$ for all $n \in \mathbb{N}$. As \widehat{X} is compact, by Theorem 4.7, we know that $({}^*P)_p$ is a countably additive probability measure on $(\widehat{X}, \mathcal{B}[\widehat{X}])$.

Claim 5.17. *The sequence $\{P_n\}_{n \in \mathbb{N}}$ converges weakly to $({}^*P)_p$ in $(\widehat{X}, \mathcal{B}[\widehat{X}])$.*

PROOF: Let f be a bounded continuous function from \widehat{X} to \mathbb{R} . Let g be the restriction of f to X . Then g is a bounded uniformly continuous function from X to \mathbb{R} . By assumption, we have $\int_X g \, dP_n \rightarrow \int_X g \, dP$. Thus, we have $\int_{\widehat{X}} f \, dP_n \rightarrow \int_{\widehat{X}} f \, dP$. By Corollary 5.5, we have $\int_{\widehat{X}} f \, dP = \int_{\widehat{X}} f \, d({}^*P)_p$, completing the proof. \square

By Theorem 5.15, we have $({}^*P_{N_1})_p = ({}^*P_{N_2})_p$ for all infinite $N_1, N_2 \in {}^*\mathbb{N}$.

Now suppose we have $({}^*P_{N_1})_p = ({}^*P_{N_2})_p$ for all infinite $N_1, N_2 \in {}^*\mathbb{N}$. For every $n \in {}^*\mathbb{N}$, *P_n can be extended to an internal probability measure on $({}^*\widehat{X}, {}^*\mathcal{B}[\widehat{X}])$ by letting ${}^*P_n(\{a_0\}) = 0$. By Theorem 5.15, we know that the sequence $\{P_n\}_{n \in \mathbb{N}}$ converges weakly to a countably additive probability measure μ on $(\widehat{X}, \mathcal{B}[\widehat{X}])$. Pick an element $y \in {}^*X$ such that y is in the monad of a_0 . Define an internal probability measure ν on $({}^*\widehat{X}, {}^*\mathcal{B}[\widehat{X}])$ as following:

- (1) $\nu(A) = {}^*\mu(A)$ for all $A \in {}^*\mathcal{B}[\widehat{X}]$ such that $\{a_0, y\} \cap A = \emptyset$;
- (2) $\nu(A) = {}^*\mu(A)$ for all $A \in {}^*\mathcal{B}[\widehat{X}]$ such that $\{a_0, y\} \subset A$;
- (3) $\nu(A) = {}^*\mu(A) - {}^*\mu(\{a_0\})$ for all $A \in {}^*\mathcal{B}[\widehat{X}]$ such that $y \notin A$ and $a_0 \in A$;
- (4) $\nu(A) = {}^*\mu(A) + {}^*\mu(\{a_0\})$ for all $A \in {}^*\mathcal{B}[\widehat{X}]$ such that $y \in A$ and $a_0 \notin A$.

By the internal definition principle and the fact that $\nu(\{a_0\}) = 0$, ν defines an internal probability measure on $({}^*X, {}^*\mathcal{B}[X])$. Thus, the internal push-down ν^p defines a charge on $(X, \mathcal{B}[X])$.

Claim 5.18. $\nu_p = \mu$.

PROOF: Pick a set $E \in \mathcal{B}[\widehat{X}]$. Suppose that $y \notin \text{st}^{-1}(E)$. Then we know that $a_0 \notin \text{st}^{-1}(E)$. By the definition of ν , we have $\nu(B) = {}^*\mu(B)$ for all $B \in {}^*\mathcal{B}[\widehat{X}]$ such that $B \subset \text{st}^{-1}(E)$. By Lemma 4.9, we have

$$(5.14) \quad \mu(E) = \overline{{}^*\mu}(\text{st}^{-1}(E))$$

$$(5.15) \quad = \sup\{{}^*\mu(B) : B \in {}^*\mathcal{B}[\widehat{X}] \wedge B \subset \text{st}^{-1}(E)\}$$

$$(5.16) \quad = \sup\{\nu(B) : B \in {}^*\mathcal{B}[\widehat{X}] \wedge B \subset \text{st}^{-1}(E)\} = \overline{\nu}(\text{st}^{-1}(E)).$$

Now suppose that $y \in \text{st}^{-1}(E)$. Then we have $a_0 \in E \subset \text{st}^{-1}(E)$. Thus, we have $\nu(B) = {}^*\mu(B)$ for all $B \in {}^*\mathcal{B}[\widehat{X}]$ such that $B \supset \text{st}^{-1}(E)$. By Lemma 4.9, we have

$$(5.17) \quad \mu(E) = \overline{{}^*\mu}(\text{st}^{-1}(E))$$

$$(5.18) \quad = \inf\{{}^*\mu(B) : B \in {}^*\mathcal{B}[\widehat{X}] \wedge B \supset \text{st}^{-1}(E)\}$$

$$(5.19) \quad = \inf\{\nu(B) : B \in {}^*\mathcal{B}[\widehat{X}] \wedge B \supset \text{st}^{-1}(E)\} = \underline{\nu}(\text{st}^{-1}(E)).$$

As E is arbitrary, we have the desired result. □

Pick a bounded uniformly continuous function f from X to \mathbb{R} . We can extend f to a bounded continuous function \hat{f} from \widehat{X} to \mathbb{R} . By assumption, we have

$$(5.20) \quad \int_X f \, dP_n = \int_{\widehat{X}} \hat{f} \, dP_n \rightarrow \int_{\widehat{X}} \hat{f} \, d\mu.$$

By Corollary 5.5, we have

$$(5.21) \quad \int_{\widehat{X}} \hat{f} \, d\mu = \int_{\widehat{X}} \hat{f} \, d\nu^p = \int_X f \, d\nu^p.$$

Hence we have shown that $\{P_n\}_{n \in \mathbb{N}}$ converges weakly to ν^p which is a charge on $(X, \mathcal{B}[X])$, completing the proof. □

REFERENCES

- [1] Aldaz J. M., *Representation of Measures via the Standard Part Map*, Ph.D. Thesis, University of Illinois at Urbana-Champaign, Champaign, 1991.
- [2] Aldaz J. M., *Representing abstract measures by Loeb measures: a generalization of the standard part map*, Proc. Amer. Math. Soc. **123** (1995), no. 9, 2799–2808.
- [3] Anderson R. M., *Star-finite representations of measure spaces*, Trans. Amer. Math. Soc. **271** (1982), no. 2, 667–687.
- [4] Anderson R. M., Rashid S., *A nonstandard characterization of weak convergence*, Proc. Amer. Math. Soc. **69** (1978), no. 2, 327–332.
- [5] Arkeryd L. O., Cutland N. J., Henson C. W., eds., *Nonstandard Analysis*, Theory and Applications, Proc. of the NATO Advanced Study Institute on Nonstandard Analysis and Its

- Applications, Edinburgh, 1996, NATO Advanced Science Institutes Series C: Mathematical and Physical Sciences, 493, Kluwer Academic Publishers Group, Dordrecht, 1997.
- [6] Bhaskara R. K. P. S., Bhaskara R. M., *Theory of Charges*, A study of finitely additive measures, Pure and Applied Mathematics, 109, Academic Press, Harcourt Brace Jovanovich Publishers, New York, 1983.
 - [7] Cutland N. J., Neves V., Oliveira F., Sousa-Pinto J., eds., *Developments in Nonstandard Mathematics*, Pitman Research Notes in Mathematics Series, 336, Papers from the Int. Col. (CIMNS94), Aveiro, 1994, Longman, Harlow, 1995.
 - [8] Duanmu H., Rosenthal J. S., Weiss W., *Ergodicity of Markov Processes via Non-standard Analysis*, available at <http://probability.ca/jeff/ftplib/duanmu1.pdf>.
 - [9] Duanmu H., Roy D. M., *On Extended Admissible Procedures and Their Nonstandard Bayes Risk*, available at arXiv:1612.09305v2 [math.ST].
 - [10] Kadane J. B., Schervish M. J., Seidenfeld T., *Statistical implications of finitely additive probability*, Stud. Bayesian Econometrics Statist., 6, North-Holland, Amsterdam, 1986, pages 59–76.
 - [11] Keisler H. J., *An infinitesimal approach to stochastic analysis*, Mem. Amer. Math. Soc. **48** (1984), no. 297, 184 pages.
 - [12] Lindström T., *Pushing Down Loeb-Measures*, Pure Mathematics: Preprint Series, Matematisk Institutt Universitetet i Oslo, Oslo, 1981.
 - [13] Loeb P. A., *Conversion from nonstandard to standard measure spaces and applications in probability theory*, Trans. Amer. Math. Soc. **211** (1975), 113–122.
 - [14] McShane E. J., *Extension of range of functions*, Bull. Amer. Math. Soc. **40** (1934), no. 12, 837–842.
 - [15] Render H., *Pushing down Loeb measures*, Math. Scand. **72** (1993), no. 1, 61–84.
 - [16] Rosenthal J. S., *A First Look at Rigorous Probability Theory*, World Scientific Publishing, Hackensack, 2006.
 - [17] Ross D., *Compact measures have Loeb preimages*, Proc. Amer. Math. Soc. **115** (1992), no. 2, 365–370.
 - [18] Ross D. A., *Pushing down infinite Loeb measures*, Math. Scand. **104** (2009), no. 1, 108–116.
 - [19] Yosida K., Hewitt E., *Finitely additive measures*, Trans. Amer. Math. Soc. **72** (1952), 46–66.

H. Duanmu, W. Weiss:

DEPARTMENT OF STATISTICS, UNIVERSITY OF TORONTO, SIDNEY SMITH HALL,
100 ST GEORGE ST, TORONTO, ON M5S 3G3, ONTARIO, CANADA

E-mail: duanmuhaosui@hotmail.com

E-mail: weiss@math.utoronto.ca

(Received March 19, 2018, revised August 24, 2018)