

John Alexander Cruz Morales; Alexander Torres-Gomez
On F-algebroids and Dubrovin's duality

Archivum Mathematicum, Vol. 55 (2019), No. 2, 109–122

Persistent URL: <http://dml.cz/dmlcz/147750>

Terms of use:

© Masaryk University, 2019

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON F -ALGEBROIDS AND DUBROVIN'S DUALITY

JOHN ALEXANDER CRUZ MORALES AND ALEXANDER TORRES-GOMEZ

ABSTRACT. In this note we introduce the concept of F -algebroid, and give its elementary properties and some examples. We provide a description of the almost duality for *Frobenius manifolds*, introduced by Dubrovin, in terms of a composition of two anchor maps of a unique cotangent F -algebroid.

1. INTRODUCTION

The notion of *Frobenius manifolds* was introduced by Dubrovin [5, 6, 7] in order to give a geometrical origin to the WDVV associativity equations discovered by E. Witten, R. Dijkgraaf, E. Verlinde, and H. Verlinde, in their study of topological quantum field theories from the physical point of view. For alternative approaches to introduce the notion of Frobenius manifolds see [1, 12].

Less than 20 years ago, C. Hertling and Y. Manin [11] defined what is known today as an F -manifold where they described a geometrical structure very close related to a *Frobenius manifold* but which does not need the introduction of a metric tensor and underlies any *Frobenius manifold*.

In this note, inspired by the notion of a *Lie algebroid* [2, 8, 9, 14, 20] and F -manifolds [10, 11], we define an F -algebroid which, roughly speaking, is a vector bundle whose fiber is an F -algebra and it has a map (called the *anchor*) mapping sections of the vector bundle to sections of the tangent bundle satisfying certain conditions, see section 4 below. In terms of this new geometrical structure, an F -manifold is just an F -algebroid structure over a tangent bundle whose *anchor* is the identity map.

Both *Frobenius manifolds* and F -manifolds, as can be seen below, are structures defined over the tangent bundle, then a natural question to ask is if these structures can be generalised to be defined over a general vector bundle. This kind of reasoning was what took us to the notion of an F -algebroid.

In the theory of *Frobenius manifolds* Dubrovin introduced the idea of almost duality [6], which also was extended to F -manifolds by Manin [18]. This duality relates two different multiplicative structures on the tangent bundle. Using the new structure

2010 *Mathematics Subject Classification*: primary 53C15; secondary 53D45.

Key words and phrases: F -manifolds, Frobenius manifolds, Lie algebroids.

Received January 11, 2019. Editor J. Slovák.

DOI: 10.5817/AM2019-2-109

proposed in this note we are able to “explain” this duality as the result of two different anchor maps over a unique cotangent F -algebroid, see below for details.

In sections 2 and 3 we remind the reader the notion of *Frobenius manifold* and F -manifold. Next, we define what we mean by an F -algebroid. Then, we describe the notion of almost dual *Frobenius manifolds* in terms of F -algebroids. Finally, we give some remarks where we briefly describe different questions about this new structure of F -algebroids which we are currently studying. In the appendix we provide the definition of *Lie algebroid* and give some examples.

Acknowledgement. We want to thank Yuri Manin for reading a draft of this note and his valuable comments. In particular, he drew our attention to the paper [4] by V. Dotsenko and pointed out the possibility of finding relations between the approach in that paper and ours. A. Torres-Gomez is partially supported by Universidad del Norte grant number 2018-17 “Agenda I+D+I”. J.A. Cruz Morales wants to thank Max Plank Institute for Mathematics for its hospitality and support during his visit in January 2018 where part of this work was done.

2. FROBENIUS MANIFOLD

2.1. Definition. Let M be a smooth manifold and TM its tangent bundle with the structures (\bullet, e, g, C) define on it where, for $X, Y, Z \in \Gamma(TM)$,

- (1) \bullet is a multiplicative structure on sections of the tangle bundle which is commutative and associative, that is, $\bullet: \Gamma(TM) \times \Gamma(TM) \rightarrow \Gamma(TM)$ and $X \bullet Y = Y \bullet X$, $(X \bullet Y) \bullet Z = X \bullet (Y \bullet Z)$;
- (2) $e \in \Gamma(TM)$ is a unit vector field with respect to the multiplicative structure \bullet , that is, $e \bullet X = X$;
- (3) g is a Riemannian (or semi-Riemannian metric) on M compatible with \bullet , that is, $g: \Gamma(TM) \times \Gamma(TM) \rightarrow C^\infty(M)$ and $g(X \bullet Y, Z) = g(X, Y \bullet Z)$;
- (4) C is a 3-tensor field defined by $C(X, Y, Z) := g(X \bullet Y, Z)$.

The manifold M is called a Frobenius manifold if the 6-tuple $(TM, \bullet, e, g, C, \mathcal{E})$, where $\mathcal{E} \in \Gamma(TM)$, satisfies:

- a) the metric g is flat and the unit vector field e is covariantly constant, that is, $\text{Riemann}(g) = 0$ and $\nabla e = 0$ where ∇ is the Levi-Civita connection of g ;
- b) the covariant derivative of C is a symmetric 4-tensor field, that is, $\nabla_W C(X, Y, Z)$ must be symmetric in $X, Y, Z, W \in \Gamma(TM)$.
- c) the Euler vector field \mathcal{E} must be linear, $\nabla \mathcal{E} = 0$, and for $X, Y \in \Gamma(TM)$ it obeys

$$\begin{aligned} \mathcal{L}_{\mathcal{E}}(X \bullet Y) - (\mathcal{L}_{\mathcal{E}}X) \bullet Y - X \bullet (\mathcal{L}_{\mathcal{E}}Y) &= X \bullet Y, \\ \mathcal{L}_{\mathcal{E}}g(X, Y) &= (2 - d)g(X, Y), \end{aligned}$$

where \mathcal{L} stands for the Lie derivative and d is called the charge.

Note that for any $p \in M$ the 4-tuple $(T_p M, \bullet_p, g_p, e_p)$ defines a Frobenius algebra.

2.2. WDVV Equations. On a Frobenius manifold M there exists a local system of flat coordinates (t^1, \dots, t^n) , because of the flatness condition on the metric g . In a coordinate basis $(\partial/\partial t^1, \dots, \partial/\partial t^n)$ the components of the metric g and the tensor C are given by

$$(2.1) \quad \eta_{ij} = g\left(\frac{\partial}{\partial t^i}, \frac{\partial}{\partial t^j}\right),$$

and

$$(2.2) \quad \frac{\partial}{\partial t^i} \bullet \frac{\partial}{\partial t^j} = \sum_k C_{ij}^k(t) \frac{\partial}{\partial t^k},$$

where $i, j, k = 1, \dots, n$. Note that the components η_{ij} and its inverse η^{ij} are constants. Usually this flat coordinates are chosen in such a way that the unit vector field is $e = \partial/\partial t^1$. Moreover, the Euler vector field can be written as

$$(2.3) \quad \mathcal{E} = \sum_i \left(\sum_j a_j^i t^j + b^i \right) \frac{\partial}{\partial t^i},$$

for some constants a_j^i, b^i obeying $a_1^i = \delta_1^i$ and $b^1 = 0$.

The structure functions $C_{ij}^k(t)$ can be written locally as third derivatives of a function $F: M \rightarrow \mathbb{R}$ called the potential, that is,

$$(2.4) \quad C_{ij}^k(t) = \sum_l \eta^{kl} \frac{\partial^3 F(t)}{\partial t^l \partial t^i \partial t^j},$$

obeying the following equations

$$(2.5) \quad \frac{\partial^3 F(t)}{\partial t^1 \partial t^i \partial t^j} = \eta_{ij},$$

$$(2.6) \quad \sum_{m,n} \frac{\partial^3 F(t)}{\partial t^i \partial t^j \partial t^m} \eta^{mn} \frac{\partial^3 F(t)}{\partial t^n \partial t^k \partial t^l} = \sum_{m,n} \frac{\partial^3 F(t)}{\partial t^l \partial t^j \partial t^m} \eta^{mn} \frac{\partial^3 F(t)}{\partial t^n \partial t^k \partial t^i},$$

$$(2.7) \quad \mathcal{E}F = (3-d)F + \frac{1}{2}A_{ij} t^i t^j + B_i t^i + c,$$

where A_{ij}, B_i, c are constants. The last two equations are known as WDVV associative equation and quasi-homogeneity condition, respectively.

2.3. Duality. The notion of duality between the Frobenius manifold $(M, \bullet, e, \mathcal{E}, \eta)$ and the almost-Frobenius manifold $(M, *, \mathcal{E}, g)$ is the statement that the multiplicative structure $*$ is given in terms of the multiplicative structures \bullet and the inverse of the Euler vector field \mathcal{E}^{-1} as

$$(2.8) \quad X * Y = X \bullet Y \bullet \mathcal{E}^{-1},$$

and the metric g is written in terms of the metric η and \mathcal{E}^{-1} as

$$(2.9) \quad g(X, Y) = \eta(\mathcal{E}^{-1} \bullet X, Y).$$

3. F-MANIFOLD

Let M be a manifold and denote by $[X, Y]$ the Lie bracket between $X, Y \in \Gamma(TM)$. Define $P_X(Y, Z)$, measuring the deviation of the structure $(\Gamma(TM), \bullet, [,])$ from being a Poisson algebra on $(\Gamma(TM), \bullet)$, as

$$(3.1) \quad P_X(Y, Z) := [X, Y \bullet Z] - [X, Y] \bullet Z - Y \bullet [X, Z].$$

An F-manifold is a triple (M, \bullet, e) , where

- (1) \bullet is a multiplicative structure on sections of the tangent bundle which is commutative and associative, that is, $\bullet: \Gamma(TM) \times \Gamma(TM) \rightarrow \Gamma(TM)$ and $X \bullet Y = Y \bullet X$, $(X \bullet Y) \bullet Z = X \bullet (Y \bullet Z)$;
- (2) $e \in \Gamma(TM)$ is a unit vector field with respect to the multiplicative structure \bullet , that is, $e \bullet X = X$;

satisfying the following identity

$$(3.2) \quad P_{X \bullet Y}(Z, W) = X \bullet P_Y(Z, W) + Y \bullet P_X(Z, W),$$

for any $X, Y, Z, W \in \Gamma(TM)$.

Theorem (Hertling and Manin, Weak Frobenius Manifolds, 1999 [11]).

(a) *Let (M, e, g, C) be a Frobenius manifold with multiplication \bullet . Then (M, \bullet, e) is an F-manifold.*

(b) *Let (M, \bullet) be an F-manifold, whose multiplication law is semisimple on an open dense subset. Assume that it admits an invariant flat metric η that defines the cubic tensor $C(X, Y, Z) = \eta(X \bullet Y, Z)$. Then (M, g, C) is a Frobenius manifold.*

4. F-ALGEBROIDS

4.1. Definition and examples of F-algebroids. Let TM be the tangent bundle of a manifold M ; $[,]$ a Lie bracket between section of TM ; \bullet denote a multiplicative commutative and associative structure on section of TM ; and $e \in \Gamma(TM)$ a unit vector field with respect to the multiplicative structure \bullet . An F-algebroid is a 5-tuple $(E, [,]_E, \diamond, \mathcal{U}, \rho)$ where

- (1) E is a vector bundle over a manifold M ;
- (2) $[,]_E : \Gamma(E) \times \Gamma(E) \rightarrow \Gamma(E)$ is a Lie bracket on sections of the vector bundle, that is, it is anti-symmetric and satisfies Jacobi identity $[\alpha, [\beta, \gamma]_E]_E + [\beta, [\gamma, \alpha]_E]_E + [\gamma, [\alpha, \beta]_E]_E = 0$;
- (3) $\diamond : \Gamma(E) \times \Gamma(E) \rightarrow \Gamma(E)$ is a commutative and associative multiplicative structure on sections of the vector bundle, that is, $\alpha \diamond \beta = \beta \diamond \alpha$ and $(\alpha \diamond \beta) \diamond \gamma = \alpha \diamond (\beta \diamond \gamma)$;
- (4) $\mathcal{U} \in \Gamma(E)$ is a unit with respect to the multiplicative structure \diamond , that is, $\mathcal{U} \diamond \alpha = \alpha$;
- (5) $\rho: \Gamma(E) \rightarrow \Gamma(TM)$, called the anchor map, relates sections of the vector bundle to sections of the tangent bundle;

satisfying the following conditions for $\alpha, \beta, \gamma, \tau \in \Gamma(E)$ and $f \in C^\infty(M)$:

a) defining $P_\alpha(\beta, \gamma)$ as

$$P_\alpha(\beta, \gamma) := [\alpha, \beta \diamond \gamma]_E - [\alpha, \beta]_E \diamond \gamma - \beta \diamond [\alpha, \gamma]_E,$$

it satisfies

$$P_{\alpha \diamond \beta}(\gamma, \tau) = \alpha \diamond P_\beta(\gamma, \tau) + \beta \diamond P_\alpha(\gamma, \tau);$$

b) homomorphism $\rho(\alpha \diamond \beta) = \rho(\alpha) \bullet \rho(\beta)$;

c) Leibniz rule, $[\alpha, f \beta]_E = (\rho(\alpha) f) \beta + f [\alpha, \beta]_E$;

d) Lie algebra homomorphism, $\rho([\alpha, \beta]_E) = [\rho(\alpha), \rho(\beta)]$.

That is we have

$$(E, \diamond, [\ , \]_E, \mathcal{U}) \xrightarrow{\rho} (TM, \bullet, [\ , \], e)$$

satisfying a), b), c) and d) above. Note that the triple (TM, \bullet, e) is an F-manifold.

Some examples of F-algebroids are the following:

- (1) **F-algebras and Frobenius algebras.** Every F-algebra and Frobenius algebra is an F-algebroid over the one point manifold. For the case of Frobenius algebra we introduce a symmetric bilinear form compatible with the product.
- (2) **Tangent F-algebroid.** In this case the vector bundle is the tangent bundle of M , the anchor map is the identity map, and we recover the definition of F-manifold.
- (3) In the following three examples, related to the theory of Frobenius manifolds, let the multiplicative structure \bullet be semisimple on an open dense subset and assume that M admits an invariant flat metric η that defines the cubic tensor $C(X, Y, Z) = \eta(X \bullet Y, Z)$.
 - (a) **Tangent Frobenius F-algebroid.** Take the vector bundle as the tangent bundle $E = TM$ and the anchor map as the identity $\rho = \text{Id}$. This would correspond to a Frobenius manifold.
 - (b) **Cotangent Frobenius F-algebroid.** Here the vector bundle is the cotangent bundle of the manifold M , $E = T^*M$. In flat coordinates $\{t^i\}$, the metric η is written locally as

$$(4.1) \quad \eta = \sum_{i,j} \eta_{ij}(t) dt^i \otimes dt^j,$$

and its inverse

$$(4.2) \quad \eta^{\text{inv}} = \sum_{i,j} \eta^{ij}(t) \frac{\partial}{\partial t^i} \otimes \frac{\partial}{\partial t^j}.$$

The anchor is given by

$$(4.3) \quad \rho_1 = \eta^{\text{inv}\#} = \sum_{i,j} \eta^{ij}(t) \frac{\partial}{\partial t^i} \otimes \frac{\partial}{\partial t^j},$$

and acts on a base of sections $dt^\alpha \in \Gamma(T^*M)$ as

$$(4.4) \quad \rho_1(dt^i) = \eta^{\text{inv}\#}(dt^i) = \sum_j \eta^{ij}(t) \frac{\partial}{\partial t^j}.$$

The symmetric 3-tensor C is written in components as $C = \sum_{ijk} C_{ijk}(t) dt^i \otimes dt^j \otimes dt^k$.

We have:

$$(T^*M, \diamond, \mathcal{U}) \xrightarrow{\rho_1 = \eta^{\text{inv}\#}} (TM, \bullet, e).$$

Define the product \diamond as

$$(4.5) \quad dt^i \diamond dt^j = \sum_k C_k^{ij}(t) dt^k,$$

where $C_k^{ij}(t) = \eta^{il} \eta^{jm} C_{lmk}$. From the homomorphism $\rho_1(\alpha \diamond \beta) = \rho_1(\alpha) \bullet \rho_1(\beta)$ we obtain

$$(4.6) \quad \begin{aligned} \rho_1(dt^i) \bullet \rho_1(dt^j) &= \rho_1(dt^i \diamond dt^j), \\ \sum_{k,l} \eta^{ik} \eta^{jl} \frac{\partial}{\partial t^k} \bullet \frac{\partial}{\partial t^l} &= \sum_k C_k^{ij} \rho_1(dt^k), \\ \sum_{k,l} \eta^{ik} \eta^{jl} \frac{\partial}{\partial t^k} \bullet \frac{\partial}{\partial t^l} &= \sum_{k,m} C_k^{ij} \eta^{km} \frac{\partial}{\partial t^m}, \end{aligned}$$

and thus the product \bullet is given by

$$(4.7) \quad \frac{\partial}{\partial t^k} \bullet \frac{\partial}{\partial t^l} = \sum_m C_{kl}^m(t) \frac{\partial}{\partial t^m}$$

The F-manifold (TM, \bullet, e) with $C(X, Y, Z) = \eta(X \bullet Y, Z)$ is a Frobenius manifold.

- (c) **Cotangent almost Frobenius F-algebroid.** Here we have a similar situation as in the case before where the vector bundle $E = T^*M$ is the cotangent bundle of a manifold M . Moreover, we have an Euler vector field \mathcal{E} . This time the anchor map is given by

$$(4.8) \quad \rho_2 = \sum_{i,j,k} \mathcal{E}^{-1k}(t) C_k^{ij}(t) \frac{\partial}{\partial t^i} \otimes \frac{\partial}{\partial t^j},$$

and acts on a base of sections dt^i as

$$(4.9) \quad \rho_2(dt^i) = \left(\sum_{j,k,l} \mathcal{E}^{-1k} C_k^{jl} \frac{\partial}{\partial t^j} \otimes \frac{\partial}{\partial t^l} \right) (dt^i) = \sum_{j,k} \mathcal{E}^{-1k} C_k^{ij} \frac{\partial}{\partial t^j}.$$

Note that the objet $\sum_k \mathcal{E}^k(t) C_k^{ij}(t) = g^{ij}(t)$ can be considered as a symmetric bilinear form on T^*M . We have:

$$(T^*M, \diamond, \mathcal{U}) \xrightarrow{\rho_2 = \sum_{i,j,k} \mathcal{E}^{-1k} C_k^{ij} \frac{\partial}{\partial t^i} \otimes \frac{\partial}{\partial t^j}} (T(M \setminus \Sigma), *, \mathcal{E})$$

where $\Sigma \subset M$ is the discriminant¹.

Again, define the product \diamond in sections of T^*M as

$$(4.10) \quad dt^i \diamond dt^j = \sum_k C_k^{ij}(t) dt^k .$$

Then, from the homomorphism $\rho_2(\alpha \diamond \beta) = \rho_2(\alpha) * \rho_2(\beta)$ we obtain that the product $*$ on section of TM now is given by

$$(4.11) \quad \begin{aligned} \rho_2(dt^i) * \rho_2(dt^j) &= \rho_2(dt^i \diamond dt^j) , \\ \rho_2(dt^i) * \rho_2(dt^j) &= \sum_k C_k^{ij} \rho_2(dt^k) , \\ \rho_2(dt^i) * \rho_2(dt^j) &= \sum_{k,l,m} C_k^{ij} \mathcal{E}^{-1m} C_m^{kl} \frac{\partial}{\partial t^l} . \end{aligned}$$

5. DUALITY AND F-ALGEBROIDS

The notion of duality between the Frobenius manifold $(M, \bullet, e, \mathcal{E}, \eta)$ and the almost-Frobenius manifold $(M, *, \mathcal{E}, g)$ can be understood as the composition between two anchor maps, as the following theorem shows.

Theorem 1. *Let*

$$(T^*M, \diamond, \mathcal{U}) \xrightarrow{\rho_1 = \eta^{\text{inv}} \#} (TM, \bullet, e, g, C, \mathcal{E})$$

be a cotangent Frobenius F-algebroid and

$$(T^*M, \diamond, \mathcal{U}) \xrightarrow{\rho_2 = \sum_{i,j,k} \mathcal{E}^{-1k} C_k^{ij} \frac{\partial}{\partial t^i} \otimes \frac{\partial}{\partial t^j}} (T(M \setminus \Sigma), *, \mathcal{E}, g)$$

a cotangent almost Frobenius F-algebroid. Then, there exists a map called \mathcal{D} given by

$$(5.1) \quad \mathcal{D} = \rho_2 \circ \rho_1^{-1}$$

relating the two multiplicative structures \bullet and $$ by*

$$(5.2) \quad X * Y = X \bullet Y \bullet \mathcal{E}^{-1} ,$$

and the two metrics η and g by

$$(5.3) \quad g(X, Y) = \eta(X \bullet \mathcal{E}^{-1}, Y) .$$

¹See [6] for details about the discriminant.

In other words, we have the following commuting diagram

$$\begin{array}{ccc} (T^*M, \diamond, \mathcal{U}) & \xrightarrow{\rho_1} & (TM, \bullet, e, g, C, \mathcal{E}) \\ & \searrow \rho_2 & \downarrow D = \rho_2 \circ \rho_1^{-1} \\ & & (T(M \setminus \Sigma), *, \mathcal{E}, g) \end{array}$$

The map $\mathcal{D} = \rho_2 \circ \rho_1^{-1}$ is what is called a duality in the theory of Frobenius manifolds.

Proof. Let $\{t^i\}$ be flat coordinates and $\{dt^i\}$ and $\{\frac{\partial}{\partial t^i}\}$ a base of section of T^*M and TM , respectively. In the coordinates $\{t^i\}$, we can write, locally,

$$(5.4) \quad \rho_1 = \sum_{i,j} \eta^{ij}(t) \frac{\partial}{\partial t^i} \otimes \frac{\partial}{\partial t^j},$$

$$(5.5) \quad \rho_1^{-1} = \sum_{i,j} \eta_{ij}(t) dt^i \otimes dt^j,$$

and

$$(5.6) \quad \rho_2 = \sum_{i,j,k} \mathcal{E}^{-1k}(t) C_k^{ij}(t) \frac{\partial}{\partial t^i} \otimes \frac{\partial}{\partial t^j}.$$

Let $X, Y \in TM$, then locally we have

$$(5.7) \quad X = \sum_i X^i(t) \frac{\partial}{\partial t^i}, \quad Y = \sum_j Y^j(t) \frac{\partial}{\partial t^j}.$$

Let us divide the proof in two parts corresponding to the map between the multiplicative structures and the metrics.

Multiplicative structures. Locally, $(X \bullet Y) \in TM$ is written as

$$(5.8) \quad X \bullet Y = \sum_{i,j} X^i Y^j \frac{\partial}{\partial t^i} \bullet \frac{\partial}{\partial t^j} = \sum_{i,j,k} C_{ij}^k X^i Y^j \frac{\partial}{\partial t^k}.$$

Acting with ρ_1^{-1} on $(X \bullet Y) \in TM$ we obtain

$$(5.9) \quad \rho_1^{-1}(X \bullet Y) = \sum_{i,j,k} C_{ij}^k X^i Y^j \rho_1^{-1} \left(\frac{\partial}{\partial t^k} \right) = \sum_{i,j,k} C_{ijk} X^i Y^j dt^k.$$

Now, acting with ρ_2 on $\rho_1^{-1}(X \bullet Y) \in T^*M$ we find

$$(5.10) \quad \begin{aligned} \rho_2(\rho_1^{-1}(X \bullet Y)) &= \sum_{i,j,k} C_{ijk} X^i Y^j \rho_2(dt^k) = \sum_{i,j,k,l,m} C_{ijk} C_l^{km} X^i Y^j \mathcal{E}^{-1l} \frac{\partial}{\partial t^m}, \\ &= X \bullet Y \bullet \mathcal{E}^{-1}. \end{aligned}$$

Thus, acting with the map $\mathcal{D} = \rho_2 \circ \rho_1^{-1}$ on $(X \bullet Y) \in TM$ we find

$$(5.11) \quad \mathcal{D}(X \bullet Y) = \rho_2 \circ \rho_1^{-1}(X \bullet Y) = X \bullet Y \bullet \mathcal{E}^{-1} = X * Y,$$

proving the relation between the multiplicative structures \bullet and \ast .
Metrics. Acting with ρ_1^{-1} on $X \in TM$ we get

$$(5.12) \quad \rho_1^{-1}(X) = \sum_i X^i \rho_1^{-1}\left(\frac{\partial}{\partial t^i}\right) = \sum_{i,j} \eta_{ij} X^i dt^j.$$

Now, acting with ρ_2 on $\rho_1^{-1}(X)$ we find

$$(5.13) \quad \rho_2 \circ \rho_1^{-1}(X) = \sum_{i,j} \eta_{ij} X^i \rho_2(dt^j) = \sum_{i,j,k} C_{ij}^k X^i \mathcal{E}^{-1j} \frac{\partial}{\partial t^k} = X \bullet \mathcal{E}^{-1}.$$

Then, the two metrics η and g are related as

$$(5.14) \quad g(X, Y) = \eta(X \bullet \mathcal{E}^{-1}, Y).$$

□

Remark 1. This theorem also holds when we consider the tangent F -algebroid where the underlying F -manifold algebra is semisimple. In this case the proof is similar to that of Theorem 1.

6. FINAL REMARKS

In this final section we will consider some questions and perspectives for future work.

So far, we have been discussing how different multiplicative structures can be related and the role played by the anchor maps in order to establish a duality between them. However, we can consider more than two products and it makes sense to ask how they are related. This question is motivated by studying how the anchor maps behave under composition (we only consider the semisimple case).

Let us consider the following situation. Let (M, \star_0) and (M, \star_1) be two dual F -manifold algebras such that $X \star_1 Y = X \star_0 Y \star_0 \mathcal{E}_0^{-1}$. Now consider an F -manifold algebra (M, \star_2) dual to (M, \star_1) , meaning $X \star_2 Y = X \star_1 Y \star_1 \mathcal{E}_1^{-1}$. A natural question is: Is there a relation between (M, \star_0) and (M, \star_2) ? A simple computation leads to

$$X \star_2 Y = X \star_0 Y \star_0 (\mathcal{E}_0^{-1})^2 \star_0 \mathcal{E}_1^{-1}.$$

Note that the invertible element \mathcal{E}_1^{-1} for the product \star_1 is not necessarily an invertible element for \star_0 , so \star_0 and \star_2 are not dual in the sense we have been discussing above. This motivates the following definition.

Definition 1. Let \mathcal{E}_0 be an eventual identity² for \star_0 and \mathcal{E}_1 be an eventual identity for a product \star_1 dual to \star_0 . An element \mathcal{I} is a pseudo-eventual identity for the product \star_0 if $\mathcal{I} = \mathcal{E}_0 \star_0 \mathcal{E}_1^{-1}$ and the product $X \star_2 Y = X \star_0 Y \star_0 \mathcal{I}$ defines an F -manifold algebra structure.

In this case we will say that (M, \star_0) and (M, \star_2) are pseudo-dual structures.

²See [3] for the definition of eventual identities.

It is easy to see that the notion of pseudo-duality can be defined as composition of anchor maps for the corresponding F -algebroids in the same way that the notion of duality. That composition defines a map \mathcal{P} called pseudo-duality map.

Proposition 1. *The composition of the maps \mathcal{D}_0 and \mathcal{D}_1 in Theorem 1 is the map \mathcal{P}_{01} relating the pseudo-dual products \star_0 and \star_2 .*

Proof. Let us consider the following diagram

$$\begin{array}{ccc}
 (T^*M_0, \diamond, \mathcal{U}) & \xrightarrow{\rho_0} & (TM_0, \star_0, e_0, \mathcal{E}_0) \\
 & \searrow \rho_1 & \downarrow D_0 = \rho_1 \circ \rho_0^{-1} \\
 & & (TM_1, \star_1, e_1, \mathcal{E}_1) \\
 & \searrow \rho_2 & \downarrow D_1 = \rho_2 \circ \rho_1^{-1} \\
 & & (TM_2, \star_2, e_2, \mathcal{E}_2)
 \end{array}$$

Observe that $\mathcal{P}_{01} = \rho_2 \circ \rho_0^{-1}$. This follows from an argument similar to that of Theorem 1. Since $\rho_2 = \mathcal{D}_1 \circ \rho_1$ we have that $\mathcal{P}_{01} = \mathcal{D}_1 \circ \mathcal{D}_0$ □

In general, we can have an enumerable family of products and then we can ask how the products in this family relate to each other. We have the following proposition.


Proposition 2. *Consider the family of products $\{\star_i\}_{i \in I}$ where I is an ordered set such that \star_i is dual to \star_{i+1} . Let \mathcal{E}_i^{-1} be an eventual identity for \star_i and \mathcal{I} be a power of a pseudo-eventual identity for \star_i . If $j \geq i + 3$ then the product $X \star_j Y = X \star_i Y \star_i \mathcal{I} \star_i \mathcal{E}_i^{-1}$ defines an F -manifold algebra structure.*

Proof. Without loss of generality we will consider four products $\star_0, \star_1, \star_2$ and \star_3 , such that \star_i is dual to \star_{i+1} for $i = 0, 1, 2, 3$. Since \star_2 and \star_3 are dual then $X \star_3 Y = X \star_2 Y \star_2 \mathcal{E}_2^{-1}$. For the previous proposition \star_0 and \star_2 are pseudo-dual, so

$$X \star_3 Y = (X \star_0 Y \star_0 \mathcal{I}) \star_2 \mathcal{E}_2^{-1} = (X \star_0 Y \star_0 \mathcal{I}) \star_0 \mathcal{E}_2^{-1} \star_0 \mathcal{I} = X \star_0 Y \star_0 \mathcal{I}^2 \star_0 \mathcal{E}_2^{-1},$$

where \mathcal{I} is a pseudo-eventual identity for \star_0 . The same argument works for more than four products. □

Now consider the following diagram



From the diagram it is easy to see that the relation between the product \star_i and \star_j can be understood as composition of the associated anchor maps. This relation is weaker than the duality and the pseudo-duality defined above, so we will call it weak duality. An analogue of proposition 1 is true for weak dualities. This means, that weak duality is a composition of dualities (or pseudo-dualities).

The notion of weak duality and the family of products behind it lead to the following questions: Is it possible to consider a chain of F -algebroids connected by duality, pseudo-duality and weak duality maps? Which properties this chain will have?

In mathematics the notion of horizontal categorification, also known as “oidification”, is a procedure giving a structure X the structure X -oid. For example, starting with a Hopf algebra to construct a Hopf algebroid. In our work, we have been constructing a F -algebroid by mimicking the construction of Lie algebroids, so a natural question is whether the object we have defined corresponds to the horizontal categorification of F -manifold algebras. Let us point out that, at least for us, it is not clear how Lie algebroids are horizontal categorification of Lie algebras, so in that sense our question is also valid in the Lie theoretic setting. We think that this question could be related to some discussions in [19].

Another further direction to work in the formalism of F -algebroids is the study of the analogies between F -geometry and Poisson geometry. We do not have a precise picture here but we believe that this question is related to a bigger question asking for studying the geometry of F -algebroids. This question comes with the necessity of constructing more examples of F -algebroids with interesting mathematical structures. Note that in the theory of Lie algebroids there is a 1-1 correspondence between Lie algebroid structures on a vector bundle and homogeneous Poisson structures on the total space of the dual bundle. We believe that we should have an analogue result in the context of F -algebroids.

Following ideas in [16], we consider that F -algebroids might be an adequate context to study deformations of Frobenius manifolds and F -manifolds in analogy to the theory

of deformations of complex manifolds [13, 15]. On the other hand, since the duality of Dubrovin can be seen, in some cases, as instance of mirror symmetry phenomenon, we also think that F -algebroids can play a role in mirror symmetry.

Finally, we think that giving an algebraic definition of our F -algebroid (similar to the Lie-Rinehart pair description of a Lie algebroid) we could make contact with the construction of the operad $FMan$ in [4], this is work in progress.

A. LIE ALGEBROID

Some reviews on Lie algebroids can be read in [2, 8, 9, 14, 20].

A.1. Definition and some examples. A Lie algebroid is a triple $(E, [\cdot, \cdot]_E, \rho)$ where

- (1) E is a vector bundle over a manifold M ,
- (2) $[\cdot, \cdot]_E : \Gamma(E) \times \Gamma(E) \rightarrow \Gamma(E)$ is a Lie bracket on sections of the vector bundle, that is, it is anti-symmetric and satisfies Jacobi identity $[\alpha, [\beta, \gamma]_E]_E + [\beta, [\gamma, \alpha]_E]_E + [\gamma, [\alpha, \beta]_E]_E = 0$, and
- (3) $\rho : \Gamma(E) \rightarrow \Gamma(TM)$, called the anchor map, maps sections of the vector bundle to sections of the tangent bundle.

The Lie bracket $[\cdot, \cdot]_E$ and the anchor ρ satisfy the following conditions for $\alpha, \beta, \gamma \in \Gamma(E)$ and $f \in C^\infty(M)$:

- a) Leibniz rule, $[\alpha, f\beta]_E = (\rho(\alpha)f)\beta + f[\alpha, \beta]_E$;
- b) Lie algebra homomorphism, $\rho([\alpha, \beta]_E) = [\rho(\alpha), \rho(\beta)]$;

A few examples of Lie algebroids are the following

- **Lie algebra.** Every Lie algebra is a Lie algebroid over the one point manifold.
- **Tangent Lie algebroid.** In this case the vector bundle is the tangent bundle of M , the anchor map is the identity map and the bracket is the usual Lie bracket of vector fields.
- **Cotangent Lie algebroid.** Here the vector bundle is the cotangent bundle of a Poisson manifold M with Poisson bivector Π . We define the anchor map as follows:

- Let $\alpha \in \Gamma(T^*M)$ be a section of the cotangent bundle, then the anchor map is $\rho(\alpha) = \Pi^\#(\alpha)$.

In a coordinate basis $\{X^i\}$, with $\Pi = \sum_{i,j} \frac{1}{2} \Pi^{ij}(X) \frac{\partial}{\partial X^i} \wedge \frac{\partial}{\partial X^j}$ and $\alpha = \sum_i \alpha_i dX^i$, we have $\Pi^\#(\alpha) = \sum_{i,j} \alpha_i \Pi^{ij} \frac{\partial}{\partial X^j}$; and

- Let α and $\beta \in \Gamma(T^*M)$, then the bracket is

$$[\alpha, \beta] = \mathcal{L}_{\Pi^\#(\alpha)}\beta - \mathcal{L}_{\Pi^\#(\beta)}\alpha + d\Pi(\alpha, \beta),$$

where $\mathcal{L}_{\Pi^\#(\alpha)}$ denotes the Lie derivative of the 1-form β in the direction of the vector field $\Pi^\#(\alpha)$ and $\Pi(\alpha, \beta)$ the natural pairing between the vector $\Pi^\#(\alpha)$ and the 1-form β . Cotangent Lie algebroids can be understood from the perspective of Poisson geometry.

REFERENCES

- [1] Audin, M., *Symplectic geometry in Frobenius manifolds and quantum cohomology*, J. Geom. Phys. **25** (1–2) (1998), 183–204.
- [2] Crainic, M., Fernandes, R.L., *Lectures integrability Lie brackets*, Geom. Topol. Monogr. **17** (2011), 1–107.
- [3] David, L., Strachan, I.A.B., *Dubrovin's duality for F-manifolds with eventual identities*, Adv. Math. **226** (4) (2011), 4031–4060.
- [4] Dotsenko, V., *Algebraic structures of F-manifolds via pre-Lie algebras*, Ann. Mat. Pura Appl. (4) **198** (2019), 517–527.
- [5] Dubrovin, B., *Geometry of 2D topological field theories*, Lecture Notes in Math., vol. 1620, Springer, 1996.
- [6] Dubrovin, B., *On almost duality for Frobenius manifolds*, Amer. Math. Soc. Transl. **212** (2004), 75–132.
- [7] Dubrovin, B., *WDVV Equations and Frobenius Manifolds*, Encyclopedia of Mathematical Physics, vol. 1, Elsevier, 2006, pp. 438–447.
- [8] Dufour, J.P., Zung, N.T., *Poisson Structures and Their Normal Forms*, Birkhauser, 2000.
- [9] Fernandes, R.L., *Lie algebroids, holonomy and characteristic classes*, Adv. Math. **170** (1) (2002), 119–179.
- [10] Hertling, C., *Frobenius manifolds and moduli spaces for singularities*, Cambridge University Press, 2004.
- [11] Hertling, C., Manin, Y., *Weak Frobenius manifolds*, Internat. Math. Res. Notices **6** (1999), 277–286.
- [12] Hitchin, N., *Frobenius manifolds*, Gauge Theory and Symplectic Geometry, Springer, 1997.
- [13] Kodaira, K., *Complex Manifolds and Deformation of Complex Structures*, Springer, 2005.
- [14] Mackenzie, K.C.H., *General Theory of Lie Groupoids and Lie Algebroids*, Cambridge University Press, 2005.
- [15] Manetti, M., *Lectures on deformation of complex manifolds*, Rendiconti di Matematica **24** (2004), 1–183.
- [16] Manin, Y., *Mirrors, functoriality, and derived geometry*, [arXiv:1708.02849](https://arxiv.org/abs/1708.02849).
- [17] Manin, Y., *Frobenius Manifolds, Quantum Cohomology, and Moduli Spaces*, Amer. Math. Soc. Colloq. Publ. **47** (1999), xiv+303 pp.
- [18] Manin, Y., *F-manifolds with flat structure and Dubrovin's duality*, Adv. Math. **198** (1) (2005), 5–26.
- [19] Manin, Y., *Grothendieck-Verdier duality patterns in quantum algebra*, Izv. Ross. Akad. Nauk Ser. Mat. **81** (4) (2017), 158–166.
- [20] Weinstein, A., *Linearization problems Lie algebroids and Lie groupoids*, Lett. Math. Phys. **52** (2000), 93–102.

JOHN ALEXANDER CRUZ MORALES,
 DEPARTAMENTO DE MATEMÁTICAS,
 UNIVERSIDAD NACIONAL DE COLOMBIA,
 BOGOTÁ, COLOMBIA,
E-mail: jacruzmo@unal.edu.co

ALEXANDER TORRES-GOMEZ,
DEPARTAMENTO DE MATEMÁTICAS Y ESTADÍSTICA,
UNIVERSIDAD DEL NORTE,
BARRANQUILLA, COLOMBIA,
E-mail: alexander.torres.gomez@gmail.com