

Higher Structures

David Jaz Myers

Good Fibrations through the Modal Prism


Higher Structures, Vol. 6 (2022), No. 1, 212–255

Persistent URL: <http://dml.cz/dmlcz/153448>

Terms of use:

© Institute of Mathematics, Czech Academy of Sciences, 2022

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library*
<http://dml.cz>

Good Fibrations through the Modal Prism

David Jaz Myers^a

^aDepartment of Mathematics, 404 Krieger Hall 3400 N. Charles Street, Baltimore, MD 21218

Abstract

Homotopy type theory is a formal language for doing abstract homotopy theory — the study of identifications. But in unmodified homotopy type theory, there is no way to say that these identifications come from identifying the path-connected points of a space. In other words, we can do abstract homotopy theory, but not algebraic topology. Shulman’s *Real Cohesive HoTT* remedies this issue by introducing a system of modalities that relate the spatial structure of types to their homotopical structure. In this paper, we develop a theory of *modal fibrations* for a general modality, and apply it in particular to the shape modality of real cohesion. We then give examples of modal fibrations in Real Cohesive HoTT, and develop the theory of covering spaces.

Communicated by: Richard Garner.

Received: 12th February, 2020. Accepted: 12th April, 2022.

MSC: 55U35.

Keywords: Homotopy Type Theory, Modality, Fibration, Axiomatic Cohesion.

1. Introduction

While homotopy theory — the study of identifications — has been well developed in homotopy type theory, algebraic topology — the study of the connectivity of space — has been somewhat lacking. This is because Book HoTT (the homotopy type theory of the HoTT Book [13]) has no way of saying that a type is *the homotopy type* of another type. While we can define both the homotopy circle S^1 as a higher inductive type and the topological circle

$$\mathbb{S}^1 := \{(x, y) : \mathbb{R}^2 \mid x^2 + y^2 = 1\},$$

in Book HoTT alone we do not have the tools to say that S^1 is the homotopy type of \mathbb{S}^1 .

In his Real Cohesive Homotopy Type Theory [12], Shulman solves this issue by adding a system of modalities which includes the *shape* modality \int that takes a type X to its homotopy

Email address: davidjaz@jhu.edu (David Jaz Myers)

© David Jaz Myers, 2022, under a Creative Commons Attribution 4.0 International License.

DOI: 10.21136/HS.2022.04

type $\int X$.¹ In Real Cohesive HoTT, every type has a spatial structure and every map is continuous with respect to this spatial structure. This spatial structure is distinct from the *homotopical* structure of identifications that every type has in homotopy type theory. But these two structures are brought together by the \int modality, which allows us to identify points by giving spatial paths between them. Formally, the \int modality is given by localizing at the type of Dedekind real numbers \mathbb{R} — in other words, by identifying points which are connected by paths $\gamma : \mathbb{R} \rightarrow X$.²

As with any modality, there is a *modal unit* $(-)^{\int} : X \rightarrow \int X$, a quotient map of sorts, which is the universal map from X to a *discrete* type — one with only homotopical and no spatial structure.³ For any map $f : X \rightarrow Y$, we have a naturality square which induces a map from the fiber of f over $y : Y$ to its *homotopy fiber*, the fiber of $\int f$:

$$\begin{array}{ccc}
 \text{fib}_f(y) & \overset{\delta}{\dashrightarrow} & \text{fib}_{\int f}(y^{\int}) \\
 \downarrow & & \downarrow \\
 X & \xrightarrow{(-)^{\int}} & \int X \\
 f \downarrow & & \downarrow \int f \\
 Y & \xrightarrow{(-)^{\int}} & \int Y
 \end{array}$$

The fibers of maps between discrete types are themselves discrete, so the map $\delta : \text{fib}_f(y) \rightarrow \text{fib}_{\int f}(y^{\int})$ factors uniquely through $(-)^{\int} : \text{fib}_f(y) \rightarrow \int \text{fib}_f(y)$ by the universal property of the unit. This gives us a useful diagram (Figure 1) which I like to call *the modal prism*.

$$\begin{array}{ccc}
 \text{fib}_f(y) & \xrightarrow{\delta} & \text{fib}_{\int f}(y^{\int}) \\
 \searrow^{(-)^{\int}} & & \nearrow^{\gamma} \\
 & \int \text{fib}_f(y) &
 \end{array}$$

Figure 1: The Modal Prism.

Looking through the modal prism, we see a rainbow of different possibilities for a function $f : X \rightarrow Y$.

Definition 1.1. Let $f : X \rightarrow Y$ and consider the modal prism as in Figure 1. Then f is

- \int -*modal* if its fibers are discrete, that is, if $(-)^{\int}$ is an equivalence for all $y : Y$,
- \int -*connected* if its fibers are homotopically contractible, that is, if $\int \text{fib}_f(y)$ is contractible for all $y : Y$,
- \int -*étale* if its fibers are its homotopy fibers, that is, if δ is an equivalence for all $y : Y$.
- a \int -*equivalence* if its homotopy fibers are contractible, that is, if $\text{fib}_{\int f}(y^{\int})$ is contractible for all $y : Y$,

¹The symbol “ \int ” is an *esh*, the IPA symbol for the voiceless palato-alveolar fricative phoneme /sh/ that begins the word “shape”. It is not an integral sign.

²In this paper, we reserve the term *path* (in X) for function $\gamma : \mathbb{R} \rightarrow X$, while we use the term *identification* for points of the type $x = y$ (for $x, y : X$). This conflicts with the terminology of the HoTT Book, in which “path” is used for what we call identifications. But, in our setting, the shape modality \int takes a path $\gamma : \mathbb{R} \rightarrow X$ and gives an identification $\gamma(0)^{\int} = \gamma(1)^{\int}$ in the homotopy type $\int X$. So, when one is working with homotopy types $\int X$, the difference between our terminology and the terminology of the HoTT Book is blurred.

³That is, every path is constant in a discrete type, but there may still be non-trivial identifications between its points.

- a \mathfrak{f} -fibration if the homotopy type of its fibers are its homotopy fibers, that is, if γ is an equivalence for all $y : Y$.

For the shape modality, a map is modal when it has discrete fibers, and is a modal equivalence, or (weak) homotopy equivalence, when it induces an equivalence on homotopy types. It is modally connected when it has the stronger property that its fibers are homotopically contractible; for comparison, consider the inclusion $x : \mathbb{R} \rightarrow \mathbb{R}^2$ of the x -axis, which is clearly a homotopy equivalence but is not \mathfrak{f} -connected since some of its fibers are empty. Finally, a \mathfrak{f} -étale map is a weak relative of a covering map; it has a *unique* lifting against any homotopy equivalence.

The notions of modal maps, connected maps, and modal equivalences appear in the HoTT Book ([13]). For the n -truncation modality, these are n -truncated and n -connected maps respectively, with modal equivalences not given a specific name. The notion of modal étale map is due to Wellen as a “formally étale map” in [15], building on work of Schreiber in the setting of higher topos theory [10]. In the case of \mathfrak{f} , it appears as a “modal covering” in [14].

The notion of modality has also made its way into the ∞ -categorical literature through the work of Anel, Biederman, Finster, and Joyal (see [2] and [1]). In these papers, they define a modality as a *stable orthogonal factorization system* (one of the equivalent ways of defining a modality in HoTT), and translate a homotopy type theoretic generalized Blakers-Massey Theorem into the language of ∞ -categories and apply it to the Goodwillie calculus of functors. As Shulman has proven that every ∞ -topos models HoTT ([11]), the results in this paper concerning modal fibrations (in Section 3) apply in any ∞ -topos as well.

The notion of modal fibration is, as far as I know, novel to this paper. It gives a good notion of fibration in real cohesion which works not just for set level spaces (e.g. manifolds) but also spaces with both topological and homotopical content (e.g. orbifolds and Lie groupoids). A map is a \mathfrak{f} -fibration when the homotopy type of its fibers are the fibers of its action on homotopy types; this gives us the long fiber sequence on homotopy groups we expect from a fibration in real cohesion. This definition closely resembles the classical notion of *quasi-fibration* due to Dold and Thom [6], though it is much better behaved (see Remark 3.1).

In Section 2, we will refresh ourselves on modalities and look through the modal prism to see the different kinds of functions associated with a modality. Then, in Section 3 we will develop the basic theory of \diamond -fibrations for an arbitrary modality \diamond , and justify the name. In summary, the \diamond -fibrations are closed under composition and pullback and may be characterized in any one of the following ways.

Theorem 1.2. *For a map $f : X \rightarrow Y$, the following are equivalent:*

1. f is a \diamond -fibration.
2. \diamond preserves all fibers of f .
3. \diamond preserves all pullbacks along f .
4. The \diamond -connected/ \diamond -modal and \diamond -equivalence/ \diamond -étale factorizations of f agree.
5. The \diamond -modal factor of f is \diamond -étale.
6. The \diamond -equivalence factor of f is \diamond -connected.
7. The \diamond -naturality square of f is \diamond -cartesian.
8. The connecting map $\mathbf{tot}(\gamma)$ between the two factorizations of f is a \diamond -fibration.
9. f has \diamond -locally constant \diamond -fibers in the sense that $\diamond \mathbf{fib}_f : Y \rightarrow \mathbf{Type}_\diamond$ factors through $\diamond Y$.
10. (If \diamond -units are surjective:) For every $x : X$, the induced map $\mathbf{fib}_{(-)\diamond}(x^\diamond) \rightarrow \mathbf{fib}_{(-)\diamond}((fx)^\diamond)$ is \diamond -connected.

In particular, we will prove in Theorem 3.14 that a map $f : X \rightarrow Y$ is an \diamond -fibration if and only if the type family $\diamond \mathbf{fib}_f : Y \rightarrow \mathbf{Type}$ factors through the modal unit $(-)^{\diamond} : Y \rightarrow \diamond Y$. For the modality \mathfrak{f} , this means that a map is a \mathfrak{f} -fibration if and only if the homotopy type of its fiber over $y : Y$ is locally constant in y ; that is, a map is a \mathfrak{f} -fibration if and only if its fibers form a *local system* on its codomain.

We will also characterize the $\|-\|_n$ -fibrations as those maps which are surjective on π_{n+1} in Corollary 3.19.

In Section 4, we give a brief review of Shulman’s Real Cohesive HoTT. We then prove in Section 5 that the classifying types of bundles of discrete structures are themselves discrete (see Theorem 5.9 for the precise statement). As a corollary, we find in Theorem 6.1 that maps whose fibers have a merely constant homotopy type are \mathfrak{f} -fibrations. Morally, this result says that if all the fibers of a map have the same homotopy type so that one can comfortably write

$$F \rightarrow E \xrightarrow{p} B$$

with F well defined up to homotopy, then p is a \mathfrak{f} -fibration.

In the remaining sections, we will show how this theory can be applied to synthetic algebraic topology. Because the homotopy type of the fibers of a \mathfrak{f} -fibration are its homotopy fibers, whenever

$$F \rightarrow E \xrightarrow{p} B$$

is a fiber sequence with p a \mathfrak{f} -fibration, $\mathfrak{f}F \rightarrow \mathfrak{f}E \xrightarrow{\mathfrak{f}p} \mathfrak{f}B$ is also a fiber sequence. Using the fact that the fibers of the map $(\cos, \sin) : \mathbb{R} \rightarrow \mathbb{S}^1$ are merely equivalent to \mathbb{Z} , Theorem 6.1 implies that this map is a \mathfrak{f} -fibration, and that therefore,

$$\mathbb{Z} \rightarrow \mathfrak{f}\mathbb{R} \rightarrow \mathfrak{f}\mathbb{S}^1$$

is a fiber sequence. Since $\mathfrak{f}\mathbb{R} \simeq *$ is contractible, this calculates the loop space of the topological circle \mathbb{S}^1 without passing through the higher inductive circle S^1 . We consider this and other examples of \mathfrak{f} -fibrations, including:

- The map $(\cos, \sin) : \mathbb{R} \rightarrow \mathbb{S}^1$ (in Section 6.1).
- The homogeneous coordinates $\mathbb{S}^n \rightarrow \mathbb{R}P^n$, $\mathbb{S}^{2n+1} \rightarrow \mathbb{C}P^n$, and $\mathbb{S}^{4n+3} \rightarrow \mathbb{H}P^n$, including as special cases the Hopf fibration $\mathbb{S}^3 \rightarrow \mathbb{C}P^1$ and the quaternionic Hopf fibration $\mathbb{S}^7 \rightarrow \mathbb{H}P^1$ (in Section 6.2).
- The rotation map $\mathbf{SO}(n+1) \rightarrow \mathbb{S}^n$ (in Section 7.1).
- The homotopy quotient $\mathbb{R} \vee \mathbb{R} \rightarrow (\mathbb{R} \vee \mathbb{R}) // C_2$, and many other homotopy quotients (in Section 7.2).

After this, we prove some corollaries for the theory of higher groups in Sections 7 and 8. We begin by reviewing the definition of higher groups, and then show that the homotopy quotient $X \rightarrow X // G$ of a type by the action of a crisp higher group is always a \mathfrak{f} -fibration. We then prove that \mathfrak{f} preserves the connectedness of crisp types, and conclude that the homotopy type of a higher group is itself a higher group.

Finally, in Section 9, we turn to the theory of covering spaces. We define the notion of covering following Wellen [14], and show that the type of coverings on a type is equivalent to the type of actions of its fundamental groupoid on discrete sets. We then show that every pointed type has a universal cover, and prove that this universal cover has the expected universal property. We end by showing that the universal cover of a higher group is a higher group.

Notation. In this paper, we will use Agda-inspired notation for the dependent pair and dependent function types. For a type family $E : B \rightarrow \mathbf{Type}$, we write

$$(b : B) \times E(b) \equiv \sum_{b:B} E(b), \text{ and}$$

$$(b : B) \rightarrow E(b) \equiv \prod_{b:B} E(b)$$

for the dependent pair (or dependent sum) type and the dependent function (or product) type respectively. The elements of $(b : B) \times E(b)$ are pairs (b, e) with $b : B$ and $e : E(b)$. The elements of $(b : B) \rightarrow E(b)$ are functions $b \mapsto f(b)$ with $f(b) : E(b)$ for $b : B$.

2. Modalities and the Modal Prism

A modality is a way of changing what it means for two elements of a type to be identified. To each type X , we associate a new type $\diamond X$ and a function $(-)^{\diamond} : X \rightarrow \diamond X$. For two points $x, y : X$ to be identified by the modality then means that $x^{\diamond} = y^{\diamond}$ as elements of $\diamond X$. Here are a few examples of modalities, with emphasis on those we will focus on in this paper.

- With the trivial modality $\diamond X = *$, any two points are uniquely identified.
- With the n -truncation modality $\|- \|_n$, two points are identified by giving an $(n - 1)$ -truncated identification between them. The base case is $\|X\|_{-2} = *$, the trivial modality.
- With the *shape* modality \int , two points may be identified by giving a path between them (that is, a map from the real line \mathbb{R} which sends 0 to one point and 1 to the other). We call $\int X$ the *homotopy type* of a type X .⁴
- With the *crystalline* modality \mathfrak{J} , two points may be identified by giving an *infinitesimal* path between them. We call $\mathfrak{J}X$ the *de Rham stack* of a type X .⁵

While the elementary theory of modalities appeared in the HoTT Book [13], the notion was developed more fully by Rijke, Shulman, and Spitters in [9]. In that paper, they give equivalences between four different notions of modality and prove a number of useful lemmas along the way. We will take our modalities to be “higher modalities”, one of the many equivalent notions of modality.

Definition 2.1. A *higher modality* consists of a *modal operator* $\diamond : \mathbf{Type} \rightarrow \mathbf{Type}$ together with:

- For each type X , a *modal unit*

$$(-)^{\diamond} : X \rightarrow \diamond X$$

- For every $A : \mathbf{Type}$ and $P : \diamond A \rightarrow \mathbf{Type}$, an *induction principle*

$$\mathbf{ind}_A^{\diamond} : ((a : A) \rightarrow \diamond P(a^{\diamond})) \rightarrow ((u : \diamond A) \rightarrow \diamond P(u)),$$

- For every $A : \mathbf{Type}$, $P : \diamond A \rightarrow \mathbf{Type}$, $f : (a : A) \rightarrow \diamond P(a^{\diamond})$ and $x : A$, a *computation rule*

$$\mathbf{comp}_A^{\diamond} : \mathbf{ind}_A^{\diamond}(f)(x^{\diamond}) = f(x),$$

- For any $u, v : \diamond A$, a witness that the modal unit $(-)^{\diamond} : u = v \rightarrow \diamond(u = v)$ is an equivalence.

⁴The modality \int appears as Definition 9.6 of [12], and we review it in Section 4.

⁵The crystalline modality appears formally as Axiom 3.4.1 in [15], and in the higher categorical setting in Definition 4.2.1 of [10], where it is called the *infinitesimal shape* modality

We say a type X is \diamond -modal if $(-)^{\diamond} : X \rightarrow \diamond X$ is an equivalence, and we define

$$\mathbf{Type}_{\diamond} \equiv (X : \mathbf{Type}) \times \text{is}\diamond\text{Modal}(X)$$

to be the universe of \diamond -modal types. A type X is \diamond -separated if for all $x, y : X$, the type of identifications $x = y$ is \diamond -modal.

A modality is in particular a *reflective subuniverse*: pre-composition by $(-)^{\diamond}$ gives an equivalence

$$(\diamond X \rightarrow Z) \xrightarrow{\sim} (X \rightarrow Z)$$

whenever Z is \diamond -modal (see Theorem 1.13 of [9]). Any map $\eta : X \rightarrow K$ from X to a modal type K which satisfies the same property is called a \diamond -unit, since from this property it can be show that $K \simeq \diamond X$ and $\eta = (-)^{\diamond}$ under this equivalence.

Modal types are closed under the basic operations of dependent type theory in the following way.

Lemma 2.2. Let X be a type and $P : X \rightarrow \mathbf{Type}$ a family of types.

- If X is modal and for all $x : X$, Px is modal, then $(x : X) \times Px$ is modal.
- If for all $x : X$, Px is modal, then $(x : X) \rightarrow Px$ is modal.

Proof. See Theorem 1.32 and Lemma 1.26 of [9]. □

As a corollary, a number of useful properties of modal types are also modal.

Corollary 2.3. Let A be a modal type. Then

$$\text{isContractible}(A) \equiv (a : A) \times ((a' : A) \rightarrow (a = a'))$$

is modal. If B is also a modal type and $f : A \rightarrow B$, then

$$\text{isEquiv}(f) \equiv (b : B) \rightarrow \text{isContractible}(\text{fib}_f(b))$$

is modal.

When we use the induction principle of a modality, it often makes sense to think of it “backwards”. That is, we think of the induction principle as saying that in order to map out of $\diamond A$ into a modal type, it suffices to map out of A . Or, with variables, in order to define $T(u) : \diamond P(u)$ for $u : \diamond A$, it suffices to assume that $u \equiv a^{\diamond}$ for $a : A$. In prose, we will just say that \diamond -induction lets us assume u is of the form a^{\diamond} .

We can extend the operation of \diamond to a functor using the induction principle. If $f : X \rightarrow Y$, then define $\diamond f : \diamond X \rightarrow \diamond Y$ by $\diamond f(x^{\diamond}) \equiv f(x)^{\diamond}$, or explicitly by

$$\diamond f \equiv \mathbf{ind}_X^{\diamond}((-)^{\diamond} \circ f).$$

Using the computation rule, we get a *naturality square*

$$\begin{array}{ccc} X & \xrightarrow{(-)^{\diamond}} & \diamond X \\ f \downarrow & & \downarrow \diamond f \\ Y & \xrightarrow{(-)^{\diamond}} & \diamond Y \end{array}$$

Any commuting square induces a map from the fiber of the left map to the fiber of the right. Therefore, we get the map $\delta : \mathbf{fib}_f(y) \rightarrow \mathbf{fib}_{\diamond f}(y^\diamond)$ for any $y : Y$ given by

$$\delta((x : X), (p : fx = y)) := (x^\diamond, \mathbf{comp}^\diamond \cdot (\mathbf{ap}(-)^\diamond p)).$$

As the sum of modal types is modal, $\mathbf{fib}_{\diamond f}(y^\diamond) \equiv (u : \diamond X) \times (\diamond f(u) = y^\diamond)$ is modal. Therefore, this map factors through $\diamond \mathbf{fib}_f(y)$ uniquely, giving us the *modal prism*.

$$\begin{array}{ccc} \mathbf{fib}_f(y) & \xrightarrow{\delta} & \mathbf{fib}_{\diamond f}(y^\diamond) \\ & \searrow^{(-)^\diamond} & \nearrow^\gamma \\ & \diamond \mathbf{fib}_f(y) & \end{array}$$

The modal prism divides functions in 5 possible kinds. Four of these possibilities arrange themselves into orthogonal factorization systems; the other gives a mediating notion which is the focus of this paper.

Definition 2.4. Let $f : X \rightarrow Y$ and consider the modal prism as in Figure 1. Then f is

- \diamond -modal if $(-)^{\diamond}$ is an equivalence for all $y : Y$,
- \diamond -connected if $\diamond \mathbf{fib}_f(y)$ is contractible for all $y : Y$,
- \diamond -étale if δ is an equivalence for all $y : Y$.
- a \diamond -equivalence if $\mathbf{fib}_{\diamond f}(y^\diamond)$ is contractible for all $y : Y$,
- a \diamond -fibration if γ is an equivalence for all $y : Y$.

Remark 2.5. By a quick application of \diamond -induction, we see that f is a \diamond -equivalence if and only if $\diamond f$ is an equivalence. And, by the lemma that a square is a pullback if and only if the induced map on fibers is an equivalence, f is \diamond -étale if and only if its naturality square is a pullback.

We can see relations between these definitions right off the bat.

Lemma 2.6. Let $f : X \rightarrow Y$. Then:

- f is \diamond -étale if and only if it is \diamond -modal and a \diamond -fibration.
- f is \diamond -connected if and only if it is a \diamond -equivalence and a \diamond -fibration.

Proof. Since the modal prism commutes, if f is \diamond -modal and a \diamond -fibration, then it is \diamond -étale. On the other hand, since $\mathbf{fib}_{\diamond f}(y^\diamond)$ is modal, if f is \diamond -étale then $\mathbf{fib}_f(y)$ is \diamond -modal and so $(-)^{\diamond}$ is an equivalence and hence so is γ .

If f is a \diamond -equivalence and a \diamond -fibration, then $\diamond \mathbf{fib}_f(y)$ is contractible as it is equivalent to the contractible $\mathbf{fib}_{\diamond f}(y^\diamond)$. On the other hand, if f is \diamond -connected, then it is a \diamond -equivalence by Lemma 1.35 of [9], and so γ is a map between contractible types and is therefore an equivalence. □

Recall that any function $f : X \rightarrow Y$ gives an equivalence $X \simeq (y : Y) \times \mathbf{fib}_f(y)$ over Y . Therefore, by totalizing the modal prism, we can find two factorizations of any map f , connected in the middle by $\mathbf{tot}(\gamma)$:

$$\begin{array}{ccccc} & & X & & \\ & \swarrow^{\mathbf{tot}((-)^{\diamond})} & \downarrow f & \searrow^{\mathbf{tot}(\delta)} & \\ (y : Y) \times \diamond \mathbf{fib}_f(y) & \dashrightarrow & & \dashrightarrow & (y : Y) \times \mathbf{fib}_{\diamond f}(y^\diamond) \\ & \searrow^{\mathbf{fst}} & \downarrow f & \swarrow_{\mathbf{fst}} & \\ & & Y & & \end{array}$$

In [9], Rijke, Shulman, and Spitters prove that the left factorization is a *stable orthogonal factorization system*. In particular, $\mathbf{tot}((-)^\diamond)$ is \diamond -connected, and $\mathbf{fst} : (y : Y) \times \diamond \mathbf{fib}_f(y) \rightarrow Y$ is \diamond -modal, and these give the unique \diamond -connected/ \diamond -modal factorization of f . The connected/modal factorization of a map f is also preserved under pullback; if $y : A \rightarrow Y$ is any map, then the factorization of the pullback y^*f is the pullback of the factorization of f along y .

This can be seen most clearly by viewing the factorization system from the point of view of type families. A map $f : X \rightarrow Y$ corresponds to the type family $\mathbf{fib}_f : Y \rightarrow \mathbf{Type}$, and its modal factor corresponds to the type family $\diamond \mathbf{fib}_f : Y \rightarrow \mathbf{Type}$. On type families, pullback along $y : A \rightarrow Y$ corresponds to composition, so y^*f corresponds to $\lambda a : A. \mathbf{fib}_f(ya) : A \rightarrow \mathbf{Type}$. The modal factorization of the pullback y^* is then $\lambda a : A. \diamond \mathbf{fib}_f(ya)$, which is precisely the pullback of the modal factorization of f .

In his thesis [8], Rijke proves that the right factorization is an *orthogonal factorization system*. In particular, $\mathbf{tot}(\delta)$ is a \diamond -equivalence and $\mathbf{fst} : (y : Y) \times \mathbf{fib}_{\diamond f}(y^\diamond) \rightarrow Y$ is \diamond -étale, and this is the unique \diamond -equivalence/ \diamond -étale factorization of f . This is, however, not a stable factorization system because the \diamond -equivalences are not in general preserved under pullback (see Remark 3.8 for an example).

Another important concept in the theory of modalities is that of a \diamond -cartesian square (see, for example, Definition 3.7.1 of [2]). We will make use of \diamond -cartesian squares in developing the theory of modal fibrations, so we will establish a few lemmas here.

Definition 2.7. A commuting square

$$\begin{array}{ccc} A & \xrightarrow{g} & B \\ f \downarrow & & \downarrow h \\ C & \xrightarrow{k} & D \end{array}$$

is \diamond -cartesian if the cartesian gap map $A \rightarrow B \times_D C$ is \diamond -connected.

Note that a id-cartesian square for the identity modality \mathbf{id} is simply a pullback. Before proving our lemmas concerning \diamond -cartesian squares,

Lemma 2.8. Consider a square

$$\begin{array}{ccc} A & \xrightarrow{g} & B \\ f \downarrow & & \downarrow h \\ C & \xrightarrow{k} & D \end{array}$$

commuting via $S : (x : A) \rightarrow (k(f(x)) = h(g(x)))$. Let $c : C$, and define the map $G : \mathbf{fib}_f(c) \rightarrow \mathbf{fib}_h(kc)$ by

$$G(x : A, w : fx = c) := (gx, S(x)^{-1} \cdot k_* w).$$

Then for any $(b, p) : \mathbf{fib}_h(kc)$, we have an equivalence $\mathbf{fib}_G((b, p)) = \mathbf{fib}_{\mathbf{gap}}((c, bp))$ with the fiber of the gap map $A \rightarrow B \times_D C$.

Proof. We find the equivalence as the following composite:

$$\begin{aligned} \mathbf{fib}_G((b, p)) &: \equiv ((x, w) : \mathbf{fib}_g(c)) \times (G(x, w) = (b, p)) \\ &= (x : A) \times (w : fx = c) \times ((gx, S(x)^{-1} \cdot k_* w) = (b, p)) \\ &= (x : A) \times ((gx, fx, S(x)^{-1}) = (b, c, p)) \\ &= \mathbf{fib}_{\mathbf{gap}}((b, c, p)). \end{aligned}$$

□

Using this, we can give a characterization of \diamond -cartesian maps which resembles the usual characterization of pullbacks as fiberwise equivalences.

Lemma 2.9. A commuting square

$$\begin{array}{ccc} A & \xrightarrow{g} & B \\ f \downarrow & & \downarrow h \\ C & \xrightarrow{k} & D \end{array}$$

is \diamond -cartesian if and only if for every $c : C$, the induced map

$$G : \mathbf{fib}_f(c) \rightarrow \mathbf{fib}_h(kc)$$

induced on fibers is \diamond -connected.

Proof. By Lemma 2.8, the fibers of the gap map are the fibers of G ; so, the fibers of the gap map are \diamond -connected if and only if the fibers of G are. \square

The following lemmas may be found in [2] as Lemmas 3.7.4 and 3.7.3 respectively. We will prove them in HoTT.

Lemma 2.10. Consider a pair of commuting squares:

$$\begin{array}{ccccc} A & \longrightarrow & B & \longrightarrow & E \\ \downarrow & & \downarrow & & \downarrow \\ C & \xrightarrow{k} & D & \longrightarrow & F \end{array}$$

Then

1. If the left square and the right square are \diamond -cartesian, then so is the composite square.
2. If the left square and the composite square are \diamond -cartesian, and k is surjective, then the right square is \diamond -cartesian.
3. If the right square is a pullback and the composite square is \diamond -cartesian, then the left square is \diamond -cartesian.

Proof. We will appeal to Lemma 2.9 a number of times. To prove the first fact, let $c : C$ and consider the following diagram:

$$\begin{array}{ccccc} \mathbf{fib}_f(c) & \longrightarrow & \mathbf{fib}_h(kc) & \longrightarrow & \mathbf{fib}_\ell(jkc) \\ \downarrow & & \downarrow & & \downarrow \\ A & \longrightarrow & B & \longrightarrow & E \\ f \downarrow & & \downarrow h & & \downarrow \ell \\ C & \xrightarrow{k} & D & \xrightarrow{j} & F \end{array}$$

The squares are \diamond -cartesian when the maps on fibers are \diamond -connected, and \diamond -connected maps are closed under composition, so the outer square is also \diamond -cartesian.

With a modification of the above argument, we can prove the third fact. Suppose instead that the right square is a pullback, so that $\mathbf{fib}_h(kc) \rightarrow \mathbf{fib}_\ell(jkc)$ is an equivalence. Then since the composite map $\mathbf{fib}_f(c) \rightarrow \mathbf{fib}_\ell(jkc)$ is \diamond -connected, so is $\mathbf{fib}_f(c) \rightarrow \mathbf{fib}_h(kc)$.

To prove the second fact, suppose that $d : D$; then, since k is assumed to be surjective and we are trying to prove a proposition, we may suppose we have a $c : C$ with $kc = d$. Then we

can consider the above diagram again with $\text{fib}_f(c) \rightarrow \text{fib}_h(d)$ and $\text{fib}_f(c) \rightarrow \text{fib}_\ell(jd)$ modally connected. By right cancellability of modally connected maps (Lemma 1.33 of [9]), we see that therefore $\text{fib}_h(d) \rightarrow \text{fib}_\ell(jd)$ is \diamond -connected. □

Lemma 2.11. Suppose that

$$\begin{array}{ccc} A & \xrightarrow{g} & B \\ f \downarrow & & \downarrow h \\ C & \xrightarrow{k} & D \end{array}$$

is a \diamond -cartesian square. In its modal factorization

$$\begin{array}{ccccc} A & \longrightarrow & (b : B) \times \diamond \text{fib}_g(b) & \longrightarrow & B \\ \downarrow & & \downarrow & & \downarrow \\ C & \longrightarrow & (d : D) \times \diamond \text{fib}_k(d) & \longrightarrow & D \end{array} \tag{1}$$

the right square is a pullback.

Proof. Here we will use the proof of this fact from Lemma 3.7.3 of [2]. Consider the following diagram:

$$\begin{array}{ccccccc} A & \longrightarrow & B \times_D C & \xrightarrow{x} & B \times_D ((d : D) \times \diamond \text{fib}_k(d)) & \xrightarrow{y} & B \\ & \searrow f & \downarrow & & \downarrow & & \downarrow h \\ & & C & \xrightarrow{\ell} & (d : D) \times \diamond \text{fib}_k(d) & \xrightarrow{r} & D \end{array}$$

where we have taken two pullbacks. By construction, ℓ is \diamond -connected and r is \diamond -modal. By stability of the \diamond -connected / \diamond -modal factorization system, x is also \diamond -connected and y is \diamond -modal. Since by hypothesis the gap map $A \rightarrow B \times_D C$ is \diamond -connected, the composite $A \rightarrow B \times_D ((d : D) \times \diamond \text{fib}_k(d))$ is \diamond -connected, so by the uniqueness of \diamond -connected / \diamond -modal factorizations, we see that $B \times_D ((d : D) \times \diamond \text{fib}_k(d))$ must be equivalent to \diamond -factorization $(b : B) \times \diamond \text{fib}_g(b)$. Therefore, the right hand pullback square in the above diagram is equivalent to the right hand square in Diagram 1, showing that it is a pullback. □

Using these lemmas, we can prove a slight improvement of the Proposition 5.1 of [4], using essentially the same proof.

Theorem 2.12. Suppose that

$$\begin{array}{ccc} A & \xrightarrow{g} & B \\ f \downarrow & & \downarrow h \\ C & \xrightarrow{k} & D \end{array}$$

is a \diamond -cartesian square, and that B and D are \diamond -modal. Then the square

$$\begin{array}{ccc} \diamond A & \xrightarrow{\tilde{g}} & B \\ \diamond f \downarrow & & \downarrow h \\ \diamond C & \xrightarrow{\tilde{k}} & D \end{array}$$

is a pullback, where the maps $\tilde{g} : \diamond A \rightarrow B$ and $\tilde{k} : \diamond C \rightarrow D$ are the unique factorizations of g and k respectively.

Proof. Consider the following diagram:

$$\begin{array}{ccccc}
 A & \xrightarrow{r} & B \times_D \diamond C & \xrightarrow{x} & B \\
 f \downarrow & & \downarrow & & \downarrow h \\
 C & \xrightarrow{(-)^\diamond} & \diamond C & \xrightarrow{\tilde{k}} & D
 \end{array} \tag{2}$$

We will start by showing that the map $r : A \rightarrow B \times_D \diamond C$ is \diamond -connected. Let $c : C$, and extend the diagram as follows:

$$\begin{array}{ccccc}
 \text{fib}_f(c) & \xrightarrow{z} & \text{fib}_{\text{snd}}(c^\diamond) & \xrightarrow{\sim} & \text{fib}_h(kc) \\
 \downarrow & & \downarrow & & \downarrow \\
 A & \xrightarrow{r} & B \times_D \diamond C & \xrightarrow{x} & B \\
 f \downarrow & & \downarrow & & \downarrow h \\
 C & \xrightarrow{(-)^\diamond} & \diamond C & \xrightarrow{\tilde{k}} & D
 \end{array}$$

Since the square on the bottom right is a pullback, we get an equivalence between the map $z : \text{fib}_f(c) \rightarrow \text{fib}_{\text{snd}}(c^\diamond)$ and the composite $G : \text{fib}_f(c) \rightarrow \text{fib}_h(kc)$. Since, by Lemma 2.9, G is \diamond -connected, we see for all $c : C$ the map $z : \text{fib}_f(c) \rightarrow \text{fib}_{\text{snd}}(c^\diamond)$ is \diamond -connected. Since $(-)^\diamond$ is always \diamond -connected, we may conclude by Lemma 1.39 of [9] that the map $r : A \rightarrow B \times_D \diamond C$ is \diamond -connected.

Now, as the pullback of maps between modal types, $B \times_D \diamond C$ is modal. Therefore, r is a \diamond -connected map into a \diamond -modal type, which makes it a \diamond -unit. Therefore, the square on the right in Diagram 2 is the square we are trying to show is a pullback. \square

Remark 2.13. We can also see Theorem 2.12 as a corollary of Lemma 2.11 by noting that the right square in that lemma will be the square in the conclusion of Theorem 2.12 when B and D are modal.

3. Modal Fibrations

Recall that a map $f : X \rightarrow Y$ is a \diamond -fibration if and only if the induced map $\gamma : \diamond \text{fib}_f(y) \rightarrow \text{fib}_{\diamond f}(y^\diamond)$ is an equivalence for all $y : Y$. In other words, $f : X \rightarrow Y$ is a \diamond -fibration if \diamond preserves its fibers in the sense that whenever

$$F \rightarrow X \xrightarrow{f} Y$$

is a fiber sequence (for any pointing of Y), so is

$$\diamond F \rightarrow \diamond X \xrightarrow{\diamond f} \diamond Y.$$

In other words, a \diamond -fibration is a map f whose fibers “correctly represent” the fibers of $\diamond f$.

For example, consider the shape modality \mathbb{J} . A \mathbb{J} -fibration is a map $f : X \rightarrow Y$ whose fibers have the same homotopy type as its homotopy fibers, the fibers of its induced map $\mathbb{J}f : \mathbb{J}X \rightarrow \mathbb{J}Y$ on homotopy types. A simple example of a \mathbb{J} -fibration is the projection $\pi_1 : \mathbb{R}^3 \rightarrow \mathbb{R}^2$; all the fibers of this map are identifiable with \mathbb{R} whose shape is contractible, and the fibers of its induced map on homotopy types are contractible. An example of a map which isn’t a fibration is the inclusion $i : * \rightarrow \mathbb{R}^2$ of the origin into the real plane. Over the point $(1, 1) : \mathbb{R}^2$, the fiber of i is empty, and so its homotopy type is empty. But the induced map $\mathbb{J}i : \mathbb{J}* \rightarrow \mathbb{J}\mathbb{R}^2$ is an equivalence since $\mathbb{J}\mathbb{R}^2$ is contractible, and so all the fibers of $\mathbb{J}i$ are equivalent to $*$ which is not empty.

Remark 3.1. This is the sense in which a \diamond -fibration is a “fibration”. It most closely resembles the notion of *quasi-fibration* of topological spaces introduced by Dold and Thom in [6], which is a continuous map $f : X \rightarrow Y$ such that for all $y \in Y$, the canonical map from the inverse image $f^{-1}(y)$ to the homotopy fiber $\text{fib}_f(y)$ is a weak equivalence. If, seeking analogy, we take “weak equivalence” to be \diamond -equivalence (which, for \mathbb{J} , means that a map is a weak equivalence if it induces an equivalence on homotopy types), then a \diamond -fibration is map f whose fibers are weakly equivalent to its “modal fibers”, the fibers of $\diamond f$.

However, the notion of \diamond -fibration is somewhat more robust than the notion of quasi-fibration, even in the case of \mathbb{J} . As we will see, \diamond -fibrations are closed under pullback, while quasi-fibrations are not. In this sense, \diamond -fibrations more closely resemble the *universal quasi-fibrations* introduced by Goodwillie in an email to the ALGTOP mailing list [7]. Intuitively, this is because universal quantification in type theory says more than it does in set theory — it implies a sort of continuity. We will come back to this subtle point in the next section when we introduce the notion of a crisp variable from Shulman’s real hohesion [12] in order to give a trick for showing a map is a \mathbb{J} -fibration.

Before we get there, let’s develop the basic theory of \diamond -fibrations for a general modality. First, we will characterize \diamond -fibrations as those maps on which the two factorization systems of \diamond agree.

Lemma 3.2. For $f : X \rightarrow Y$, the following are equivalent:

1. f is a \diamond -fibration.
2. The \diamond -modal factor of f is \diamond -étale.
3. The \diamond -equivalence factor of f is \diamond -connected.
4. The \diamond -connected/ \diamond -modal and \diamond -equivalence/ \diamond -étale factorizations of f are equal as factorizations of f .
5. The \diamond -naturality square for f is \diamond -cartesian.

Proof. We will first show that the first two conditions are equivalent; then we will argue that the next three are all equivalent by the uniqueness of each factorization. Finally, we note that the last condition is immediately equivalent to the third, since the \diamond -equivalence factor of f is the gap map of the \diamond -naturality square.

By Lemma 1.24 of [9], the unique factorization of the map

$$\lambda(y, x). (y, x^\diamond)^\diamond : (y : Y) \times \text{fib}_f(y) \rightarrow \diamond((y : Y) \times \text{fib}_f(y))$$

through $\diamond((y : Y) \times \text{fib}_f(y))$ is an equivalence. Therefore, the composite

$$(y : Y) \times \diamond \text{fib}_f(y) \xrightarrow{(-)^\diamond} \diamond((y : Y) \times \diamond \text{fib}_f(y)) \xrightarrow{\sim} \diamond((y : Y) \times \text{fib}_f(y))$$

is a \diamond -unit. So, for any $y : Y$, we get a diagram

$$\begin{array}{ccccc} \text{fib}_f(y) & \longrightarrow & \diamond \text{fib}_f(y) & \xrightarrow{\gamma} & \text{fib}_{\diamond f}(y^\diamond) \\ \downarrow & & \downarrow & & \downarrow \\ X & \longrightarrow & (y : Y) \times \diamond \text{fib}_f(y) & \longrightarrow & \diamond X \\ f \downarrow & & \downarrow & & \downarrow \\ Y & \xrightarrow{\text{id}} & Y & \longrightarrow & \diamond Y \end{array}$$

in which the bottom right square is a \diamond -naturality square. The map f is a \diamond -fibration if and only if the connecting map γ is an equivalence for all $y : Y$, and this happens if and only if the bottom right square is a pullback. But the bottom right square is a pullback precisely when $\text{fst} : (y : Y) \times \diamond \text{fib}_f(y) \rightarrow Y$ is \diamond -étale.

On the other hand, the fourth condition implies the second and third by simply transporting the properties. Each of the second and third also imply the fourth by the uniqueness of each factorization. Without loss of generality, consider the second condition. The \diamond -connected factor of f is always a \diamond -equivalence, so if the modal factor of f is \diamond -étale then the \diamond -connected/ \diamond -modal factorization is a \diamond -equivalence/ \diamond -étale factorization and so is equal to the canonical one by the uniqueness of such factorizations. \square

As a corollary, we can prove that \diamond -fibrations are closed under pullback, and give a descent theorem for \diamond -fibrations.

Corollary 3.3. Let

$$\begin{array}{ccc} A & \xrightarrow{x} & X \\ g \downarrow & & \downarrow f \\ B & \xrightarrow{y} & Y \end{array}$$

be a \diamond -cartesian square. If f is a fibration, then so is g . In particular, \diamond -fibrations are closed under pullback.

Proof. Consider the following cube:

$$\begin{array}{ccccc} & & \diamond A & \xrightarrow{\quad} & \diamond X \\ & \nearrow & \downarrow & \nearrow & \downarrow \diamond f \\ A & \xrightarrow{\quad} & X & & \\ g \downarrow & & \downarrow & & \\ B & \xrightarrow{\quad} & Y & & \\ & \nearrow & \downarrow & \nearrow & \\ & & \diamond B & \xrightarrow{\quad} & \diamond Y \end{array} \tag{3}$$

By hypothesis, the front face is \diamond -cartesian and, since f is a \diamond -fibration, so is the rightmost face. Therefore, by Lemma 2.10, the diagonal square is \diamond -cartesian. Then, by Theorem 2.12, the back face is a pullback. Then, by Lemma 2.10 again, the leftmost face is \diamond -cartesian, which shows that g is a \diamond -fibration. \square

Remark 3.4. It is at this point that we require a full modality, rather than just a *reflective subuniverse*. The proof of Theorem 2.12 uses the fact that \diamond -units are \diamond -connected, a fact which characterizes modalities amongst localizations (also known as reflective subuniverses). However, if one could prove Theorem 2.12 without using this fact, or prove that the pullback of a \diamond -étale map is \diamond -étale for \diamond a reflective subuniverse, then we could prove the pullback stability of \diamond -fibrations and so the rest of the theory of \diamond -fibrations would go through as well.

Using Lemma 2.10 and the characterization of \diamond -fibrations as those maps whose naturality squares are \diamond -cartesian, we can show that \diamond -fibrations have the same closure properties as \diamond -cartesian squares.

Theorem 3.5. *Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be maps.*

1. *If f and g are \diamond -fibrations, then $g \circ f$ is a \diamond -fibration.*
2. *If f and $g \circ f$ are \diamond -fibrations, and $\diamond f$ is surjective, then g is a \diamond -fibration.*
3. *If g is \diamond -étale and $g \circ f$ is a \diamond -fibration, then f is a \diamond -fibration.*

Proof. We apply Lemma 2.10 to the squares

$$\begin{array}{ccccc} X & \xrightarrow{f} & Y & \xrightarrow{g} & Z \\ \downarrow & & \downarrow & & \downarrow \\ \diamond X & \xrightarrow{\diamond f} & \diamond Y & \xrightarrow{\diamond g} & \diamond Z \end{array}$$

For the third part, remember that g is \diamond -étale precisely when its naturality square is a pullback. \square

We now have the tools to characterize \diamond -fibrations in another way. A modality is called *lex* if it preserves all pullbacks. Not all modalities are lex; for example, the truncation modalities are not, and nor is \mathbb{J} . The \diamond -fibrations are precisely the maps along which \diamond is lex. That is, \diamond preserves all pullbacks of a map f if and only if that map is a \diamond -fibration.

Theorem 3.6. *A map $f : X \rightarrow Y$ is a \diamond -fibration if and only if \diamond preserves every pullback of it in the sense that whenever the square on the left is a pullback, so is the square on the right.*

$$\begin{array}{ccc} A & \xrightarrow{x} & X \\ g \downarrow & & \downarrow f \\ B & \xrightarrow{y} & Y \end{array} \qquad \begin{array}{ccc} \diamond A & \xrightarrow{\diamond x} & \diamond X \\ \diamond g \downarrow & & \downarrow \diamond f \\ \diamond B & \xrightarrow{\diamond y} & \diamond Y \end{array}$$

Remark 3.7. For the case of \mathbb{J} , Theorem 3.6 gives us a sufficient condition for a pullback to be a *homotopy pullback* (that is, a pullback on homotopy types): if one of the legs is a \mathbb{J} -fibration, then the pullback is a homotopy pullback.

Proof. If \diamond preserves all pullbacks of f , then by taking $B \equiv *$, we see that \diamond preserves all fibers of f which by definition makes it a \diamond -fibration.

On the other hand, suppose that f is a \diamond -fibration and that the square on the left above is a pullback. Then the connecting map $\alpha : \mathbf{fib}_g(a) \rightarrow \mathbf{fib}_f(ya)$ is an equivalence for all $a : A$. Furthermore, g is also a \diamond -fibration by Corollary 3.3 and therefore the maps $\gamma_f : \diamond \mathbf{fib}_f(ya) \rightarrow \mathbf{fib}_{\diamond f}((ya)^\diamond)$ and $\gamma_g : \diamond \mathbf{fib}_g(a) \rightarrow \mathbf{fib}_{\diamond g}(a^\diamond)$ are equivalences for all $a : A$. These maps fit together into a commuting square:

$$\begin{array}{ccc} \diamond \mathbf{fib}_g(a) & \xrightarrow{\diamond \alpha} & \diamond \mathbf{fib}_f(ya) \\ \gamma_g \downarrow & & \downarrow \gamma_f \\ \mathbf{fib}_{\diamond g}(a^\diamond) & \longrightarrow & \mathbf{fib}_{\diamond f}((ya)^\diamond) \end{array}$$

Since the sides and top are equivalences, the bottom is also an equivalence.

Now, in order to show that the square on the right is a pullback, we need for the induced map $\zeta : \mathbf{fib}_{\diamond g}(u) \rightarrow \mathbf{fib}_{\diamond f}(\diamond y(u))$ to be an equivalence for all $u : \diamond B$. But we have only shown it for $u \equiv a^\diamond$, since $\diamond y(a^\diamond) = (ya)^\diamond$ by naturality. Luckily, as both $\mathbf{fib}_{\diamond g}(u)$ and $\mathbf{fib}_{\diamond f}(\diamond y(u))$ are \diamond -modal, $\mathbf{isEquiv}(\zeta)$ is also \diamond -modal for all $u : \diamond B$. We may therefore assume that $u \equiv a^\diamond$ by \diamond -induction. \square

As a corollary of this, we can prove a partial stability of the \diamond -equivalence/ \diamond -étale factorization system. A factorization system is stable if the left class is stable under pullback.

Remark 3.8. The class of \diamond -equivalences is not stable under pullback in general. For example, consider the following pullback

$$\begin{array}{ccc} \emptyset & \longrightarrow & * \\ \downarrow & & \downarrow 1 \\ * & \xrightarrow{0} & \mathbb{R} \end{array}$$

Though the bottom map is a \mathfrak{J} -equivalence since \mathbb{R} is homotopically contractible, the top map is not a \mathfrak{J} -equivalence.

On the other hand, \diamond -equivalences are preserved by pullback along \diamond -fibrations.

Corollary 3.9. Suppose that the following square is a pullback. If f is a \diamond -fibration and y a \diamond -equivalence, then x is a \diamond -equivalence.

$$\begin{array}{ccc} A & \xrightarrow{x} & X \\ g \downarrow & & \downarrow f \\ B & \xrightarrow{y} & Y \end{array}$$

Proof. Since f is a \diamond -fibration, the square

$$\begin{array}{ccc} \diamond A & \xrightarrow{\diamond x} & \diamond X \\ \diamond g \downarrow & & \downarrow \diamond f \\ \diamond B & \xrightarrow{\diamond y} & \diamond Y \end{array}$$

is also a pullback. But $\diamond y$ is an equivalence by hypothesis, and therefore so is $\diamond x$. □

All of this pullback preserving lets us add a few more conditions to the long list of equivalent conditions for lexness in Theorem 3.1 of [9].

Proposition 3.10. The following are equivalent:

1. The modality \diamond is lex.
2. Every map is a \diamond -fibration.
3. If every map $f_i : A_i \rightarrow B_i$ is a \diamond -fibration in a family of maps f , then the total map $\text{tot}(f) : (i : I) \times A_i \rightarrow (i : I) \times B_i$ is a \diamond -fibration.
4. For any map $f : X \rightarrow Y$, the connecting map $\text{tot}(\gamma) : (y : Y) \times \diamond \text{fib}_f(y) \rightarrow (y : Y) \times \text{fib}_{\diamond f}(y^\diamond)$ between factorizations of f is a \diamond -fibration.
5. The universal map $\mathbf{Type}_* \rightarrow \mathbf{Type}$ is a \diamond -fibration.

Proof. Conditions 1 and 2 are equivalent by the characterization of \diamond -fibrations in terms of pullback preservation, and condition 2 trivially implies conditions 3, 4, and 5. Every map between \diamond -modal types is \diamond -étale since for \diamond -modal types the modal units are equivalences. Therefore, the connecting map $\gamma : \diamond \text{fib}_f(y) \rightarrow \text{fib}_{\diamond f}(y^\diamond)$ is \diamond -étale and in particular a \diamond -fibration for any map $f : X \rightarrow Y$ and $y : Y$. This means that condition 3 implies condition 4. On the other hand, since \diamond -fibrations are closed under composition, if $\text{tot}(\gamma)$ is a \diamond -fibration then the \diamond -modal factor of any map $f : X \rightarrow Y$ is a \diamond -fibration, as it is the composite of $\text{tot}(\gamma)$ and the \diamond -étale factor of f . Therefore, by Lemma 3.2, f is a \diamond -fibration, so that condition 4 implies condition 2.

Finally, the last condition implies the second since \diamond -fibrations are closed under pullback. □

All objects are “fibrant” with respect to \diamond -fibrations in the sense that the terminal map is always a \diamond -fibration. We can say something more — every projection map $\text{fst} : A \times B \rightarrow A$ is a \diamond -fibration.

Lemma 3.11. For any types A and B , the projection map $\text{fst} : A \times B \rightarrow A$ is a \diamond -fibration.

Proof. This follows directly from the fact that \diamond preserves products. The map $(-)^{\diamond} \times (-)^{\diamond} : A \times B \rightarrow \diamond A \times \diamond B$ is a \diamond -unit by Lemma 1.27 of [9], and so for any $a : A$ we get a map of fiber sequences:

$$\begin{array}{ccc} B & \xrightarrow{(-)^{\diamond}} & \diamond B \\ \downarrow & & \downarrow \\ A \times B & \xrightarrow{(-)^{\diamond} \times (-)^{\diamond}} & \diamond A \times \diamond B \\ \text{fst} \downarrow & & \downarrow \text{fst} \\ A & \xrightarrow{(-)^{\diamond}} & \diamond A \end{array}$$

where the bottom square is a \diamond -naturality square. The induced map $\gamma : \diamond \text{fib}_{\text{fst}}(a) \rightarrow \text{fib}_{\diamond \text{fst}}(a^{\diamond})$ is therefore equal to the identity map of $\diamond B$, and so is an equivalence. \square

A map $f : X \rightarrow Y$ is equal to a projection $\text{fst} : Y \times Z \rightarrow Y$ if and only if $\text{fib}_f : Y \rightarrow \mathbf{Type}$ is constant, that is, if it factors through the point.

$$\begin{array}{ccc} Y & \xrightarrow{\text{fib}_f} & \mathbf{Type} \\ \downarrow & \nearrow Z & \\ * & & \end{array}$$

We have just shown that such maps are \diamond -fibrations, but we can do better. We can show that a map is a \diamond -fibration if and only if it has \diamond -locally constant \diamond -fibers in the sense made precise in the upcoming Theorem 3.14. First, we prove a similar characterization of \diamond -étale maps. This is the *modal descent* theorem of [4].

Lemma 3.12. Let $E : Y \rightarrow \mathbf{Type}_{\diamond}$ be a family of modal types. Then E factors through the modal unit of Y if and only if $\text{fst} : (y : Y) \times Ey \rightarrow Y$ is \diamond -étale. In particular, the type of such factorizations is a proposition.

Proof. If fst is \diamond -étale, then $\gamma : Ey \rightarrow \text{fib}_{\diamond \text{fst}}(y^{\diamond})$ is an equivalence; therefore, $\text{fib}_{\diamond \text{fst}} : \diamond Y \rightarrow \mathbf{Type}_{\diamond}$ is such a factorization.

On the other hand, suppose that $\tilde{E} : \diamond Y \rightarrow \mathbf{Type}_{\diamond}$ with $w : (y : Y) \rightarrow (Ey \simeq \tilde{E}y^{\diamond})$ is a factorization. Then the square

$$\begin{array}{ccc} (y : Y) \times Ey & \xrightarrow{\text{tot}(w)} & (u : \diamond Y) \times \tilde{E}u \\ \text{fst} \downarrow & & \downarrow \text{fst} \\ Y & \xrightarrow{\quad} & \diamond Y \end{array}$$

is a pullback. Since the unit $Y \rightarrow \diamond Y$ is \diamond -connected and \diamond -connected maps are closed under pullback, $\text{tot}(w)$ is \diamond -connected. As $(u : \diamond Y) \times \tilde{E}u$ is a sum of modal types over a modal type, it is modal, and therefore $\text{tot}(w)$ is a \diamond -unit and this square is a \diamond -naturality square. But then $\text{fst} : (y : Y) \times Ey \rightarrow Y$ is \diamond -étale since its \diamond -naturality square is a pullback.

To show that the type of such factorizations is a proposition, we just need to show that any factorization equals $(\text{fib}_{\diamond \text{fst}}, \gamma)$. This follows immediately from the uniqueness of \diamond -units. \square

As a corollary, we can characterize the \diamond -étale maps into a type Y .

Corollary 3.13. For any type Y , the type

$$\mathring{\text{Ét}}_{\diamond}(Y) := (X : \mathbf{Type}) \times (f : X \rightarrow Y) \times \text{is}\diamond\text{étale}(f)$$

is equivalent to the type $\diamond Y \rightarrow \mathbf{Type}_{\diamond}$ of families of modal types varying over $\diamond Y$.

Proof. Consider the following equivalence:

$$\begin{aligned} \mathring{\text{Ét}}_{\diamond}(Y) &:= (X : \mathbf{Type}) \times (f : X \rightarrow Y) \times \text{is}\diamond\text{étale}(f) \\ &\simeq (X : \mathbf{Type}) \times (f : X \rightarrow Y) \times (\tilde{E} : \diamond Y \rightarrow \mathbf{Type}_{\diamond}) \times \text{fib}_f = \tilde{E} \circ (-)^{\diamond} \\ &\simeq (E : Y \rightarrow \mathbf{Type}_{\diamond}) \times (\tilde{E} : \diamond Y \rightarrow \mathbf{Type}_{\diamond}) \times (E = \tilde{E} \circ (-)^{\diamond}) \\ &\simeq \diamond Y \rightarrow \mathbf{Type}_{\diamond} \square \end{aligned}$$

We may now prove the main theorem of this section, characterizing \diamond -fibrations as those maps with \diamond -locally constant \diamond -fibers.

Theorem 3.14. Let $E : Y \rightarrow \mathbf{Type}$ be a family of types. Then $\text{fst} : (y : Y) \times Ey \rightarrow Y$ is a \diamond -fibration if and only if there is a type family $\tilde{E} : \diamond Y \rightarrow \mathbf{Type}_{\diamond}$ making the following square commute:

$$\begin{array}{ccc} Y & \xrightarrow{E} & \mathbf{Type} \\ \downarrow & & \downarrow \diamond \\ \diamond Y & \xrightarrow{\tilde{E}} & \mathbf{Type}_{\diamond} \end{array}$$

Remark 3.15. In the case of the \mathfrak{J} modality, Theorem 3.14 can be understood as characterizing the \mathfrak{J} -fibrations as those maps whose fibers form a *local system* on their codomain. The factorization $\tilde{E} : \mathfrak{J}Y \rightarrow \mathbf{Type}_{\mathfrak{J}}$ of $\mathfrak{J}E : Y \rightarrow \mathbf{Type}_{\mathfrak{J}}$ shows that the homotopy types of the fibers Ey are *locally constant* in y . Moreover, the usual transport of identifications in $\mathfrak{J}Y$ gives rise to a *monodromy* action of the homotopy type $\mathfrak{J}Y$ on the homotopy types $\mathfrak{J}Ey$ of the fibers Ey .

Proof. By Lemma 3.2, fst is a fibration if and only if its modal factor $\mathcal{R}(\text{fst}) : (y : Y) \times \diamond(Ey) \rightarrow Y$ is \diamond -étale. By Lemma 3.12, $\mathcal{R}(\text{fst})$ is \diamond -étale if and only if $\diamond E : Y \rightarrow \mathbf{Type}_{\diamond}$ factors through $\diamond Y$. But this is exactly what we are asking for! \square

What is a $\|\!-\!\|_n$ -fibration? A map is a $\|\!-\!\|_n$ -equivalence exactly when it induces an equivalence on the homotopy groups π_k for $0 \leq k \leq n$ (see Theorem 8.8.3 of [13]), and is $\|\!-\!\|_n$ -connected when it furthermore induces a surjection on π_{n+1} (see Corollary 8.8.6 of [13]). Since a map is a $\|\!-\!\|_n$ -fibration if and only if its $\|\!-\!\|_n$ -equivalence factor is $\|\!-\!\|_n$ -connected, we might expect that a map is a $\|\!-\!\|_n$ -fibration if it induces a surjection on π_{n+1} . We can prove this naive conjecture by giving one more equivalent characterization of \diamond -fibrations — this time with a small caveat.

We first need an elementary lemma concerning fibers.

Lemma 3.16. Consider a square

$$\begin{array}{ccc} A & \xrightarrow{g} & B \\ f \downarrow & & \downarrow h \\ C & \xrightarrow{k} & D \end{array}$$

commuting via $S : (x : A) \rightarrow (k(f(x)) = h(g(x)))$. Let $a : A$, and define $F : \text{fib}_g(ga) \rightarrow \text{fib}_k(kfa)$ by

$$F(x : A, p : gx = ga) := (fx, S(x) \cdot h_*p \cdot S(a)^{-1}).$$

For $(c, q) : \text{fib}_k(kfa)$, define $G : \text{fib}_f(c) \rightarrow \text{fib}_h(kfa)$ by

$$G(x : A, w : fx = c) := (gx, S(x)^{-1} \cdot k_*w \cdot q).$$

Then we have an equivalence $\text{fib}_F(c, q) = \text{fib}_G(ga, S(a)^{-1})$ giving a (judgementally) commuting square

$$\begin{array}{ccc} \text{fib}_F(c, q) (= \text{fib}_G(ga, S(a)^{-1})) & \longrightarrow & \text{fib}_f(c) \\ \downarrow & & \downarrow \\ \text{fib}_g(ga) & \longrightarrow & A \end{array}$$

Proof. We find the equivalence as the following composite:

$$\begin{aligned} \text{fib}_F(c, q) &:= ((x, p) : \text{fib}_g(ga)) \times (F(x, p) = (c, q)) \\ &= (x : A) \times (p : gx = ga) \times ((fx, S(x) \cdot h_*p \cdot S(a)^{-1}) = (c, q)) \\ &= (x : A) \times (p : gx = ga) \times (w : fx = c) \times (k_*w^{-1} \cdot S(x) \cdot h_*p \cdot S(a)^{-1} = q) \\ &= (x : A) \times (w : fx = c) \times (p : gx = ga) \times (h_*p^{-1} \cdot S(x) \cdot k_*w \cdot q = S(a)^{-1}) \\ &= (x : A) \times (w : fx = c) \times (G(x, w) = (ga, S(a)^{-1})) \\ &= \text{fib}_G(ga, S(a)^{-1}). \end{aligned}$$

Note that throughout this equivalence, $x : A$ is not affected by the equivalences. Therefore, we end up with the judgementally commuting square as desired. \square

Theorem 3.17. *Let $f : X \rightarrow Y$.*

1. *If f is a \diamond -fibration, then for all $x : X$ the induced map $\text{fib}_{(-)\diamond}(x^\diamond) \rightarrow \text{fib}_{(-)\diamond}((fx)^\diamond)$ is \diamond -connected.*
2. *If the modal unit $(-)\diamond : X \rightarrow \diamond X$ is surjective, and for all $x : X$ the induced map $\text{fib}_{(-)\diamond}(x^\diamond) \rightarrow \text{fib}_{(-)\diamond}((fx)^\diamond)$ is \diamond -connected, then f is a \diamond -fibration.*

Proof. First, suppose that f a \diamond -fibration, and let $x : X$ seeking to show that the induced map $\text{fib}_{(-)\diamond}(x^\diamond) \rightarrow \text{fib}_{(-)\diamond}((fx)^\diamond)$ is \diamond -connected. By Lemma 3.16, the fiber of the induced map over $(y, p) : \text{fib}_{(-)\diamond}((fx)^\diamond)$ is equivalent to the fiber of $\delta : \text{fib}_f(y) \rightarrow \text{fib}_{\diamond f}(y^\diamond)$ over $(x^\diamond, S(x)^{-1})$ where $S : (x : X) \rightarrow (fx)^\diamond = \diamond f(x^\diamond)$ is witness to the commutativity of the naturality square. Since f is a \diamond -fibration, this δ is a \diamond -equivalence; but it is a \diamond -equivalence landing in a modal type, and is therefore a \diamond -unit, which is to say it is \diamond -connected.

Conversely, suppose that the modal unit $(-)\diamond : X \rightarrow \diamond X$ is surjective. We aim to show that $f : X \rightarrow Y$ is a \diamond -fibration, so it suffices to prove that the maps $\delta : \text{fib}_f(y) \rightarrow \text{fib}_{\diamond f}(y^\diamond)$ are \diamond -connected for all $y : Y$. So, suppose we have $(u, p) : \text{fib}_{\diamond f}(y^\diamond)$, seeking to show that $\text{fib}_\delta(u, p)$ is \diamond -connected. By the surjectivity of $(-)\diamond : X \rightarrow \diamond X$, we may assume u is of the form x^\diamond . Then Lemma 3.16 tells us that $\text{fib}_\delta(x^\diamond, p)$ is equivalent to the fiber of the induced map $\text{fib}_{(-)\diamond}(x^\diamond) \rightarrow \text{fib}_{(-)\diamond}((fx)^\diamond)$ over $(fx, S(x))$. But by hypothesis, this fiber was \diamond -connected. \square

Remark 3.18. The condition that $(-)\diamond : X \rightarrow \diamond X$ be surjective is often trivially satisfied. For many modalities — the n -truncation modalities and the shape modality included — *all* modal

units are surjective. In this case, Theorem 3.17 characterizes the \diamond -fibrations with no caveats. We might refer to modalities whose units are surjective as *global* modalities; they are counterposed to topological modalities, which are given by a nullification at a family of propositions, since any global topological modality is trivial. More specifically, any global modality is cotopological in the sense of Theorem 3.22 of [9].

Corollary 3.19. A map $f : X \rightarrow Y$ is a $\|-\|_n$ -fibration if and only if for all $y : Y$ and $(x, p) : \text{fib}_f(y)$, the induced map $\pi_{n+1}(X, x) \rightarrow \pi_{n+1}(Y, y)$ is surjective.

Proof. By Theorem 3.17, f is a $\|-\|_n$ -fibration if and only if the induced map $\text{fib}_{|-|_n}(x) \rightarrow \text{fib}_{|-|_n}(y)$ is $\|-\|_n$ -connected. As the fibers of $\|-\|_n$ -units, $\text{fib}_{|-|_n}(x)$ and $\text{fib}_{|-|_n}(y)$ are $\|-\|_n$ -connected, so the induced map is $\|-\|_n$ -connected if and only if the induced map

$$\pi_{n+1}(\text{fib}_{|-|_n}(x), (x, \text{refl})) \rightarrow \pi_{n+1}(\text{fib}_{|-|_n}(y), (y, \text{refl}))$$

is a surjection. But this map is equivalent to the induced map $\pi_{n+1}(X, x) \rightarrow \pi_{n+1}(Y, y)$. \square

Before moving on, let’s briefly consider a pair of modalities $\diamond \leq \blacklozenge$, where every \diamond -modal type is \blacklozenge -modal. For example, $\|-\|_n \leq \|-\|_{n+1}$. In particular, $\diamond X$ is \blacklozenge -modal, and so the unit $(-)^{\diamond} : X \rightarrow \diamond X$ factors uniquely through $(-)^{\blacklozenge} : X \rightarrow \blacklozenge X$, giving us a commuting diagram:

$$\begin{array}{ccc} X & \xrightarrow{(-)^{\blacklozenge}} & \blacklozenge X \\ & \searrow (-)^{\diamond} & \downarrow c \\ & & \diamond X \end{array}$$

Lemma 3.20. Suppose that every \diamond -modal type is \blacklozenge -modal. Then the connecting map $c : \blacklozenge X \rightarrow \diamond X$ is a \diamond -unit. As a corollary, for any $f : X \rightarrow Y$, we get a \diamond -naturality square

$$\begin{array}{ccc} \blacklozenge X & \longrightarrow & \diamond X \\ \blacklozenge f \downarrow & & \downarrow \diamond f \\ \blacklozenge Y & \longrightarrow & \diamond Y \end{array}$$

Proof. Let Z be a \diamond -modal type. It is therefore also \blacklozenge -modal. Precomposing by the above commutative triangle gives us a commutative diagram:

$$\begin{array}{ccc} (X \rightarrow Z) & \xleftarrow{\sim} & (\blacklozenge X \rightarrow Z) \\ & \swarrow \sim & \uparrow \\ & & (\diamond X \rightarrow Z) \end{array}$$

Because Z is both \diamond -modal and \blacklozenge -modal, the two horizontal maps are equivalences, and therefore the vertical map is an equivalence, as desired. \square

We aim to demonstrate the following relations between the different kinds of maps associated to these modalities.

Theorem 3.21. Suppose that every \diamond -modal type is \blacklozenge -modal, and that $f : X \rightarrow Y$. Then:

1. If f is \diamond -modal, then it is \blacklozenge -modal.
2. If f is \diamond -étale, then it is \blacklozenge -étale.

3. If f is a \blacklozenge -equivalence, then it is a \diamond -equivalence.
4. If f is \blacklozenge -connected, then it is \diamond -connected.
5. If f is a \blacklozenge -fibration and $\blacklozenge f$ is a \diamond -fibration, then f is a \diamond -fibration.

Proof of Theorem 3.21.

1. If f is \diamond -modal, then its fibers are \diamond -modal and so by hypothesis \blacklozenge -modal, so that f is \blacklozenge -modal.
2. If f is \diamond -étale, then by Lemma 3.12, fib_f factors through $\diamond X$ as $E : \diamond X \rightarrow \mathbf{Type}$. But then $E \circ c : \blacklozenge X \rightarrow \mathbf{Type}$ is a factorization of fib_f through $\blacklozenge X$, so that f is \blacklozenge -étale.
3. If f is a \blacklozenge -equivalence, then $\blacklozenge f$ is an equivalence. But then since $\diamond \blacklozenge f$ is equivalent to $\diamond f$ by Lemma 3.20, $\diamond f$ is an equivalence.
4. If f is \blacklozenge -connected, then $\blacklozenge \text{fib}_f(y)$ is contractible for all $y : Y$. But then $\diamond \text{fib}_f(y) = \diamond \blacklozenge \text{fib}_f(y)$ is contractible for all $y : Y$, so f is \diamond -connected.
5. Consider the following diagram.

$$\begin{array}{ccc}
 Y & \xrightarrow{\text{fib}_f} & \mathbf{Type} \\
 \downarrow & & \downarrow \blacklozenge \\
 \blacklozenge Y & \xrightarrow{\text{fib}_{\blacklozenge f}} & \mathbf{Type}_{\blacklozenge} \\
 \downarrow & & \downarrow \diamond \\
 \diamond Y & \xrightarrow{\text{fib}_{\diamond \blacklozenge f}} & \mathbf{Type}_{\diamond}
 \end{array}$$

If f is a \blacklozenge -fibration then the upper square commutes, and if $\blacklozenge f$ is a \diamond -fibration then the lower square commutes. If the outer square commutes, then fib_f factors through $\diamond Y$, and so is a \diamond -fibration. \square

4. A Brief Review of Cohesive HoTT

In this section, we review Mike Shulman’s Real Cohesive Homotopy Type Theory (as found in [12]). The *shape* modality \mathfrak{f} which sends a type to its homotopy type is defined in the context of Real Cohesive HoTT. It is the interplay of this modality with the *comodality* \mathfrak{b} that defines real cohesion, and that we will exploit to give a trick for showing that a map is a \mathfrak{f} -fibration.

For the reader who isn’t too familiar with real cohesion and doesn’t feel like getting too familiar with it, worry not. The details in this section revolve around the notion of *crisp* objects, which will be explained below. But every object (type or element) which appears in the empty context — that is to say, with no free variables in its definition — is crisp. Therefore, if you need a heuristic for understanding what it means to, say, have a crisp type $Z :: \mathbf{Type}$, just imagine that this means that Z has no free variables in its definition. For example, \mathbb{N} , \mathbb{Z} , \mathbb{R} , and \mathbf{Type} are all crisp types, while $0 : \mathbb{N}$, $\pi : \mathbb{R}$, and $\lambda x. x^2 + 2 : \mathbb{R} \rightarrow \mathbb{R}$ are all crisp elements since they have no free variables. Furthermore, any natural number may be assumed to be crisp, so that types like \mathbb{R}^n may be taken as crisp even though they involve a free variable $n : \mathbb{N}$.

In type theory, if you can argue that for all $x : X$, there is an $f(x) : Y$, then you have given a function $f : X \rightarrow Y$ in the process. In Shulman’s Real Cohesive HoTT, all functions will be *continuous* in a topological sense. So, saying that for $x : X$ we have a $f(x) : Y$ means that $f(x)$ must depend *continuously* on x . But not all dependencies are continuous. What if we want to express a discontinuous dependence?

To address this concern, Shulman introduces the notion of a “crisp variable”

$$a :: A$$

to express a discontinuous dependence. Hypothesizing $a :: A$ means that we can use a in a discontinuous manner; one way this is realized is in the crisp Law of Excluded middle.

Axiom 1 (Crisp excluded middle). For any crisp $P :: \mathbf{Prop}$, we have $P \vee \neg P$.

This axiom lets us use case analysis when assuming a crisp element of a set, even if the set has a native topology that wouldn’t admit case analysis constructively (such as the Dedekind real numbers \mathbb{R} , which cannot constructively be separated into two disjoint parts).

Any variable appearing in the type of a crisp variable must also be crisp, and a crisp variable may only be substituted by expressions that *only* involve crisp variables. When all the variables in an expression are crisp, we say that that expression is crisp; so, we may only substitute crisp expressions in for crisp variables. Constants — like $0 : \mathbb{N}$ or $\mathbb{N} : \mathbf{Type}$ — appearing in an empty context are therefore always crisp. This means that one cannot give a closed form example of a term which is *not* crisp; all terms with no free variables are crisp. For emphasis, we will say that a term which is not crisp is *cohesive*. The rules for crisp type theory can be found in Section 2 of [12].

One way to think of the difference between a cohesive dependence — for all $x : X$, $f(x) : Y$ — and a crisp dependence — for all $x :: X$, $f(x) : Y$ — is that the former expresses that $f(x)$ depends on a generic $x : X$, whereas in the latter we are saying that for *each* individual x , there is an $f(x)$.⁶

Given a *crisp* type X , we can remove its spatial structure to get a type $\flat X$. If X is a set, $\flat X$ can be thought of as its set of points.⁷ The rules for \flat can be found in Section 4 of [12]. They may be summed up by saying that $\flat X$ is inductively generated by elements of the form x^\flat for *crisp* $x :: X$. In particular, whenever we have a type family $C : \flat X \rightarrow \mathbf{Type}$, an $x : \flat X$, and an element $f(u) : C(u^\flat)$ depending on a *crisp* $u :: X$, we get an element

$$(\text{let } u^\flat := x \text{ in } f(u)) : C(x)$$

and if $x \equiv v^\flat$, then $(\text{let } u^\flat := x \text{ in } f(u)) \equiv f(v)$. This allows us to think of $\flat X$ as “the type of *crisp* points of X ”.

We have an inclusion $(-)_\flat : \flat X \rightarrow X$ given by $x_\flat \equiv \text{let } u^\flat := x \text{ in } u$. Since we are thinking of a dependence on a crisp variable as a *discontinuous* dependence, if this map $(-)_\flat : \flat X \rightarrow X$ is an equivalence then every *discontinuous* dependence on $x :: X$ underlies a *continuous* dependence on x . This leads us to the following definition:

Definition 4.1. A crisp type $X :: \mathbf{Type}$ is *crisply discrete* if the counit $(-)_\flat : \flat X \rightarrow X$ is an equivalence.⁸

We would like our formal notion of continuity coming from crisp types to match our topological notion of continuity as measured by continuous paths. We have a notion of discreteness coming from crisp variables — *crisply discrete* — but we also need a topological notion of discreteness.

⁶In particular, by the crisp excluded middle axiom, we may deal with each $x :: X$ on a case by case basis.

⁷This intuition really only works for sets, since if G is a group then $\flat BG$ behaves like the moduli stack of principal G -bundles with flat connection, and not “the type of points of BG ”.

⁸See Remark 6.13 of [12] for a discussion on some of the subtleties in the notion of crisp discreteness.

Definition 4.2. A type X is *discrete* if every path in it is constant in the sense that the inclusion of constant paths $X \rightarrow (\mathbb{R} \rightarrow X)$ is an equivalence.

Remark 4.3. The real numbers \mathbb{R} in Definition 4.2 — and throughout this paper — are the *Dedekind* real numbers and not the Cauchy real numbers. It can be proven in real cohesion (with a form of the axiom of choice) that the Cauchy real numbers are discrete, and that indeed they are equivalent to $\flat\mathbb{R}$ — see Corollary 8.28 of [12].

Note that we can form the proposition “is discrete” for any type, while we can only form the proposition “is crisply discrete” for crisp types, since to form $\flat X$, X must be crisp. The main axiom of real cohesion, which ties the liminal sort of topology implied by the use of crisp variables to the concrete topology of the real numbers, is that for crisp types being discrete and being crisply discrete coincide.

Axiom 2 ($\mathbb{R}\flat$). A crisp type $X :: \mathbf{Type}$ is crisply discrete if and only if it is discrete.

We can now define the shape modality as a localization.

Definition 4.4. The *shape* or *homotopy type* $\int X$ of a type X is defined to be the localization of X at the type of Dedekind real numbers \mathbb{R} (see Definition 9.6 of [12]). By construction, a type is \int -modal if and only if it is discrete.

Since \int is given by localization at a small type,⁹ it is accessible in the sense of [9]. Therefore, by Lemma 2.24 of [9], it may be extended canonically to any larger universe. For this reason, and because \flat is universe polymorphic, we will elide the size issues in the use of \int and, for example, consider the type of discrete types \mathbf{Type}_\flat to be \int -separated.

In the upcoming sections, we will need not only the shape modality \int , but the *n-truncated* shape modality \int_n .

Definition 4.5. Let \int_n be the modality whose modal types are discrete, *n-truncated* types. It can be constructed by localizing at the real line \mathbb{R} and the homotopy *n-sphere* S^n .

It may be tempting to define $\int_n X$ as $\|\int X\|_n$, but it is not currently known whether $\|D\|_n$ of a discrete type D is discrete; the author suspects that it is not true in general. However, for crisp types, this is true.

Proposition 4.6. Let $X :: \mathbf{Type}$ be a crisp type. Then $\int_n X = \|\int X\|_n$.

Proof. Since X is crisp, so is $\int X$. Since $\int X$ is crisp, $\|\int X\|_n$ is crisply an *n-type*. Then, by Corollary 6.7 of [12], $\flat\|\int X\|_n = \|\flat\int X\|_n$. But $\int X$ is discrete, so by Axiom $\mathbb{R}\flat$, $\flat\int X = \int X$. Therefore, $\|\int X\|_n$ is a discrete *n-type* and so the canonical map $\|\int X\|_n \rightarrow \int_n X$ is an equivalence. \square

We can think of $\int_n X$ as the “fundamental *n-groupoid*” of X . In particular,

- $\int_0 X$ is the set of connected components of X .
- $\int_1 X$ is the fundamental groupoid of X .

We can prove that $\int_0 X$ is the set of connected components of X in a naive sense.

Definition 4.7. Let X be a type. A *connected component* of X is a subtype $C : X \rightarrow \mathbf{Prop}$ of X which is

⁹Assuming propositional resizing, \mathbb{R} is as small as \mathbb{N} ; without propositional resizing, \mathbb{R} has the size of the universe of \mathbb{N} . We will assume propositional resizing here, as is common in homotopy type theory and valid in any ∞ -topos.

1. Inhabited: there is merely an $x : X$ such that $C(x)$.
2. Connected: If $C \subseteq P \cup \neg P$, then $C \subseteq P$ or $C \subseteq \neg P$.¹⁰
3. Detachable: For any $x : X$, either $C(x)$ or $\neg C(x)$.¹¹

We denote the set of connected components of X by $\pi_0 X$.

Connected components are quite rigid; if two connected components have non-empty intersection, then they are equal.

Lemma 4.8. Suppose that C and D are connected components of X . Then $C = D$ if and only if $C \cap D$ is non-empty.

Proof. If $C = D$, then $C \cap D$ is C and so is inhabited.

Since D is detachable, we have that $X \subseteq D \cup \neg D$, and therefore $C \subseteq D \cup \neg D$. Now, C is connected, so $C \subseteq D$ or $C \subseteq \neg D$; but it can't be the latter because then their intersection would be empty. So, $C \subseteq D$ and symmetrically $D \subseteq C$. \square

Intuitively, $\int_0 X$ should be the set of connected components of X and $(-)^{\int_0} : X \rightarrow \int_0 X$ should send $x : X$ to the connected component x^{\int_0} it is contained in. We can justify this intuition with the following theorem.

Lemma 4.9. Let $u : \int_0 X$, and let $C_u : X \rightarrow \mathbf{Prop}$ be defined by

$$C_u(x) := u = x^{\int_0}$$

Then C_u is a connected component of X , giving us a map $C : \int_0 X \rightarrow \pi_0 X$.

Proof. We need to prove that C_u is inhabited, connected, and detachable.

1. C_u is inhabited because $(-)^{\int_0}$ is surjective (by the same proof as that of Corollary 9.12 of [12]).
2. Suppose that $C_u \subseteq P \cup \neg P$. Consider the map $\chi : (x : X) \times C_u(x) \rightarrow \{0, 1\}$ sending x to 0 if $P(x)$ and x to 1 if $\neg P(x)$. As $\{0, 1\}$ is a discrete set (by Theorems 6.19 and 6.21 of [12], noting that $\{0, 1\} = \{0\} + \{1\}$), χ factors uniquely through $\int_0((x : X) \times C_u(x))$. But $(x : X) \times C_u(x) \equiv \mathbf{fib}_{(-)^{\int_0}}$ is a fiber of a \int_0 -unit, and so is \int_0 -connected. Therefore χ is constant, and so either all x in C_u satisfy P , or they all satisfy $\neg P$.
3. Since $\int_0 X$ is a discrete set, it has decidable equality by Lemma 8.15 of [12]. So, for any $x : X$, either $u = x^{\int_0}$ or not. But that exactly means that $C_u(x)$ or not. \square

Theorem 4.10. Let X be a type. Then the map $C : \int_0 X \rightarrow \pi_0 X$ of Lemma 4.9 is an equivalence.

Proof. We will show that the map C is surjective and injective.

1. To show that C is surjective, suppose that U is a connected component of X , seeking to witness $\|\mathbf{fib}_C(U)\|$. Since we are seeking a proposition and U is inhabited, we may assume that $x : X$ is in U . Then x is in $C_{x^{\int_0}} \cap U$, so that $C_{x^{\int_0}} = U$ by Lemma 4.8.
2. To show that C is injective, suppose that $C_u = C_v$ seeking to show that $u = v$. If $C_u = C_v$, then $C_u \cap C_v = C_u$ is merely inhabited. Since we are seeking a proposition, let x be an element in the intersection. But then $u = x^{\int_0}$ and $v = x^{\int_0}$, so $u = v$. \square

¹⁰This expresses the connectivity of C because it says that if C is contained in a *disjoint* union, it is contained wholly in one part.

¹¹This says that C is a *component* of X in the sense that X is the disjoint union of C and its complement.

Remark 4.11. Though we have framed this paper as taking place in the setting of Real Cohesion, it will in fact mostly use the “locally contractible” part of the theory — namely, crisp variables, the comodality \flat , the modality \flat , and the axiom relating them for crisp types. The only extra condition is that \flat commute with propositional truncation, which, as proven in [12], uses the codiscrete modality $\#$. It also follows from the fact (Proposition 8.8 of [12]) that propositions are discrete which only uses that \flat is given by localization at a family of pointed types.

In particular, Theorem 5.9 replies only on crisp type theory, while Theorem 6.1 relies on the adjoint relationship of \flat and \flat (namely, that crisp types are \flat -modal if and only if they are \flat -comodal). Theorems 7.7 and 8.6 relies only on Theorem 6.1, and are therefore also valid in general cohesion. On the other hand, the specific examples in Sections 6, 7, 8 and 9 take place in real cohesion.

Therefore, the theory of \flat -fibrations and coverings in the coming sections should work equally well in other settings that have an adjoint $\diamond \dashv \square$ modality/comodality pair implemented using crisp variables in which \square preserves propositional truncation. A likely example of such a situation would be the adjoint pair $\mathfrak{J} \dashv \&$ between the crystalline modality \mathfrak{J} which is given by localizing at a family of infinitesimal types, and the infinitesimal flat modality $\&$ which appears (in the language of ∞ -toposes, rather than type theory) in Schreiber’s [10]. Since \mathfrak{J} is the localization at a family of pointed types, propositions are crystalline and so $\&$ commutes with propositional truncation. In this setting, Theorem 6.1 would be used with Lemma 3.12 to show that the projections of certain bundles are \mathfrak{J} -étale (that is, formally étale or locally diffeomorphic).

The modality \mathfrak{J} is left exact, and so every map is an \mathfrak{J} -fibration. However, \mathfrak{J} -étale maps include the *formally étale* maps, or *local diffeomorphisms*. So the applications to covering theory of Section 9 can be interpreted in this setting as well.

5. Classifying Types of Discrete Structures are Discrete

In this section, we will show that the classifying types of bundles of crisply discrete structures are themselves discrete. As a corollary, the fibers of such a bundle depend only on the homotopy type of the base space. We will use this fact to show that maps whose fibers have a merely constant homotopy type — merely equivalent to some crisply discrete type — are \flat -fibrations.

First, we need a good notion of “type of discrete objects”. We will call these types *locally discrete*.

Definition 5.1. A type X is *locally discrete* if it is \flat -separated, that is, for all $x, y : X$, $x = y$ is discrete. A crisp type X is *locally crisply discrete* if for all crisp $x, y :: X$, $x = y$ is crisply discrete; more explicitly, for all $x, y : \flat X$, $x_{\flat} = y_{\flat}$ is crisply discrete.

Remark 5.2. We can’t explicitly quantify over crisp elements $x, y :: X$ in Shulman’s crisp type theory, but we can quantify over cohesive elements $x, y : \flat X$. These amount to the same thing, since if x and y are crisp elements of X , then $x_{\flat} = y_{\flat}$ is the same type as $x = y$.

In Agda, which has incorporated the \flat modality since version 2.6, we can quantify over crisp variables.

That we can think of locally discrete types as being types *of* discrete objects is justified by the following lemma.

Lemma 5.3. The type \mathbf{Type}_{\flat} of discrete types is locally discrete.

Proof. For any modality, the types of identifications between modal types are equivalent to modal types. In particular, \mathbf{Type}_f is separated relative to the canonical extension of f to any universe containing \mathbf{Type} . \square

In [5], Christensen, Opie, Rijke, and Scoccola show that if a modality \diamond is given by localization at a type X , then the \diamond -separated types also form a modality whose operator is given by localization at the suspension ΣX (see Lemma 2.15 and Remark 2.16 of [5]). As a corollary, by Lemma 2.2 we get that locally discrete types are closed under dependent sums.

Lemma 5.4. If X is locally discrete and $P : X \rightarrow \mathbf{Type}$ is a family of locally discrete types, then $(x : X) \times Px$ is locally discrete.

We can package this result into a useful extension of the idea that a locally discrete type is a type of discrete objects. Many structured objects are captured by the notion of a *standard notion of structure*, which appears in the HoTT Book [13] in Section 9.8 as a tool to prove the structure identity principle. A standard notion of structure on a category \mathcal{C} is a pair (P, H) where $P : \mathcal{C}_0 \rightarrow \mathbf{Type}$ assigns to each object of \mathcal{C} its type of (P, H) -structures (and H gives a notion of homomorphism between such structures). For example, a group is a standard notion of structure on the category of sets by letting P take each set to the set of group structures on it. We can read the previous lemma as saying that discretely structured discrete objects are also discrete, in the following way.

Corollary 5.5. Let \mathcal{C} be a category whose type of objects \mathcal{C}_0 is locally discrete type, and (P, H) be a standard notion of structure on \mathcal{C} such that for all $x : \mathcal{C}_0$, Px is discrete. Then the type of (P, H) structures is locally discrete.

Proof. The type of structures is just the dependant sum $(x : \mathcal{C}_0) \times Px$, which is locally discrete by the above corollary. \square

There are two ways to say a crisp type $X :: \mathbf{Type}$ is discrete: either $(-)_b : \flat X \rightarrow X$ is an equivalence or $(-)^f : X \rightarrow \flat X$ is an equivalence. Correspondingly, there are two ways to say that a crisp type is locally discrete, which we have given the names of locally discrete and locally crisply discrete. Though a crisp type which is locally discrete will always be locally crisply discrete, these two notions are likely *not* equivalent in general since the latter only quantifies over *crisp* elements of X . We can, however, give another characterization of locally crisply discrete types.

Lemma 5.6. A crisp type X is locally crisply discrete if and only if $(-)_b : \flat X \rightarrow X$ is an embedding.

Proof. Recall the left exactness of \flat (Theorem 6.1 of [12]); we have an equivalence $\flat(x = y) \simeq (x^b = y^b)$ for all crisp $x, y :: X$ making the following diagram commute:

$$\begin{array}{ccc}
 \flat(x = y) & \xrightarrow{\simeq} & x^b = y^b \\
 \searrow^{(-)_b} & & \swarrow^{\mathbf{ap}(-)_b} \\
 & x = y &
 \end{array}$$

Now, X is locally crisply discrete if and only if the downwards map on the left is an equivalence, and $(-)_b$ is an embedding if and only if the downwards map on the right is an equivalence. \square

Let's turn our attention to classifying types. In general, any type X can be seen as “classifying” the maps into it. This rather abstract way of thinking is more useful the more readily the objects of X can be turned into types, since maps into **Type** correspond to arbitrary bundles of types. For an $x : X$, the following general definition gives a classifying type for “bundles of x s”.

Definition 5.7. For a type X and a term $x : X$, we define

$$\mathbf{BAut}_X(x) := (y : X) \times \|x = y\|$$

This notation is inspired by the notation for the classifying space BG of principal G -bundles for a topological group G . If $G \simeq \mathbf{Aut}_X(x)$ is the group of automorphisms of some object (as, for example, $\mathbf{GL}_n(\mathbb{R}) \simeq \mathbf{Aut}_{\mathbf{Vect}_{\mathbb{R}}}(\mathbb{R}^n)$), then $\mathbf{BAut}_X(x)$ as defined above does classify principal G -bundles. If $\mathbf{Aut}_X(x)$ has a recognizable name G , we will write BG for $\mathbf{BAut}_X(x)$.

We will now show that if X is crisply locally discrete, and $x :: X$ is a crisp element, then $\mathbf{BAut}_X(x)$ is discrete.

Lemma 5.8. For any crisp type X and crisp $x :: X$, we have an equivalence $\flat \mathbf{BAut}_X(x) \simeq \mathbf{BAut}_{\flat X}(x^\flat)$ making the following triangle commute:

$$\begin{array}{ccc} \flat \mathbf{BAut}_X(x) & \xrightarrow{\simeq} & \mathbf{BAut}_{\flat X}(x^\flat) \\ & \searrow^{(-)_\flat} & \swarrow_{(y,p) \mapsto y_\flat, \dots} \\ & \mathbf{BAut}_X(x) & \end{array}$$

Proof. Consider the following equivalence:

$$\begin{aligned} \flat \mathbf{BAut}_X(x) &::= \flat((y : X) \times \|x = y\|) \\ &\simeq (u : \flat X) \times \text{let } y^\flat := u \text{ in } \flat \|x = y\| \\ &\simeq (u : \flat X) \times \text{let } y^\flat := u \text{ in } \|\flat(x = y)\| \\ &\simeq (u : \flat X) \times \text{let } y^\flat := u \text{ in } \|x^\flat = y^\flat\| \\ &\simeq \mathbf{BAut}_{\flat X}(x^\flat). \end{aligned}$$

The first equivalence follows from Lemma 6.8, the second from Corollary 6.7, and the third from Theorem 6.1 of [12]. The final equivalence follows from Lemma 4.4 of [12], which says that $(\text{let } y^\flat := u \text{ in } f(y^\flat)) = f(u)$.

On $(y, p)^\flat : \flat \mathbf{BAut}_X(x)$, this equivalence yields $(y^\flat, \dots) : \mathbf{BAut}_{\flat X}(x^\flat)$, and so when applying $(-)_\flat$ to either side, we find that the result is the same. \square

Theorem 5.9. *Suppose X is locally crisply discrete and $x :: X$. Then $\mathbf{BAut}_X(x)$ is (crisply) discrete.*

Proof. By the above lemma, it suffices to prove that $(y, \cdot) \mapsto (y_\flat, \cdot) : \mathbf{BAut}_{\flat X}(x^\flat) \rightarrow \mathbf{BAut}_X(x)$ is an equivalence. Now, $(-)_\flat : \flat X \rightarrow X$ is an embedding because X is locally crisply discrete, so the map in question is an embedding as well. We just need to show it is surjective.

Suppose $y : \mathbf{BAut}_X(x)$. To prove surjectivity, we need to inhabit $\|\text{fib}(y)\|$. Because we are trying to prove a proposition, we may assume that $p : x = y$; but then $(x^\flat, p) : \text{fib}(y)$. \square

6. Examples of \mathfrak{f} -Fibrations

By using Theorem 5.9 together with Theorem 3.14, we get a nice trick for showing that a map $f : X \rightarrow Y$ is a \mathfrak{f} -fibration. We just need give a crisply discrete type $F :: \mathbf{Type}_{\mathfrak{f}}$ such that $\int \mathfrak{fib}_f(y)$ is merely equivalent to F for all $y : Y$.

Theorem 6.1. *Let $f : X \rightarrow Y$. If there is a crisp type $F :: \mathbf{Type}_{\mathfrak{f}}$ such that for all $y : Y$, $\|F = \int \mathfrak{fib}_f(y)\|$, then f is a \mathfrak{f} -fibration. If furthermore we have that $\|F = \mathfrak{fib}_f(y)\|$ for all $y : Y$, then f is \mathfrak{f} -étale. If F is an n -type, then f is a \int_{n+1} -fibration (resp. \int_{n+1} -étale).*

Proof. By hypothesis, $\int \mathfrak{fib}_f$ factors through $\mathbf{BAut}(F)$. Since F is a crisp element of a locally discrete type, $\mathbf{BAut}(F)$ is discrete by Theorem 5.9 and therefore $\int \mathfrak{fib}_f$ factors through $\mathfrak{f}Y$. But then, by Theorem 3.14, f is a \mathfrak{f} -fibration. The second claim follows in the same way from Lemma 3.12. If F is an n -type, then $\mathbf{BAut}(F)$ is an $(n + 1)$ -type, and so the maps factor further through $\int_{n+1}X$. □

With a little effort, we can extend this trick to classify fibrations over disconnected spaces whose fibers over each part are different. A little care must be taken around crispness.

Corollary 6.2. Let $X, Y :: \mathbf{Type}$ and $f :: X \rightarrow Y$. Assuming the crisp axiom of choice, f is a \mathfrak{f} -fibration if and only if there is a $F :: \|\mathfrak{f}Y\|_0 \rightarrow \mathbf{Type}$ such that for all $y : Y$, $\|F(|y^{\mathfrak{f}}|_0) = \int \mathfrak{fib}_f(y)\|$.

Proof. First, if there is an $F :: \|\mathfrak{f}Y\|_0 \rightarrow \mathbf{Type}$ such that for all $y : Y$, $\|F(|y^{\mathfrak{f}}|_0) = \int \mathfrak{fib}_f(y)\|$, then $\int \mathfrak{fib}_f : Y \rightarrow \mathbf{Type}$ factors through $(u : \|\mathfrak{f}Y\|_0) \times \mathbf{BAut}(F(u))$. Since $\|\mathfrak{f}Y\|_0$ is crisply discrete (by Proposition 4.6) and for all $z : \mathfrak{b} \|\mathfrak{f}Y\|_0$ we have that (let $v^{\mathfrak{b}} := z$ in $\mathbf{isdiscrete}(\mathbf{BAut}(F(v)))$) by Theorem 5.9, we find that $(u : \|\mathfrak{f}Y\|_0) \times \mathbf{BAut}(F(u))$ is crisply discrete by Theorem 6.20 of [12]. Therefore, $\int \mathfrak{fib}_f$ factors through $(-)^{\mathfrak{f}}$, proving that f is an \mathfrak{f} -fibration.

On the other hand, suppose that f is a fibration. Assuming the crisp axiom of choice (Theorem 6.30 of [12]), there is a crisp section $s :: \|\mathfrak{f}Y\|_0 \rightarrow Y$ of $|(-)^{\mathfrak{f}}|_0 : Y \rightarrow \|\mathfrak{f}Y\|_0$; that is, we may choose an element in every fiber. Define $F(u) ::= \int \mathfrak{fib}_f(su)$. It remains to show that $\|F(|y^{\mathfrak{f}}|_0) = \int \mathfrak{fib}_f(y)\|$ for all $y : Y$. Since f is a fibration, we have that $\int \mathfrak{fib}_f = \mathfrak{fib}_{\mathfrak{f}f} \circ (-)^{\mathfrak{f}}$ and so

$$\left\| F(|y^{\mathfrak{f}}|_0) = \int \mathfrak{fib}_f(y) \right\| \simeq \left\| \mathfrak{fib}_{\mathfrak{f}f}((s|y^{\mathfrak{f}}|_0)^{\mathfrak{f}}) = \mathfrak{fib}_{\mathfrak{f}f}(y^{\mathfrak{f}}) \right\|$$

It will suffice to show that $\|s|y^{\mathfrak{f}}|_0^{\mathfrak{f}} = y^{\mathfrak{f}}\|$. But this is equivalent to $|s|y^{\mathfrak{f}}|_0|_0 = |y^{\mathfrak{f}}|_0$, which holds since s is a section. □

We can now use Theorem 6.1 to give a number of examples of \mathfrak{f} -fibrations. In this section, we will be working in real cohesion, assuming that \mathfrak{f} is given by localization at the type \mathbb{R} of Dedekind real numbers. We will add two more examples later, in Sections 7.1 and 7.2.

6.1 The Universal Cover of the Circle We will now show that the map $(\cos, \sin) : \mathbb{R} \rightarrow \mathbb{S}^1$ is a \mathfrak{f} -fibration, where \mathbb{S}^1 is the unit circle in \mathbb{R}^2 . In Section 9, we will show that it is indeed the universal cover of the circle \mathbb{S}^1 .

Lemma 6.3. The map $(\cos, \sin) : \mathbb{R} \rightarrow \mathbb{S}^1$ is \int_1 -étale, and so in particular is a \mathfrak{f} -fibration.

Proof. Let $r \equiv (\cos, \sin)$. Over $(x, y) : \mathbb{S}^1$, the fiber of r is $r^*(x, y) ::= \{\theta : \mathbb{R} \mid \cos \theta = x, \sin \theta = y\}$. We will show that $r^*(x, y)$ is merely equivalent to \mathbb{Z} .

For any $\theta : r^*(x, y)$ and $k : \mathbb{Z}$, we have that $\theta + 2\pi k$ is in $r^*(x, y)$. This gives map $\lambda k. \theta + 2\pi k : \mathbb{Z} \rightarrow r^*(x, y)$. Moreover, given any other $\varphi : r^*(x, y)$, the difference $\varphi - \theta$ is an integral multiple of 2π , which gives us a map $\lambda\varphi. \frac{\varphi - \theta}{2\pi} : r^*(x, y) \rightarrow \mathbb{Z}$. These maps are clearly inverse, and since r is merely surjective there is always some θ we may choose to make this equivalence.

We have therefore shown that $r^* : \mathbb{S}^1 \rightarrow \mathbf{Type}$ factors through $\mathbf{BAut}(\mathbb{Z})$.¹² But \mathbb{Z} is a crisply discrete set, so by Theorem 6.1, r is a fibration. \square

We can now use the fact that (\cos, \sin) is a fibration to calculate the fundamental group of the circle.

Theorem 6.4. *Let \mathbb{S}^1 be the unit circle in \mathbb{R}^2 . Then $\Omega \int \mathbb{S}^1 \simeq \mathbb{Z}$.*

Proof. Since

$$\mathbb{Z} \rightarrow \mathbb{R} \rightarrow \mathbb{S}^1$$

is a fiber sequence and (\cos, \sin) is a \int -fibration,

$$\mathbb{Z} \rightarrow * \rightarrow \int \mathbb{S}^1$$

is a fiber sequence, showing that $\Omega \int \mathbb{S}^1 \simeq \mathbb{Z}$. \square

6.2 Hopf Fibrations In the following, let \mathbb{K} be the real numbers \mathbb{R} , the complex numbers \mathbb{C} , or the quaternions \mathbb{H} . We will denote the apartness relation on any of these number systems by $x \# y$; for real numbers this means $|x - y| > 0$, and for the other two number systems this means $\|x - y\| > 0$. If X is a set with an apartness relation and $x : X$, we will denote by $X \# \{x\}$ the set of elements $y : X$ with $x \# y$.

Remark 6.5. In the presence of Shulman’s Axiom T of [12], the notions of apartness and non-equality in \mathbb{R} , \mathbb{C} , and \mathbb{H} coincide (see Theorem 8.32 of that paper). In this case, we could replace all instances of apartness by non-equality. Otherwise, we make no use of Axiom T.

Definition 6.6. A *line* in \mathbb{K}^{n+1} is a proposition $\mathcal{L} : \mathbb{K}^{n+1} \rightarrow \mathbf{Prop}$ satisfying:

1. There is (merely) an $x \# 0$ element in \mathcal{L} which is apart from 0.
2. For any element x in \mathcal{L} and $c : \mathbb{K}$, the scaled element cx is in \mathcal{L} .
3. For any elements x and y in \mathcal{L} , there is a unique $c : \mathbb{K}$ such that $cx = y$.

For a line \mathcal{L} , we define $\{\mathcal{L}\} := (x : \mathbb{K}^{n+1}) \times \mathcal{L}(x)$ to be its extent. We denote the type of lines in \mathbb{K}^{n+1} by $\mathbb{K}P^n$.

Quite obviously, every line is somehow identifiable with \mathbb{K} .

Lemma 6.7. Let $\mathcal{L} : \mathbb{K}P^n$ be a line. Then

$$\|\{\mathcal{L}\} = \mathbb{K}\|.$$

Proof. Since we are proving a proposition and since there exists a element apart from zero on \mathcal{L} , we may assume we have such an element x . Then the map $y \mapsto c$ where c is the unique element of \mathbb{K} such that $cx = y$ determines a map $\{\mathcal{L}\} \rightarrow \mathbb{K}$. Since for any $c : \mathbb{K}$, cx is on \mathcal{L} , this map is surjective. It is injective by the uniqueness condition (3). \square

¹²In fact, since the fibers are actually \mathbb{Z} -torsors, r^* factors through $\mathbf{B}\mathbb{Z}$, which would work just as well.

For any $x : \mathbb{K}^{n+1} \# \{0\}$, we get the line $\mathbb{K}x$ in the direction of x defined as

$$\mathbb{K}x(y) := \exists c : \mathbb{K}, cx = y.$$

We have a function $\tilde{h} : \mathbb{K}^{n+1} \# \{0\} \rightarrow \mathbb{K}P^n$, sending x to $\mathbb{K}x$. We refer to its restriction $h : \mathbb{S}_{\mathbb{K}^{n+1}} \rightarrow \mathbb{K}P^n$ to the unit sphere of \mathbb{K}^{n+1} as the *generalized Hopf map*.

Suppose that $\mathcal{L} : \mathbb{K}P^n$ is a line and consider the fiber $\text{fib}_{\tilde{h}}(\mathcal{L})$. By definition, this is the type of all elements $x : \mathbb{K}^{n+1} - \{0\}$ such that $\mathbb{K}x = \mathcal{L}$.

Lemma 6.8. For any line $\mathcal{L} : \mathbb{K}P^n$,

$$\text{fib}_{\tilde{h}}(\mathcal{L}) = \{\mathcal{L}\} \# 0$$

And, as a corollary,

$$\text{fib}_h(\mathcal{L}) = (x : \{\mathcal{L}\}) \times (\|x\| = 1)$$

consists of the elements on the line \mathcal{L} of unit length.

Proof. Suppose that x is in \mathcal{L} . By property 2, cx is in \mathcal{L} for any $c : \mathbb{K}$, and by property 3, every element of \mathcal{L} may be so expressed in a unique way. Therefore, $\mathbb{K}x = \mathcal{L}$.

On the other hand, if $\mathbb{K}x = \mathcal{L}$, then in particular $1 \cdot x = x$ is in \mathcal{L} . □

Putting together these two lemmas, we conclude that for all $\mathcal{L} : \mathbb{K}P^n$, the fiber of h over \mathcal{L} is merely equivalent to the unit sphere of \mathbb{K} :

$$\|\text{fib}_h(\mathcal{L}) = \mathbb{S}_{\mathbb{K}}\|.$$

In particular, their homotopy types are merely equivalent, and so by Theorem 6.1,

$$\mathbb{S}_{\mathbb{K}} \rightarrow \mathbb{S}_{\mathbb{K}^{n+1}} \rightarrow \mathbb{K}P^n$$

is a \mathfrak{J} -fibration.

Substituting \mathbb{R} , \mathbb{C} , and \mathbb{H} back in for \mathbb{K} , we see that:

Theorem 6.9.

- $\mathbb{S}^0 \rightarrow \mathbb{S}^n \rightarrow \mathbb{R}P^n$ is a \mathfrak{J} -fibration.¹³
- $\mathbb{S}^1 \rightarrow \mathbb{S}^{2n+1} \rightarrow \mathbb{C}P^n$ is a \mathfrak{J} -fibration. This includes the original Hopf fibration $\mathbb{S}^1 \rightarrow \mathbb{S}^3 \rightarrow \mathbb{C}P^1$.
- $\mathbb{S}^3 \rightarrow \mathbb{S}^{4n+3} \rightarrow \mathbb{H}P^n$ is a \mathfrak{J} -fibration. This includes the quaternionic Hopf fibration $\mathbb{S}^3 \rightarrow \mathbb{S}^7 \rightarrow \mathbb{H}P^1$.

6.3 A \mathfrak{J} -Fibration which is not a Hurewicz Fibration In this example we will prove that the projection of the x and y -axes onto the x -axis is a \mathfrak{J} -fibration. This is a classic example of a quasi-fibration which is not a Hurewicz fibration, since the x -axis cannot be lifted to a path going through a point $y \neq 0$ in the fiber over $x = 0$.

First, we need a useful and straightforward lemma.

Lemma 6.10. Let X be a type with a point $x_0 : X$ and suppose that for every $x : X$, we have a path $\gamma_x : \mathbb{R} \rightarrow X$ with $\gamma_x(0) = x$ and $\gamma_x(1) = x_0$. Then $\mathfrak{J}X$ is contractible.

¹³We will see in the next section that it is a covering map.

Proof. Define the map $\tilde{\gamma} : \mathbb{R} \rightarrow (X \rightarrow X)$ by $\tilde{\gamma}(t)(x) = \gamma_x(t)$ and note that $\tilde{\gamma}(0) = \text{id}_X$ and $\tilde{\gamma}(1) = \text{const}_{x_0}$, the constant map at x_0 . This gives us an identification $\text{id}_x^f = \text{const}_{x_0}^f$ in $\mathfrak{f}(X \rightarrow X)$. It remains to show that such an identification implies that $\mathfrak{f}X$ is contractible.

The functorial action of \mathfrak{f} gives a map $(X \rightarrow X) \rightarrow (\mathfrak{f}X \rightarrow \mathfrak{f}X)$, and since the latter is \mathfrak{f} -modal this factors uniquely through $\mathfrak{f}(X \rightarrow X)$. By construction, the map $\mathfrak{f}(X \rightarrow X) \rightarrow (\mathfrak{f}X \rightarrow \mathfrak{f}X)$ sends id_X^f to $\mathfrak{f}\text{id}_X$, which equals $\text{id}_{\mathfrak{f}X}$ by functoriality. Furthermore, $\text{const}_{x_0}^f$ gets sent to $\mathfrak{f}(\text{const}_{x_0}) = \mathfrak{f}(x_0 \circ !)$ where $! : X \rightarrow *$ is the terminal morphism. By functoriality, this equals the composite $\mathfrak{f}X \xrightarrow{!} \mathfrak{f}* \xrightarrow{\mathfrak{f}x_0} \mathfrak{f}X$, which is the constant map at x_0^f . Therefore, the identity of $\mathfrak{f}X$ factors through a constant map, and so $\mathfrak{f}X$ is contractible. \square

Remark 6.11. We can think of the function $\gamma_{(-)}(-) : X \rightarrow (\mathbb{R} \rightarrow X)$ of Lemma 6.10 as a weak form of multiplicative action of \mathbb{R} on X . If we write $t \cdot x \equiv \gamma_x(t)$, then the assumptions $\gamma_x(0) = x_0$ and $\gamma_x(1) = x$ read as $0 \cdot x = x_0$ and $1 \cdot x = x$. Seen this way, Lemma 6.10 shows us that any type with such a multiplicative action of \mathbb{R} — say, a vector space — is \mathfrak{f} -connected.

As a corollary, we find that the projection

$$\{(x, y) : \mathbb{R}^2 \mid xy = 0\} \rightarrow \{x : \mathbb{R}\}$$

is \mathfrak{f} -connected (and is therefore in particular a \mathfrak{f} -fibration). The fiber of this projection over $x : \mathbb{R}$ is $\{y : Y \mid xy = 0\}$, and for every y in the fiber we have the path $t \mapsto ty$ from 0 to y .

Remark 6.12. We shouldn't expect all quasi-fibrations to be \mathfrak{f} -fibrations. The closest analogue of a quasi-fibration in real hoheision would be a map $f : X \rightarrow Y$ such that for every *crisp* $y :: Y$, $\gamma : \mathfrak{f}\text{fib}_f(y) \rightarrow \text{fib}_{\mathfrak{f}f}(y^f)$ is an equivalence. This is strictly weaker than our definition of \mathfrak{f} -fibration; it amounts to the claim that the pullback of f along $(-)_b : bY \rightarrow Y$ is a \mathfrak{f} -fibration.

7. Homotopy Quotients are \mathfrak{f} -Fibrations.

In this section, we show that the quotient map $X \rightarrow X // G$ from a type X to the homotopy quotient $X // G$ of X by an action of the ∞ -group G is a fibration whenever G is *crisp*. If the action is *crisp* and *transitive*, then for any *crisp* point $x :: X$, the map $G \rightarrow X$ given by acting on x is a fibration as well. We will then give two more examples of \mathfrak{f} -fibrations.

Before we prove these things, we should review the definition of ∞ -group and ∞ -group action. These notions can be found in [3], which develops the basic theory of ∞ -groups and proves a stabilization theorem about them.

Definition 7.1. An ∞ -group is a type G identified with the loop space ΩBG of a pointed, 0-connected type BG (called the *delooping* of G). Since singleton types are contractible, the type of ∞ -groups is equivalent to the type of pointed, 0-connected types.

$$\begin{aligned} \infty\text{-Grp} &: \equiv (G : \mathbf{Type}) \times (BG : \mathbf{Type}_*^{>0}) \times (G = \Omega BG) \\ &\simeq \mathbf{Type}_*^{>0}. \end{aligned}$$

For this reason, we will often identify G with ΩBG .

We may think of the elements of BG as G -torsors, and the point $\text{pt}_{BG} : BG$ as G acting on itself. Indeed, for any group G in the axiomatic sense (a set equipped with operations satisfying laws), we may construct its delooping BG as the type of G -torsors, pointed at G .

In particular, for all $x : X$, $\text{Orbit}(x) \simeq G$.

Now we can prove our main theorem for this section.

Theorem 7.7. *Let G be a crisp ∞ -group, and $X^{(-)} : \mathbf{BG} \rightarrow \mathbf{Type}$ an action of G . Then the quotient map $[-] : X \rightarrow X // G$ is a \mathfrak{f} -fibration.*

If furthermore $X^{(-)}$ is crisp, then the classifying map $\text{fst} : X // G \rightarrow \mathbf{BG}$ is a \mathfrak{f} -fibration, and if the action is transitive and $x :: X$, then the map $g \mapsto gx : G \rightarrow X$ is a \mathfrak{f} -fibration.

Proof. Each fact follows quickly from Proposition 7.6 and Theorem 6.1.

Since \mathbf{BG} is 0-connected, the map $x \mapsto [x] := (\text{pt}_{\mathbf{BG}}, x)$ is surjective. Since by Proposition 7.6 the fiber $\text{fib}_{[-]}([x]) \simeq G$ for all $x : X$; in particular for all $(t, y) : X // G$ we have a term of $\|\text{fib}_{[-]}((t, y)) = G\|$. Since G is crisp, we may take the homotopy type of each side to discover (by Theorem 6.1) that $[-] : X \rightarrow X // G$ is a \mathfrak{f} -fibration.

If $X^{(-)}$ is crisp, then so is $X := X^{\text{pt}_{\mathbf{BG}}}$ (since the ∞ -group G , and hence its delooping \mathbf{BG} and its basepoint $\text{pt}_{\mathbf{BG}}$ are assumed crisp). Since \mathbf{BG} is 0-connected, all the fibers of $\text{fst} : X // G \rightarrow \mathbf{BG}$ are merely equivalent to X , and therefore their homotopy types are merely equivalent to its homotopy type. So, by Theorem 6.1, the classifying map $\text{fst} : X // G \rightarrow \mathbf{BG}$ is a \mathfrak{f} -fibration.

Suppose that $x :: X$. If the action is transitive, then for any $y : X$, $\|\text{Stab}(y) = \text{Stab}(x)\|$. Since x is crisp, so is $\text{Stab}(x)$, so by Theorem 6.1 this proves that the map $g \mapsto gx : G \rightarrow X$ (whose fiber over $y : X$ is $\text{Stab}(y)$ by Proposition 7.6) is a \mathfrak{f} -fibration. \square

We can use Theorem 7.7 to give two more examples of \mathfrak{f} -fibrations.

7.1 $\mathbf{SO}(n) \rightarrow \mathbf{SO}(n + 1) \rightarrow \mathbb{S}^n$ We will first construct a delooping $\mathbf{BSO}(n)$ of the special orthogonal group, and then define the action of $\mathbf{SO}(n + 1)$ on the n -sphere as a map $\mathbf{BSO}(n + 1) \rightarrow \mathbf{Type}$ (with $n \geq 1$). We will prove that the fiber of the map $\mathbf{SO}(n + 1) \rightarrow \mathbb{S}^n$ given by acting on the base point has fiber $\mathbf{SO}(n)$. Finally, by Theorem 7.7, we will conclude that the map $\mathbf{SO}(n + 1) \rightarrow \mathbb{S}^n$ is a \mathfrak{f} -fibration.

Definition 7.8. An *orientation* on a normed real n -dimensional vector space V is a unit length element of its exterior power $\Lambda^n V$, equipped with the norm

$$\langle v_1 \wedge \cdots \wedge v_n, w_1 \wedge \cdots \wedge w_n \rangle := \det[\langle v_i, w_j \rangle_V]$$

We define $\mathbf{BSO}(n)$ to be the type of normed real n -dimensional vector spaces V equipped with an orientation that are merely isomorphic to \mathbb{R}^n with its standard norm and orientation. We point $\mathbf{BSO}(n)$ at \mathbb{R}^n with its standard norm and orientation.

We need to justify this definition of $\mathbf{BSO}(n)$.

Lemma 7.9. $\Omega \mathbf{BSO}(n) = \mathbf{SO}(n)$.

Proof. A linear automorphism of \mathbb{R}^n which preserves the norm is given by an orthogonal matrix. If this furthermore preserves the standard orientation on \mathbb{R}^n , that means its n^{th} -exterior power is the identity; but this is given by multiplying by its determinant, so its determinant must be 1. \square

We can now define the action of $\mathbf{SO}(n + 1)$ on the n -sphere \mathbb{S}^n .

Definition 7.10. For $(V, \langle -, - \rangle)$ a normed vector space, let $\mathbb{S}_V := \{v : V \mid \|v\| = 1\}$ be its unit sphere. Note that $\mathbb{S}_{\mathbb{R}^n} \equiv \mathbb{S}^{n-1}$ by definition.

The map $(V, \langle -, - \rangle, \omega) \mapsto \mathbb{S}_V : \mathbf{BSO}(n+1) \rightarrow \mathbf{Type}$ induces the action of $\mathbf{SO}(n+1)$ on \mathbb{S}^n .

Lemma 7.11. The action of $\mathbf{SO}(n+1)$ on \mathbb{S}^n is transitive, and the stabilizer of the basepoint $1 : \mathbb{S}^n$ may be identified with $\mathbf{SO}(n)$.

Proof. For $v : \mathbb{S}^n$, consider v as a unit vector in \mathbb{R}^{n+1} . Then v may be merely extended to a orthonormal basis of \mathbb{R}^{n+1} by the Gram-Schmidt process. The resulting matrix will have determinant either 1 or -1 , but since $\{-1, 1\}$ has decidable equality, we can choose to swap two of these basis vectors to get a special orthogonal matrix that sends $(1, 0, \dots, 0) : \mathbb{S}^n$ to v .

The stabilizer of the basepoint $1 : \mathbb{S}^n$ may be identified with the special orthogonal matrices whose first column has its first entry 1 and all other entries 0. Since the matrix is orthogonal, there can be nothing but 0s in the first row as well. Therefore, the bottom minor given by removing the first row and first column is also special orthogonal, and this gives an identification of the stabilizer with $\mathbf{SO}(n)$. □

Finally, by Theorem 7.7, we may conclude that

$$\mathbf{SO}(n) \rightarrow \mathbf{SO}(n+1) \rightarrow \mathbb{S}^n$$

is a \mathfrak{f} -fibration.

7.2 A \mathfrak{f} -fibration over a 1-type So far we have only seen \mathfrak{f} -fibrations over sets. But with Cohesive HoTT, we can work directly with topological stacks as well. In this example, we will see an example of a \mathfrak{f} -fibration over a 1-type — a stacky version of the real numbers.

Often, a map will fail to be a fibration at a few points because it is ramified there. For example, the map $\mathbb{R} \vee \mathbb{R} \rightarrow \mathbb{R}$ induced by the identity maps

$$\begin{array}{ccc}
 * & \xrightarrow{0} & \mathbb{R} \\
 0 \downarrow & & \downarrow \\
 \mathbb{R} & \longrightarrow & \mathbb{R} \vee \mathbb{R} \\
 & \searrow & \downarrow \text{dashed} \\
 & & \mathbb{R}
 \end{array}
 \begin{array}{l}
 \text{id} \\
 \text{id}
 \end{array}$$

is almost a \mathfrak{f} -fibration (indeed, almost a covering), but it is ramified over 0. However, when such a “ramified fibration” appears as the quotient of a group action, it can be rectified into a \mathfrak{f} -fibration by replacing the base by the homotopy quotient.

In the above example, note that we can also see this map as the quotient

$$\mathbb{R} \vee \mathbb{R} \rightarrow \mathbb{R} \vee \mathbb{R} / C_2$$

of the action of the cyclic group C_2 of order 2 on $\mathbb{R} \vee \mathbb{R}$ given by permuting the factors. The homotopy quotient $\mathbb{R} \vee \mathbb{R} // C_2$ will be a stacky version of the reals where 0 has automorphism group C_2 . Now the fiber over 0 consists of both a point over 0 (of which there is just one), together with an identification of its image with 0, of which there are now two. So the fibers have become locally constant; they are in fact merely equivalent to the group C_2 .

This can be made formal by appealing to the upcoming Theorem 7.7. We will construct the example above.

Definition 7.12. Let \mathbf{BC}_2 be the type of 2-element sets pointed at $\{0, 1\}$, noting that $C_2 = \Omega\mathbf{BC}_2$.

For $T : \mathbf{BC}_2$, let X^T be the cofiber of $(\text{id}, 0) : T \rightarrow T \times \mathbb{R}$. Note that $X := X^{\text{pt}\mathbf{BC}_2}$ may be identified with $\mathbb{R} \vee \mathbb{R}$. This gives the action of C_2 on $\mathbb{R} \vee \mathbb{R}$ by permuting the factors.

Theorem 7.7 then tells us that

$$C_2 \rightarrow \mathbb{R} \vee \mathbb{R} \rightarrow \mathbb{R} \vee \mathbb{R} // C_2$$

is a \mathfrak{f} -fibration. Explicitly $\mathbb{R} \vee \mathbb{R} // C_2$ is the type of pairs $(T : \mathbf{BC}_2) \times X^T$ of 2-element sets T and elements of the cofiber of the inclusion $(\text{id}, 0) : T \rightarrow T \times \mathbb{R}$.

A map can be a “ramified fibration” even if each fiber¹⁵ is the same. An example of this is the Mobius band given by rotating $[-1, 1]$ around a circle with a half turn mapping down onto $[-1, 1]/\text{sgn}$ sending each longitudinal circle to the set of points it intersects in a fixed copy of $[-1, 1]$ in the Mobius band.

Each fiber of this map is a circle, but as one travels from $[1]$ to $[0]$ in $[-1, 1]/\text{sgn}$, the fibers double over. So while each fiber is the same, they do not have a well defined transport along paths as a \mathfrak{f} -fibration would. The trick here is the word “each”; it is true that every fiber is a circle over each *crisp* point of $[-1, 1]/\text{sgn}$, but not over a generic point as Theorem 6.1 requires.

This ramification can be fixed by considering the map to $[-1, 1] // \text{sgn}$, a stacky version of $[0, 1]$ in which 0 has an automorphism group C_2 .

8. The Shape of a Crisp n -Connected Type is n -Connected

One might expect that if X is $\|-\|_n$ -connected, then its homotopy type $\mathfrak{f}X$ would also be $\|-\|_n$ -connected. While we do not know whether this is true in general, we can prove it for crisp types $X :: \mathbf{Type}$. To do this, we need to recall a bit of the theory of *separated* types for a modality from [5].

Definition 8.1. A type X is \diamond -separated if for all $x, y : X$, the type of identifications $x = y$ is \diamond -modal. By Theorem 2.26 of [5], the \diamond -separated types form a modality \diamond' , and we may inductively define

$$\begin{aligned} \diamond^{(0)} &:: \equiv \diamond \\ \diamond^{(n+1)} &:: \equiv \diamond^{(n)'} \end{aligned}$$

We now need to import a few lemmas from [5].

Lemma 8.2. Any \diamond -modal type is $\diamond^{(n)}$ -modal, and the canonical factorization $\diamond^{(n)}X \rightarrow \diamond X$ of the \diamond -unit through the $\diamond^{(n)}$ -unit is a \diamond -unit.

Proof. By hypothesis, the identification types in $\diamond X$ are \diamond -modal, so that $\diamond X$ is \diamond' -modal, and so on. This proves the first statement.

The second statement now follows by Lemma 3.20. □

Lemma 8.3. For any modality \diamond and any pointed type X , there is an equivalence

$$\Omega^n \diamond^{(n)} X \simeq \diamond \Omega^n X$$

¹⁵That is, over each *crisp* point.

Proof. This follows immediately from Proposition 2.27 of [5] by induction. □

Lemma 8.4. Suppose that \diamond is given by localization at a map $A \rightarrow *$. Then $\diamond^{(n)}$ is given by localization at $\Sigma^n A \rightarrow *$.

Proof. This follows immediately from Lemma 2.15 of [5] by induction. □

As a corollary, we find that the n -fold locally discrete modalities $\mathfrak{J}^{(n)}$ are given by localization at $\Sigma^n \mathbb{R} \rightarrow *$. Since \mathbb{R} is inhabited, as a corollary we find that $\mathfrak{J}^{(n)}$ preserves n -connected types.

Lemma 8.5. Suppose that $-1 \leq k \leq n$. If X is k -connected, then $\mathfrak{J}^{(n)} X$ is k -connected.

Proof. This follows immediately from Corollary 3.13 of [5] by induction. In particular, since \mathbb{R} is (-1) -connected, by Theorem 8.2.1 of [13] $\Sigma^n \mathbb{R}$ is $(n - 1)$ -connected and so $(k - 1)$ -connected. Corollary 3.13 of [5] then applies to the map $\Sigma^n \mathbb{R} \rightarrow *$. □

We are now ready to prove that \mathfrak{J} preserves n -connected crisp types.

Theorem 8.6. Let $X :: \mathbf{Type}$ be a crisp, n -connected type for $n \geq -1$. Then the canonical map $\mathfrak{J}^{(n+1)} X \rightarrow \mathfrak{J} X$ induced by factoring the \mathfrak{J} -unit through the $\mathfrak{J}^{(n+1)}$ -unit is an equivalence, and so in particular $\mathfrak{J} X$ is n -connected.

Proof. For $n \equiv -1$, the statement follows tautologically. It remains to show that assuming the statement for n implies $n + 1$. We note here that since \mathbb{N} is crisply discrete, we may assume all natural numbers are crisp.

First, we argue that we may assume that X is crisply pointed. Since X is $(n + 1)$ -connected and $n \geq -1$, in particular $\|X\|$ is contractible and so also $\flat \|X\|$ is contractible. By Corollary 6.7 of [12], $\flat \|X\| \simeq \|\flat X\|$ so that $\|\flat X\|$ is also contractible. Since we are trying to prove that a map is an equivalence, which is a proposition, we may assume that we have a $u : \flat X$, and therefore assume that we have $u \equiv x^\flat$ for a crisp $x :: X$.

Now, assume that $x :: X$ is a crisp point of X and that X is $(n + 1)$ -connected. Then ΩX is a crisp, n -connected type and therefore $\mathfrak{J}^{(n+1)} \Omega X \rightarrow \mathfrak{J} \Omega X$ is an equivalence by hypothesis; in particular $\mathfrak{J}^{(n+1)} \Omega X$ is discrete. Therefore, $\mathfrak{J}^{(n+1)} \Omega X \simeq \Omega \mathfrak{J}^{(n+2)} X$ is discrete. By Lemma 8.5, $\mathfrak{J}^{(n+2)} X$ is $(n + 1)$ -connected and therefore in particular 0-connected; therefore, it is locally crisply discrete. Since it is pointed and 0-connected, it is also equivalent to $\mathbf{BAut}_{\mathfrak{J}^{(n+2)} X}(x^{\mathfrak{J}^{(n+1)}})$ and so by Theorem 5.9, it is discrete. But then the canonical map $\mathfrak{J}^{(n+2)} X \rightarrow \mathfrak{J} X$ is an equivalence by Lemma 8.2. □

Using Theorem 8.6, we can show that the homotopy type of a higher group is a higher group.

Definition 8.7. A k -commutative ∞ -group is a type G identified with $\Omega^{k+1} \mathbf{B}^{k+1} G$ for a pointed, k -connected type $\mathbf{B}^{k+1} G$.¹⁶ A homomorphism of k -commutative ∞ -groups is a pointed map $\mathbf{B}^{k+1} G \rightarrow \mathbf{B}^{k+1} H$.

¹⁶In [3], k -commutative ∞ -groups are called $(k + 1)$ -tuply groupal, but I couldn't bear to subject the reader to such terminology.

Lemma 8.8. The equivalence $\diamond\Omega^{(n)} = \Omega^{(n)}\diamond^{(n)}$ of Lemma 8.3 is natural. Let $f : X \cdot \rightarrow Y$ be a pointed map between pointed types. Then the following square commutes:

$$\begin{array}{ccc} \diamond\Omega^n X & \xrightarrow{\diamond\Omega^n f} & \diamond\Omega^n Y \\ \sim \downarrow & & \downarrow \sim \\ \Omega^n \diamond^{(n)} X & \xrightarrow{\Omega^n \diamond^{(n)} f} & \Omega^n \diamond^{(n)} Y \end{array}$$

Proof. Since $\Omega^n \diamond^{(n)} Y$ is modal, we may check that this commutes on $p : \Omega^n X$. When restricted to $\Omega^n X$, the square becomes Ω^n applied to the $\diamond^{(n)}$ -naturality square, which commutes. \square

Theorem 8.9. Suppose that G is a crisp, k -commutative ∞ -group with $(k + 1)$ -fold delooping $\mathbb{B}^{k+1}G$. Then $\int G$ is a k -commutative ∞ -group with delooping $\int \mathbb{B}^{k+1}G$ and the unit $(-)^{\int} : G \rightarrow \int G$ is a homomorphism.

Proof. By Theorem 8.6, $\int \mathbb{B}^{k+1}G$ is k -connected and may be pointed at $\mathbf{pt}_{\mathbb{B}^{k+1}G}^{\int}$. By the same theorem,

$$\begin{aligned} \Omega^{k+1} \int \mathbb{B}^{k+1}G &\simeq \Omega^{k+1} \int^{(k+1)} \mathbb{B}^{k+1}G \\ &\simeq \int \Omega^{k+1} \mathbb{B}^{k+1}G \\ &\simeq \int G. \end{aligned}$$

By Lemma 8.8 and the fact that the composite $\mathbb{B}^{k+1}G \rightarrow \int^{(k+1)} \mathbb{B}^{k+1}G \xrightarrow{\sim} \int \mathbb{B}^{k+1}G$ is equal to the unit $\mathbb{B}^{k+1}G \rightarrow \int \mathbb{B}^{k+1}G$, this unit deloops the unit $G \rightarrow \int G$, showing that the latter is a k -commutative homomorphism. \square

As a corollary, we can understand the homotopy type of some classifying types.

- Let $\mathbf{BGL}_1(\mathbb{R})$ be the type of 1-dimensional real vector spaces. Since $\int \mathbf{BGL}_1(\mathbb{R}) = \{-1, 1\}$ may be identified with the group of signs, we get find that $\int \mathbf{BGL}_1(\mathbb{R}) = \mathbb{B}\mathbb{Z}/2$. We can call the \int -unit $w_1 : \mathbf{BGL}_1(\mathbb{R}) \rightarrow \mathbb{B}\mathbb{Z}/2$ the *first Stiefel-Whitney class*, since pushing forward by it sends a real line bundle to a first degree cocycle in $\mathbb{Z}/2$ cohomology. Since this is a \int -unit, we see that the first Stiefel-Whitney class is the universal discrete cohomological invariant of a real line bundle.
- Let $\mathbf{BU}(1)$ be the type of 1-dimensional normed complex vector spaces. Since $\int \mathbf{U}(1) = \mathbb{B}\mathbb{Z}$ is a pointed, connected type whose loop space is \mathbb{Z} , we find that $\int \mathbf{BU}(1) = \mathbb{B}^2\mathbb{Z}$. We can call the \int -unit $c_1 : \mathbf{BU}(1) \rightarrow \mathbb{B}^2\mathbb{Z}$ the *first Chern class*, since pushing forward by it sends a Hermitian line bundle to a second degree cocycle in integral cohomology. Since this is a \int -unit, we see that the first Chern class is the universal discrete cohomological invariant of a complex line bundle.

We can now show, with a quick modal argument, that the first Chern class of the Hopf fibration generates $H^2(\mathbb{S}^2; \mathbb{Z})$.

Proposition 8.10. The first Chern class $c_1(h)$ of the Hopf fibration $h : \mathbb{S}^3 \rightarrow \mathbb{S}^2$ generates $H^2(\mathbb{S}^2; \mathbb{Z})$.

Proof. For the purpose of this proof, we make an identification of \mathbb{S}^2 with $\mathbb{C}P^1$ and so take the points of \mathbb{S}^2 to be complex lines in \mathbb{C}^2 . We will show that the \int_2 -unit $\mathbb{S}^2 \rightarrow \int_2 \mathbb{S}^2$ generates $H^2(\mathbb{S}^2; \mathbb{Z})$, and then that $c_1(h)$ factors uniquely through this unit.

Consider the long exact sequence of homotopy groups associated to the Hopf fibration. Since we have calculated (in Lemma 6.3) that $\Omega\mathbb{S}^1 \simeq \mathbb{Z}$, we see that $\pi_2(\mathbb{S}^2) \simeq \pi_1(\mathbb{S}^1) = \mathbb{Z}$. Therefore, \mathbb{S}^2 is a $\mathbb{B}^2\mathbb{Z}$, and the \mathbb{S}^2 -unit $(-)^{\mathbb{S}^2} : \mathbb{S}^2 \rightarrow \mathbb{S}^2$ induces the identity on π_2 and so generates $H^2(\mathbb{S}^2; \mathbb{Z})$.

It remains to show that $c_1(h) : \mathbb{S}^2 \rightarrow \mathbb{B}^2\mathbb{Z}$ is an \mathbb{S}^2 -unit. Let $\chi : \mathbb{S}^2 \rightarrow \mathbb{B}\mathbb{U}(1)$ send a line $\mathcal{L} : \mathbb{S}^2$ in \mathbb{C}^2 to $\{\mathcal{L}\}$, the normed 1-dimensional complex vector space that it is as a subspace of \mathbb{C}^2 . This classifies the Hopf fibration by Lemma 6.8 and because a unitary isomorphism with \mathbb{C} is determined by an element of unit norm:

$$\text{fib}_\chi(\mathbb{C}) \equiv (\mathcal{L} : \mathbb{S}^2) \times (\{\mathcal{L}\} = \mathbb{C}) \simeq (\mathcal{L} : \mathbb{S}^2) \times (\ell : \{\mathcal{L}\}) \times (\|\ell\| = 1) \simeq (\mathcal{L} : \mathbb{S}^2) \times \text{fib}_h(\mathcal{L})$$

In other words, $c_1(h) \equiv c_1 \circ \chi$. Now, the fibers of χ are merely equivalent to \mathbb{S}^3 , and $\mathbb{S}^3 = *$, so it is \mathbb{S}^2 -connected. But c_1 is an \mathbb{S}^2 -unit and so also \mathbb{S}^2 -connected. Therefore, $c_1 \circ \chi$ is a \mathbb{S}^2 -connected map into a \mathbb{S}^2 -modal type; by Lemma 1.38 of [9], it is therefore a \mathbb{S}^2 -unit. □

9. A Bit of Covering Space Theory

In this section, we'll see a bit of modal covering theory and get a sense of how working with coverings using modalities feels. In his *Cohesive Covering Theory* extended abstract [14], Wellen defines a *modal covering* map $\pi : E \rightarrow B$ for a modality \diamond to be a \diamond -étale map. He then specializes to the modality \mathbb{J}_1 to recover the usual covering theory. Here, in light of further conversation with Wellen, we will make a slightly less general definition of covering map which relates more closely to the traditional theory.

Definition 9.1. A map $\pi : E \rightarrow B$ is a *cover* if it is \mathbb{J}_1 -étale and its fibers are sets.

Recall from Section 2 that \diamond -equivalences lift uniquely against \diamond -étale maps. In particular, in any square

$$\begin{array}{ccc} * & \longrightarrow & E \\ 0 \downarrow & & \downarrow \pi \\ \mathbb{R} & \longrightarrow & B \end{array}$$

there is a unique filler since \mathbb{R} is \mathbb{J}_1 -connected. Therefore, covers satisfy the *unique* path lifting property.

We can quickly prove the classical theorem that coverings of a space X correspond to actions of the fundamental groupoid of X on discrete sets.

Theorem 9.2. Let X be a type and let $\text{Cov}(X)$ denote the type of covers of X . Then

$$\text{Cov}(X) \simeq (\mathbb{J}_1 X \rightarrow \mathbf{Type}_{\mathbb{J}_0}).$$

Proof. This follows immediately from Corollary 3.13, applied to the modality \mathbb{J}_1 . This corollary says that \mathbb{J}_1 -étale maps into X correspond to maps from $\mathbb{J}_1 X$ to $\mathbf{Type}_{\mathbb{J}_1}$. If furthermore the fibers are sets, then the maps go from $\mathbb{J}_1 X$ to $\mathbf{Type}_{\mathbb{J}_0}$. □

Classically, the universal cover is just any simply connected cover. We can let this characterization lead us to a definition of the universal cover of a pointed, homotopically connected

space. Let X be a space and $\pi : \tilde{X} \rightarrow X$ a covering with \tilde{X} simply connected in the sense that $\int_1 \tilde{X} = *$. Since π is a covering, and hence \int_1 -étale, the \int_1 -naturality square

$$\begin{array}{ccc} \tilde{X} & \longrightarrow & \int_1 \tilde{X} \\ \pi \downarrow & & \downarrow \int_1 \pi \\ X & \longrightarrow & \int_1 X \end{array}$$

is a pullback. But $\int_1 X = *$, so this shows us that $\tilde{X} = \text{fib}_{(-)\int_1}(u)$ for some $u : \int_1 X$. This leads us to the following definition.

Definition 9.3. Let X be a type and $\text{pt}_X : X$ a base point. Suppose further that X is homotopically connected in the sense that $\|\int_1 X\|_0 = *$. Then the *universal cover* $\pi : \tilde{X} \cdot \rightarrow X$ is defined to be $\text{fst} : \text{fib}_{(-)\int_1}(\text{pt}_X^{\int_1}) \rightarrow X$, with $\text{pt}_{\tilde{X}} \equiv (\text{pt}_X, \text{refl})$ and $\text{pt}_\pi \equiv \text{refl}$:

$$\begin{array}{ccc} \tilde{X} & \longrightarrow & * \\ \pi \downarrow & & \downarrow \text{pt}_X^{\int_1} \\ X & \longrightarrow & \int_1 X \end{array}$$

Theorem 9.4. *The universal cover $\pi : \tilde{X} \rightarrow X$ is the initial pointed cover of X . That is, for any pointed cover $c : C \cdot \rightarrow X$, there is a unique pointed cover $\chi_c : \tilde{X} \cdot \rightarrow C$ such that $c \circ \chi_c = \pi$ as pointed maps.*

Proof. We need to show that the universal cover is a cover with the correct universal property.

First, note that as the fiber of a \int_1 -unit, \tilde{X} is \int_1 -connected (that is, simply connected). Therefore, the naturality square

$$\begin{array}{ccc} \tilde{X} & \longrightarrow & \int_1 \tilde{X} \\ \pi \downarrow & & \downarrow \int_1 \pi \\ X & \longrightarrow & \int_1 X \end{array}$$

is equal to the square

$$\begin{array}{ccc} \tilde{X} & \longrightarrow & * \\ \pi \downarrow & & \downarrow \text{pt}_X^{\int_1} \\ X & \longrightarrow & \int_1 X \end{array}$$

which is a pullback. As the \int_1 -naturality square of π is a pullback, π is \int_1 -étale. The fiber of π over any point $x : X$ is equivalent to $x^{\int_1} = \text{pt}_X^{\int_1}$, which is a type of identifications in the 1-type $\int_1 X$ and is therefore a set. This proves that π is a cover.

Now for the universal property. Note that since $\pi(\text{pt}_{\tilde{X}}) \equiv \text{pt}_X$, the data of a pointed cover $c : C \cdot \rightarrow X$ can be expressed as a square

$$\begin{array}{ccc} * & \xrightarrow{\text{pt}_C} & C \\ \text{pt}_{\tilde{X}} \downarrow & & \downarrow c \\ \tilde{X} & \xrightarrow{\pi} & X \end{array}$$

in which the map c is a cover. A filler of that square is precisely a pointed map $\tilde{X} \rightarrow C$ over X . But \tilde{X} is \int_1 -connected and therefore the map $\text{pt}_{\tilde{X}} : * \rightarrow \tilde{X}$ is an \int_1 -equivalence. And since c is

a \int_1 -étale map and \int_1 -equivalences are orthogonal to \int_1 -étale maps by Lemma 6.1.23 of [8], the type of fillers of this square is contractible.

It remains to show that the unique filler of the square is a cover. Since c and π are \int_1 -étale, it is \int_1 -étale. And since c and π have set fibers, it does as well. Therefore, it is a cover. \square

As promised, Lemma 6.3 does prove that $(\cos, \sin) : \mathbb{R} \rightarrow \mathbb{S}^1$ is the universal cover of the circle. This map is \int_1 -étale, its fibers are sets, and \mathbb{R} is simply connected.

Theorem 6.1 provides us with a simple trick for showing that a map is a cover.

Corollary 9.5. Let $\pi : E \rightarrow B$. If there is a crisply discrete set F such that $\|\text{fib}_\pi(b) = F\|$ for all $b : B$, then π is a cover.

Remark 9.6. As promised in Section 6.2, the map $\mathbb{S}^{n+1} \rightarrow \mathbb{R}P^n$ is a covering map, and since \mathbb{S}^{n+1} is simply connected for $n \geq 0$, this is the universal cover of $\mathbb{R}P^n$.

We can prove a seemingly suspect proposition with this trick: any map with finite fibers is a cover. To do this, we need to prove a bit of folklore.

Lemma 9.7. Let $\text{Fin} := (X : \mathbf{Type}) \times \|(n : \mathbb{N}) \times X = \{1, \dots, n\}\|$ be the type of finite types (types X for which there exists an n such that $X = \{1, \dots, n\}$). There is an equivalence

$$\text{Fin} \simeq (n : \mathbb{N}) \times \mathbf{BAut}(n)$$

between the type of finite types and the sum over $n : \mathbb{N}$ of the classifying types $\mathbf{BAut}(n) := (X : \mathbf{Type}) \times \|X = \{1, \dots, n\}\|$ of the symmetric group $\mathbf{Aut}(n)$.

Proof. Note that

$$\begin{aligned} (n : \mathbb{N}) \times \mathbf{BAut}(n) &\equiv (n : \mathbb{N}) \times (X : \mathbf{Type}) \times \|X = \{1, \dots, n\}\| \\ &\simeq (X : \mathbf{Type}) \times (n : \mathbb{N}) \times \|X = \{1, \dots, n\}\|. \end{aligned}$$

Therefore, it will suffice to show that $(n : \mathbb{N}) \times \|X = \{1, \dots, n\}\| \simeq \|(n : \mathbb{N}) \times X = \{1, \dots, n\}\|$ assuming that $X : \mathbf{Type}$. But the obvious map $(n, |p|) \mapsto |(n, p)|$ is a $\|-\|$ -unit by Lemma 1.24 of [9], so it will suffice to show that $(n : \mathbb{N}) \times \|X = \{1, \dots, n\}\|$ is a proposition.

Suppose that (n, p) and (m, q) are of type $(n : \mathbb{N}) \times \|X = \{1, \dots, n\}\|$, seeking $(n, p) = (m, q)$. From p and q , we get $\|\{1, \dots, n\} = \{1, \dots, m\}\|$. A simple induction shows that this occurs if and only if $n = m$. \square

Proposition 9.8. Let $\pi : E \rightarrow B$ be a map whose fibers are finite in the sense that for every $b : B$, there exists an $n : \mathbb{N}$ such that $\|\text{fib}_\pi(b) = \{1, \dots, n\}\|$. Then π is a cover.

Proof. Note that this condition says that the map $\text{fib}_\pi : B \rightarrow \mathbf{Type}$ factors through $\text{Fin} \hookrightarrow \mathbf{Type}$. But by Lemma 9.7, $\text{Fin} \simeq (n : \mathbb{N}) \times \mathbf{BAut}(n)$, and since \mathbb{N} is crisply discrete, we have an equivalence

$$(n : \mathbb{N}) \times \mathbf{BAut}(n) \simeq (n : \flat \mathbb{N}) \times \text{let } n := m^b \text{ in } \mathbf{BAut}(m).$$

Now, in the inner expression, $m :: \mathbb{N}$ is *crisp*, and so Theorem 5.9 applies and $\mathbf{BAut}(m)$ is discrete. Therefore, Fin is a discretely indexed sum of discrete types, and so it is also discrete. It is, furthermore, a 1-type since it is a set indexed sum of 1-types.

Therefore, fib_b factors through $\int_1 B$ and so by Lemma 3.12, is \int_1 -étale. By hypothesis, its fibers are finite and therefore sets, so it is a cover. \square

Remark 9.9. What is strange about this theorem is that there appear to be counterexamples. Consider the map $\mathbb{R} \vee \mathbb{R} \rightarrow \mathbb{R}$ we looked at in Example 7.2. It seems like its fibers are finite. By a quick application of descent, we can see that its fiber over $r : \mathbb{R}$ is equivalent to the suspension $\Sigma(r = 0)$ of the proposition that $r = 0$. The inclusion of the endpoints of the suspension are always jointly surjective, so there is a *surjection* $\{0, 1\} \rightarrow \Sigma(r = 0)$. But we cannot prove this is a bijection, or that there is a bijection from $\Sigma(r = 0)$ to $\{0\}$ without deciding the proposition $r = 0$. We can't decide whether a real number is 0 (since the reals are connected), so we can't find a precise cardinality for the fiber. This example emphasizes the difference between cardinal finiteness (being equivalent to some $\{1, \dots, n\}$) and Kuratowski finiteness (admitting a surjection from some $\{1, \dots, n\}$) in real cohesion.

Remark 9.10. While the map $\mathbb{R} \vee \mathbb{R} \rightarrow \mathbb{R}$ we considered in Example 7.2 is not a covering, the homotopy quotient $\mathbb{R} \vee \mathbb{R} \rightarrow \mathbb{R} \vee \mathbb{R} // C_2$ is a cover, and is in fact the universal cover of $\mathbb{R} \vee \mathbb{R} // C_2$. To see this, note that $\mathbb{R} \vee \mathbb{R}$ is contractible since it is given as a crisp pushout and \int preserves crisp pushouts. The fibers of the homotopy quotient are merely equivalent to C_2 , which is a discrete set, so the map is a covering. This gives an example of the universal cover of a space which is not a set.

For a particular example of these results, consider an n -fold cover of the circle \mathbb{S}^1 .

Definition 9.11. An n -fold cover $\pi : E \rightarrow B$ is a map whose fibers have n elements. By Corollary 9.5, an n -fold cover is indeed a cover.

Theorem 9.12. Let $n : \mathbb{N}$. The type of n -fold covers of \mathbb{S}^1 whose fiber over $(1, 0)$ is identified with a fixed n -element set $\{1, \dots, n\}$ is equivalent to the type $\mathbf{Aut}(n)$ of permutations of n elements.

Proof. First, we note that since \mathbb{N} is crisply discrete, we may assume without loss of generality that n is crisp and that the fixed n -element set $\{1, \dots, n\}$ is also crisp. The type in question is

$$(f : \mathbb{S}^1 \rightarrow \mathbf{BAut}(n)) \times (f(1, 0) = \{1, \dots, n\})$$

the type of *pointed* maps from the circle to $\mathbf{BAut}(n)$. But Theorem 5.9, $\mathbf{BAut}(n)$ is discrete and so this is equivalent to the type

$$(f : \int \mathbb{S}^1 \rightarrow \mathbf{BAut}(n)) \times (f(1, 0)^\int = \{1, \dots, n\}).$$

By Theorem 9.5 of [12], $(\mathbb{S}^1 \rightarrow X) \simeq (S^1 \rightarrow X)$ for any discrete X , and so the above type is equivalent to

$$(f : S^1 \rightarrow \mathbf{BAut}(n)) \times (f(\mathbf{pt}) = \{1, \dots, n\})$$

which, by the universal proposty of S^1 , is equivalent to $\Omega \mathbf{BAut}(n) \simeq \mathbf{Aut}(n)$. □

Looking at some examples of n -fold coverings (such as Figure 2), we might get the idea that the set of connected components of the total space corresponds to the cycle type of its induced permutation. Somewhat more objectively, we might expect that the set of connected components of the total space should correspond to the set of orbits of the action of the induced permutation on the elements of a fiber. We can prove this using a nice modal argument.

Theorem 9.13. Let $\pi : E \rightarrow B$ be a cover over a pointed base B with fiber F which is connected in the sense that $\int_1 B$ is 0-connected. Then

$$\int_1 E = F // \pi_1(B)$$

where $\pi_1(B) := \Omega(\int_1 B, \mathbf{pt}_B^1)$ is the fundamental group of B .




Figure 2: A 5-fold cover of the circle corresponding to the permutation (12)(354). It has cycle type (2, 3), corresponding to the 2 elements of the fiber in the top connected component, and the 3 elements in the bottom.

Proof. Since $\pi : E \rightarrow B$ is a cover, $\mathbf{fib}_\pi : B \rightarrow \mathbf{Type}$ factors through $\int_1 B$ as $\mathbf{fib}_{\int_1 \pi}$:

$$\begin{array}{ccc}
 B & \xrightarrow{\mathbf{fib}_\pi} & \mathbf{Type} \\
 (-)^{\int_1} \downarrow & \nearrow \mathbf{fib}_{\int_1 \pi} & \\
 \int_1 B & &
 \end{array}$$

witnessed by $\delta : \mathbf{fib}_\pi(b) \xrightarrow{\sim} \mathbf{fib}_{\int_1 \pi}(b^{\int_1})$. Taking total spaces, we find that the following square is a pullback:

$$\begin{array}{ccc}
 E & \xrightarrow{\mathbf{tot}(\delta)} & (t : \int_1 B) \times \mathbf{fib}_{\int_1 \pi}(t) \\
 \pi \downarrow & & \downarrow \mathbf{fst} \\
 B & \xrightarrow{(-)^{\int_1}} & \int_1 B
 \end{array}$$

Since $(-)^{\int_1} : B \rightarrow \int_1 B$ is \int_1 -connected (by Theorem 1.32 of [9]) and \int_1 -connected maps are preserved under pullback (by Theorem 1.34 of [9]), the top map $\mathbf{tot}(\delta)$ is also \int_1 -connected.

Now, since $\int_1 B$ is 0-connected, when pointed at $\mathbf{pt}_B^{\int_1}$ it can be considered as the delooping $\mathbf{B}\pi_1(B)$ of the fundamental group of B . Then, the homotopy quotient $\mathbf{fib}_\pi(\mathbf{pt}_B) // \pi_1(B)$ can be constructed as the pair type

$$F // \pi_1(B) := (t : \int_1 B) \times \mathbf{fib}_{\int_1 \pi}(t).$$

See Section 7 for a brief introduction to the theory of higher groups and Lemma 7.5 for a justification of this construction.

So, the canonical map $E \rightarrow F // \pi_1(B)$ is \int_1 -connected and therefore in particular a \int_1 -equivalence. But as a \int_1 -modally indexed sum of \int_1 -modal types, $\mathbf{fib}_\pi(\mathbf{pt}_B) // \pi_1(B)$ is \int_1 -modal, so we find that $\int_1 E = F // \pi_1(B)$. □

Corollary 9.14. Let $\pi : E \rightarrow \mathbb{S}^1$ be an n -fold covering of the circle whose fiber over $(1, 0)$ is identified with $\{1, \dots, n\}$, and let $\varphi : \mathbf{Aut}(n)$ be the corresponding permutation. Then the set of connected components of the total space E is equivalent to the set of orbits of the action of φ on $\{1, \dots, n\}$.

Proof. The set of connected components of the total space may be constructed as $\|\int_1 E\|_0$, which by Theorem 9.13 is equivalent to $\|\mathbf{fib}_\pi((1, 0)) // \pi_1(\mathbb{S}^1)\|_0$. As we calculated in Theorem 6.4,

$\pi_1(\mathbb{S}^1) = \mathbb{Z}$, and by hypothesis $\text{fib}_\pi((1, 0)) = \{1, \dots, n\}$. So the connected components of E is equivalent to $\|\{1, \dots, n\} // \mathbb{Z}\|_0$ with the action given by $1 \mapsto \varphi$. By Lemma 7.5, two elements of $\|\{1, \dots, n\} // \mathbb{Z}\|_0$ are equal if and only if there is an integer that sends one to the other; in other words, this is the set of orbits of the action of φ , as desired. \square

We can extend the definition of a cover naturally to an “ n -cover” using the modality \int_n .

Definition 9.15. A map $\pi : E \rightarrow B$ is an n -cover if it is \int_n -étale and its fibers are $(n - 1)$ -types.

The theory of n -covers follows just as smoothly as the theory of covers. For every fact above about covers, there is an analogous fact about n -covers proved in the same way. In particular, a universal n -cover is just a \int_n -connected n -cover. We can describe the universal 2-cover of the 2-sphere.

Theorem 9.16. Let $h : \mathbb{S}^3 \rightarrow \mathbb{S}^2$ be the Hopf fibration. Then the \int -modal factor $\text{fst} : (s : \mathbb{S}^2) \times \int \text{fib}_h(s) \rightarrow \mathbb{S}^2$ of the Hopf fibration is the universal 2-cover of the 2-sphere.

Proof. Let $\pi : E \rightarrow \mathbb{S}^2$ denote the \int -modal factor of the Hopf fibration. Note that $\text{fib}_\pi(s) = \int \text{fib}_h(s)$ is merely equivalent to the crisply discrete 1-type $\int \mathbb{S}^1$ for all $s : \mathbb{S}^2$, and is therefore by Theorem 6.1 is \int_2 -étale and so a 2-cover. Furthermore, $\int E \simeq \int \mathbb{S}^3$, so it is \int_2 -connected (since $\int \mathbb{S}^3 = S^3$ is 2-connected), and therefore the universal 2-cover. \square

The theory of n -covers seems related to the theory of Whitehead towers, but the precise relationship between these notions in Cohesive HoTT is not yet clear to the author.

We can show that the universal cover of a crisp ∞ -group is also an ∞ -group. If G is a crisp ∞ -group, then so is $\int_1 G \simeq \|\int G\|_1$ by Theorem 8.9 and so we get a long fiber sequence:

$$\begin{array}{ccccccc}
 & & \dots & \longrightarrow & \pi_1(G) & & \\
 & & & & \searrow & & \\
 & & & & \curvearrowright & & \\
 & & \tilde{G} & \longrightarrow & G & \longrightarrow & \int_1 G \\
 & & & & \searrow & & \\
 & & & & \curvearrowright & & \\
 & & \mathbf{B}\tilde{G} & \longrightarrow & \mathbf{B}G & \longrightarrow & \int_2 \mathbf{B}G
 \end{array}$$

The delooping of \tilde{G} is defined to be the fiber of $(-)^{\int_2} : \mathbf{B}G \rightarrow \int_2 \mathbf{B}G$, and it is 0-connected since the unit $(-)^{\int_1} : G \rightarrow \int_1 G$ is surjective. Note that $\mathbf{B}\tilde{G}$ is the universal 2-cover of $\mathbf{B}G$.

We can continue this fiber sequence on as long as G can be delooped, taking $\int_{k+1} \mathbf{B}^k G$ as the delooping of $\int_k \mathbf{B}^{k-1} G$ and taking $\mathbf{B}^k \tilde{G}$ to be the universal $(k + 1)$ -cover of $\mathbf{B}^k G$. In particular, we get a long fiber sequence:

$$\begin{array}{ccccccc}
 & & \dots & \longrightarrow & \mathbb{Z} & & \\
 & & & & \searrow & & \\
 & & & & \curvearrowright & & \\
 & & \mathbb{R} & \longrightarrow & \mathbf{U}(1) & \longrightarrow & \mathbf{B}\mathbb{Z} \\
 & & & & \searrow & & \\
 & & & & \curvearrowright & & \\
 & & \mathbf{B}\mathbb{R} & \longrightarrow & \mathbf{B}\mathbf{U}(1) & \longrightarrow & \mathbf{B}^2 \mathbb{Z} \\
 & & & & \searrow & & \\
 & & & & \curvearrowright & & \\
 & & \dots & & & &
 \end{array}$$

This gives us a long exact sequence $H^*(-; \mathbb{Z}) \rightarrow H^*(-; \mathbb{R}) \rightarrow H^*(-; \mathbb{U}(1)) \rightarrow H^{*+1}(-; \mathbb{Z})$ in *continuous* cohomology.

In this paper, we have defined a notion of *modal fibration* and explored the fibrations for the *shape* modality of Real Cohesive HoTT. We have seen that it is often quite easy to prove a map is a \mathfrak{J} -fibration — indeed, if you know what the fiber is ahead of time, it is often trivial. After a fibration is found, many simple calculations can be done with purely modal arguments.

Though we only briefly discussed them in this paper, the author hopes that this framework can make calculations in the theory of orbifolds and Lie groupoids more approachable and more conceptual.

Acknowledgements

I would like to thank Felix Wellen for introducing me to the modal covering story, and for many interesting conversations on the topic. I would also like to thank Egbert Rijke for his work on modalities and for a fruitful discussion on modal fibrations. I would also like to thank the reviewer whose helpful comments have improved numerous parts of the exposition and cleaned up the proofs of a few lemmas. And, crucially, I would like to thank Emily Riehl for her helpful comments and guidance during the drafting of this paper.

References

- [1] M. Anel et al. “Goodwillie’s calculus of functors and higher topos theory”. In: *Journal of Topology* 11.4 (2018), 1100–1132. ISSN: 1753-8416. DOI: [10.1112/topo.12082](https://doi.org/10.1112/topo.12082). URL: <http://dx.doi.org/10.1112/topo.12082>.
- [2] Mathieu Anel et al. *A Generalized Blakers-Massey Theorem*. 2017. arXiv: [1703.09050](https://arxiv.org/abs/1703.09050) [[math.AT](#)].
- [3] Ulrik Buchholtz, Floris van Doorn, and Egbert Rijke. *Higher Groups in Homotopy Type Theory*. 2018. arXiv: [1802.04315](https://arxiv.org/abs/1802.04315) [[cs.LG](#)].
- [4] Felix Cherubini and Egbert Rijke. “Modal descent”. In: *Mathematical Structures in Computer Science* (2020), 1–29. ISSN: 1469-8072. DOI: [10.1017/S0960129520000201](https://doi.org/10.1017/S0960129520000201). URL: <http://dx.doi.org/10.1017/S0960129520000201>.
- [5] J. Daniel Christensen et al. “Localization in Homotopy Type Theory”. In: *arXiv e-prints* (2018). arXiv: [1807.04155](https://arxiv.org/abs/1807.04155) [[math.AT](#)].
- [6] Albrecht Dold and Rene Thom. “Quasifaserungen und Unendliche Symmetrische Produkte”. In: *Annals of Mathematics* 67.2 (1958), pp. 239–281. ISSN: 0003486X. URL: <http://www.jstor.org/stable/1970005>.
- [7] Tom Goodwillie. *quasifibrations*. Email to ALGTOP mailing list. May 2001.
- [8] Egbert Rijke. “Classifying Types: Topics in synthetic homotopy theory”. PhD thesis. Carnegie Mellon University, Aug. 2018.
- [9] Egbert Rijke, Michael Shulman, and Bas Spitters. “Modalities in homotopy type theory”. In: *arXiv e-prints* (2017). arXiv: [1706.07526](https://arxiv.org/abs/1706.07526) [[math.CT](#)].
- [10] Urs Schreiber. “Differential cohomology in a cohesive infinity-topos”. In: (2013). arXiv: [1310.7930](https://arxiv.org/abs/1310.7930) [[math-ph](#)].

- [11] Michael Shulman. *All $(\infty, 1)$ -toposes have strict univalent universes*. 2019. arXiv: 1904.07004 [math.AT].
- [12] Michael Shulman. “Brouwer’s fixed-point theorem in real-cohesive homotopy type theory”. In: *Mathematical Structures in Computer Science* 28.6 (2018), 856–941. DOI: 10.1017/S0960129517000147.
- [13] The Univalent Foundations Program. *Homotopy Type Theory: Univalent Foundations of Mathematics*. Institute for Advanced Study: <https://homotopytypetheory.org/book>, 2013.
- [14] Felix Wellen. “Cohesive Covering Theory”. In: *Proceedings of HoTT/UF 2018*. Ed. by Huber Ahrens and Mörtberg. Aug. 2018. URL: https://hott-uf.github.io/2018/abstracts/HoTTUF18_paper_15.pdf.
- [15] Felix Wellen. “Formalizing Cartan Geometry in Modal Homotopy Type Theory”. PhD thesis. Karlsruher Institut für Technologie, July 2017.