

Higher Structures

Arnaud Mayeux
Dilatations of categories


Higher Structures, Vol. 9 (2025), No. 2, 62–75

Persistent URL: <http://dml.cz/dmlcz/153492>

Terms of use:

© Institute of Mathematics, Czech Academy of Sciences, 2025

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library*
<http://dml.cz>

Dilatations of categories

Arnaud Mayeux^a

^a*Einstein Institute of Mathematics, Edmond J. Safra Campus, The Hebrew University of Jerusalem, Givat Ram. Jerusalem, 9190401, Israel*

Abstract

Dilatations modify categories by imposing that some morphisms factorize through some others. This is formalized by a universal property. This text is devoted to introduce and study this construction. Examples of dilatations of categories include localizations of categories and dilatations of rings.

Communicated by: Andrey Lazarev.

Received: July 12th 2023. Accepted: July 22nd, 2025.

MSC: 18A05, 18E35.

Keywords: dilatations of categories, factorizations of morphisms, fractions of morphisms, categories of fractions, localizations of categories, algebraic dilatations, universal property of dilatations.

1. Introduction

Given a subset S of a commutative ring A , one has the localization $S^{-1}A$ of A relatively to S . Needless to insist on the fact that this is a fundamental construction. The localization process of a commutative ring is extended in several ways, among them:

1. localization of categories [7] (cf. also [2, 8]), a basic construction used in many branches of mathematics;
2. dilatation of rings [9, 11], the building blocks of dilatations of schemes [9] (cf. also [4, 5, 10] and the large number of references therein).

In this text, we provide a construction unifying all of these constructions in a single construction: dilatations of categories. Intuitively, localization is a process that adds and imposes inverses of elements or morphisms. In other words, localization adds some "fractions" with prescribed denominators. Intuitively, dilatations do the same thing except that it only adds some fractions where both numerators and denominators are prescribed. Let \mathcal{C} be a category, and let Σ be a collection of morphisms of \mathcal{C} . Let us recall some properties of the localization $\mathcal{C}[\Sigma^{-1}]$. We have a functor $L : \mathcal{C} \rightarrow \mathcal{C}[\Sigma^{-1}]$. The objects of the localization $\mathcal{C}[\Sigma^{-1}]$ coincide with the objects of \mathcal{C} ,

Email address: arnaud.mayeux@mail.huji.ac.il (Mayeux)

© Mayeux, 2025, under a Creative Commons Attribution 4.0 International License.

DOI: [10.21136/HS.2025.10](https://doi.org/10.21136/HS.2025.10)

i.e. L is the identity on objects. Given a morphism d in Σ , $L(d)$ is an isomorphism in $\mathcal{C}[\Sigma^{-1}]$. If $F : \mathcal{C} \rightarrow \mathcal{D}$ is another functor such that $F(d)$ is an isomorphism for all d in Σ , then F factors through L . Now assume that for any d in Σ we have a sieve N_d , in \mathcal{C} over the codomain of d . The dilatation process will provide a category \mathcal{C}' and a functor $\Theta : \mathcal{C} \rightarrow \mathcal{C}'$. The objects of \mathcal{C}' will also coincide with the objects of \mathcal{C} . For any $d \in \Sigma$ and any $n \in N_d$, there exists a unique arrow b in \mathcal{C}' such that the diagram

$$\begin{array}{ccc} X & \xrightarrow{\Theta(n)} & Y \\ & \searrow b & \nearrow \Theta(d) \\ & & Z \end{array}$$

commutes. The element b is thought of as a non-commutative fraction $d \setminus n = d^{-1} \circ n$. Dilatations also satisfy a natural universal property (Theorem 3.10), namely Θ is universal among all functors $F : \mathcal{C} \rightarrow \mathcal{D}$ such that

1. the sieve generated by $F(N_d)$ is included in the sieve generated by $F(d)$, for all morphisms d in the collection Σ ,
2. the localization $\mathcal{D} \rightarrow \mathcal{D}[\Sigma^{-1}]$ is faithful.

Condition (1) allows existence of the factorization morphisms "b" as before and condition (2) allows unicity. To prove the existence of dilatations in this note, we use directed graphs.

We now discuss the structure of this paper. Definition 2.13 and antecedent material provide the definition of dilatations of categories. It holds in a very general framework. Proposition 3.1, Theorem 3.10 and Proposition 3.12 are the main results. Facts 2.15 and 5.1 make connections with localizations of categories and dilatations of rings. We also discuss a related notion in §4, that of codilatations of categories. After that, using the fact that algebraic structures (sets with composition laws and axioms) are in particular categories with a single objects, we introduce dilatations of some non-commutative structures (e.g. monoids). Finally, Fact 5.2 shows that the "natural generalization" of another characterization of dilatations of rings, in terms of localizations, does not hold for categories.

2. Dilatations of categories: definition via fractions

We now fix a category \mathcal{C} with objects $Ob\mathcal{C}$ and morphisms $Mor\mathcal{C}$.

2.1 Sieves Recall that if $X \xrightarrow{a} Y$ belongs to $Mor\mathcal{C}$, $dom(a) := X$ is called the domain and $cod(a) := Y$ the codomain (note that dom and cod also make sense for directed graphs). Recall that a sieve in \mathcal{C} is a collection of morphisms stable by precompositions. Note that a union of sieves is a sieve. A sieve over an object Y is a sieve made of morphisms with codomain Y . Similarly, a cosieve is a collection of morphisms stable by postcompositions.

Definition 2.1. Let E be a collection of morphisms of \mathcal{C} . The collection $e \circ n$ with $n \in Mor\mathcal{C}$, $e \in E$ and $cod(n) = dom(e)$ is a sieve called the sieve generated by E . This sieve is denoted $S_E^{\mathcal{C}}$. If $E = \{d\}$ is a singleton, we put $S_E^{\mathcal{C}} = S_d^{\mathcal{C}}$. Note that if $d = Id_X$, $S_{Id_X}^{\mathcal{C}}$ is the sieve made of all morphisms with codomain X . Similarly, we use the notation $CoS_E^{\mathcal{C}}$ to denote the cosieve generated by E .

2.2 Directed graphs and localizations In this section, we discuss localizations of categories. The content of this subsection is classical and does not claim originality (cf. e.g. [7], [8] and [2]), however we provide a self-contained description of this construction. Let Σ be a collection of morphisms of \mathcal{C} .

Definition 2.2. Let \mathcal{G} be the oriented graph defined as follows. The vertices of \mathcal{G} are equal to the objects of \mathcal{C} . The directed lines of \mathcal{G} are made of

1. for each morphism a of \mathcal{C} , a directed line $dom(a) \xrightarrow{a} cod(a)$ of \mathcal{G} ,
2. for each morphism d in Σ a directed line $cod(d) \xrightarrow{l_d} dom(d)$ of \mathcal{G} (in particular $dom(l_d) = cod(d)$ and $cod(l_d) = dom(d)$).

Definition 2.3. A Σ -sequence of directed lines in \mathcal{G} is a sequence, finite and possibly empty, of directed lines x_1, x_2, \dots, x_n of \mathcal{G} such that $cod(x_i) = dom(x_{i+1})$ ($n \geq 0$ is an integer). By convention, an empty sequence is just an element in the collection $Ob\mathcal{C}$. For a non-empty sequence $s = (x_1, \dots, x_n)$, we define the domain as $dom(s) = dom(x_1)$ and the codomain as $cod(s) = cod(x_n)$. If a sequence s is empty and given by an object X , then we put $dom(s) = cod(s) = X$. Note that Σ -sequences of directed lines with compatible domains and codomains can be composed (by convention, composing a sequence s with an empty sequence e gives s).

Definition 2.4. We say that two Σ -sequences of directed lines s and s' in \mathcal{G} are equivalent if $dom(s) = dom(s')$, $cod(s) = cod(s')$ and if one can be obtained from the other by a chain of elementary equivalences of the following types:

1. a sequence x_1, \dots, x_n such that x_i, x_{i+1} are equal to $a, a' \in Mor\mathcal{C}$ and $cod(a) = dom(a')$, for some $1 \leq i, i+1 \leq n$, is equivalent to the sequence $x_1, \dots, x_{i-1}, x', x_{i+2}, \dots, x_n$ where x' is the composition of a and a' in \mathcal{C} ,
2. a sequence x_1, \dots, x_n such that x_i, x_{i+1} are equal to l_d, d with $d \in \Sigma$, for some $1 \leq i, i+1 \leq n$, is equivalent to the sequence $x_1, \dots, x_{i-1}, x_{i+2}, \dots, x_n$,
3. a sequence x_1, \dots, x_n such that x_i, x_{i+1} are equal to d, l_d with $d \in \Sigma$, for some $1 \leq i, i+1 \leq n$, is equivalent to the sequence $x_1, \dots, x_{i-1}, x_{i+2}, \dots, x_n$,
4. for any object X , the sequence Id_X is equivalent to the empty sequence at X .

In other words and informally, two Σ -sequences of directed lines are equivalent if one can be obtained from the other by the operations consisting in exchanging parts of sequences as follows:

$$\begin{aligned} X &\xrightarrow{a} Y \xrightarrow{a'} Z \rightsquigarrow X \xrightarrow{a' \circ a} Z \\ Y &\xrightarrow{l_d} Z \xrightarrow{d} Y \rightsquigarrow Y \text{ (the empty sequence at } Y) \\ Z &\xrightarrow{d} Y \xrightarrow{l_d} Z \rightsquigarrow Z \text{ (the empty sequence at } Z) \\ X &\xrightarrow{Id_X} X \rightsquigarrow X \text{ (the empty sequence at } X). \end{aligned}$$

The equivalence class of a sequence s is denoted by $[s]$. Note that equivalence classes of Σ -sequences of directed lines with compatible domains and codomains can be composed associatively.

Definition 2.5. A Σ -fraction is an equivalence class of Σ -sequence.

Fact 2.6. Let d, d' in Σ such that $cod(d) = dom(d')$ and let d'' be their composite. Assume that d'' belongs to Σ . Then

$$[cod(d') \xrightarrow{l_{d'}} dom(d') \xrightarrow{l_d} dom(d)] = [cod(d') \xrightarrow{l_{d''}} dom(d)].$$

Proof. This follows from the equalities

$$\begin{aligned}
&= [\text{cod}(d') \xrightarrow{l_{d'}} \text{dom}(d') \xrightarrow{l_d} \text{dom}(d)] \\
&= [\text{cod}(d') \xrightarrow{l_{d'}} \text{dom}(d') \xrightarrow{l_d} \text{dom}(d) \xrightarrow{d''} \text{cod}(d') \xrightarrow{l_{d''}} \text{dom}(d)] \\
&= [\text{cod}(d') \xrightarrow{l_{d'}} \text{dom}(d') \xrightarrow{l_d} \text{dom}(d) \xrightarrow{d} \text{cod}(d) \xrightarrow{d'} \text{cod}(d') \xrightarrow{l_{d''}} \text{dom}(d)] \\
&= [\text{cod}(d') \xrightarrow{l_{d''}} \text{dom}(d)].
\end{aligned}$$

□

Definition 2.7. The localization of \mathcal{C} relatively to Σ is the category $\mathcal{C}[\Sigma^{-1}]$ whose objects are the objects of \mathcal{C} and whose morphisms are Σ -fractions. We have a canonical functor $L : \mathcal{C} \rightarrow \mathcal{C}[\Sigma^{-1}]$.

Proposition 2.8 (Universal property of localization). *Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor such that $F(d)$ is invertible for all d in Σ . Then there exists a unique functor $F' : \mathcal{C}[\Sigma^{-1}] \rightarrow \mathcal{D}$ such that $F = F' \circ L$.*

Proof. We first prove unicity. Assume that F' exists. Let X be in $\text{Ob}\mathcal{C} = \text{Ob}\mathcal{C}'$, then $F'(X) = F'(L(X)) = F(X)$. Let a be in $\text{Mor}\mathcal{C}$, then $F'([a]) = F'(L(a)) = F(a)$. Let d be in Σ , then $F'([l_d]) \circ F(d) = F'([l_d] \circ [d]) = \text{Id}_{\text{dom}(F(d))}$ and $F(d) \circ F'([l_d]) = F'([d] \circ [l_d]) = \text{Id}_{\text{cod}(F(d))}$. This implies that $F'([l_d]) = F(d)^{-1}$. This proves unicity. To prove existence, it is enough to prove that the assignments $X \mapsto X, [a] \mapsto F(a), l_d \mapsto F(d)^{-1}$ (as before) provide a well-defined functor F' . For this it is enough to prove that the assignments are compatible with the elementary equivalences of Definition 2.4, which is immediate. □

2.3 Dilatations of categories: definition

Let \mathcal{C} be a category.

Definition 2.9. A center in \mathcal{C} is a collection $\{[N_i, d_i]\}_{i \in I}$ of pairs $[N_i, d_i]$, indexed by a collection I and such that, for all i in I , d_i is a morphism of \mathcal{C} and N_i is a sieve over $\text{cod}(d_i)$.

We now fix a center $\{[N_i, d_i]\}_{i \in I}$ in \mathcal{C} (we sometimes use the notation N_{d_i} to denote N_i).

Put $\Sigma = \{d_i\}_{i \in I}$.

Definition 2.10. A $\{[N_i, d_i]\}_{i \in I}$ -fraction is a Σ -fraction such that a representative can be written as

$$X_1 \xrightarrow{n_1} Y_1 \xrightarrow{l_{d_{i_1}}} X_2 \xrightarrow{n_2} Y_2 \xrightarrow{l_{d_{i_2}}} X_3 \dots X_k \xrightarrow{n_k} Y_k \xrightarrow{l_{d_{i_k}}} X_{k+1} \xrightarrow{a} X_{k+2}$$

with $a \in \text{Mor}\mathcal{C}$, $k \geq 0$ and $i_j \in I$, $n_i \in N_{i_j}$ for all $j \in \{1, \dots, k\}$.

Fact 2.11. $\{[N_i, d_i]\}_{i \in I}$ -fractions with compatible domains and codomains can be composed associatively.

Proof. This is immediate since the N'_i 's are sieves. Associativity is immediate. □

Remark 2.12. 1. If $X \xrightarrow{a} Y$ is a morphism in \mathcal{C} , the $\{d_i\}_{i \in I}$ -fraction $[X \xrightarrow{a} Y]$ is a $\{[N_i, d_i]\}_{i \in I}$ -fraction. In particular, the class $[\text{Id}_X]$ of the identity at an object X is a $\{[N_i, d_i]\}_{i \in I}$ -fraction.

2. The Σ -fraction

$$X_1 \xrightarrow{n_1} Y_1 \xrightarrow{l_{d_{i_1}}} X_2 \xrightarrow{n_2} Y_2 \xrightarrow{l_{d_{i_2}}} X_3 \dots X_k \xrightarrow{n_k} Y_k \xrightarrow{l_{d_{i_k}}} X_{k+1}$$

is a $\{[N_i, d_i]\}_{i \in I}$ -fraction since it is equivalent to

$$X_1 \xrightarrow{n_1} Y_1 \xrightarrow{l_{d_{i_1}}} X_2 \xrightarrow{n_2} Y_2 \xrightarrow{l_{d_{i_2}}} X_3 \dots X_k \xrightarrow{n_k} Y_k \xrightarrow{l_{d_{i_k}}} X_{k+1} \xrightarrow{Id_{X_{k+1}}} X_{k+1}.$$

Definition 2.13. The dilatation of \mathcal{C} with center $\{[N_i, d_i]\}_{i \in I}$ is the category \mathcal{C}' defined as follows. The objects of \mathcal{C}' are equal to the objects of \mathcal{C} . If X and Y are objects in \mathcal{C}' , morphisms between X and Y are given by $\{[N_i, d_i]\}_{i \in I}$ -fractions (cf. Definition 2.10) with domain X and codomain Y . Fact 2.11 shows that \mathcal{C}' is indeed a category, which is also denoted $\mathcal{C} \left[\{(d_i)^{-1} \circ N_i\}_{i \in I} \right]$.

Fact 2.14. We have a faithful functor $\mathcal{C} \left[\{(d_i)^{-1} \circ N_i\}_{i \in I} \right] \rightarrow \mathcal{C} \left[\{(d_i)_{i \in I}\}^{-1} \right]$.

Proof. The functor is obtained regarding $\{[N_i, d_i]\}_{i \in I}$ -fractions as Σ -fractions. \square

Fact 2.15. Assume that, for all i in I , N_i is the sieve $S_{Id_{\text{cod}(d_i)}}^{\mathcal{C}}$, then $\mathcal{C} \left[\{(d_i)^{-1} \circ N_i\}_{i \in I} \right]$ is the localization $\mathcal{C}[\{(d_i)_{i \in I}\}^{-1}]$.

Proof. In this case, the faithful functor of Fact 2.14 is also full, as any $\{(d_i)_{i \in I}\}$ -fraction is equivalent to a $\{[S_{Id_{\text{cod}(d_i)}}^{\mathcal{C}}, d_i]\}_{i \in I}$ -fraction. \square

Corollary 2.16. Assume that \mathcal{C} is small, then $\mathcal{C} \left[\{(d_i)^{-1} \circ N_i\}_{i \in I} \right]$ is small.

Proof. Combine Fact 2.14 and [2, Proposition 5.2.2]. \square

3. Universal property of dilatations and related results

We proceed with the notation from §2.3. Recall that $\mathcal{C}' = \mathcal{C} \left[\{(d_i)^{-1} \circ N_i\}_{i \in I} \right]$.

Proposition 3.1. The following assertions hold.

1. We have a canonical functor $\Theta : \mathcal{C} \rightarrow \mathcal{C}'$ defined as follows. If X is an object of \mathcal{C} , $\Theta(X) = X$. If $X \xrightarrow{a} Y$ is a morphism of \mathcal{C} , $\Theta(a) = [X \xrightarrow{a} Y]$.
2. For any i in I and any $n \in N_i$, there exists a unique morphism $\text{dom}(n) \xrightarrow{b} \text{dom}(d_i)$

in \mathcal{C}' such that the following triangle commutes

$$\begin{array}{ccc} \text{dom}(n) & \xrightarrow{[n]} & \text{cod}(n) \\ & \searrow b & \nearrow [d_i] \\ & & \text{dom}(d_i) \end{array} .$$

Intuitively, the morphism b is thought of as a non-commutative fraction $d_i \setminus n = (d_i)^{-1} \circ n$ of morphisms. Mathematically, we have

$$b = [X \xrightarrow{n} Y \xrightarrow{l_{d_i}} Z].$$

Proof. 1. Obvious.

2. Put $X = \text{dom}(n)$, $Y = \text{cod}(n)$ and $Z = \text{dom}(d_i)$. The class $[X \xrightarrow{n} Y \xrightarrow{l_{d_i}} Z]$ composed with the class $[Z \xrightarrow{d_i} Y]$ equals the class $[n]$. This shows that there exists an arrow b in \mathcal{C}' such that the triangle commutes. We now prove unicity. Assume that $m \in \text{Mor}\mathcal{C}'$ satisfies that the composition $X \xrightarrow{m} Z \xrightarrow{\Theta(d_i)} Y$ equals $X \xrightarrow{\Theta(n)} Y$. Let s be a representative sequence of m . Then the class $[X \xrightarrow{s} Z \xrightarrow{d_i} Y \xrightarrow{l_{d_i}} Z]$ is equal to the class $[X \xrightarrow{n} Y \xrightarrow{l_{d_i}} Z]$ and also to the class $[X \xrightarrow{s} Z]$. This shows that m equals $[X \xrightarrow{n} Y \xrightarrow{l_{d_i}} Z]$. \square

Recall that a bimorphism is defined as a morphism which is both a monomorphism and an epimorphism. Bimorphisms are also called regular morphisms.

Fact 3.2. *A morphism whose image under a faithful functor is a bimorphism is itself a bimorphism.*

Proof. Let $F : \mathcal{A} \rightarrow \mathcal{B}$ be a faithful functor. Let f be a morphism of \mathcal{A} such that $F(f)$ is a bimorphism. Let a, b, c, d be morphisms of \mathcal{A} such that $a \circ f = b \circ f$ and $f \circ c = f \circ d$. Then $F(a) \circ F(f) = F(b) \circ F(f)$ and $F(f) \circ F(c) = F(f) \circ F(d)$. So $F(a) = F(b)$ and $F(c) = F(d)$. Consequently $a = b$ and $c = d$. This proves that f is a bimorphism. \square

Proposition 3.3. *Let $i \in I$, then $\Theta(d_i)$ is a bimorphism in \mathcal{C}' .*

Proof. A morphism whose image under a faithful functor is a bimorphism is itself a bimorphism by Fact 3.2. Now Proposition 3.3 follows from Fact 2.14. \square

Definition 3.4. For any $i \in I$, let $S_{\Theta(N_i)}^{\mathcal{C}'}$ be the sieve over $\text{cod}(\Theta(d_i))$ in \mathcal{C}' generated by $\{\Theta(n) | n \in N_i\}$. Similarly, let $S_{\Theta(d_i)}^{\mathcal{C}'}$ be the sieve over $\text{cod}(\Theta(d_i))$ generated by $\Theta(d_i)$.

Proposition 3.5. *Let $i \in I$, then $S_{\Theta(N_i)}^{\mathcal{C}'} \subset S_{\Theta(d_i)}^{\mathcal{C}'}$.*

Proof. Let x be in $S_{\Theta(N_i)}^{\mathcal{C}'}$. Then $x = [n] \circ t$ with $n \in N_i$. We have

$$x = [n] \circ t = [d_i] \circ [dom(n) \xrightarrow{n} cod(n) \xrightarrow{l_{d_i}} dom(d_i)] \circ t.$$

This finishes the proof. \square

Definition 3.6. Let $\text{Cat}_{\mathcal{C}}^{\Sigma\text{-reg}}$ be the full subcategory of the comma category \mathcal{C}/Cat whose objects $F : \mathcal{C} \rightarrow \mathcal{D}$ are arbitrary functors out of \mathcal{C} such that the localization functor $\mathcal{D} \rightarrow \mathcal{D}[F(\Sigma)^{-1}]$ is faithful.

Fact 3.7. *The functor $\Theta : \mathcal{C} \rightarrow \mathcal{C} \left[\{(d_i)^{-1} \circ N_i\}_{i \in I} \right]$ is an object in the category $\text{Cat}_{\mathcal{C}}^{\Sigma\text{-reg}}$.*

Proof. Fact 2.14 says that $\mathcal{C} \left[\{(d_i)^{-1} \circ N_i\}_{i \in I} \right] \rightarrow \mathcal{C}[\{(d_i)_{i \in I}\}^{-1}]$ is faithful. Now we observe that $\mathcal{C} \left[\{(d_i)^{-1} \circ N_i\}_{i \in I} \right] \left[\{\Theta(d_i)\}_{i \in I}\right]^{-1}$ identifies with the localization $\mathcal{C}[\{(d_i)_{i \in I}\}^{-1}]$. This finishes the proof. \square

Fact 3.8. *Let $\Sigma' \subset \Sigma$ be a subcollection. If $L : \mathcal{C} \rightarrow \mathcal{C}[\Sigma^{-1}]$ is faithful, then $L' : \mathcal{C} \rightarrow \mathcal{C}[\Sigma'^{-1}]$ is also faithful.*

Proof. Observe that $\mathcal{C}[\Sigma^{-1}] = \mathcal{C}[\Sigma'^{-1}][L'(\Sigma)^{-1}]$, so that we have a commutative triangle of functors

$$\begin{array}{ccc} \mathcal{C} & & \\ \downarrow L' & \searrow L & \\ \mathcal{C}[\Sigma'^{-1}] & \xrightarrow{l} & \mathcal{C}[\Sigma^{-1}] \end{array} .$$

Now let a, b be two morphisms such that $L'(a) = L'(b)$. We obtain $l(L'(a)) = l(L'(b))$ and $a = b$. So L' is faithful. \square

Remark 3.9. We remark that [3, Lemma 4.4] shows that if \mathcal{C} is semi-abelian and integral (cf. [3] and references therein for precise definitions) and if B is the collection of all bimorphisms of \mathcal{C} , then the identity functor $\mathcal{C} \rightarrow \mathcal{C}$ itself belongs to $Cat_{\mathcal{C}}^{B\text{-reg}}$. Still in the above setting of [3], [3, Lemma 4.4] together with Fact 3.8 implies that the identity functor $\mathcal{C} \rightarrow \mathcal{C}$ itself belongs to $Cat_{\mathcal{C}}^{\Sigma\text{-reg}}$ for any subcollection Σ of B . Note that we do not use [3, Lemma 4.4] in the present paper.

Theorem 3.10. (*Universal property*) Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be an object in $Cat_{\mathcal{C}}^{\Sigma\text{-reg}}$ such that for any i in I , we have

$$S_{F(N_i)}^{\mathcal{D}} \subset S_{F(d_i)}^{\mathcal{D}} .$$

Then there is a unique functor $F' : \mathcal{C}' \rightarrow \mathcal{D}$ such that the triangle of functors

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{D} \\ \searrow \Theta & & \nearrow F' \\ & \mathcal{C}' & \end{array}$$

commutes (recall that $\mathcal{C}' = \mathcal{C} \left[\{(d_i)^{-1} \circ N_i\}_{i \in I} \right]$).

Proof. Assume that such a F' exists, then $F'(X) = F'(\Theta(X)) = F(X)$ for all X in $Ob\mathcal{C} = Ob\mathcal{C}'$. Now any morphism in \mathcal{C}' is a composition of morphisms of the form $[X \xrightarrow{n} Y \xrightarrow{l_{d_i}} Z]$ with $i \in I, n \in N_i$, or of the form $[dom(a) \xrightarrow{a} cod(a)]$. Necessarily we have $F'([X \xrightarrow{n} Y \xrightarrow{l_{d_i}} Z]) = t$ where

$$\begin{array}{ccc} F(X) & \xrightarrow{F(n)} & F(Y) \\ & \searrow t & \nearrow F(d_i) \\ & & F(Z) \end{array}$$

t is the unique morphism such that commutes (this t exists

since $F(n)$ belongs to $S_{F(d_i)}^{\mathcal{D}}$ by assumption and is unique because $\mathcal{D} \rightarrow \mathcal{D}[F(\Sigma)^{-1}]$ is faithful and so $F(d_i)$ is a bimorphism by Fact 3.2). Necessarily we have $F'([a]) = F'(\Theta(a)) = F(a)$. This shows that F' is unique and given by the formula above if it exists. Reciprocally, the assignment $X \mapsto F(X), [a] \mapsto F(a), [X \xrightarrow{n} Y \xrightarrow{l_{d_i}} Z] \mapsto t$ (the unique t as before), is a well-defined functor F' satisfying that $F = F' \circ \Theta$. \square

Fact 3.11. Let $H : \mathcal{A} \rightarrow \mathcal{B}$ be a functor between two categories. Let E be a collection of morphisms of \mathcal{A} . Then $S_{H(E)}^{\mathcal{B}} = S_{H(S_E^{\mathcal{A}})}^{\mathcal{B}}$.

Proof. We have $E \subset S_E^{\mathcal{A}}$, so $S_{H(E)}^{\mathcal{B}} \subset S_{H(S_E^{\mathcal{A}})}^{\mathcal{B}}$. Reciprocally, let x be in $S_{H(S_E^{\mathcal{A}})}^{\mathcal{B}}$, then there exists $b \in Mor\mathcal{B}, a \in Mor\mathcal{A}$ and $e \in E$, such that $x = H(e \circ a) \circ b$. Since H is a functor, we have $x = H(e) \circ (H(a) \circ b)$. So x belongs to $S_{H(E)}^{\mathcal{B}}$. \square

Proposition 3.12. *The functor $\Theta : \mathcal{C} \rightarrow \mathcal{C} \left[\{(d_i)^{-1} \circ N_i\}_{i \in I} \right]$ represents the covariant functor $Cat_{\mathcal{C}}^{\Sigma\text{-reg}} \rightarrow Set$ given by*

$$(\mathcal{C} \xrightarrow{F} \mathcal{D}) \mapsto \begin{cases} \{*\}, & \text{if } S_{F(N_i)}^{\mathcal{D}} \subset S_{F(d_i)}^{\mathcal{D}} \text{ for any } i \text{ in } I; \\ \emptyset, & \text{else.} \end{cases} \quad (3.1)$$

Proof. Let $(\mathcal{C} \xrightarrow{F} \mathcal{D})$ be an object of $Cat_{\mathcal{C}}^{\Sigma\text{-reg}}$. If $F' \in Hom_{\mathcal{C}}(\mathcal{C}', \mathcal{D})$, then for all i in I we have

$$S_{F(N_i)}^{\mathcal{D}} = S_{F'(\Theta(N_i))}^{\mathcal{D}} = S_{F'(S_{\Theta(N_i)}^{\mathcal{C}'})}^{\mathcal{D}} \subset S_{F'(S_{\Theta(d_i)}^{\mathcal{C}'})}^{\mathcal{D}} = S_{F'(\Theta(d_i))}^{\mathcal{D}} = S_{F(d_i)}^{\mathcal{D}}.$$

We used Proposition 3.5 and Fact 3.11. Therefore, if there exists i such that $S_{F(N_i)}^{\mathcal{D}} \not\subset S_{F(d_i)}^{\mathcal{D}}$, then $Hom_{\mathcal{C}}(\mathcal{C}', \mathcal{D}) = \emptyset$. If $S_{F(N_i)}^{\mathcal{D}} \subset S_{F(d_i)}^{\mathcal{D}}$ for all i in I , Theorem 3.10 implies that $Hom_{\mathcal{C}}(\mathcal{C}', \mathcal{D})$ is a singleton $\{*\}$. This finishes the proof. \square

Fact 3.13. *For each i in I , let M_i be a subsieve of N_i . Then we have a canonical functor*

$$\varphi : \mathcal{C} \left[\{(d_i)^{-1} \circ M_i\}_{i \in I} \right] \rightarrow \mathcal{C} \left[\{(d_i)^{-1} \circ N_i\}_{i \in I} \right].$$

The functor φ is faithful.

Proof. Any $\{[M_i, d_i]\}_{i \in I}$ -fraction is a $\{[N_i, d_i]\}_{i \in I}$ -fraction since M_i is a subsieve of N_i for all i . \square

Proposition 3.14. *Let $K \subset I$ be a subcollection. Then we have a canonical functor*

$$\Phi : \mathcal{C} \left[\{(d_i)^{-1} \circ N_i\}_{i \in K} \right] \rightarrow \mathcal{C} \left[\{(d_i)^{-1} \circ N_i\}_{i \in I} \right].$$

Moreover

1. *if $N_i = S_{d_i}^{\mathcal{C}}$ for all i in $I \setminus K$, then Φ is full,*
2. *if $\mathcal{C}[(\{d_i\}_{i \in K})^{-1}] \rightarrow \mathcal{C}[(\{d_i\}_{i \in I})^{-1}]$ is faithful, then Φ is faithful.*

Proof. Put $\Gamma = \{d_i\}_{i \in K}$. We have a natural application from Γ -sequences to Σ -sequences. This induces an application from Γ -fractions to Σ -fractions. This provides Φ by restriction. To prove (i), we observe that for each piece of Σ -sequence of the kind $l_{d_i} \circ n$ with $i \in I \setminus K$ and $n \in N_i$, by assumption $n = d_i \circ f$ for some f in \mathcal{C} , so that l_{d_i} disappears and the remaining sequence is a Γ -sequence. To prove (ii), consider the commutative diagram of functors

$$\begin{array}{ccc} \mathcal{C} \left[\{(d_i)^{-1} \circ N_i\}_{i \in K} \right] & \longrightarrow & \mathcal{C} \left[\{(d_i)^{-1} \circ N_i\}_{i \in I} \right] \\ \downarrow & & \downarrow \\ \mathcal{C}[\Gamma^{-1}] & \longleftarrow & \mathcal{C}[\Sigma^{-1}] \end{array}.$$

The vertical arrows are faithful by Fact 2.14. The lower arrow is faithful by assumption. This shows that the upper arrow is faithful and finishes (ii). \square

Proposition 3.15. *Let $\{[N_j, d_j]\}_{j \in J}$ be another center in \mathcal{C} . Put $I' = I \sqcup J$. Proposition 3.1 provides functors*

$$\Theta : \mathcal{C} \rightarrow \mathcal{C} \left[\{(d_i)^{-1} \circ N_i\}_{i \in I'} \right]$$

and

$$\Theta' : \mathcal{C} \rightarrow \mathcal{C} [\{(d_i)^{-1} \circ N_i\}_{i \in I'}].$$

We observe that $\left\{ \left[S_{\Theta(N_j)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]} , \Theta(d_j) \right] \right\}_{j \in J}$ is a center in $\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]$, so that we get a dilatation functor

$$\beta : \mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}] \rightarrow \mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}] \left[\left\{ (\Theta(d_j))^{-1} \circ S_{\Theta(N_j)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]} \right\}_{j \in J} \right].$$

We observe that Proposition 3.14 provides a canonical functor

$$\Phi : \mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}] \rightarrow \mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I'}].$$

1. The functor Φ belongs to $\text{Cat}_{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]}^{\{\Theta(d_j)\}_{j \in J}\text{-reg}}$.
2. The functor $\beta \circ \Theta$ belongs to $\text{Cat}_{\mathcal{C}}^{\{d_i\}_{i \in I'}\text{-reg}}$.
3. There exists a unique functor

$$\alpha : \mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I'}] \rightarrow \mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}] \left[\left\{ (\Theta(d_j))^{-1} \circ S_{\Theta(N_j)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]} \right\}_{j \in J} \right]$$

such that $\beta \circ \Theta = \alpha \circ \Theta'$.

4. There exists a unique functor

$$\alpha' : \mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}] \left[\left\{ (\Theta(d_j))^{-1} \circ S_{\Theta(N_j)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]} \right\}_{j \in J} \right] \rightarrow \mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I'}]$$

such that $\Phi = \alpha' \circ \beta$.

5. The functors $\alpha \circ \alpha'$ and $\alpha' \circ \alpha$ are identity functors.
6. There exists an isomorphism of categories

$$\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}] \left[\left\{ (\Theta(d_j))^{-1} \circ S_{\Theta(N_j)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]} \right\}_{j \in J} \right] \xrightarrow{\sim} \mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I'}].$$

Proof. 1. We have to prove that $\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I'}] \rightarrow \mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I'}][\{(\Phi \circ \Theta(d_j))\}_{j \in J}^{-1}]$ is faithful. It is enough to observe that $\Phi \circ \Theta = \Phi'$ and apply Fact 3.7 and Fact 3.8.

2. Applying Fact 2.14 twice, we get a faithful functor

$$\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I'}][\{(\Theta(d_j))^{-1} \circ S_{\Theta(N_j)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]} \}_{j \in J}] \rightarrow \mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I'}][\{[d_j]_{j \in J}\}^{-1}]. \quad (3.2)$$

Now we observe that

$$\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I'}][\{[d_j]_{j \in J}\}^{-1}] = \mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I'}] \quad (3.3)$$

and that

$$\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I'}][\{(\Theta(d_j))^{-1} \circ S_{\Theta(N_j)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]} \}_{j \in J}][\{(\beta \circ \Theta)(d_i)\}_{i \in I'}^{-1}] = \mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I'}]^{-1}. \quad (3.4)$$

Now (3.2), (3.3) and (3.4) together implies that the functor $\beta \circ \Theta$ belongs to $\text{Cat}_{\mathcal{C}}^{\{d_i\}_{i \in I'}\text{-reg}}$.

3. By Theorem 3.10, it is enough to prove that, for all k in I' :

$$S_{(\beta \circ \Theta)(N_k)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I'}][\{(\Theta(d_j))^{-1} \circ S_{\Theta(N_j)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]} \}_{j \in J}]} \subset S_{(\beta \circ \Theta)(d_k)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I'}][\{(\Theta(d_j))^{-1} \circ S_{\Theta(N_j)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]} \}_{j \in J}]} \quad (3.5)$$

Let us first take k in I . By Proposition 3.5, we have

$$S_{\Theta(N_k)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]} \subset S_{\Theta(d_k)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]}$$

therefore we get

$$S_{\beta\left(S_{\Theta(N_k)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]}\right)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}][\{(\Theta(d_j))^{-1} \circ S_{\Theta(N_j)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]}]\}_{j \in J}} \subset S_{\beta\left(S_{\Theta(d_k)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]}\right)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}][\{(\Theta(d_j))^{-1} \circ S_{\Theta(N_j)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]}]\}_{j \in J}}. \quad (3.6)$$

Now (3.6) and Fact 3.11 implies (3.5) for all $k \in I$. Now let $k \in J$. By Proposition 3.5, we have

$$S_{\beta\left(S_{\Theta(N_k)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]}\right)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}][\{(\Theta(d_j))^{-1} \circ S_{\Theta(N_j)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]}]\}_{j \in J}} \subset S_{\beta\left(S_{\Theta(d_k)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]}\right)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}][\{(\Theta(d_j))^{-1} \circ S_{\Theta(N_j)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]}]\}_{j \in J}}. \quad (3.7)$$

Now (3.7) and Fact 3.11 implies (3.5) for all $k \in J$. Finally, (3.5) holds for all $k \in I'$.

4. By Theorem 3.10, it is enough to prove that, for all k in J ,

$$S_{\Phi\left(S_{\Theta(N_k)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]}]\right)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I'}]} \subset S_{\Phi\left(S_{\Theta(d_k)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]}]\right)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I'}]}. \quad (3.8)$$

Using that $\Theta' = \Phi \circ \Theta$, Proposition 3.5 shows that, for all k in J ,

$$S_{\Phi\left(S_{\Theta(N_k)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I'}]}]\right)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I'}]} \subset S_{\Phi\left(S_{\Theta(d_k)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I'}]}]\right)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I'}]}. \quad (3.9)$$

Now (3.9) and Fact 3.11 implies (3.8) for all $k \in J$, as required.

5. This is an immediate consequence of Theorem 3.12.

6. This is an immediate consequence of (5). □

Remark 3.16. One can prove Proposition 3.15 (6) directly using explicit computations on fractions. However, Proposition 3.15 (3) (4) and (5) provides stronger uniqueness assertions.

Fact 3.17. Let $H : \mathcal{A} \rightarrow \mathcal{B}$ be a functor. Let N, N' be two sieves of A . Then

$$S_{H(N)}^{\mathcal{B}} \cup S_{H(N')}^{\mathcal{B}} = S_{H(N \cup N')}^{\mathcal{B}}.$$

Proof. The inclusion \subset is immediate. Let x be in $S_{H(N \cup N')}^{\mathcal{B}}$, then $x = H(n) \circ y$ with $n \in N \cup N'$. So x belongs to $S_{H(N)}^{\mathcal{B}} \cup S_{H(N')}^{\mathcal{B}}$. □

Proposition 3.18. For all $i \in I$, let N'_i be another sieve over $\text{cod}(d_i)$ and let N''_i be the union of N_i and N'_i . Then $\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}, \{(d_i)^{-1} \circ N'_i\}_{i \in I}]$ identifies with $\mathcal{C}[\{(d_i)^{-1} \circ N''_i\}_{i \in I}]$ (more precisely there are unique \mathcal{C} -functors from each category to the other and these morphisms are isomorphisms).

Proof. We observe that $\Theta_1 : \mathcal{C} \rightarrow \mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}, \{(d_i)^{-1} \circ N'_i\}_{i \in I}]$ and $\Theta_2 : \mathcal{C} \rightarrow \mathcal{C}[\{(d_i)^{-1} \circ N''_i\}_{i \in I}]$ belongs to $\text{Cat}_{\mathcal{C}}^{\Sigma\text{-reg}}$ by Fact 3.7. For $k \in I$, by Proposition 3.5 we have

$$S_{\Theta_1(N_k)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}, \{(d_i)^{-1} \circ N'_i\}_{i \in I}]} \cup S_{\Theta_1(N'_k)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}, \{(d_i)^{-1} \circ N'_i\}_{i \in I}]} \subset S_{\Theta_1(d_k)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}, \{(d_i)^{-1} \circ N'_i\}_{i \in I}]}.$$

So by Fact 3.17 we have

$$S_{\Theta_1(N_k \cup N'_k)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}, \{(d_i)^{-1} \circ N'_i\}_{i \in I}]} \subset S_{\Theta_1(d_k)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}, \{(d_i)^{-1} \circ N'_i\}_{i \in I}]}.$$

So by Theorem 3.10, we get a unique \mathcal{C} -functor

$$\alpha : \mathcal{C}[\{(d_i)^{-1} \circ N''_i\}_{i \in I}] \rightarrow \mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}, \{(d_i)^{-1} \circ N'_i\}_{i \in I}].$$

For $k \in I$, by Proposition 3.5, we have

$$S_{\Theta_2(N''_k)}^{\mathcal{C}[\{(d_i)^{-1} \circ N''_i\}_{i \in I}]} \subset S_{\Theta_2(d_k)}^{\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]}.$$

Since $N_k, N'_k \subset N''_k$, by Theorem 3.10, we get a unique \mathcal{C} -functor

$$\alpha' : \mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}, \{(d_i)^{-1} \circ N'_i\}_{i \in I}] \rightarrow \mathcal{C}[\{(d_i)^{-1} \circ N''_i\}_{i \in I}].$$

Proposition 3.12 implies that $\alpha \circ \alpha'$ and $\alpha' \circ \alpha$ are identity morphisms. \square

4. Codilatations of categories

We used fractions of morphisms $(d_i)^{-1} \circ N_i$, the construction given in this note also makes sense for fractions $V_i \circ (d_i)^{-1}$ (each V_i is now a cosieve), pictorially:

$$\begin{array}{ccc} X & \xrightarrow{v} & Y \\ & \searrow^{d_i} & \nearrow_{\exists!} \\ & & Z \end{array} \quad \forall i \in I, \forall v \in V_i.$$

In this section, we introduce this construction formally. We define codilatations using dilata-tions and opposite categories. Let \mathcal{C} be a category.

Definition 4.1. A cocenter $\{[V_i, d_i]\}_{i \in I}$ in \mathcal{C} is a collection of pairs $[V_i, d_i]$ where d_i is a morphism of \mathcal{C} and V_i is a cosieve from $\text{dom}(d_i)$.

We now fix a cocenter $\{[V_i, d_i]\}_{i \in I}$.

Fact 4.2. In \mathcal{C}^{op} , for all i in I , V_i can be regarded as sieve over $\text{cod}^{\mathcal{C}^{op}}(d_i) = \text{dom}^{\mathcal{C}}(d_i)$. In particular $\{[V_i, d_i]\}_{i \in I}$ can be regarded as a center in \mathcal{C}^{op} .

Proof. By definition, the collection of morphisms of a cosieve is a sieve in the opposite category. \square

Definition 4.3. We put $\mathcal{C}[\{V_i \circ (d_i)^{-1}\}_{i \in I}] = (\mathcal{C}^{op}[\{(d_i)^{-1} \circ V_i\}_{i \in I}])^{op}$. The category $\mathcal{C}[\{V_i \circ (d_i)^{-1}\}_{i \in I}]$ is called the codilatation of \mathcal{C} with cocenter $\{[V_i, d_i]\}_{i \in I}$.

Fact 4.4. Let \mathcal{A} be a category. Let E be a collection of morphisms of \mathcal{A} . Then $\text{CoS}_E^{\mathcal{A}} = S_E^{\mathcal{A}^{op}}$.

Proof. $\text{CoS}_E^{\mathcal{A}} = \{x \circ_{\mathcal{A}} e \mid \text{dom}^{\mathcal{A}}(x) = \text{cod}^{\mathcal{A}}(e)\} = \{e \circ_{\mathcal{A}^{op}} x \mid \text{dom}^{\mathcal{A}^{op}}(e) = \text{cod}^{\mathcal{A}}(x)\} = S_E^{\mathcal{A}^{op}}$. \square

Proposition 4.5. 1. We have a canonical functor $\Upsilon : \mathcal{C} \rightarrow \mathcal{C}[\{V_i \circ (d_i)^{-1}\}_{i \in I}]$.
 2. We have a canonical faithful functor $\mathcal{C}[\{V_i \circ (d_i)^{-1}\}_{i \in I}] \rightarrow \mathcal{C}[\{(d_i)_{i \in I}^{-1}\}]$.
 3. The functor Υ is an object in the category $\text{Cat}_{\mathcal{C}}^{\{d_i\}_{i \in I}\text{-reg}}$.
 4. The functor Υ represents the covariant functor $\text{Cat}_{\mathcal{C}}^{\{d_i\}_{i \in I}\text{-reg}}$ to Set given by

$$(\mathcal{C} \xrightarrow{F} \mathcal{D}) \longmapsto \begin{cases} \{*\}, & \text{if } \text{CoS}_{F(v_i)}^{\mathcal{D}} \subset \text{CoS}_{F(d_i)}^{\mathcal{D}} \text{ for all } i \text{ in } I; \\ \emptyset, & \text{else.} \end{cases} \quad (4.1)$$

- Proof.* 1. The canonical functor $\mathcal{C}^{op} \rightarrow \mathcal{C}^{op} [\{(d_i)^{-1} \circ V_i\}_{i \in I}]$ of Proposition 3.1 induces a canonical functor $\Upsilon : \mathcal{C} \rightarrow (\mathcal{C}^{op} [\{(d_i)^{-1} \circ V_i\}_{i \in I}])^{op}$.
2. Fact 2.14 provides a faithful functor $\mathcal{C}^{op} [\{(d_i)^{-1} \circ V_i\}_{i \in I}] \rightarrow \mathcal{C}^{op} [(\{d_i\}_{i \in I})^{-1}]$. Since $\mathcal{C}^{op} [(\{d_i\}_{i \in I})^{-1}] = (\mathcal{C} [(\{d_i\}_{i \in I})^{-1}])^{op}$ (e.g. using the explicit descriptions with fractions), we get the desired faithful functor $(\mathcal{C}^{op} [\{(d_i)^{-1} \circ V_i\}_{i \in I}])^{op} \rightarrow \mathcal{C} [(\{d_i\}_{i \in I})^{-1}]$.
3. Fact 3.7 implies that $\mathcal{C}^{op} \rightarrow \mathcal{C}^{op} [\{(d_i)^{-1} \circ V_i\}_{i \in I}]$ is an object in the category $Cat_{\mathcal{C}^{op}}^{\{d_i\}_{i \in I}\text{-reg}}$. This implies that the functor Υ is an object in the category $Cat_{\mathcal{C}}^{\{d_i\}_{i \in I}\text{-reg}}$.
4. Combine Proposition 3.12 and Fact 4.4. □

5. Some examples and remarks

5.1 Universal property of localizations of categories Let \mathcal{C} be a category and let Σ be a class of morphisms of \mathcal{C} . Fact 2.15 shows that the localization $\mathcal{C}[\Sigma^{-1}]$ is equal to the dilatation $\mathcal{C}[\{d^{-1} \circ N_d\}_{d \in \Sigma}]$ where $N_d = S_{Id_{cod(d)}}^{\mathcal{C}}$. Obviously, $\mathcal{C}[\Sigma^{-1}][[\Sigma^{-1}]] = \mathcal{C}[\Sigma^{-1}]$ and so $\mathcal{C}[\Sigma^{-1}]$ belongs to $Cat_{\mathcal{C}}^{\Sigma\text{-reg}}$. Now if $F : \mathcal{C} \rightarrow \mathcal{D}$ is another \mathcal{C} -category, such that $F(\Sigma)$ is made of isomorphisms, again $\mathcal{D} = \mathcal{D}[F(\Sigma)^{-1}]$ so that \mathcal{D} also belongs to $Cat_{\mathcal{C}}^{\Sigma\text{-reg}}$. Moreover in this case it is obvious that for any $d \in \Sigma$, we have

$$S_{F(N_d)}^{\mathcal{D}} \subset S_{F(d)}^{\mathcal{D}}.$$

So by Theorem 3.10, F factors uniquely through \mathcal{C}' . So the universal property of dilatations generalizes the universal property of localizations.

5.2 Dilatations of commutative rings and semirings

Proposition 5.1. *Let A be a commutative unital ring. Let I be a set and let $\{[M_i, a_i]\}_{i \in I}$ be a center in A in the sense of [9]. Let $A[\{\frac{M_i}{a_i}\}_{i \in I}]$ be the dilatation of A with center $\{[M_i, a_i]\}_{i \in I}$ as in [9]. Let \mathcal{C} be the category attached to A . In particular a_i is a morphism of \mathcal{C} and M_i is a sieve for all i in I . We have $Ob\mathcal{C} = \{\bullet\}$, a singleton. Let $\mathcal{C}[\{\frac{M_i}{a_i}\}_{i \in I}]$ be the category attached to the ring $A[\{\frac{M_i}{a_i}\}_{i \in I}]$. We have an identification of \mathcal{C} -functors $\mathcal{C}[\{(a_i)^{-1} \circ M_i\}_{i \in I}] = \mathcal{C}[\{\frac{M_i}{a_i}\}_{i \in I}]$.*

Proof. Let Φ be the functor $\mathcal{C}[\{(a_i)^{-1} \circ M_i\}_{i \in I}] \rightarrow \mathcal{C}[\{\frac{M_i}{a_i}\}_{i \in I}]$ given by $[\bullet \xrightarrow{m} \bullet \xrightarrow{a_i} \bullet] \mapsto \frac{m}{a_i}$. It is well-defined, surjective, injective and provides the desired identification. □

This shows that dilatations of categories generalize dilatations of rings. As noticed in [9], dilatations of commutative semirings make sense. The same argument as before shows that dilatations of semirings also provide examples of dilatations of categories.

5.3 Dilatations of monoids Let \mathcal{C} be a small category with a single object, i.e. a (not necessarily commutative) monoid M . Let $\{[N_i, d_i]\}_{i \in I}$ be a center of \mathcal{C} . Then $\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]$ is a category with a single object endowed with a functor $\mathcal{C} \rightarrow \mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]$. In other words, $\mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]$ is an M -monoid M' . We refer to this construction as dilatations of monoids. This generalizes localizations of monoids. Codilatations of monoids also make sense.

5.4 Dilatations of pre-additive categories and non-commutative rings Let \mathcal{C} be a pre-additive category, in general a dilatation of \mathcal{C} is not pre-additive as we know that this fails already for localization, cf. e.g. [1]. Nevertheless, as in the case of localizations, it should be possible to study dilatations of pre-additive categories once we have an adapted calculus of fractions. This is related to investigate dilatations of non-commutative rings.

5.5 A characterization of dilatations of rings that does not hold for categories In the context of dilatations of rings, [9, §2] shows that for any sub- A -algebra B of a localization $A[(\{a_i\}_{i \in I})^{-1}]$, there is a multi-center in A such that B identifies with the associated dilatation of A . Here we explain that this characterization does not extend to dilatations of categories.

Fact 5.2. *There exists a category \mathcal{C} , a collection of morphisms Γ , a functor $\mathcal{C} \rightarrow \mathcal{D}$ that is the identity on objects, and a faithful functor $\mathcal{D} \rightarrow \mathcal{C}[\Gamma^{-1}]$ such that*

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{\quad} & \mathcal{C}[\Gamma^{-1}] \\ & \searrow & \nearrow \\ & \mathcal{D} & \end{array}$$

commutes and such that $\mathcal{C} \rightarrow \mathcal{D}$ is not isomorphic to $\mathcal{C} \rightarrow \mathcal{C}[\{(d_i)^{-1} \circ N_i\}_{i \in I}]$ for all centers $\{[N_i, d_i]\}_{i \in I}$.

Proof. Let \mathcal{C} be the category with two objects X and Y and whose morphisms are as follows

$$\text{Hom}_{\mathcal{C}}(X, X) = \text{Id}_X, \text{Hom}_{\mathcal{C}}(Y, Y) = \text{Id}_Y, \text{Hom}_{\mathcal{C}}(Y, X) = \emptyset \text{ and } \text{Hom}_{\mathcal{C}}(X, Y) = \{a, b\}$$

Let Γ be the collection of morphisms of \mathcal{C} given by

$$\Gamma = \{b\}.$$

Then $\mathcal{C}[\Gamma^{-1}]$ is the category whose objects are X and Y and whose morphisms are as follows

$$\begin{aligned} \text{Hom}_{\mathcal{C}[\Gamma^{-1}]}(X, X) &= \{\text{Id}_X, b^{-1}a, b^{-1}ab^{-1}a, \dots, (b^{-1}a)^n, \dots (n \in \mathbb{N})\}, \\ \text{Hom}_{\mathcal{C}[\Gamma^{-1}]}(Y, Y) &= \{\text{Id}_Y, ab^{-1}, ab^{-1}ab^{-1}, \dots, (ab^{-1})^n, \dots (n \in \mathbb{N})\}, \\ \text{Hom}_{\mathcal{C}[\Gamma^{-1}]}(X, Y) &= \{b, a, ab^{-1}a, ab^{-1}ab^{-1}a, \dots, a(b^{-1}a)^n, \dots (n \in \mathbb{N})\}, \\ \text{Hom}_{\mathcal{C}[\Gamma^{-1}]}(Y, X) &= \{b^{-1}, b^{-1}ab^{-1}, b^{-1}ab^{-1}ab^{-1}, \dots, b^{-1}(ab^{-1})^n, \dots (n \in \mathbb{N})\}. \end{aligned}$$

Now let \mathcal{D} be the subcategory of $\mathcal{C}[\Gamma^{-1}]$ whose objects are X and Y and whose morphisms are as follows

$$\begin{aligned} \text{Hom}_{\mathcal{D}}(X, X) &= \text{Id}_X, \\ \text{Hom}_{\mathcal{D}}(Y, Y) &= \text{Id}_Y, \\ \text{Hom}_{\mathcal{D}}(X, Y) &= \{b, a, ab^{-1}a\}, \\ \text{Hom}_{\mathcal{D}}(Y, X) &= \emptyset. \end{aligned}$$

Then we have canonical functors $\mathcal{C} \rightarrow \mathcal{D}$ and $\mathcal{D} \rightarrow \mathcal{C}[\Gamma^{-1}]$. We claim that there is no center $\{[N_i, d_i]\}_{i \in I}$ such that \mathcal{D} identifies with the associated dilatation. To prove this claim, we chose a center $\{[N_i, d_i]\}_{i \in I}$ in \mathcal{C} , put $\Sigma = \{d_i\}_{i \in I}$ and exhaustively distinguish two cases:

1. if $(a \in \Sigma \text{ and } (\text{Id}_Y \in N_a \text{ or } b \in N_a)) \text{ or } (b \in \Sigma \text{ and } (\text{Id}_Y \in N_b \text{ or } a \in N_b))$, then $\#\text{Hom}_{\mathcal{C}'}(X, Y)$ is infinite,
2. if $(a \in \Sigma \Rightarrow N_a \subset \{a\}) \text{ and } (b \in \Sigma \Rightarrow N_b \subset \{b\})$, then $\mathcal{C}' = \mathcal{C}$.

In all cases $\mathcal{C}' \neq \mathcal{D}$. □

Acknowledgements

I am grateful to the referee for his suggestion to remove an unnecessary assumption, which led to many improvements. The author has received funding from the ISF grant No. 1577/23 and ERC grant agreement No. 101002592.

References

- [1] M. Artin. *Non commutative rings*. Class notes Math 251 Berkeley, 1999. https://math.mit.edu/~zyun/Artin_notes.pdf
- [2] F. Borceux. *Handbook of categorical algebra I Basic category theory*. Encyclopedia of Mathematics and its Applications, 50. Cambridge University Press, Cambridge, 1994.
- [3] A. Buan, B. Marsh. *From triangulated categories to module categories via localization II: calculus of fractions*. J. London Math. Soc.(2) 86 (2012) 152–170.
- [4] A. Dubouloz, A. Mayeux. *A polyptych of multi-centered deformation spaces*, [ArXiv:2411.15606](https://arxiv.org/abs/2411.15606).
- [5] A. Dubouloz, A. Mayeux, J.P. dos Santos. *A survey on algebraic dilatations*, Proceedings of the workshop “Langlands Program: Number Theory and Representation Theory”, BIRS-CMO, 2023
- [6] N. D. Duong, P. H. Hai, J. P. dos Santos. *On the structure of affine flat group schemes over discrete valuation rings, I*. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 18, no. 3, 977–1032, 2018
- [7] P. Gabriel, M. Zisman. *Calculus of fractions and homotopy theory*. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 35. New York: Springer-Verlag, 1967
- [8] S. Gelfand, Y. Manin. *Methods of homological algebra*. Second edition. Springer Monographs in Mathematics. Springer-Verlag, 2003.
- [9] A. Mayeux. *Multi-centered dilatations, congruent isomorphisms and Rost double deformation space*, Transformation Groups (2024). <https://doi.org/10.1007/s00031-024-09894-9>
- [10] A. Mayeux, T. Richarz, M. Romagny. *Néron blowups and low-degree cohomological applications*, Algebr. Geom. 10 (2023), no. 6, 729–753.
- [11] The Stacks project authors. *The Stacks Project*. <https://stacks.math.columbia.edu>, 2018