

Jan Mařík

On pseudo-compact spaces

Proc. Japan Acad. 35 (1959), 120–121

Persistent URL: <http://dml.cz/dmlcz/502110>

Terms of use:

© The Japan Academy, 1959

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

25. On Pseudo-compact Spaces

By Jan MARÍK

(Comm. by K. KUNUGI, M.J.A., March 12, 1959)

Notations. Let S be a topological space. Let Z be the family of all sequences $\{f_n(x)\}_{n=1,2,\dots}$, where f_n are (finite real) continuous functions on S such that $f_n(x) \rightarrow 0$ for each $x \in S$. Let Z_0 be the family of all bounded sequences $\{f_n\} \in Z$; let N (resp. E , resp. U) be the family of all non-increasing (resp. equi-continuous, resp. uniformly convergent) sequences $\{f_n\} \in Z$. Further we put $N_0 = N \cap Z_0$, $E_0 = E \cap Z_0$.

If b is a real number, we write $b_+ = \max(b, 0)$.

Lemma 1. $U \cup N \subset E$.

Proof. If $\{f_n\} \in U \cup N$, $x \in S$, $\varepsilon = 2\eta > 0$, then there exist an index p and a neighbourhood V of x such that $|f_n(y)| < \eta$ for each $n > p$ and each $y \in V$. Further we can find a neighbourhood W of x such that $|f_n(x) - f_n(y)| < \varepsilon$ for $n = 1, \dots, p$ and for each $y \in W$. Obviously $|f_n(x) - f_n(y)| < \varepsilon$ for each n and each $y \in V \cap W$.

Lemma 2. If $\{f_n\} \in E$, then the function $f(x) = \sum_{n=1}^{\infty} (|f_n(x)| - \varepsilon)_+$ is continuous for each $\varepsilon > 0$.

Proof. Suppose that $x \in S$ and that $\varepsilon = 2\eta > 0$. There exist an index p and a neighbourhood V of x such that $|f_n(x)| < \eta$ for each $n > p$ and that $|f_n(x) - f_n(y)| < \eta$ for each $y \in V$ and each n . Now, if $n > p$ and if $y \in V$, we have $|f_n(y)| < \varepsilon$, whence $(|f_n(y)| - \varepsilon)_+ = 0$. It follows that $f(y) = \sum_{n=1}^p (|f_n(y)| - \varepsilon)_+$ for each $y \in V$, which completes the proof.

Lemma 3. If S is pseudo-compact, then $N \subset U$.

Proof. If $\{f_n\} \in N$ and if $\varepsilon = 2\eta > 0$, then, by Lemmas 1 and 2, the function $f(x) = \sum_{n=1}^{\infty} (f_n(x) - \eta)_+$ is continuous. Since S is pseudo-compact, there exists a number A such that $f(x) < A$ for each $x \in S$. If $f_n(x) \geq \varepsilon$, then $(f_n(x) - \eta)_+ \geq \eta$ for $k = 1, 2, \dots, n$, so that $n\eta \leq f(x) < A$, $n < A\eta^{-1}$. We see that $f_n(x) < \varepsilon$ for each $n \geq A\eta^{-1}$ and for each $x \in S$.

Lemma 4. If S is pseudo-compact, then $E \subset U$.

Proof. Let $\{f_n\}$ be a sequence of E and let $\varepsilon = 2\eta$ be a positive number. Lemma 2 implies that the functions $g_n(x) = \sum_{k=n}^{\infty} (|f_k(x)| - \eta)_+$ are continuous; obviously $\{g_n\} \in N$ and so, by Lemma 3, $\{g_n\} \in U$. There exists an index p such that $g_p(x) < \eta$ for each $x \in S$. If $n \geq p$, then we have $|f_n(x)| \leq (|f_n(x)| - \eta)_+ + \eta \leq g_p(x) + \eta < 2\eta = \varepsilon$ for each $x \in S$, which proves the lemma.

Lemma 5. If $N_0 \subset U$, then S is pseudo-compact.

Proof. Let f be an unbounded continuous function on S ; put $f_n(x) = \arctg(n^{-1}|f(x)|)$. Then $\{f_n\} \in N_0 - U$.

Theorem. If some of the families E, E_0, N, N_0 is contained in U , then S is pseudo-compact. If, conversely, S is pseudo-compact, then $U = E = E_0, U \ni N = N_0$.

Proof. Lemma 1 implies that $N \subset E$; now it is obvious that $N_0 \subset N, N_0 \subset E_0 \subset E$. If some of the families E, E_0, N, N_0 is contained in U , then we have $N_0 \subset U$ and, by Lemma 5, S is pseudo-compact.

Now let S be pseudo-compact. From Lemmas 1 and 4 we see that $E = U$, so that $E = E_0$. Lemma 3 implies that $N \subset U$; obviously $N = N_0$ and the proof is complete.

Remark. This theorem is a slight generalization of Theorem 3 of [1] and of Theorem 3 of [2].

References

- [1] K. Iséki: A characterisation of pseudo-compact spaces, Proc. Japan Acad., **33**, 320-322 (1957).
- [2] K. Iséki: Pseudo-compactness and strictly continuous convergence, Proc. Japan Acad., **33**, 424-428 (1957).